



Relationships between predicted moonlighting proteins, human diseases, and comorbidities from a network perspective

Andreas Zanzoni, Charles E. Chapple, Christine Brun

► To cite this version:

Andreas Zanzoni, Charles E. Chapple, Christine Brun. Relationships between predicted moonlighting proteins, human diseases, and comorbidities from a network perspective. *Frontiers in Physiology*, 2015, 10.3389/fphys.2015.00171 . hal-01202109

HAL Id: hal-01202109

<https://amu.hal.science/hal-01202109>

Submitted on 18 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Relationships between predicted moonlighting proteins, human diseases, and comorbidities from a network perspective

Andreas Zanzoni^{1,2}, Charles E. Chapple^{1,2} and Christine Brun^{1,2,3*}

¹ INSERM, UMR_S1090 TAGC, Marseille, France, ² Aix-Marseille Université, UMR_S1090, TAGC, Marseille, France, ³ Centre National de la Recherche Scientifique, Marseille, France

OPEN ACCESS

Edited by:

Rafael Tabarés-Seisdedos,
University of Valencia, Spain

Reviewed by:

Scott H. Harrison,
North Carolina A&T State University,
USA

Sylvie Ricard-Blum,
University Claude Bernard Lyon 1,
France

*Correspondence:

Christine Brun,
INSERM, Aix-Marseille Université,
TAGC, UMR_S1090, Parc Scientifique
de Luminy, Case 928, Marseille,
F-13009, France
brun@tagc.univ-mrs.fr

Specialty section:

This article was submitted to
Systems Biology,
a section of the journal
Frontiers in Physiology

Received: 12 November 2014

Accepted: 20 May 2015

Published: 23 June 2015

Citation:

Zanzoni A, Chapple CE and Brun C
(2015) Relationships between
predicted moonlighting proteins,
human diseases, and comorbidities
from a network perspective.
Front. Physiol. 6:171.
doi: 10.3389/fphys.2015.00171

Moonlighting proteins are a subset of multifunctional proteins characterized by their multiple, independent, and unrelated biological functions. We recently set up a large-scale identification of moonlighting proteins using a protein-protein interaction (PPI) network approach. We established that 3% of the current human interactome is composed of predicted moonlighting proteins. We found that disease-related genes are over-represented among those candidates. Here, by comparing moonlighting candidates to non-candidates as groups, we further show that (i) they are significantly involved in more than one disease, (ii) they contribute to complex rather than monogenic diseases, (iii) the diseases in which they are involved are phenotypically different according to their annotations, finally, (iv) they are enriched for diseases pairs showing statistically significant comorbidity patterns based on Medicare records. Altogether, our results suggest that some observed comorbidities between phenotypically different diseases could be due to a shared protein involved in unrelated biological processes.

Keywords: moonlighting proteins, human disease, protein-protein interactions, multifunctional proteins, disease comorbidity

Introduction

Moonlighting proteins are a special sub-class of multifunctional proteins performing multiple autonomous, often unrelated functions (Jeffery, 1999; Huberts et al., 2010; Copley, 2012). For instance, the human aconitase, encoded by the *ACO1* gene, is an enzyme of the tricarboxylic acid cycle (TCA cycle) which turns into a translation regulator upon a conformational change, when the iron concentration is low (Volz, 2008). Such switch between functions can molecularly depend on changes in substrates, post-translational modifications, the action of effector molecules, and other physiochemical factors. They may, in some cases, modify the oligomerization or conformational status of the protein. Moreover, additional functions may arise from a shift in the sub-cellular localization, the physicochemical environment or the pathological status of the cell, or a combination thereof (Jeffery, 2014).

In disease states, the switch of function of a moonlighting protein may occur in different ways. First, moonlighting proteins can be “neomorphic” (Jeffery, 2011), meaning that the protein does not play the same role in healthy and disease conditions. Therefore, the secondary function of the protein is solely revealed during the progression of a pathological state. This is illustrated by HMMR, a nuclear microtubule-associated protein in normal cells, which is exported to the

extracellular matrix in certain tumors, where it binds the CD44 antigen, activates the ERK pathway, and ultimately promotes metastasis (reviewed in Maxwell et al., 2008; Jiang et al., 2010). Second, the different functions performed by a moonlighting protein in a healthy organism can be impaired by the effect of mutations in the disease state, often leading to complex phenotypes. For example, the phosphoglucose isomerase (PGI) is a moonlighting protein in healthy conditions. It converts Glucose 6-phosphate to Fructose 6-phosphate and also functions as a neurotrophic factor (Haga et al., 2000). Mutation leading to the inactivation of both molecular functions provokes a hemolytic anemia with neurological defects whereas a mutation impairing only the enzymatic function causes solely hemolytic anemia (Kugler et al., 1998). This naturally raises the question of a possible relationship between moonlighting proteins and the co-occurrence of different diseases within a single individual, i.e., disease comorbidity. Interestingly, recent works suggest that disease comorbidity generally occurs due to the dys-regulation of a common underlying cellular process through shared genes or protein interactions (Goh et al., 2007; Park et al., 2009). Hence, in the case of diseases that share a moonlighting protein, we hypothesize their co-occurrence may be caused by the dysregulation of distinct cellular processes.

As a first step toward the large-scale identification of moonlighting proteins, we have recently developed a pipeline that can detect what we have termed “extreme multifunctional proteins” (EMFs) from a protein-protein interaction (PPI) network (Chapple et al., 2015). These are proteins whose multiple functions are very different to one another. Although, EMFs will not all necessarily adhere to the strict definition of a moonlighting protein in term of protein organization, that is performing several unrelated functions without partitioning them between different several protein domains, (for further discussions see Jeffery, 1999; Huberts et al., 2010; Copley, 2012; Chapple et al., 2015), they will be however enriched in moonlighting candidates. Our analysis showed that 30% of the human interactome consists of multifunctional proteins, 10% of which are candidate EMFs. We demonstrated that they form a distinct sub-group of proteins characterized by specific features, constituting a signature of extreme multifunctionality (Chapple et al., 2015).

Here, we further investigate their involvement in human disorders and their possible role in disease comorbidities. Indeed, we found that EMF candidates appear to be involved in more than one disease, to contribute to complex rather than monogenic diseases, and to be involved in diseases leading to different phenotypes according to their annotations, when they are compared to other non-candidates proteins. Finally, they are implicated in diseases showing statistically significant comorbidity patterns based on MediCare records (Park et al., 2009). Overall, these results suggest that a protein involved in unrelated cellular processes can be implicated in phenotypically different diseases, therefore providing a possible molecular explanation for some known comorbidities between diseases observed at the level of individuals and populations.

Materials and Methods

A Dataset of EMF Proteins

A dataset of 412 EMF candidates (Chapple et al., 2015) have been identified from a human protein-protein interaction network containing 74388 interactions between 12865 proteins, constructed by gathering interaction data from several databases through the PSICQUIC query interface (Aranda et al., 2011) (February 2013). Overlapping clusters have been extracted from the network using OCG (Becker et al., 2012), and annotated with Gene Ontology terms (Biological Process) using a majority rule. The EMF candidates are selected as proteins belonging to clusters annotated to dissimilar GO terms. GO term dissimilarity was assessed using two metrics of GO term co-occurrence, one based on the frequency of two terms annotating the same protein and the other on the frequency of interactions between proteins annotated to one term and those annotated to the other. These metrics allow the identification of functions that are very rarely performed by (i) a single protein and (ii) by interacting proteins, two proxies we consider indicators of *unrelated functions* (Chapple et al., 2015).

Disease Annotations

OMIM (Amberger et al., 2009) disease-protein associations and non-synonymous mutations were obtained from the UniprotKB database (Magrane and Consortium, 2011). OMIM disease phenotypes were mapped to Human Phenotype (Köhler et al., 2014) (HPO, “organ abnormality” branch) and Disease Ontology (Osborne et al., 2009) (DO) terms using their respective OBO ontology files. The broad HPO phenotype categories were defined using SimCT, a tool for ontology clustering (Herrmann et al., 2009).

Phenotypic Similarity Measures

For each disease pair a - b associated with every protein of interest, we computed the phenotypic similarity of their HPO annotations using the Sørensen–Dice coefficient defined as:

$$PhenoSim_{S-D} = \frac{2(A \cap B)}{A \cup B}$$

and the Jaccard index defined as:

$$PhenoSim_j = \frac{A \cap B}{A \cup B}$$

where A is the list of HPO terms associated to disease a , and B the list of HPO terms associated to disease b .

Disease Semantic Similarity

We used the DOSE package for R (Yu et al., 2015) for computing semantic similarities among Disease Ontology (DO) terms associated with every disease pair by applying both Lin’s information content-based method and Wang’s graph-based method (for a review, see Gan et al., 2013).

Comorbidity Data

The list of disease pairs with statistically significant comorbidity patterns and linked by gene sharing was taken from Park et al. (2009).

Pfam Domain Detection

We used the Pfamscan tool to identify Pfam domains (version 26.0) (Punta et al., 2012). We kept only Pfam-A matches with an $E < 10^{-5}$.

Results

A Dataset of EMF Candidates in the Human Interactome

EMFs, including moonlighting proteins, are expected to perform their independent functions through different interaction partners. In a recent work (Chapple et al., 2015), we have identified EMF protein candidates from the human interactome using OCG (Becker et al., 2012), an algorithm that identifies overlapping protein clusters in an interaction network. The cellular functions in which protein clusters are involved have been defined with regards to the Gene Ontology Biological Process annotations of their constituent proteins. Then, EMF candidates are proteins found at the intersection of clusters involved in very dissimilar cellular functions (see Methods section for details). Using this approach, we have identified two distinct categories of multifunctional proteins in the human interactome: 412 EMF candidates (MF-CAN) and 3434 non-EMF multifunctional proteins (MCNC for multi-clustered non-candidate). A third category consists of the remaining 9019 proteins in the interactome, which belong to only one cluster (MONO for mono-clustered). The proteins belonging to the three categories are listed in Supplementary Data Sheet 1. We have shown that the EMF candidates form a distinct sub-group of proteins displaying specific features distinguishing them from non-candidates proteins and constituting a signature of extreme multifunctionality. Among the striking features, EMF candidates are more connected in the network, enriched in short linear motifs (SLiMs) and in disease-related proteins compared to non-candidates, and are less intrinsically disordered than network hubs (Chapple et al., 2015).


EMF Candidates Are Enriched in Proteins Involved in more than One Disease

Using OMIM disease annotations, we found that the multifunctional proteins of the human interactome (i.e., proteins belonging to more than one network cluster, corresponding to MF-CAN and MCNC taken together) are enriched in proteins involved in diseases (1.33-fold, $P < 2.2 \times 10^{-16}$, Fisher's exact test, two-sided). We still observed this enrichment when considering MF-CAN and MCNC separately (Table 1). In addition, MF-CAN and MCNC are both significantly enriched in proteins involved in more than two diseases, with MF-CAN showing a higher proportion of such proteins compared to MCNC (2.04-fold and 1.25-fold, respectively). On the other hand, the MONO category is significantly depleted in disease proteins (Table 1). Given these results, we decided to focus our attention on those proteins of the three categories that are involved in more than two diseases. Interestingly, MF-CAN proteins are associated with a higher number of diseases compared to the other categories (3.6 compared to 3 and 2.9, on average, for MCNC and MONO respectively, $P = 2.6 \times 10^{-3}$ and

TABLE 1 | Proteins in the human interactome involved in OMIM diseases.

Set	Total number	At least one disease	At least two diseases
MULTI	3846	830 (1.33, $P < 2.2 \times 10^{-16}$)	298 (1.34, $P = 3.8 \times 10^{-7}$)
MF-CAN	412	113 (1.84, $P = 8.2 \times 10^{-8}$)	52 (2.04, $P = 1.8 \times 10^{-4}$)
MCNC	3434	717 (1.28, $P = 9.1 \times 10^{-12}$)	246 (1.25, $P = 6.8 \times 10^{-4}$)
MONO	9019	1365 (0.87, $P < 2.2 \times 10^{-16}$)	349 (0.82, $P = 3.8 \times 10^{-7}$)

Enrichment ratios (ER) and P -values for over- ($ER > 1$) and under-representation ($ER < 1$) are reported between parentheses (Fisher's Exact test, two-sided, $\alpha = 0.05$). See main text for details.



$P = 1.3 \times 10^{-4}$, Mann-Whitney U test, one-sided) (Figure 1). This suggests that MF-CAN proteins are particularly associated to multiple diseases.

EMF Protein Candidates Contribute to Complex Diseases

We next sought to verify whether MF-CAN, MCNC, and MONO proteins are associated with the same diseases. Figure 2A shows that most of the diseases (89%) in which the MF-CAN are involved, are also associated with at least one protein belonging to one of the other categories (compared to only 34% and 26% for MCNC and MONO, respectively). On the other hand, 86% of the disease pairs associated with MF-CAN are specific to this category (compared to ~68% for the other two categories) (Figure 2B). This, therefore, suggests that our EMF candidates are mainly involved in specific combinations of diseases. Moreover, the fact that diseases associated to MF-CAN are associated with several other genes/proteins as well (Figure 2A) suggests that EMF candidates are implicated in combinations of complex diseases rather than monogenic ones.




FIGURE 2 | Shared disease phenotypes and pairs among MF-CAN, MCNC, and MONO proteins. (A) Most of the MF-CAN proteins are annotated with OMIM disease annotations also present in the other categories, whereas **(B)** they are associated with specific combination of diseases.

EMF Candidates Are Involved in Phenotypically Dissimilar Diseases

Different OMIM diseases caused by the dysregulation of a same gene can affect the same body organs or have similar phenotypic manifestations (Goh et al., 2007), or not. We therefore tested this on the disease pairs associated with MF-CAN, MCNC, and MONO proteins.

To do so, we mapped OMIM diseases to the Human Phenotype Ontology (HPO) (Köhler et al., 2014). For each disease, HPO provides a list of annotation terms describing the phenotypes associated to the disease. We clustered them in 19 broad classes of organ abnormalities. We found that proteins belonging to MF-CAN, MCNC, and MONO are over-represented in almost all of the HPO broad classes (**Figure 3**). Particularly, MF-CAN proteins are over-represented in all HPO classes except “abnormality of prenatal development or birth” (**Figure 3A**), MCNC proteins are under-represented in the “abnormality of metabolism/homeostasis” class (**Figure 3B**) whereas MONO proteins are under-represented in three HPO classes: “abnormality of the breast,” “abnormality of the blood and blood-forming tissue,” and “abnormality of head and neck” (**Figure 3C**). More interestingly, we observed a significantly higher fraction of shared MF-CAN proteins, 32% on average, between HPO classes compared to MCNC (21%, $P = 2.3 \times 10^{-13}$, Mann-Whitney U test, one-sided) and MONO (19%, $P < 2.2 \times 10^{-16}$). This result shows that MF-CAN proteins are involved in disease affecting distinct body parts. Therefore, this led us to evaluate the phenotypic similarity between disease pairs by computing the Sorensen-Dice distance based on their HPO annotation terms (see Methods for details). For each pair of diseases in which a same gene is involved, we have compared their phenotypic descriptions to assess their similarity aiming at grasping possible differences. We indeed found that those pairs associated with MF-CAN are significantly different (**Figure 4**, Supplementary Figure 1) compared to the other categories ($P = 4.7 \times 10^{-5}$ and $P = 1.3 \times 10^{-11}$ for MCNC and MONO, respectively, Mann-Whitney U test, one-sided). We obtained similar results using semantic similarity measures (Supplementary Figure 2) based on DO annotations (Schriml et al., 2012). Indeed, MONO disease pairs are significantly more similar than MF-CAN and MCNC pairs ($P = 2.7 \times 10^{-3}$ and 1.5×10^{-5} , respectively, Mann-Whitney U test, one-sided). These

results could be due to the fact that multifunctional candidates (MF-CAN and MCNC) are acting in several biological processes, and are therefore involved in different diseases and phenotypes.

Diseases leading to different phenotypes could be due to mutations located in different protein regions. Given the strict definition applied to moonlighting protein which should perform their multiple functions without partitioning these into multiple domains (Jeffery, 1999), we sought to verify whether mutations causing different diseases are located within the same functional domains. For this, we compared the phenotypic similarities of disease pairs whose mutations are either found in the same Pfam domains or outside. Interestingly, MF-CAN proteins show a lower phenotypic similarity than MCNC and MONO in both situations (Supplementary Figure 3). Therefore, diseases in which MF-CAN are involved, display different phenotypes irrespective to the location of the corresponding mutations.

Overall, these results show that MF-CAN are involved in diseases leading to different phenotypes and affecting different organs.


EMF Candidates are Involved in Comorbid Diseases

In order to estimate whether moonlighting candidates could explain some comorbidity patterns, we used the list of comorbid diseases established by Park et al. (2009), based on Medicare records. We found (**Table 2**, Supplementary Data Sheet 2) that EMF candidates are enriched in proteins involved in at least one disease pair showing statistically significant comorbidity patterns ($P = 3.5 \times 10^{-6}$, Fisher’s exact test, two-sided), whereas MONO are significantly depleted in such pairs ($P = 1.1 \times 10^{-5}$).

Our results therefore indicate that EMF proteins involved in several diseases can contribute to comorbidity, probably through the unrelated biological processes in which they participate.

An Example of EMF Candidate Involved in a Pair of Comorbid Diseases

Androgen receptor (AR) is one of the MF-CAN proteins associated to a disease pair showing a comorbidity pattern (Supplementary Data Sheet 2): prostate cancer (PC) and spinal and bulbar muscular atrophy (SBMA, ICD-9: 335.1), an illness leading to progressive muscle weakness and atrophy, known also as Kennedy’s disease. SBMA is a late-onset neurodegenerative



disease, caused by an extreme expansion of trinucleotide CAG repeats in the AR gene first exon, which encodes for a stretch of glutamines (polyQ) in the N-terminal transactivation domain of the AR protein (Kumar et al., 2011). Interestingly, *in vivo* studies on mice harboring humanized AR alleles showed that long polyQ may be involved in late-onset PC and influence hormone sensitivity (Albertelli et al., 2008; Simanainen et al., 2011). Recently, a case of SBMA and PC comorbidity has been reported in the literature (Kosaka et al., 2012). Altogether, these observations are consistent with the comorbidity reported in Medicare, since it collects hospital visit reports of elderly patients (age equal or above 65). Interestingly, polyQ stretches induce the formation of AR aggregates within neuronal cells, thus impairing AR interaction ability. We can therefore speculate that AR aggregates in prostate cell can have a similar effect, probably by perturbing a different set of AR interactions.

Discussion

Network approaches recently suggested that the co-occurrence of different diseases within the same individual, i.e., comorbidity, correlates with the sharing or the interactions of their underlying molecular components in cellular networks (Park et al., 2009; Zhou et al., 2014). Nevertheless, these molecular components can also be involved in distinct cellular processes. Since moonlighting, and by extension EMF, proteins are implicated in unrelated cellular processes (Jeffery, 1999; Copley, 2012), often taking places in different subcellular localizations, we postulate that disease comorbidities could be due not only to the dysregulation of shared underlying sub-networks as previously proposed, but also to the dysregulation of different ones in which a shared protein, able to moonlight, is taking part. To test this hypothesis, we investigated the potential involvement of a particular group of proteins, the EMF candidates, in




TABLE 2 | Number of proteins associated with at least one disease pair showing comorbidity patterns.

Set	Total number	Disease comorbidity
MF-CAN	52	27 (1.95, $P = 3.5 \times 10^{-6}$)
MCNC	246	67 (1.25, $P = 0.0858$)
MONO	349	58 (0.52, $P = 1.1 \times 10^{-5}$)

Enrichment ratios (ER) and *P*-values for over- ($ER > 1$) and under-representation ($ER < 1$) are reported between parentheses (Fisher's Exact test, two-sided, $\alpha = 0.05$).

diseases. We show that EMF candidates are involved in multiple diseases that are phenotypically dissimilar and affect distinct organs.

Moreover, we found that a significant fraction of these EMF candidates represents a possible molecular link between comorbid diseases. Nevertheless, although a gene/protein may be involved in different diseases, these diseases are not necessarily co-occurring within a single individual. Our finding that comorbid diseases identified from Medicare records are enriched among EMF candidates therefore reinforces (i) the fact that diseases in which an EMF is involved can occur in a same individual and (ii) our hypothesis that comorbidity can be due their involvement.

Diseases associated to EMF proteins display different phenotypes according to HPO, independently of the location of the mutations in the protein sequence, within or outside known protein domains. This raises several possibilities. First, these mutations may occur in existing domains not yet discovered. Indeed, current tools may not have detected some domains, since they are below the detection thresholds, or differing from the model used by the prediction tools. Second, this lack of difference

may also be explained by the fact that disease mutations occur in SLiMs, short stretches of amino acids sequences that represent potential regulatory sites and are located mostly in intrinsically disordered regions (van Roey et al., 2013). As (i) we previously showed that EMFs are enriched in SLiMs compared to MCNC (Chapple et al., 2015) and (ii) a recent report showed that disease mutations occur more likely within SLiMs than neutral missense mutations in intrinsically disordered regions (Uyar et al., 2014), the second hypothesis mentioned above is thus supported.

In an early study on co-occurring metabolic diseases, Barabasi and colleagues (Lee et al., 2008) proposed that the lack of observed comorbidities between 69% of the metabolic disease pairs they investigated, could be due to moonlighting enzymes involved in non-metabolic functions. Indeed, the corresponding subnetworks were missing from the metabolic network they were investigating. Interestingly, for 95% of the EMF, one of the functions is related to various metabolic processes whereas the other is non-metabolic, such as signaling, transport, or localization (Table 1 in Chapple et al., 2015). Therefore, given the link between EMFs and moonlighting explained in the Introduction, our results support the hypothesis that moonlighting proteins could indeed provide the missing connections between metabolic diseases and the rest of the disease network (Goh et al., 2007).

It should be noted that our observations cross and link several integrated functional levels (Brun et al., 2004; Barabási, 2007). First, the disease pairs in which EMF candidates play a role are the most dissimilar in terms of phenotype. Since candidates have been selected as involved in unrelated cellular functions, this observation directly links cellular processes to disease phenotypes, based on the analysis of independent annotations (Gene Ontology and Human Phenotype Ontology). It thus appears that for these proteins, noticeable differences are observed from the molecular level (mutations) to the cellular level (unrelated processes) up to the tissular/organismal level (phenotypes), thus emphasizing the complexity of the link between genotype and phenotype.

Finally, grounded on our computational identification of moonlighting candidates, this work represents, to our knowledge, the first effort toward the elucidation of the convoluted relationships between moonlighting protein functions, disease phenotypes and comorbidities. Overall, although much remains to be learned, our analysis underlined the necessity of identifying moonlighting proteins and to better understand their role in human diseases, particularly in the context of drug development to avoid side effects. Indeed, knowing that the different diseases in which a protein is involved, are due to the dysregulation of different processes may help choosing specific interactions to inhibit in order to act on one disease and not the other. In addition, a drug inhibiting an unsuspected moonlighting protein may cause side effects if it also impairs the secondary function of the protein. Therefore, the identification of candidate moonlighting proteins can be useful to gain a deeper understanding of disease

molecular details and to design more effective therapeutic strategies.

Acknowledgments

The authors received financial support from the French “Plan Cancer 2009–2013” (Systems Biology call, A12171AS), Institut National de la Santé et de la Recherche Médicale (Inserm),

Aix-Marseille University, Centre National de la Recherche Scientifique (CNRS).

Supplementary Material

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fphys.2015.00171>

References

- Albertelli, M. A., O'Mahony, O. A., Brogley, M., Tosoian, J., Steinkamp, M., Daignault, S., et al. (2008). Glutamine tract length of human androgen receptors affects hormone-dependent and -independent prostate cancer in mice. *Hum. Mol. Genet.* 17, 98–110. doi: 10.1093/hmg/ddm287
- Amberger, J., Bocchini, C. A., Scott, A. F., and Hamosh, A. (2009). McKusick's online mendelian inheritance in man (OMIM). *Nucleic Acids Res.* 37, D793–D796. doi: 10.1093/nar/gkn665
- Aranda, B., Blankenburg, H., Kerrien, S., Brinkman, F. S. L., Ceol, A., Chataud, E., et al. (2011). PSICQUIC and PSISCORE: accessing and scoring molecular interactions. *Nat. Methods* 8, 528–529. doi: 10.1038/nmeth.1637
- Barabási, A.-L. (2007). Network medicine—from obesity to the “diseasome.” *N. Engl. J. Med.* 357, 404–407. doi: 10.1056/NEJMe078114
- Becker, E., Robisson, B., Chapple, C. E., Guénoche, A., and Brun, C. (2012). Multifunctional proteins revealed by overlapping clustering in protein interaction network. *Bioinforma. Oxf. Engl.* 28, 84–90. doi: 10.1093/bioinformatics/btr621
- Brun, C., Baudot, A., Guénoche, A., and Jacq, B. (2004). “The use of protein-protein interaction networks for genome wide protein function comparisons and predictions,” in *Methods in Proteome and Protein Analysis Principles and Practice*, eds R. M. Kamp, J. J. Calvete, and T. Choli-Papadopoulou (Heidelberg: Springer-Verlag), 103–124.
- Chapple, C. E., Robisson, B., Spinelli, L., Guien, C., Becker, E., and Brun, C. (2015). Extreme multifunctional proteins identified from a human protein interaction network. *Nat. Commun.* 6:7412. doi: 10.1038/ncomms8412
- Copley, S. D. (2012). Moonlighting is mainstream: paradigm adjustment required. *Bioessays News Rev. Mol. Cell. Dev. Biol.* 34, 578–588. doi: 10.1002/bies.201100191
- Gan, M., Dou, X., and Jiang, R. (2013). From ontology to semantic similarity: calculation of ontology-based semantic similarity. *Sci. World J.* 2013:793091. doi: 10.1155/2013/793091
- Goh, K. I., Cusick, M. E., Valle, D., Childs, B., Vidal, M., and Barabási, A. L. (2007). The human disease network. *Proc. Natl. Acad. Sci. U.S.A.* 104, 8685–8690. doi: 10.1073/pnas.0701361104
- Haga, A., Niinaka, Y., and Raz, A. (2000). Phosphohexose isomerase/autocrine motility factor/neuroleukin/maturation factor is a multifunctional phosphoprotein. *Biochim. Biophys. Acta* 1480, 235–244. doi: 10.1016/S0167-4838(00)00075-3
- Herrmann, C., Berard, S., and Tichit, L. (2009). SimCT: a generic tool to visualize ontology-based relationships for biological objects. *Bioinformatics* 25, 3197–3198. doi: 10.1093/bioinformatics/btp553
- Huberts, D. H. E. W., Venselaar, H., Vriend, G., Veenhuis, M., and van der Klei, I. J. (2010). The moonlighting function of pyruvate carboxylase resides in the non-catalytic end of the TIM barrel. *Biochim. Biophys. Acta* 1803, 1038–1042. doi: 10.1016/j.bbamer.2010.03.018
- Jeffery, J. (1999). Moonlighting proteins. *Trends Biochem. Sci.* 24, 8–11. doi: 10.1016/S0968-0004(98)01335-8
- Jeffery, C. J. (2011). Proteins with neomorphic moonlighting functions in disease. *Iubmb Life* 63, 489–494. doi: 10.1002/iub.504
- Jeffery, C. J. (2014). An introduction to protein moonlighting. *Biochem. Soc. Trans.* 42, 1679–1683. doi: 10.1042/BST20140226
- Jiang, J., Casalegno-Garduno, R., Chen, H., Schmitt, A., Schmitt, M., and Maxwell, C. A. (2010). Multifunctional proteins bridge mitosis with motility and cancer with inflammation and arthritis. *Sci. World J.* 10, 1244–1257. doi: 10.1100/tsw.2010.141
- Köhler, S., Doelken, S. C., Mungall, C. J., Bauer, S., Firth, H. V., Bailleul-Forestier, I., et al. (2014). The human phenotype ontology project: linking molecular biology and disease through phenotype data. *Nucleic Acids Res.* 42, D966–D974. doi: 10.1093/nar/gkt1026
- Kosaka, T., Miyajima, A., Kikuchi, E., Takahashi, S., Suzuki, N., and Oya, M. (2012). A case of spinal and bulbar muscular atrophy with high-stage and high-Gleason Score prostate cancer responded to maximal androgen blockade therapy. *J. Androl.* 33, 563–565. doi: 10.2164/jandrol.111.013698
- Kugler, W., Breme, K., Laspe, P., Muirhead, H., Davies, C., Winkler, H., et al. (1998). Molecular basis of neurological dysfunction coupled with haemolytic anaemia in human glucose-6-phosphate isomerase (GPI) deficiency. *Hum. Genet.* 103, 450–454. doi: 10.1007/s004390050849
- Kumar, R., Atamna, H., Zakharov, M. N., Bhasin, S., Khan, S. H., and Jasuja, R. (2011). Role of the androgen receptor CAG repeat polymorphism in prostate cancer, and spinal and bulbar muscular atrophy. *Life Sci.* 88, 565–571. doi: 10.1016/j.lfs.2011.01.021
- Lee, D.-S., Park, J., Kay, K. A., Christakis, N. A., Oltvai, Z. N., and Barabási, A.-L. (2008). The implications of human metabolic network topology for disease comorbidity. *Proc. Natl. Acad. Sci. U.S.A.* 105, 9880–9885. doi: 10.1073/pnas.0802208105
- Magrane, M., and Consortium, U. (2011). UniProt Knowledgebase: a hub of integrated protein data. *Database* 2011:bar009. doi: 10.1093/database/bar009
- Maxwell, C. A., McCarthy, J., and Turley, E. (2008). Cell-surface and mitotic-spindle RHAMM: moonlighting or dual oncogenic functions? *J. Cell Sci.* 121, 925–932. doi: 10.1242/jcs.022038
- Osborne, J. D., Flatow, J., Holko, M., Lin, S. M., Kibbe, W. A., Zhu, L. J., et al. (2009). Annotating the human genome with disease ontology. *BMC Genomics* 10(Suppl. 1):S6. doi: 10.1186/1471-2164-10-S1-S6
- Park, J., Lee, D.-S., Christakis, N. A., and Barabási, A.-L. (2009). The impact of cellular networks on disease comorbidity. *Mol. Syst. Biol.* 5, 262. doi: 10.1038/msb.2009.16
- Punta, M., Coghill, P. C., Eberhardt, R. Y., Mistry, J., Tate, J., Boursnell, C., et al. (2012). The Pfam protein families database. *Nucleic Acids Res.* 40, D290–D301. doi: 10.1093/nar/gkr1065
- Schriml, L. M., Arze, C., Nadendla, S., Chang, Y.-W. W., Mazaitis, M., Felix, V., et al. (2012). Disease ontology: a backbone for disease semantic integration. *Nucleic Acids Res.* 40, D940–D946. doi: 10.1093/nar/gkr972
- Simanainen, U., Brogley, M., Gao, Y. R., Jimenez, M., Harwood, D. T., Handelsman, D. J., et al. (2011). Length of the human androgen receptor glutamine tract determines androgen sensitivity *in vivo*. *Mol. Cell. Endocrinol.* 342, 81–86. doi: 10.1016/j.mce.2011.05.011
- Uyar, B., Weatheritt, R. J., Dinkel, H., Davey, N. E., and Gibson, T. J. (2014). Proteome-wide analysis of human disease mutations in short linear motifs: neglected players in cancer? *Mol. Biosyst.* 10, 2626–2642. doi: 10.1039/C4MB00290C
- van Roey, K., Dinkel, H., Weatheritt, R. J., Gibson, T. J., and Davey, N. E. (2013). The switches.ELM resource: a compendium of conditional

- regulatory interaction interfaces. *Sci. Signal.* 6, rs7. doi: 10.1126/scisignal.2003345
- Volz, K. (2008). The functional duality of iron regulatory protein 1. *Curr. Opin. Struct. Biol.* 18, 106–111. doi: 10.1016/j.sbi.2007.12.010
- Yu, G., Wang, L. G., Yan, G. R., and He, Q. Y. (2015). DOSE: an R/Bioconductor package for disease ontology semantic and enrichment analysis. *Bioinformatics* 31, 608–609. doi: 10.1093/bioinformatics/btu684
- Zhou, X., Menche, J., Barabási, A.-L., and Sharma, A. (2014). Human symptoms-disease network. *Nat. Commun.* 5, 4212. doi: 10.1038/ncomms5212

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Zanzoni, Chapple and Brun. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.