



**HAL**  
open science

## Synthesis and characterization of [Fe(BPMEN)-ACC]SbF 6: a structural and functional mimic of ACC-oxidase †

Y Roux, W Ghattas, F Avenier, R Guillot, Ariane Jalila Simaan, J.-P Mahy

► **To cite this version:**

Y Roux, W Ghattas, F Avenier, R Guillot, Ariane Jalila Simaan, et al.. Synthesis and characterization of [Fe(BPMEN)-ACC]SbF 6: a structural and functional mimic of ACC-oxidase †. Dalton Transactions, 2015, 44 (13), pp.5966-5968. 10.1039/c5dt00347d . hal-01258299

**HAL Id: hal-01258299**

**<https://amu.hal.science/hal-01258299>**

Submitted on 18 Jan 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Cite this: *Dalton Trans.*, 2015, 44, 5966

Received 26th January 2015,  
Accepted 24th February 2015

DOI: 10.1039/c5dt00347d

www.rsc.org/dalton

## Synthesis and characterization of [Fe(BPMEN)-ACC]SbF<sub>6</sub>: a structural and functional mimic of ACC-oxidase†

Y. Roux,‡<sup>a</sup> W. Ghattas,‡<sup>a</sup> F. Avenier,<sup>a</sup> R. Guillot,<sup>a</sup> A. J. Simaan\*<sup>b</sup> and J.-P. Mahy\*<sup>a</sup>


A mononuclear Fe(II) complex bearing 1-aminocyclopropane-1-carboxylic acid (ACCH) was synthesized and characterized. X-ray crystallography demonstrated that ACC binds to the Fe(II) ion in a bidentate mode constituting the first structural mimic of the expected binding of ACC to the Fe(II) center of the ethylene forming enzyme ACC-oxidase (ACCO). [Fe(BPMEN)ACC]SbF<sub>6</sub> also constitutes a functional biomimetic complex of ACCO, as it reacts with hydrogen peroxide producing ethylene.

The final step in the biosynthesis of the phytohormone ethylene<sup>1</sup> is the oxidation of 1-aminocyclopropane-1-carboxylic acid (ACCH)<sup>2</sup> catalyzed by ACC-oxidase (ACCO).<sup>3</sup> The X-ray crystal structure of substrate-free ACCO<sup>4</sup> has confirmed the anticipated makeup of its active site *i.e.* a non-heme Fe(II) cation coordinated by the classical N,N,O facial triad.<sup>5,6</sup> In contrast, the ambiguous binding of 1-aminocyclopropane-1-carboxylate (ACC) in the active site of ACCO has only been probed by spectroscopic studies, which have nonetheless concluded that ACC binds to the Fe(II) ion of the active site in a bidentate mode *via* both its amine and carboxylate functions.<sup>7</sup> Accordingly, the structural characterization of a dinuclear ACC-containing Fe(III) complex, [Fe<sub>2</sub>(TACN)<sub>2</sub>(μ-O)(μ-ACCH)<sub>2</sub>]<sup>4+</sup> (TACN = 1,4,7-triazacyclononane) has shown that ACC can bind iron centres. In this case however, two ACCH fragments are bridging two Fe(III) cations by their carboxylate functions.<sup>8</sup> Bidentate ACC has only been reported in a few Cu(II)-ACC complexes,<sup>9</sup> and has never been observed in the case of mononuclear Fe(II) complexes. Indeed, the good water solubility of amino acids is inappropriate for the synthesis of Fe(II) complexes, which are best

obtained in aprotic poorly-coordinating solvents that conversely do not dissolve amino acids.<sup>10</sup> To the best of our knowledge, structural characterization of amino acid-containing Fe(II) complexes has only been reported in the case of a proline-containing complex.<sup>11</sup> Proline is structurally distinguished from other natural α-amino acids by a secondary amine function engaged in a 5-membered ring and therefore is inappropriate to structurally mimic the coordination of other natural α-amino acids. Furthermore, the synthesis of the proline-containing Fe(II) complex relied on the solubilization of proline in DMSO, which did not allow to get the structural information when either phenylalanine, tryptophan or valine was used instead of proline.<sup>11</sup>

Here, in order to overcome the solubility limitations, we treated an aqueous solution of ACCH with one equivalent of tetra-*n*-butylammonium hydroxide (N(*n*-Bu)<sub>4</sub>OH). Subsequent water evaporation provided an ionic liquid fully miscible with acetonitrile, which allowed its combination with an acetonitrile solution of the previously described [Fe(BPMEN)-(CH<sub>3</sub>CN)<sub>2</sub>](SbF<sub>6</sub>)<sub>2</sub> complex (1) (BPMEN = *N,N'*-dimethyl-*N,N'*-bis(pyridylmethyl)ethane-1,2-diamine) (Scheme 1).<sup>12</sup>

When one equivalent of N(*n*-Bu)<sub>4</sub>ACC was added to an acetonitrile solution of complex 1, the solution turned from purple to a pale yellow color. The monitoring of the UV-vis absorbance as a function of the increasing amounts of



Scheme 1 Preparation of the [Fe(BPMEN)ACC]SbF<sub>6</sub> complex (2) from [Fe(BPMEN)(CH<sub>3</sub>CN)<sub>2</sub>](SbF<sub>6</sub>)<sub>2</sub> (1) under an inert atmosphere.

<sup>a</sup>Institut de Chimie Moléculaire et des Matériaux d'Orsay, Université Paris Sud, CNRS (UMR 8182), Orsay 91405 CEDEX, France.

E-mail: jean-pierre.mahy@u-psud.fr

<sup>b</sup>Aix Marseille Université, Centrale Marseille, CNRS, iSm2 UMR 7313, 13397, Marseille, France. E-mail: jalila.simaan@univ-amu.fr

† Electronic supplementary information (ESI) available: Crystallographic data, UV-Vis measurements, magnetization measurements. CCDC 1029372 and 1029373. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5dt00347d

‡ Equally contributed to the work reported.






Fig. 1 Evolution of the UV-vis spectrum of a 0.5 mM acetonitrile solution of  $[\text{Fe}(\text{BPMEN})(\text{CH}_3\text{CN})_2](\text{SbF}_6)_2$  (**1**) (bold black line) upon successive additions of up to 1 equiv. (red line) of  $\text{N}(n\text{-Bu})_4\text{ACC}$ .

$\text{N}(n\text{-Bu})_4\text{ACC}$  added to **1** (Fig. 1) showed the progressive evolution of the spectrum of **1** ( $\lambda_{\text{max}} = 373 \text{ nm}$ ,  $\epsilon = 3340 \text{ M}^{-1} \text{ cm}^{-1}$ , MLCT band)<sup>13</sup> into a new spectrum ( $\lambda_{\text{max}} = 395 \text{ nm}$ ,  $\epsilon = 860 \text{ M}^{-1} \text{ cm}^{-1}$ ). The occurrence of an isosbestic point at 410 nm clearly indicated a single transformation of the starting material into a new species. The transformation was optimal for one equivalent of  $\text{N}(n\text{-Bu})_4\text{ACC}$  added. Further addition of  $\text{N}(n\text{-Bu})_4\text{ACC}$  led to a decrease of the characteristic MLCT band at 395 nm that completely disappeared after the addition of three equivalents of  $\text{N}(n\text{-Bu})_4\text{ACC}$  (Fig. S1†).

High resolution electrospray ionization mass spectrometry (HR ESI-MS) analysis was carried out on the pale yellow solution obtained after the addition of one equivalent of the amino acid (Fig. 2). The results revealed the formation of a single new compound characterized by a peak at  $m/z$  426.1604, which is in agreement with the complexation of one ACC molecule to the  $\text{Fe}(\text{II})$  ion of complex **1** in the place of two acetonitrile molecules observed in the X-ray crystal structure (Fig. S2†).




Fig. 2 HR ESI-MS spectrum obtained upon additions of 1 equiv. of  $\text{N}(n\text{-Bu})_4\text{ACC}$  onto  $[\text{Fe}(\text{BPMEN})(\text{CH}_3\text{CN})_2](\text{SbF}_6)_2$  (**1**).

The new pale yellow species was then precipitated by the addition of ether in the acetonitrile solution and recrystallized from slow ether diffusion in acetonitrile to afford monocrystals suitable for X-ray diffraction analysis. The resulting diffraction pattern was in agreement with a  $[\text{Fe}(\text{BPMEN})\text{ACC}]\text{SbF}_6$  molecular formula for complex **2** and a structure in which ACC is bound to the  $\text{Fe}(\text{II})$  center in a bidentate mode *via* both its amine and carboxylate functions, as projected for the enzymatic active site (Fig. 3, Tables S1 and S2†).<sup>5</sup> In addition, both the UV-vis spectrum and the mass spectrum of complex **2** obtained from the solid state matched those of the species formed in solution. In comparison with the X-ray structure of complex **1**, the distorted octahedral geometry is maintained in complex **2**, but the average metal–ligand distance has increased from 1.98 Å in **1** to 2.18 Å in **2**. It is noteworthy that average Fe–N distances below 2.0 Å in **1** and above 2.1 Å in **2** suggest a low spin to high spin transition upon complexation of the amino acid to the  $\text{Fe}(\text{II})$  center.<sup>14</sup>

Bulk magnetization data were collected from the crystalline samples of complex **2**. The corresponding  $\chi_{\text{m}}T$  vs.  $T$  plot (Fig. S3†) showed an initial sharp increase (upto *ca.* 50 K) followed by a slight monotonic increase of  $\chi_{\text{m}}T$  with the increasing temperature. Both the overall shape of  $\chi_{\text{m}}T$  vs.  $T$  and a  $\chi_{\text{m}}T$  value reaching  $3.48 \text{ cm}^3 \text{ K mol}^{-1}$  at 400 K concur with a high spin mononuclear  $\text{Fe}(\text{II})$  center ( $S = 2$ ,  $g = 2.1$ ). Therefore, complex **2** is in the high spin state as suggested by the bond lengths obtained from the crystal structure. The coordination of the amino acid on the  $\text{Fe}(\text{II})$  ion stabilizes the high spin state whereas, complex **1** is known to be in the low spin state at low temperature, in spin transition at room temperature and at the high spin state only above 400 K.<sup>15</sup>

Although the enzymatic system contains an  $\text{Fe}(\text{II})$  ion in its active site, no functional mimic of ACCO reported so far involves  $\text{Fe}(\text{II})$ . Therefore, we tested complex **2** in the oxidation




Fig. 3 X-ray crystal structure of the  $[\text{Fe}(\text{BPMEN})\text{ACC}]\text{SbF}_6$  complex (**2**) (thermal ellipsoids are set at 50% probability). The hydrogen atoms and the counter ion are omitted for the sake of clarity. Selected bond lengths: Fe–N1 2.203(2) Å, Fe–N2 2.232(2) Å, Fe–N3 2.242(3) Å, Fe–N4 2.203(2) Å, Fe–N5 2.211(2) Å, Fe–O1 2.0164(19) Å.



of ACC into ethylene, first using O<sub>2</sub> and then in the presence of H<sub>2</sub>O<sub>2</sub>. The UV-vis spectrum of acetonitrile solutions of complex 2 did not change when O<sub>2</sub> was introduced. In contrast, when 10 equivalents of H<sub>2</sub>O<sub>2</sub> were added to an acetonitrile solution of complex 2, its UV-vis spectrum changed drastically, however, no clean transformation with isosbestic points could be observed (Fig. S4†). The addition of up to 100 equivalents of H<sub>2</sub>O<sub>2</sub> to complex 2 was then performed in sealed tubes and GC analysis of the resulting gas revealed that the formation of ethylene reached ca. 23% yield when 5 to 10 equivalents of H<sub>2</sub>O<sub>2</sub> were added, compared to a 15% yield in the blank experiment using a 1:1 mixture of iron(II) triflate and N(*n*-Bu)<sub>4</sub>ACC. The rather low ACC oxidation yield is not surprising considering the fact that complex 2 is hexacoordinated and thus, a direct interaction between the iron cation and hydrogen peroxide requires the de-coordination of one of the six ligands of iron. The formation of ethylene suggests that one of these six ligands is indeed labile enough to allow hydrogen peroxide activation at the metal center.

In summary, our work describes the synthesis, the reactivity and the characterization in solution and in the solid state of the first mononuclear Fe(II) complex bearing an ACC ligand. This complex demonstrates that ACC can bind to the Fe(II) ion in a bidentate mode, constituting a structural mimic of the binding of ACC to the Fe(II) center of ACCO.

We are grateful to Dr Eric Rivière for SQUID measurements and to the 'Agence Nationale de la Recherche' (ANR BIOXICAT and REBAR) and 'Région île de France' (DIM développement soutenable) for financial support.

## Notes and references

- (a) G.-P. L. Girardin, *Fortschr. Gesamtgeb. Agrikulturchem.*, 1864, 7, 199–200; (b) D. Neljubow, *Beih. Bot. Zentralbl.*, 1901, 10, 128–139.
- (a) T. Boller, R. Herner and H. Kende, *Planta*, 1979, 145, 293–303; (b) D. O. Adams and S. F. Yang, *Proc. Natl. Acad. Sci. U. S. A.*, 1979, 76, 170–174.
- (a) M. Costas, M. P. Mehn, M. P. Jensen and L. Que Jr., *Chem. Rev.*, 2004, 104, 939–986; (b) S. F. Yang and N. E. Hoffman, *Annu. Rev. Plant Physiol.*, 1984, 35, 155–189; (c) D. J. McGarvey and R. E. Christoffersen, *J. Biol. Chem.*, 1992, 267, 5964–5967.
- Z. Zhang, J.-S. Ren, I. J. Clifton and C. J. Schofield, *Chem. Biol.*, 2004, 11, 1383–1394.
- (a) V. J. Lay, A. G. Prescott, P. G. Thomas and P. John, *Eur. J. Biochem.*, 1996, 242, 228–234; (b) Z. Zhang, J. N. Barlow, J. E. Baldwin and C. J. Schofield, *Biochemistry*, 1997, 36, 15999–16007.
- (a) E. L. Hegg and L. Que Jr., *Eur. J. Biochem.*, 1997, 250, 625–629; (b) K. Koehntop, J. Emerson and L. Que Jr., *J. Biol. Inorg. Chem.*, 2005, 10, 87–93.
- (a) A. M. Rocklin, D. L. Tierney, V. Kofman, N. M. W. Brunhuber, B. M. Hoffman, R. E. Christoffersen, N. O. Reich, J. D. Lipscomb and L. Que Jr., *Proc. Natl. Acad. Sci. U. S. A.*, 1999, 96, 7905–7909; (b) A. Rocklin, K. Kato, H. Liu, L. Que Jr. and J. Lipscomb, *JBIC, J. Biol. Inorg. Chem.*, 2004, 9, 171–182; (c) J. Zhou, A. M. Rocklin, J. D. Lipscomb, L. Que Jr. and E. I. Solomon, *J. Am. Chem. Soc.*, 2002, 124, 4602–4609; (d) D. L. Tierney, A. M. Rocklin, J. D. Lipscomb, L. Que Jr. and B. M. Hoffman, *J. Am. Chem. Soc.*, 2005, 127, 7005–7013.
- W. Ghattas, Z. Serhan, N. El Bakkali-Taheri, M. Réglier, M. Kodera, Y. Hitomi and A. J. Simaan, *Inorg. Chem.*, 2009, 48, 3910–3912.
- (a) W. Ghattas, C. Gaudin, M. Giorgi, A. Rockenbauer, A. J. Simaan and M. Reglier, *Chem. Commun.*, 2006, 1027–1029; (b) W. Ghattas, M. Giorgi, C. Gaudin, A. Rockenbauer, M. Reglier, Y. Hitomi and A. J. Simaan, *Bioinorg. Chem. Appl.*, 2007, 43324; (c) W. Ghattas, M. Giorgi, Y. Mekmouche, T. Tanaka, A. Rockenbauer, M. Réglier, Y. Hitomi and A. J. Simaan, *Inorg. Chem.*, 2008, 47, 4627–4638; (d) N. Judaš and N. Raos, *Inorg. Chem.*, 2006, 45, 4892–4894.
- (a) M. Costas, K. Chen and L. Que Jr., *Coord. Chem. Rev.*, 2000, 200–202, 517–544; (b) E. P. Talsi and K. P. Bryliakov, *Coord. Chem. Rev.*, 2012, 256, 1418–1434.
- C. P. Magill, C. Floriani, A. Chiesi-Villa and C. Rizzoli, *Inorg. Chem.*, 1994, 33, 1928–1933.
- (a) K. Chen and L. Que Jr., *Chem. Commun.*, 1999, 1375–1376; (b) M. C. White, A. G. Doyle and E. N. Jacobsen, *J. Am. Chem. Soc.*, 2001, 123, 7194–7195.
- G. J. P. Britovsek, J. England and A. J. P. White, *Inorg. Chem.*, 2005, 44, 8125–8134.
- A. J. Simaan, S. Poussereau, G. Blondin, J.-J. Girerd, D. Defaye, C. Philouze, J. Guilhem and L. Tchertanov, *Inorg. Chim. Acta*, 2000, 299, 221–230.
- (a) K. P. Bryliakov, E. A. Duban and E. P. Talsi, *Eur. J. Inorg. Chem.*, 2005, 299, 72–76; (b) G. J. P. Britovsek, J. England and A. J. P. White, *Inorg. Chem.*, 2005, 44, 8125–8134.

