


***trans*-1,2-Disiloxybenzocyclobutene + O<sub>2</sub>  
&  
pseudopotentials for hybridized carbon atoms**


Yannick Carissan

*trans*-1,2-Disiloxybenzocyclobutene + O<sub>2</sub>

## Secondary reaction in lactonamycinone synthesis



### Reaction with O<sub>2</sub>





With benzocyclobutene same reaction at 190-240 °C. [1]

---

[1]. W. R. Roth, T. Ebbrecht, A. Breit *Chem.Ber.* **1988** *121*, 1357-1358

## Suggested pathway





## Theory : opening and color




-20 °C



T amb.




Ground and first excited state :




Absorbs blue : appears yellow


## Exp : UV analysis




## Suggested pathway



Intermediate exists



## Intermediate : proof of existence




Energy	kJ/mol	cm <sup>-1</sup>
singlet	0.35	30
triplet	0	0

Transition state of O<sub>2</sub> approach at 15kcal/mol.




Spin trapping : the two radical centers are characterized.

Theory : Singlet and triplet states are quasi degenerated.

## Suggested pathway




ISC




CAS(8,8) + CISD along the closing of C-O.

## Complete pathway



[1]. Phys. Chem. Chem. Phys., 2014, 16, 7513


## Photoactivation




## Photoactivation

Theory

Experiment




Emission with excitation at 400nm




## Photoactivation

Theory




Experiment

Emission with excitation at 312nm



## Photoactivation



## Pseudopotentials for hybridized carbon atoms


## General aim

- A new model :

  - Treat explicitly electrons of interest only ( $\pi$ )
  - Replace part of a molecule by a simpler object

- Constraint :

  - Use any quantum chemistry package
  - Use any quantum chemistry method



J. Am. Chem. Soc. 2009, 131, 12218-12229

## Atomic pseudopotentials

A non local atomic pseudopotential reads :

$$\hat{P} = a|\varphi\rangle\langle\varphi| \quad (1)$$

$$\text{with } \varphi = R(r)Y_m^l(\theta, \phi) \quad (2)$$

$\hat{P}$  is a shift operator on orbital  $\varphi$  : it modifies  $\epsilon_\varphi$ .

$R(r)$  is usually a gaussian function  $\exp(-\alpha r^2)$ .

$Y_m^l$  defines the angular momentum influenced by the potential.

## Atomic pseudopotentials : Perturbation analysis

$$\hat{F}^{(0)} \Phi_n^{(0)} = \epsilon_n^{(0)} \Phi_n^{(0)} \quad (3)$$

$$\hat{F} = \hat{F}^{(0)} + a|\varphi\rangle\langle\varphi| \quad (4)$$

$$\epsilon_n = \underbrace{\langle \Phi_n^{(0)} | \hat{F}^{(0)} | \Phi_n^{(0)} \rangle}_{\epsilon_n^{(0)}} + \underbrace{\langle \Phi_n^{(0)} | a | \varphi \rangle \langle \varphi | \Phi_n^{(0)} \rangle}_{\Delta\epsilon_n} \quad (5)$$

$$\Delta\epsilon_n = a \langle \varphi | \Phi_n^{(0)} \rangle^2 \quad (6)$$

Two parameters to modify the original orbitals :


1.  $a$  (eigenvalue)
2. shape of  $\varphi$  (angular momentum and spatial extent)


## From atomic to molecular shaped pseudopotentials

We aim at treating the  $\pi$  electrons *only*. For carbon atoms :

- 1 electron in 1  $p_z$  atomic orbital ;
- effective atomic charge  $Z=1$ .

Ring a bell ?








(a)  $\hat{F} = \hat{F}^{(0)}$

$z$




$z=1$




(a)  $\hat{F} = \hat{F}^{(0)}$



(b)  $\hat{F} = \hat{F}^{(0)} + \hat{P}_s$




## The $\pi$ interaction




- Shape of orbitals OK
- Excitation energy WRONG
- Ionization potential WRONG

## The $\pi$ interaction




- Shape of orbitals OK
- Excitation energy OK
- Ionization potential OK


## Combination of CC bonds : all trans even polyenes



## Fused rings




$\sigma$  bond



Extraction of  $C_{\sigma,\pi}$  :


- $Z=2$
- 1 electron in  $p_z$  and 1 electron in  $sp$



$$\hat{P}_{s_{\sigma-}} = c|Y_0^0\rangle\langle Y_0^0|$$

$$\hat{P}_{s_{\sigma+}} = c|Y_0^0\rangle\langle Y_0^0|$$

## Bonding to an active part



X	Y	dCC	dCX
H	C <sub>6</sub> H <sub>5</sub>	133.1	107.2
	(C) <sub>5</sub> C <sub>\sigma,\pi</sub>	132.0	107.3
Cl	C <sub>6</sub> H <sub>5</sub>	132.2	180.7
	(C) <sub>5</sub> C <sub>\sigma,\pi</sub>	131.6	181.1
Br	C <sub>6</sub> H <sub>5</sub>	132.4	193.2
	(C) <sub>5</sub> C <sub>\sigma,\pi</sub>	131.7	193.5
Me	C <sub>6</sub> H <sub>5</sub>	133.2	150.9
	(C) <sub>5</sub> C <sub>\sigma,\pi</sub>	132.4	151.1

## Conclusion

- We could create pseudopotentials for hybridized carbon atoms
- Use any QC package and any QC method
- Spectroscopy and geometry optimization
- Extension to other atoms with  $\pi$  electrons
- Other kind of hybridization tricky ( $\sigma$   $\pi$  orthogonality used)

Merci

- UV analysis
  - Jalila Simaan
  - Bruno Faure
- Co-workers
  - Béatrice Tuccio
  - Laurent Commeiras
  - Jean-Luc Parrain
- iSm2
- You for your attention

