



HAL
open science

Seasonal influenza vaccine uptake among people with disabilities: A nationwide population study of disparities by type of disability and socioeconomic status in France

Aurélie Bocquier, Lisa Fressard, Alain Paraponaris, Bérengère Davin, Pierre Verger

► To cite this version:

Aurélie Bocquier, Lisa Fressard, Alain Paraponaris, Bérengère Davin, Pierre Verger. Seasonal influenza vaccine uptake among people with disabilities: A nationwide population study of disparities by type of disability and socioeconomic status in France. *Preventive Medicine*, 2017, 101, pp.1 - 7. 10.1016/j.ypmed.2017.05.014 . hal-01683988

HAL Id: hal-01683988

<https://amu.hal.science/hal-01683988>

Submitted on 13 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Seasonal influenza vaccine uptake among people with disabilities: a nationwide population study of disparities by type of disability and socioeconomic status in France

People with disabilities and flu vaccine

Aurélie Bocquier^{1,2}, MS; Lisa Fressard^{1,2}, MS; Alain Paraponaris^{2,3}, PhD; Bérengère Davin^{1,2}, PhD; Pierre Verger^{1,2}, MD, MS.

¹ Aix Marseille Univ, INSERM, IRD, SESSTIM, Economics and Social Sciences Applied to Health & Analysis of Medical Information, Marseille, France.

² ORS PACA, Southeastern Health Regional Observatory, Marseille, France.

³ Aix Marseille Univ, CNRS, EHESS, Centrale Marseille, GREQAM, Marseille, France.

Address correspondence to: Aurélie Bocquier, ORS PACA, 27 bd Jean Moulin - 13385 Marseille cedex 5, France. E-mail address: aurelie.bocquier@inserm.fr

Conflict of interest statement: None

Abstract

People with disabilities use various preventive health services less frequently than others, notably because of a lower socioeconomic status (SES). We examined variations of seasonal influenza vaccine uptake according to type/severity of disability and SES. We analyzed (in 2016) data from the 2008 French national cross-sectional survey on health and disability (n=12,396 adults living in the community and belonging to target groups for seasonal influenza vaccination). We defined seasonal influenza vaccine uptake during the 2007-2008 season by the self-reporting of a flu shot between September 2007 and March 2008. We built scores of mobility, cognitive, and sensory limitations, and an SES score based on education, occupation, and income. We performed bivariate analyses and then multiple log-binomial regressions. The prevalence of vaccine uptake was 23% in the 18-64 group and 63% in the ≥ 65 group. In bivariate analyses, it was higher among people in both age groups who had mobility and/or cognitive limitations and in the ≥ 65 group among those with sensory limitations. In the multiple regression analyses, only the presence of major mobility limitations in the 18-64 group remained significant. The probability of vaccine uptake was higher in the highest SES category than in the lowest. Among at-risk groups, people with disabilities were more frequently vaccinated than others, mainly because of their higher levels of morbidity and healthcare use. Socioeconomic inequalities in access to vaccination persist in France. Future research is needed to monitor the trend in vaccine uptake in institutions.

Keywords: Influenza Vaccines; Disabled Persons; Socioeconomic Factors; Social Determinants of Health; France

Introduction

People with disabilities, especially neurological and neurodevelopmental conditions and disabilities related to problems with muscle tone, weakness, or handling of secretions, are at high risk of influenza-related complications.¹⁻³ Many countries thus recommend vaccination against seasonal influenza for these populations;^{1,4,5} since 2011, this includes France.⁶

Evidence from the literature – mainly from the United States (US) – shows that use of various preventive health services is less frequent among people with, compared to without, disabilities.⁷⁻¹⁴ Apart from physical barriers to access to health care, major contributors to these inequities include attitudinal barriers associated with healthcare providers' perceptions of people with disabilities.^{15,16} They also include socioeconomic barriers; lower socioeconomic status (SES) is more common among people with disabilities¹⁶ and is independently associated with less frequent use of several preventive care services (e.g., cancer screening).^{14,17,18}

The few studies that have addressed seasonal influenza vaccine uptake among people with disabilities took place in the US,^{10,11,19} Australia,²⁰ and Taiwan;²¹ their results suggest that disability is associated with a higher probability of vaccine uptake. This question has not been studied in Europe, and in particular not in France (see Appendix Table S1 for the bibliographic search). Findings about the association between seasonal influenza vaccine uptake and SES are more controversial.²²⁻²⁴ Lower SES has been found to be correlated with lower vaccination uptake in some countries, but a reverse gradient has been found in others (e.g., among older people in Italy).²⁵ In France, a study in the general population found no significant SES gradient in vaccine uptake²⁶ while among people treated for diabetes, those with a low income were less likely to be vaccinated regularly against seasonal influenza than more affluent individuals.²⁷

In France as in many other developed countries, reducing social health inequalities (i.e., avoidable and unfair differences in health status/health determinants between different population groups) has become a priority.²⁸ According to the conceptual framework set forth by the World Health Organization Commission on Social Determinants of Health, these inequalities arise from: 1) structural determinants: macro-level social and political mechanisms that generate and maintain social hierarchies and stratification by income, education, occupation, and other factors; and 2) intermediary determinants (e.g., health system, material circumstances, behaviors, beliefs) through which the structural factors shape health outcomes.²⁸ In France, the influenza vaccine is available free of charge for people in at-risk groups through vouchers, but a significant fraction of the at-risk population escapes this system.²⁷ Nevertheless, little is known about the social health inequalities in seasonal influenza and vaccination against it.

Using data from the French national survey on health and disability among people living in the community (*Enquête Handicap Santé-Ménages*, HSM survey), we sought to examine associations between seasonal influenza vaccine uptake and 1) the presence of disabilities, while distinguishing their type and severity; and 2) individual SES.

Materials and Methods

Sampling and data source

The HSM cross-sectional survey was conducted from April through September 2008 by the French national institute of statistics and economic studies (INSEE) and the Ministry of Social Affairs office of research, studies, evaluation and statistics (DREES).^{29,30} This survey aimed to measure the prevalence of various forms of disabling situations, assessed according to the concepts of disability developed by the World Health Organization (WHO).³¹

It was preceded in 2007 by a filter survey – the Everyday Life and Health Survey (*Enquête Vie Quotidienne et Santé*, VQS survey) – that allowed the HSM sampling frame to be built.²⁹

During the VQS survey, 101,930 households and around 260,000 individuals (for an 80% response rate from the initial sample of 127,176 households) answered a 26-item questionnaire, either by mail (about half) or in telephone or face-to-face interviews to assess the presence or absence of recognized disabilities and difficulties in accomplishing certain tasks. Respondents were then classified into four levels of disability severity, previously used in French studies.^{13,30}

Among these respondents, 39,065 were randomly selected to participate in the HSM survey. Because people with disabilities are relatively rare in the general population, they were intentionally overrepresented in the HSM sample, through the use of a sampling coefficient that increased with the presence and severity of disabilities.³² Professionals conducted face-to-face interviews at respondents' homes, using a computer-assisted personal interviewing (CAPI) system. A proxy respondent (e.g., a relative or friend or neighbor) could answer for HSM participants not able to answer alone. Finally, 29,931 subjects participated in the HSM survey (response rate: 76.6%).

The national commission for computer data and individual freedom (Commission Nationale de l'Informatique et des Libertés, CNIL) approved this study.

Study population

This study analyzed data from the HSM participants aged 18 or more at the time of the interview who lived in metropolitan France and belonged to target groups for seasonal influenza vaccination in France for the 2007-2008 season.³³ The French guidelines³³ recommend this vaccination for all people aged 65 years or over, people with specific chronic diseases, and health and other professionals in regular and extended contact with them (see Appendix Box S1). The HSM survey used a standardized list of 51 diseases presented to the participants during the interview to record the presence of diseases. The study population

included 12,396 respondents (see Appendix Table S2 for the number of persons per target group).

Dependent variable: seasonal influenza vaccine uptake

Participants were asked “*Have you ever had the flu vaccine?*” and “*If yes, when did you have your last flu shot?*” (*year, month*).²⁰ As the survey took place from April through September 2008, we defined the dependent variable, *seasonal influenza vaccine uptake during the 2007-2008 season (yes/no)*, hereafter referred as vaccine uptake, by the report of a seasonal influenza vaccine shot between September 2007 and March 2008.³⁴

Variables of interest

Type and severity of disability

The questionnaire collected information about respondents' functional limitations in three domains (see Appendix Box S2 for more details): 1) mobility limitations (7 items, e.g., level of difficulty in walking 500 meters on flat ground); 2) sensory limitations (3 items, e.g., level of difficulty in seeing, even with eyeglasses or contact lenses, the printed letters in newspapers); and 3) cognitive limitations (8 items, e.g., level of frequency of difficulty in acquiring new knowledge or skills, in school, professional training, or leisure activities).

Following a methodology commonly used in the literature,³⁵ we built a score of limitation severity for each kind of limitation by summing respondents' answers to the corresponding items, then dividing the sum by the number of items, and finally rounding the resulting score to the nearest integer to preserve the initial form of the limitation categories. The mobility and sensory scores ranged theoretically from 1 to 4, and the cognitive score from 1 to 3. Because few respondents were in the highest categories, the first two scores were then re-coded into three categories (1: no limitation; 2: moderate limitations; 3 or 4: major limitations), and the latter into two categories (1: no limitation; 2 or 3: moderate/major limitations).

We also used one question about whether the respondent received family assistance in performing daily living activities to evaluate disability severity.

Socioeconomic status

The socioeconomic variables included were: education level (did not complete high school, completed high school, some higher education); occupation (including but not limited to: not in the labor force, such as homemakers and people with disabilities; workers, such as farmers, artisans, factory workers, and office/sales/service workers; professionals and managers, including middle management and other intellectual workers); and equivalized household income (EHI) per month, which incorporates household size and composition.³⁶ According to a methodology already published (see Appendix Box S3),^{37,38} we used these three variables to calculate an individual composite SES indicator ranging from 0 (lowest SES) to 6 (highest SES). Because only a few respondents younger than 65 had an SES score of 0, they were grouped with those in category 1.

Other variables

The HSM questionnaire also assessed the following sociodemographic characteristics: gender; age (four categories, Table 1); living with a partner (*yes/no*); and type of residence area (*urban/rural*).

Data on health status and healthcare use included self-reporting of chronic diseases (i.e., those lasting 6 months or more) and number of visits to a general practitioner in the previous 12 months (dichotomized around the median = 4).

Statistical analysis

All HSM respondents were assigned a weight that reflected their probability of selection (depending on disability severity and geographic area of residence) and response. Final weights were calculated with a calibration procedure ensuring that the data were

representative of the French population living in the community for gender, age, and geographic area of residence.³²

First, we performed bivariate analyses with chi-square tests to describe variations of the prevalence of seasonal influenza vaccine uptake according to the variables of interest (the three limitation scores, family assistance, and the SES score), and other characteristics described above. Second, we performed weighted multiple log-binomial regression models to measure associations between seasonal influenza vaccine uptake and the variables of interest. Based on Andersen's behavioral model of health services use,³⁹ multiple regression models were adjusted for various factors: predisposing (age, gender, or living with a partner), enabling (type of residence area, as a proxy for availability of health services), and need (health status, number of visits to a general practitioner, and use of a proxy respondent). All these variables may be associated with both the variables of interest and vaccine uptake.^{22,40} Associations were measured with adjusted risk ratios (aRRs) and their 95% confidence intervals (95% CI). We performed log-binomial rather than logistic regressions because vaccine uptake is not rare (>10%); in this situation, odds ratios estimate risk ratios poorly; specifically, they overestimate them. Moreover, compared to the Robust Poisson method, the log-binomial method yields slightly less biased estimates, higher power, and smaller standard errors in most common situations.⁴¹

Chronic disease and healthcare use were strongly and positively correlated with both disabilities and seasonal vaccine uptake^{19,22} and may help explain the higher probability of vaccine uptake among people with disabilities.^{10,11,19-21} To explore the role of these two variables in the associations of interest, we performed two successive multiple regression models, first without and then with adjustment for these variables. We also tested interactions between SES and each of the three limitation scores.

All analyses were stratified by age (18-64/ \geq 65 years) for the two reasons. First, influenza vaccine coverage differs very substantially between these two groups.²⁷ Second, the national health insurance fund sends all people aged 65 or more a free vaccination voucher; although at-risk people aged 18-64 are also supposed to receive this voucher, a significant fraction escapes this system.²⁷

All analyses were based on two-sided p values, with $p < 0.05$ indicating statistical significance; they were conducted in 2016 with SAS 9.4 statistical software (SAS Institute, Cary, NC).

Results

Characteristics of the study population

Among the study population, 57% of people were aged 65 years or more (Table 1). Analysis of SES distribution shows 14% of respondents aged 18-64 and 30% of those aged 65 or older in the two lowest SES categories and 25% and 10%, respectively, in the two highest. Among people aged 18-64 years, 9% reported mobility limitations, 8% sensory limitations, and 7% cognitive limitations; among older respondents, these percentages were 32%, 21%, and 13% (Table 1). In both age groups, respondents with disabilities had a significantly lower SES than the others (see Appendix Table S3).

Prevalence of seasonal influenza vaccine uptake and results from bivariate analyses

Among the study population, 12,365 (99.7%) respondents answered the questions about seasonal influenza vaccine uptake: 44% reported a shot during the 2007-2008 season (63% among those aged 65 years or more; 23% among the at-risk members of the younger group). In both age groups, vaccine uptake prevalence was higher among men, older people, and those living with a partner. Only in the younger age group did we find better coverage in those with the highest SES category. It was also higher among people who had mobility and/or cognitive limitations in both age groups, and, only among the older age group, those who had sensory limitations (Table 2).

Multiple log-binomial regression analyses

The probability of vaccine uptake was higher among people with moderate or major physical limitations in both age groups, and among those aged 18-64 years with cognitive limitations; it was slightly lower in the younger group with moderate sensory limitations (Tables 3 and 4 and Tables S4 and S5 for results for covariables). After adjustment for chronic disease and healthcare use, only major physical limitations among people aged 18-64 years remained significantly associated with a higher probability of vaccine uptake (aRR=1.37 [1.02;1.84]). Those in the highest SES category had a higher probability of vaccine uptake than those in the lowest SES category, in both age groups (18-64: aRR=1.51 [1.25-1.81]; ≥ 65 : aRR=1.11 [1.02-1.21], Tables 3 and 4).

We found no significant interaction between the SES score and any of the limitation scores.

Discussion

In this study based on a representative sample of the French population living in the community, the prevalence of seasonal influenza vaccine uptake during the 2007-2008 season was 23% among at-risk respondents aged 18-64 and 63% among the older group (all at risk, by definition). In unadjusted analyses, uptake prevalence was higher among people with mobility and/or cognitive limitations; among people aged 18-64 only, it was also higher among those with the highest SES. After adjustment, especially for health status and health care use, only severe mobility limitations among people aged 18-64 and the highest SES category in both age groups were associated with a higher uptake prevalence.

Our estimates of the prevalence of seasonal influenza vaccine uptake during the 2007-2008 season were consistent with those of a previous study based on reimbursement data for the same season in the entire French population (24% among people aged 18-64 at risk of chronic disease; 63% among those at risk because aged 65 or older).³⁴ Their results confirm that, just before the 2009 A/H1N1 pandemic,^{42,43} influenza vaccine uptake was lower than the target of

75% set by the WHO and the French Public Health Policy Act of 2004 in 2007-8, although the target was close to achievement (71%) among people aged 65 years or more with mobility limitations.

Consistently with previous studies,^{10,11,19-21} we found that rates of seasonal vaccine uptake were higher among people with physical or cognitive limitations, despite their lower SES (a risk factor for non-immunization), than among the general population. We also showed that this higher rate was due in part to their higher morbidity level and more regular health care than that of the rest of the population.^{8,11,19} After adjustment for these factors, seasonal influenza vaccine uptake remained higher only among a very small group of people aged 18-64 with major physical limitations (2% of the study population). Some of them may have severe neurological diseases or neuromuscular conditions³⁰ that entitled them to a free vaccination voucher from the French National Health Insurance Fund several years before their conditions were listed as targeted in the official French guidelines in 2011.⁴⁴ This feedforward measure may have contributed to a greater attention by physicians to the vaccination of these patients. Although flu vaccination is recommended for people with significant cognitive/intellectual disabilities in some countries (e.g., the UK),⁴ this is not the case in France.

Our study suggests that the general tendency for people with disabilities to receive fewer preventive services (e.g., cervical and breast cancer screening)⁷⁻¹⁴ does not apply to seasonal influenza vaccination. Greater awareness by physicians of these patients' risks of seasonal flu than of their needs for other preventive services might partly explain this result.⁴⁵ There may also be fewer physical barriers to influenza shots than to cancer screening and dental cleaning, which may require medical equipment specifically adapted to people with disabilities.¹¹ Our results suggest the persistence of socioeconomic inequalities in seasonal influenza vaccine uptake in France involving especially people with disabilities, due to their SES, which

is generally lower than among the general population. The international literature about the existence of these inequalities is inconsistent;²⁵ inequalities have already been reported for this vaccine in France among people with diabetes²⁷ and for pneumococcal vaccine among children (partially free of charge).⁴⁶ The French National Health Insurance Fund identifies members of at-risk groups and sends them a free vaccination voucher; they thus do not pay for their seasonal influenza vaccine. But financial barriers still remain for people in at-risk groups not eligible for a voucher and/or those who have to pay the consultation fee.²⁷ Adjustment for complementary health insurance (*yes/no*), a traditional enabling factor of health service use,³⁹ did not modify these results (results not shown). These inequalities may also reflect SES differences in other predisposing factors of health services use,³⁹ such as health-related knowledge, attitudes, and beliefs, and more specifically cognitive determinants of vaccine behaviors (e.g., beliefs in vaccine efficacy and safety, and fear of side effects). A previous French study reported greater reluctance about vaccination in general among low SES people than among the affluent.⁴⁷ Previous studies in the field of cancer prevention (e.g., cancer screening¹⁷ and sun-protection behaviors³⁸) have showed that cognitive factors play a role in mediating SES differences in health behaviors. Further research is needed to verify whether this also applies to vaccination behavior.

Limitations and strengths

This study has several potential limitations. First, as in most previous studies exploring associations between disabilities and influenza vaccination,^{7,8,11,19,20,40} we used self-reported data on vaccine uptake, without physician confirmation. The prevalence of vaccine uptake here, however, was very close to that calculated from reimbursement (thus objective) data in France for the same season.³⁴ Second, no study since 2008 has explored associations between disability and influenza vaccination. This raises a question about whether its results apply today. That study was implemented before the 2009-2010 controversy about the safety of the

A/H1N1 pandemic vaccine,⁴⁸ which has durably affected seasonal influenza vaccine uptake in several countries. In France in particular, uptake has progressively decreased to reach, in 2015-16, levels lower than those in 2007-8.⁴⁹ There is no reason to think that this has affected the association between uptake and physical limitations, however, given the absence of any recent major changes in disability trends.⁵⁰ The social differentiation of attitudes towards vaccination in general in France increased between 2005 and 2010, with low SES becoming more predictive of unfavorable attitudes in 2010.⁵¹ The long-lasting consequences of the 2008 economic crisis in France may have contributed to an increase in social inequalities. These observations suggest that our results about SES differences in vaccine uptake are likely to continue to exist and might even be greater today.

Major strengths of this study include the size of the sample and its representativeness of the French population. Moreover, its design enabled us to obtain robust results about population subgroups by type and severity of disability, which has been a weakness of previous studies on preventive care use among people with disabilities.⁸ Using a SES score combining three different measures (education, occupation, and income) is likely to have limited potential misclassification related to each separate variable.

Conclusions

Our study suggests that the poorer use of prevention among people with disabilities previously observed for several types of preventive care does not exist for seasonal influenza vaccination. Among the at-risk groups, people with disabilities were more frequently vaccinated than other people, mainly because of their greater level of morbidity and healthcare use. After adjustment for these factors, only a few people with major mobility limitations were more likely to be vaccinated than those without limitation. Nonetheless, our results highlight the persistence of socioeconomic inequalities in vaccination that affect people with disabilities despite the existence of a free voucher system. These results should

prompt public health professionals, in collaboration with other partners, to continue to combat health inequalities and to consider the different social determinants of health more globally. Future research is needed to improve our understanding of the reasons for these differences and to examine influenza vaccination uptake among people with disabilities living in institutions.

Acknowledgements

This study was conducted with the financial support of the Groupement d'intérêt scientifique Institut de recherche en santé publique (GIS-IReSP) as part of their 2013 call for research projects IReSP-CNSA-MiRe/DREES "Handicap et perte d'autonomie - session 4". The GIS-IReSP had no involvement in the study design; collection, analysis and interpretation of data; the writing of the manuscript; or the decision to submit the manuscript for publication.

Highlights

- Influenza vaccine uptake is more frequent among people with mobility limitations.
- After adjustment for morbidity, this is true only for a few with major limitations.
- Socioeconomic inequalities persist in influenza vaccine uptake.

References

1. Centers for Disease Control and Prevention. People at High Risk of Developing Flu–Related Complications. http://www.cdc.gov/flu/about/disease/high_risk.htm. Accessed March 26, 2017.
2. Keren R, Zaoutis TE, Bridges CB, et al. Neurological and neuromuscular disease as a risk factor for respiratory failure in children hospitalized with influenza infection. *JAMA* 2005;294(17):2188–94. doi:10.1001/jama.294.17.2188.
3. Illinois Department of Human Services. Protection Against the Flu: Advice for Caregivers of People with Disabilities. <http://www.dhs.state.il.us/page.aspx?item=32138>. Accessed March 26, 2017.
4. National Health System. All about flu and how to stop getting it EasyRead version for people with learning disabilities. 2016. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/530741/9833_PHE_Flu-learning-disability-A4-8pp-6-WEB.pdf. Accessed March 26, 2017.
5. Australian Government Department of Health. Immunise Australia Program. Medically at risk. Immunise Aust Program 2015. <http://www.health.gov.au/internet/immunise/publishing.nsf/content/medically-at-risk-groups>. Accessed March 26, 2017.
6. Haut Conseil de la santé publique. Le Calendrier des vaccinations et les recommandations vaccinales 2011 selon l’avis du Haut Conseil de la santé publique. *Bull Epidemiol Hebd* 2011;10–11:101–56.
7. Chan L, Doctor JN, MacLehose RF, Lawson H, Rosenblatt RA, Baldwin LM, et al. Do Medicare patients with disabilities receive preventive services? A population-based study. *Arch Phys Med Rehabil*. 1999;80(6):642-6.
8. Diab ME, Johnston MV. Relationships between level of disability and receipt of preventive health services. *Arch Phys Med Rehabil*. 2004;85(5):749-57.
9. Ramirez A, Farmer GC, Grant D, Papachristou T. Disability and preventive cancer screening: results from the 2001 California Health Interview Survey. *Am J Public Health*. 2005;95(11):2057-64.
10. Parish SL, Saville AW. Women with cognitive limitations living in the community: evidence of disability-based disparities in health care. *Ment Retard*. 2006;44(4):249-59.
11. Pharr JR, Bungum T. Health disparities experienced by people with disabilities in the United States: a Behavioral Risk Factor Surveillance System study. *Glob J Health Sci*. 2012;4(6):99-108.
12. Peterson-Besse JJ, O’Brien MS, Walsh ES, Monroe-Gulick A, White G, Drum CE, et al. Clinical preventive service use disparities among subgroups of people with disabilities: A scoping review. *Disabil Health J*. 2014;7(4):373-93.

13. Bussière C, Sicsic J, Pelletier-Fleury N. The effects of obesity and mobility disability in access to breast and cervical cancer screening in France: results from the national health and disability survey. *PloS One*. 2014;9(8):e104901.
14. Lupi-Pegurier L, Clerc-Urmes I, Abu-Zaineh M, Paraponaris A, Ventelou B. Density of dental practitioners and access to dental care for the elderly: a multilevel analysis with a view on socio-economic inequality. *Health Policy Amst Neth*. 2011;103(2-3):160-7.
15. Aulagnier M, Verger P, Ravaud JF, Souville M, Lussault PY, Garnier JP, et al. General practitioners' attitudes towards patients with disabilities: the need for training and support. *Disabil Rehabil*. 2005;27(22):1343-52.
16. World Health Organization, The World Bank. *World report on disability*. Geneva: WHO; 2011.
17. Wardle J, McCaffery K, Nadel M, Atkin W. Socioeconomic differences in cancer screening participation: comparing cognitive and psychosocial explanations. *Soc Sci Med*. 2004;59(2):249-61.
18. McCaffery K, Wardle J, Nadel M, Atkin W. Socioeconomic variation in participation in colorectal cancer screening. *J Med Screen*. 2002;9(3):104-8.
19. McGuire LC, Strine TW, Okoro CA, Ahluwalia IB, Ford ES. Healthy lifestyle behaviors among older U.S. adults with and without disabilities, Behavioral Risk Factor Surveillance System, 2003. *Prev Chronic Dis*. 2007;4(1):A09.
20. Dyda A, MacIntyre CR, McIntyre P, Newall AT, Banks E, Kaldor J, et al. Factors associated with influenza vaccination in middle and older aged Australian adults according to eligibility for the national vaccination program. *Vaccine*. 2015;33(29):3299-305.
21. Yen C-F, Hsu S-W, Loh C-H, Fang W-H, Wu C-L, Chu CM, et al. Analysis of seasonal influenza vaccine uptake among children and adolescents with an intellectual disability. *Res Dev Disabil*. 2012;33(2):704-10.
22. Nagata JM, Hernández-Ramos I, Kurup AS, Albrecht D, Vivas-Torrealba C, Franco-Paredes C. Social determinants of health and seasonal influenza vaccination in adults ≥ 65 years: a systematic review of qualitative and quantitative data. *BMC Public Health*. 2013;13:388.
23. Endrich MM, Blank PR, Szucs TD. Influenza vaccination uptake and socioeconomic determinants in 11 European countries. *Vaccine*. 2009;27(30):4018-24.
24. Ward J, Raude J. Understanding influenza vaccination behaviors: a comprehensive sociocultural framework. *Expert Rev Vaccines*. 2014;13(1):17-29.
25. Chiatti C, Di Rosa M, Barbadoro P, Lamura G, Di Stanislao F, Prospero E. Socioeconomic determinants of influenza vaccination among older adults in Italy. *Prev Med*. 2010;51(3-4):332-3.

26. Vaux S, Van Cauteren D, Guthmann J-P, Le Strat Y, Vaillant V, de Valk H, et al. Influenza vaccination coverage against seasonal and pandemic influenza and their determinants in France: a cross-sectional survey. *BMC Public Health*. 2011;11:30.
27. Verger P, Cortaredona S, Pulcini C, Casanova L, Peretti-Watel P, Launay O. Characteristics of patients and physicians correlated with regular influenza vaccination in patients treated for type 2 diabetes: a follow-up study from 2008 to 2011 in southeastern France. *Clin Microbiol Infect*. 2015;21(10):930.e1-9.
28. Solar O, Irwin A. A Conceptual Framework for Action on the Social Determinants of Health. Social Determinants of Health Discussion paper 2. Geneva: World Health Organization; 2010.
http://www.who.int/sdhconference/resources/ConceptualframeworkforactiononSDH_eng.pdf. Accessed March 26, 2017.
29. Bouvier G. L'enquête Handicap-Santé. Présentation générale. Insee; 2011.
30. Palazzo C, Ravaud J-F, Trinquart L, Dalichampt M, Ravaud P, Poiraud S. Respective contribution of chronic conditions to disability in France: results from the national Disability-Health Survey. *PloS One*. 2012;7(9):e44994.
31. World Health Organization. International Classification of Functioning, Disability and Health (ICF). <http://www.who.int/classifications/icf/en/>. Accessed March 26, 2017.
32. Bouvier G. Le volet Ménages de l'enquête Handicap-Santé. Présentation, calcul des poids. Insee; 2011.
33. Haut Conseil de la santé publique. Calendrier vaccinal 2007 - Avis du Haut conseil de la santé publique. *Bull Epidemiol Hebd* 2007;31-32:269-88.
34. Tuppin P, Samson S, Weill A, Ricordeau P, Allemand H. Seasonal influenza vaccination coverage in France during two influenza seasons (2007 and 2008) and during a context of pandemic influenza A(H1N1) in 2009. *Vaccine*. 2011;29(28):4632-7.
35. Bussière C, Le Vaillant M, Pelletier-Fleury N. Screening for cervical cancer: What are the determinants among adults with disabilities living in institutions? Findings from a National Survey in France. *Health Policy*. 2015;119(6):794-801.
36. National Institute of Statistics and Economic Studies. Definitions and methods. Definitions. Consumption unit 2014.
<http://www.insee.fr/en/methodes/default.asp?page=definitions/unite-consommation.htm>. Accessed October 19, 2016.
37. Stamatakis E, Hillsdon M, Mishra G, Hamer M, Marmot M. Television viewing and other screen-based entertainment in relation to multiple socioeconomic status indicators and area deprivation: the Scottish Health Survey 2003. *J Epidemiol Community Health*. 2009;63(9):734-40.
38. Bocquier A, Fressard L, Legleye S, Verger P, Peretti-Watel P. Social Differentiation of Sun-Protection Behaviors: The Mediating Role of Cognitive Factors. *Am J Prev Med*. 2016;50(3):e81-90.

39. Andersen RM. National health surveys and the behavioral model of health services use. *Med Care*. 2008;46(7):647-53.
40. Chiatti C, Barbadoro P, Lamura G, Pennacchietti L, Di Stanislao F, D'Errico MM, et al. Influenza vaccine uptake among community-dwelling Italian elderly: results from a large cross-sectional study. *BMC Public Health*. 2011;11:207.
41. Petersen MR, Deddens JA. A comparison of two methods for estimating prevalence ratios. *BMC Med Res Methodol*. 2008;8:9.
42. World Health Assembly. Prevention and control of influenza pandemics and annual epidemics. 2003.
43. Drees. L'état de santé de la population en France. Rapport 2015. Paris: Drees; 2015.
44. Haut Conseil de la santé publique. Avis relatif à l'actualisation de la liste des sujets éligibles à la vaccination contre la grippe saisonnière. 2010.
45. Anderson P, Kitchin R. Disability, space and sexuality: access to family planning services. *Soc Sci Med*. 2000;51(8):1163-73.
46. Guthmann J-P, Chauvin P, Le Strat Y, Soler M, Fonteneau L, Levy-Bruhl D. [Low pneumococcal conjugated vaccine immunization coverage in low-income families: a study in Ile-de-France]. *Arch Pediatr*. 2014;21(6):584-92.
47. Peretti-Watel P, Raude J, Sagaon-Teyssier L, Constant A, Verger P, Beck F. Attitudes toward vaccination and the H1N1 vaccine: poor people's unfounded fears or legitimate concerns of the elite? *Soc Sci Med*. 2014;109:10-8.
48. Black S, Rappuoli R. A crisis of public confidence in vaccines. *Sci Transl Med* 2010;2(61):61mr1. doi:10.1126/scitranslmed.3001738.
49. Santé Publique France. Couverture vaccinale grippe par saison et dans chaque groupe d'âge 2015. <http://invs.santepubliquefrance.fr/Dossiers-thematiques/Maladies-infectieuses/Maladies-a-prevention-vaccinale/Couverture-vaccinale/Donnees/Grippe>. Accessed October 19, 2016.
50. Verropoulou G, Tsimbos C. Disability trends among older adults in ten European countries over 2004–2013, using various indicators and Survey of Health, Ageing and Retirement in Europe (SHARE) data. *Ageing Soc*. 2016;1-31.
51. Peretti-Watel P, Verger P, Raude J, Constant A, Gautier A, Jestin C, et al. Dramatic change in public attitudes towards vaccination during the 2009 influenza A(H1N1) pandemic in France. *Euro Surveill*. 2013;18(44):20623.

Table 1. Characteristics of the study population, by age group (France, 2008 HSM survey, $n=12,396$)^a

| | 18-64 years ($n=5,297$) | | ≥ 65 years ($n=7,099$) | |
|--|---------------------------|------|-------------------------------|------|
| | <i>n</i> | % | <i>n</i> | % |
| Demographic and socioeconomic characteristics | | | | |
| Gender | | | | |
| Men | 2,295 | 43.3 | 2,930 | 41.3 |
| Women | 3,002 | 56.7 | 4,169 | 58.7 |
| Age (years) | | | | |
| 18-45 | 2,159 | 40.8 | - | - |
| 46-64 | 3,138 | 59.2 | - | - |
| 65-74 | - | - | 3,639 | 51.3 |
| 75 or more | - | - | 3,460 | 48.7 |
| Living with a partner | | | | |
| No | 1,440 | 27.2 | 2,752 | 38.8 |
| Yes | 3,857 | 72.8 | 4,347 | 61.2 |
| Type of residence area | | | | |
| Rural | 1,647 | 31.1 | 2,278 | 32.1 |
| Urban | 3,650 | 68.9 | 4,821 | 67.9 |
| Socioeconomic status (SES) score [0-6] ^b | | | | |
| 0-1 (lowest SES) | 710 | 13.5 | 2,113 | 29.9 |
| 2 | 1,294 | 24.6 | 2,413 | 34.3 |
| 3 | 1,305 | 24.8 | 1,241 | 17.6 |
| 4 | 654 | 12.4 | 563 | 8.0 |
| 5 | 493 | 9.4 | 336 | 4.8 |

| | | | | |
|--|-------|------|-------|------|
| 6 (highest SES) | 804 | 15.3 | 394 | 5.6 |
| Health status and healthcare use | | | | |
| Number of visits to a general practitioner in the last 12 months | | | | |
| Between 1 and 4 | 3,029 | 57.2 | 3,455 | 48.7 |
| 5 or more | 1,769 | 33.4 | 3,323 | 46.8 |
| Don't know | 499 | 9.4 | 320 | 4.5 |
| Chronic disease ^c | | | | |
| No | 1,834 | 34.6 | 1,927 | 27.2 |
| Yes | 3,462 | 65.4 | 5,168 | 72.8 |
| Disabilities | | | | |
| Mobility limitations ^d | | | | |
| No limitation | 4,837 | 91.3 | 4,805 | 67.7 |
| Moderate limitations | 353 | 6.7 | 1,503 | 21.2 |
| Major limitations | 106 | 2.0 | 791 | 11.1 |
| Sensory limitations ^e | | | | |
| No limitation | 4,878 | 92.1 | 5,616 | 79.1 |
| Moderate limitations | 392 | 7.4 | 1,242 | 17.5 |
| Major limitations | 26 | 0.5 | 241 | 3.4 |
| Cognitive limitations | | | | |
| No limitation | 4,900 | 92.5 | 6,149 | 86.6 |
| Moderate/major limitations | 397 | 7.5 | 950 | 13.4 |
| Needs family assistance for performing daily living activities | | | | |
| No | 4,731 | 89.3 | 5,307 | 74.8 |
| Yes | 566 | 10.7 | 1,792 | 25.2 |
| Answered to the question him/herself | | | | |

| | | | | |
|-----|------|------|-------|------|
| No | 266 | 5.0 | 793 | 11.2 |
| Yes | 5031 | 95.0 | 6,306 | 88.8 |

^a Weighted data items.

^b 73 missing data items.

^c 5 missing data items.

^d 1 missing data item.

^e 3 missing data items.

Accepted Version

Table 2. Prevalence of seasonal influenza vaccine uptake during the 2007-2008 season according to the respondents' characteristics and age group (France, 2008 HSM survey, $n=12,365^a$)

| | 18-64 years ($n=5,277$) | ≥ 65 years ($n =7,088$) |
|--|---------------------------|--------------------------------|
| | % [95% CI] ^b | % [95% CI] ^b |
| Total sample | 23.4 [21.0;25.7] | 63.3 [61.9;64.7] |
| Demographic and socioeconomic characteristics | | |
| Gender | | |
| Men | 25.4 [23.7;27.2] | 64.8 [63.0;66.5] |
| Women | 21.8 [20.3;23.3] | 62.3 [60.8;63.7] |
| Age (years) | | |
| 18-45 | 15.1 [13.4;16.8] | - |
| 46-64 | 29.1 [27.6;30.5] | - |
| 65-74 | - | 56.7 [54.9;58.4] |
| 75 or more | - | 70.3 [68.9;71.7] |
| Living with a partner | | |
| No | 21.0 [19.1;23.0] | 61.3 [59.6;63.0] |
| Yes | 24.3 [22.9;25.7] | 64.6 [63.1;66.0] |
| Type of residence area | | |
| Rural | 23.8 [21.5;26.1] | 63.6 [61.4;65.8] |
| Urban | 23.2 [21.9;24.5] | 63.2 [61.8;64.5] |
| Socioeconomic status (SES) score ^c | | |
| 0-1 (lowest SES) | 24.7 [22.3;27.0] | 62.4 [60.5;64.3] |
| 2 | 22.7 [20.6;24.8] | 62.7 [60.7;64.6] |
| 3 | 20.0 [17.6;22.4] | 64.3 [61.5;67.1] |

| | | |
|-----------------|------------------|------------------|
| 4 | 20.9 [17.4;24.4] | 67.0 [62.7;71.3] |
| 5 | 22.6 [18.0;27.2] | 64.4 [58.7;70.0] |
| 6 (highest SES) | 31.5 [27.3;35.7] | 62.2 [56.6;67.7] |

Health status and health care use

Number of visits to a general practitioner in the last 12 months

| | | |
|-----------------|------------------|------------------|
| Between 1 and 4 | 20.7 [19.0;22.4] | 58.2 [56.3;60.1] |
| 5 or more | 29.2 [27.5;30.9] | 71.8 [70.4;73.1] |
| Don't know | 18.9 [14.6;23.2] | 30.1 [24.3;36.0] |

Chronic disease^d

| | | |
|-----|------------------|------------------|
| No | 18.4 [16.0;20.9] | 55.3 [52.4;58.1] |
| Yes | 26.0 [24.7;27.3] | 66.3 [65.1;67.5] |

Disabilities

Mobility limitations^e

| | | |
|------------------------------|------------------|------------------|
| No limitation | 22.7 [21.3;24.0] | 60.4 [58.7;62.1] |
| Moderate limitations | 29.0 [26.2;31.7] | 68.5 [66.5;70.6] |
| Important/severe limitations | 37.9 [33.3;42.6] | 71.1 [69.0;73.1] |

Sensory limitations^f

| | | |
|------------------------------|------------------|------------------|
| No limitation | 23.4 [22.1;24.6] | 62.5 [61.1;63.8] |
| Moderate limitations | 23.2 [20.1;26.2] | 66.3 [64.0;68.6] |
| Important/severe limitations | 31.8 [22.8;40.7] | 66.7 [62.9;70.6] |

Cognitive limitations

| | | |
|--------------------------------|------------------|------------------|
| No limitation | 23.0 [21.8;24.3] | 62.6 [61.3;63.9] |
| Moderate/important limitations | 28.0 [25.2;30.8] | 67.8 [65.6;69.9] |

Needs family assistance for performing daily living activities

| | | |
|----|------------------|------------------|
| No | 22.9 [21.6;24.3] | 61.1 [59.5;62.6] |
|----|------------------|------------------|

| | | |
|---------------------------------------|------------------|------------------|
| Yes | 27.3 [25.1;29.6] | 69.9 [68.3;71.4] |
| Answered to the questions him/herself | | |
| No | 23.1 [19.1;27.2] | 68.9 [66.5;71.3] |
| Yes | 23.4 [22.2;24.6] | 62.6 [61.3;63.9] |

^a Weighted data.

^b 95% confidence intervals.

^c 72 missing data.

^d 5 missing data.

^e 1 missing data.

^f 3 missing data.

Accepted Version

Table 3. Associations between seasonal influenza vaccine uptake during the 2007-2008 season and type and severity of disability, and socioeconomic status, among people aged 18-64 years: results from multiple log-binomial regression models (France, 2008 HSM survey^a)

| | Model 1 ^b | Model 2 ^c |
|---|---------------------------|---------------------------|
| | <i>n</i> =5,239 | <i>n</i> =5,238 |
| | aRR [95% CI] ^d | aRR [95% CI] ^d |
| Socioeconomic status (SES) score [0-6] | | |
| 0-1 (lowest SES) | 1.00 (ref.) | 1.00 (ref.) |
| 2 | 0.89 [0.79;1.00] | 0.92 [0.77;1.09] |
| 3 | 0.83 [0.73;0.94] | 0.87 [0.72;1.04] |
| 4 | 0.89 [0.77;1.03] | 0.93 [0.76;1.15] |
| 5 | 1.03 [0.88;1.20] | 1.10 [0.88;1.37] |
| 6 (highest SES) | 1.39 [1.23;1.58] | 1.51 [1.25;1.81] |
| Disabilities | | |
| Mobility limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.17 [1.02;1.34] | 1.07 [0.88;1.31] |
| Major limitations | 1.48 [1.21;1.82] | 1.37 [1.02;1.84] |
| Sensory limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 0.84 [0.73;0.96] | 0.82 [0.67;1.00] |
| Major limitations | 0.93 [0.62;1.39] | 0.93 [0.51;1.68] |
| Cognitive limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate/major limitations | 1.17 [1.03;1.33] | 1.13 [0.94;1.36] |

Needs family assistance for performing daily living activities

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.02 [0.90;1.16] | 0.96 [0.80;1.14] |

^a Weighted data. Missing data were excluded from the analysis.

^b Model 1 was adjusted for gender, age, living with a partner, type of residence area, and for whether the respondent answered him/herself or not.

^c Model 2 was adjusted for gender, age, living with a partner, type of residence area, whether the respondent answered him/herself or not, number of visits to a general practitioner in the last 12 months, and chronic disease.

^d Adjusted risk ratio (aRR) and 95% confidence intervals (95% CI).

Accepted Version

Table 4. Associations between seasonal influenza vaccine uptake during the 2007-2008 season and type and severity of disability, and socioeconomic status, among people ≥ 65 years: results from multiple log-binomial regression models (France, 2008 HSM survey^a)

| | Model 1 ^b | Model 2 ^c |
|---|---------------------------|---------------------------|
| | <i>n</i> =7,051 | <i>n</i> =7,048 |
| | aRR [95% CI] ^b | aRR [95% CI] ^b |
| Socioeconomic status (SES) score [0-6] | | |
| 0-1 (lowest SES) | 1.00 (ref.) | 1.00 (ref.) |
| 2 | 1.01 [0.96;1.06] | 1.00 [0.96;1.05] |
| 3 | 1.05 [1.00;1.11] | 1.03 [0.97;1.09] |
| 4 | 1.08 [1.01;1.15] | 1.06 [0.99;1.14] |
| 5 | 1.05 [0.97;1.14] | 1.05 [0.96;1.14] |
| 6 (highest SES) | 1.06 [0.97;1.15] | 1.11 [1.02;1.21] |
| Disabilities | | |
| Mobility limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.06 [1.01;1.11] | 0.99 [0.94;1.04] |
| Major limitations | 1.08 [1.01;1.15] | 1.01 [0.94;1.08] |
| Sensory limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.01 [0.97;1.05] | 1.00 [0.95;1.04] |
| Major limitations | 0.95 [0.87;1.05] | 0.95 [0.86;1.05] |
| Cognitive limitations | | |
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate/major limitations | 0.98 [0.93;1.04] | 0.98 [0.93;1.04] |

Needs family assistance for performing daily living activities

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.04 [1.00;1.09] | 1.01 [0.97;1.07] |

^a Weighted data. Missing data were excluded from the analysis.

^b Model 1 was adjusted for gender, age, living with a partner, type of residence area, and for whether the respondent answered him/herself or not.

^c Model 2 was adjusted for gender, age, living with a partner, type of residence area, whether the respondent answered him/herself or not, number of visits to a general practitioner in the last 12 months, and chronic disease.

^d Adjusted risk ratio (aRR) and 95% confidence intervals (95% CI).

Accepted Version

Accepted Version

Table S1. Keywords used for Medline search

| |
|--|
| <p>#1 Vaccination</p> <p>vaccin*[Title/Abstract] OR immunis*[Title/Abstract] OR immuniz*[Title/Abstract]</p> |
| <p>#2 Seasonal Influenza</p> <p>Influenza Vaccines [MeSH] OR Influenza, Human/prevention and control [MeSH] OR seasonal influenza vaccine [Title/Abstract] OR seasonal influenza vaccines [Title/Abstract] OR flu [Title/Abstract] NOT pandemic NOT epidemic</p> |
| <p>#3 Disabilities, functional limitations</p> <p>Disabilit*[Title/Abstract] OR disabled persons [MeSH] OR functional limitations [Title/Abstract]</p> |
| <p>#4 Final Search</p> <p>#1 AND #2 AND #3</p> |

Accepted Version

Box S1. Target groups for seasonal influenza vaccination, according to the French recommendations for the 2007-2008 season¹

- People aged 65 years or over
- Professionals at risk of exposure
 - Health professionals and professionals in regular and extended contact with people with specific underlying diseases (see below)
 - Staff working on cruise boats and airplanes and those who work in the travel industry as guides)
- People with specific underlying diseases
 - Chronic bronchopulmonary conditions including asthma, bronchopulmonary dysplasia, and cystic fibrosis
 - Poorly tolerated congenital heart disease and severe heart failure and valvulopathies
 - Serious chronic nephropathy, pure and primary nephrotic syndromes
 - Homozygous and S/C heterozygous sickle-cell anemia, sickle-cell thalassemia
 - Diabetes mellitus, type 1 or type 2, that cannot be managed by diet alone
 - Cellular immunodeficiencies (in people with HIV infection, to be indicated the team caring for the patient)
 - Children and adolescents (from 6 months to 18 years) whose health status requires prolonged treatment aspirin treatment (essentially complicated Kawasaki syndrome and chronic juvenile arthritis).
- People living in health institutions

1. http://www.invs.sante.fr/beh/2007/31_32/beh_31_32_2007.pdf

Box S2. Items included in the questionnaire of the 2008 HSM survey to measure functional limitations

Items selected to assess functional limitations were based on the Nagi Disability Scale,¹ the WHO Disability Assessment Schedule 2.0 (WHODAS 2.0),² some disability-related questions from the U.S. National Health Interview Survey³ and also inspired by previous studies⁴⁻⁶:

- Mobility limitations: level of difficulty (4-point Likert-type scale from *no difficulty* = 1 to *unable to perform* = 4) in performing seven tasks without any aid according to a methodology already published: 1) walking 500 meters on flat ground; 2) walking up and down a flight of stairs; 3) stopping, crouching and/or kneeling; 4) lifting or carrying a bag as heavy as 5 kilograms for a distance of 10 meters; 5) raising the arms, e.g. taking an object from an elevated location; 6) grasping or holding an object in the hands; 7) using the hands and fingers with normal dexterity, e.g. turning on a faucet or using a pencil.
- Sensory limitations: level of difficulty (4-point Likert-type scale from *no difficulty* = 1 to *unable to perform* = 4) in performing three tasks: 1) seeing, even with eyeglasses or contact lenses, the block characters of a newspaper; 2) or somebody's face standing 4 meters away; 3) and hearing a conversation between several persons.
- Cognitive limitations: level of frequency (3-point Likert-type scale *never* = 1, *sometimes* = 2, *often* = 3) of eight challenges: 1) has difficulty in acquiring new knowledge or skills, in school, professional training or leisure activities; 2) has difficulty in analyzing and finding solutions to problems in day-to-day life, e.g., identifying an itinerary or counting money; 3) has difficulty in concentrating or doing something for ten minutes; 4) has difficulty in remembering or has memory

lapses; 5) has difficulty in understanding what people say; 6) has periods of confusion or loses track of time; 7) endangers themselves because of their behavior; 8) is impulsive and/or aggressive.

References

1. Nagi SZ. An epidemiology of disability among adults in the United States. *Milbank Memorial Fund Quarterly: Health and Society* 1976;54:439-67.
2. WHO. WHO disability assessment schedule 2.0 WHODAS 2.0. 2014.
<http://www.who.int/classifications/icf/whodasii/en/>
3. NHIS. National health interview survey on disability. 2014.
http://www.cdc.gov/nchs/nhis/nhis_disability.htm
4. Rasch EK, Hochberg MC, Magder L, Magaziner J, Altman BM. Health of community-dwelling adults with mobility limitations in the United States: prevalent health conditions. Part I. *Archives of Physical Medicine and Rehabilitation* 2008;89:210–8.
5. Ramirez A. Disability and preventive cancer screening: results from the 2001 California health interview survey. *American Journal of Public Health* 2005;95:2057–64.
6. Reichard A, Stolze H, Fox MH. Health disparities among adults with physical disabilities or cognitive limitations compared to individuals with no disabilities in the United States. *Disability and Health Journal* 2011;4:59–67.

Box S3. Individual socioeconomic score

The individual socioeconomic (SES) score was calculated from the respondents' educational level, occupation, and equivalized household income (EHI), according to a methodology already published.^{1,2} For each SES indicator, respondents were assigned between 0 (did not complete high school, not in the labor force, lowest EHI respectively) and 2 points (educational level higher than high school level, intellectual worker, highest EHI respectively); the SES score was calculated by summing respondents' points; it could range from 0 (lowest SES) to 6 (highest SES). Because only a few respondents had an SES score of 0 or 6, they were grouped with the categories 1 and 5 respectively.

References

1. Stamatakis E, Hillsdon M, Mishra G, Hamer M, Marmot M. Television viewing and other screen-based entertainment in relation to multiple socioeconomic status indicators and area deprivation: the Scottish Health Survey 2003. *J Epidemiol Community Health*. 2009;63(9):734-40.
2. Stamatakis E, Wardle J, Cole TJ. Childhood obesity and overweight prevalence trends in England: evidence for growing socioeconomic disparities. *Int J Obes* 2005. 2010;34(1):41-7.

Table S2. Characteristics^a of the study population according to the criteria of the French recommendations for seasonal influenza vaccination for the 2007-2008 season (France, 2008 HSM survey, $n=12,396$)

| Criteria | 18-64 years ($n=5,297$) | | ≥ 65 years ($n=7,099$) | |
|--|---------------------------|-------|-------------------------------|------|
| | <i>n</i> | % | <i>n</i> | % |
| ≥ 65 years without any target disease | / | / | 3,176 | 44.7 |
| Works in the health and social field | 1,375 | 25.96 | 13 | 0.19 |
| Has at least one target disease | | | | |
| Target cardiovascular disease | 2,149 | 40.6 | 3,071 | 43.3 |
| Target respiratory disease | 1,830 | 34.6 | 978 | 13.8 |
| Diabetes | 726 | 13.7 | 910 | 12.8 |
| Other diseases | 77 | 1.5 | 51 | 0.7 |

^a Weighted data. Each individual can meet several inclusion criteria (e.g., have a chronic respiratory disease *and* diabetes).

Table S3. Associations between the socioeconomic status (SES) score and the severity of the limitation by age group and type of functional limitation (% line) (France, 2008 HSM survey)

| | <i>n</i> ^a | SES score [0-6] | | | | | | p-value ^b |
|------------------------------|-----------------------|-----------------|-------|-------|------|------|------|----------------------|
| | | 0-1 | 2 | 3 | 4 | 5 | 6 | |
| 18-64 years (n=5,260) | | | | | | | | |
| All 18-64 years | 5,260 | 13.5 | 24.6 | 24.8 | 12.4 | 9.4 | 15.4 | |
| Mobility limitations | | | | | | | | |
| No limitation | 4,800 | 11.5 | 23.8 | 25.2 | 13.0 | 10.0 | 16.5 | <.0001 |
| Moderate limitations | 353 | 32.9 | 32.9 | 22.6 | 6.6 | 2.1 | 3.0 | |
| Major limitations | 106 | 38.7 | 34.4 | 17.0 | 5.2 | 3.4 | 1.3 | |
| Sensory limitations | | | | | | | | |
| No limitation | 4,843 | 12.4 | 24.0 | 25.2 | 12.7 | 9.7 | 16.2 | <.0001 |
| Moderate limitations | 390 | 25.2 | 32.2 | 21.2 | 10.4 | 6.1 | 5.0 | |
| Major limitations | 26 | 40.2 | 31.9 | 15.7 | 3.1 | 6.4 | 2.7 | |
| Cognitive limitations | | | | | | | | |
| No limitation | 4,867 | 11.9 | 23.9 | 24.9 | 13.1 | 9.9 | 16.2 | <.0001 |
| Moderate/major limitations | 393 | 32.9 | 32.9 | 23.9 | 4.2 | 2.8 | 3.4 | |
| ≥ 65 years (n=7,063) | | | | | | | | |
| All ≥ 65 years | 7,063 | 29.9 | 34.2 | 17.6 | 8.0 | 4.8 | 5.6 | |
| Mobility limitations | | | | | | | | |
| No limitation | 4,781 | 24.6 | 33.1 | 19.5 | 9.7 | 5.8 | 7.3 | <.0001 |
| Moderate limitations | 1,492 | 39.4 | 36.3 | 14.0 | 4.9 | 2.9 | 2.5 | |
| Major limitations | 790 | 44.22 | 36.43 | 13.15 | 3.22 | 1.69 | 1.28 | |
| Sensory limitations | | | | | | | | |
| No limitation | 5,586 | 27.7 | 33.7 | 18.5 | 8.6 | 5.2 | 6.4 | <.0001 |

| | | | | | | | | |
|----------------------------|-------|------|------|-------|-----|------|------|--------|
| Moderate limitations | 1,236 | 37.1 | 36.9 | 14.2 | 6.0 | 3.2 | 2.7 | |
| Major limitations | 240 | 44.5 | 32.0 | 14.5 | 3.4 | 3.70 | 1.95 | |
| Cognitive limitations | | | | | | | | |
| No limitation | 6,118 | 28.3 | 34.1 | 17.9 | 8.4 | 5.1 | 6.2 | <.0001 |
| Moderate/major limitations | 945 | 40.7 | 34.6 | 15.78 | 5.1 | 2.2 | 1.7 | |

^a Weighted data.

^b Chi-square test.

Accepted Version

Table S4. Associations between seasonal influenza vaccine uptake during the 2007-2008 season and type and severity of disability, and socioeconomic status, among people aged 18-64 years: results from multiple log-binomial regression models (France, 2008 HSM survey^a)

| | Model 1 <i>n</i> =5,239 | Model 2 <i>n</i> =5,238 |
|---|----------------------------|----------------------------|
| | aRR [95% CI] ^b | aRR [95% CI] ^b |
| Gender | | |
| Men | 1.00 (ref.) | 1.00 (ref.) |
| Women | 0.87 [0.81;0.93] | 0.87 [0.78;0.96] |
| Age (years) | | |
| 18-45 | 1.00 (ref.) | 1.00 (ref.) |
| 46-64 | 1.93 [1.77;2.10] | 1.83 [1.62;2.07] |
| Living with a partner | | |
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.03 [0.95;1.12] | 1.03 [0.91;1.17] |
| Type of residence area | | |
| Rural | 1.00 (ref.) | 1.00 (ref.) |
| Urban | 0.97 [0.90;1.05] | 0.97 [0.87;1.08] |
| Socioeconomic status (SES) score [0-6] | | |
| 0-1 (lowest SES) | 1.00 (ref.) | 1.00 (ref.) |
| 2 | 0.89 [0.79;1.00] | 0.92 [0.77;1.09] |
| 3 | 0.83 [0.73;0.94] | 0.87 [0.72;1.04] |
| 4 | 0.89 [0.77;1.03] | 0.93 [0.76;1.15] |
| 5 | 1.03 [0.88;1.20] | 1.10 [0.88;1.37] |
| 6 (highest SES) | 1.39 [1.23;1.58] | 1.51 [1.25;1.81] |

Disabilities

Mobility limitations

| | | |
|----------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.17 [1.02;1.34] | 1.07 [0.88;1.31] |
| Major limitations | 1.48 [1.21;1.82] | 1.37 [1.02;1.84] |

Sensory limitations

| | | |
|----------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 0.84 [0.73;0.96] | 0.82 [0.67;1.00] |
| Major limitations | 0.93 [0.62;1.39] | 0.93 [0.51;1.68] |

Cognitive limitations

| | | |
|----------------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate/major limitations | 1.17 [1.03;1.33] | 1.13 [0.94;1.36] |

Needs family assistance for performing daily living activities

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.02 [0.90;1.16] | 0.96 [0.80;1.14] |

The respondent answered him/herself

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.07 [0.91;1.26] | 1.09 [0.86;1.38] |

Number of visits to a general practitioner in the last 12 months

| | | |
|------------|---|------------------|
| 1-4 | - | 1.00 (ref.) |
| ≥5 | - | 1.32 [1.18;1.47] |
| Don't know | - | 0.95 [0.77;1.17] |

Chronic disease

| | | |
|-----|---|------------------|
| No | - | 1.00 (ref.) |
| Yes | - | 1.20 [1.06;1.36] |

^a Weighted data. Missing data were excluded from the analysis.

^b Adjusted risk ratio (aRR) and 95% confidence intervals (95% CI).

Accepted Version

Table S5. Associations between seasonal influenza vaccine uptake during the 2007-2008 season and type and severity of disability, and socioeconomic status, among people ≥ 65 years: results from multiple log-binomial regression models (France, 2008 HSM survey^a)

| | Model 1 <i>n</i> =7,051 | Model 2 <i>n</i> =7,048 |
|---|----------------------------|----------------------------|
| | aRR [95% CI] ^b | aRR [95% CI] ^b |
| Gender | | |
| Men | 1.00 (ref.) | 1.00 (ref.) |
| Women | 0.97 [0.94;1.01] | 0.96 [0.93;1.00] |
| Age (years) | | |
| 65-74 | 1.00 (ref.) | 1.00 (ref.) |
| ≥ 75 | 1.24 [1.19;1.29] | 1.21 [1.17;1.27] |
| Living with a partner | | |
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.10 [1.06;1.14] | 1.08 [1.04;1.13] |
| Type of residence area | | |
| Rural | 1.00 (ref.) | 1.00 (ref.) |
| Urban | 0.99 [0.96;1.03] | 1.00 [0.96;1.04] |
| Socioeconomic status (SES) score [0-6] | | |
| 0-1 (lowest SES) | 1.00 (ref.) | 1.00 (ref.) |
| 2 | 1.01 [0.96;1.06] | 1.00 [0.96;1.05] |
| 3 | 1.05 [1.00;1.11] | 1.03 [0.97;1.09] |
| 4 | 1.08 [1.01;1.15] | 1.06 [0.99;1.14] |
| 5 | 1.05 [0.97;1.14] | 1.05 [0.96;1.14] |
| 6 (highest SES) | 1.06 [0.97;1.15] | 1.11 [1.02;1.21] |

Disabilities

Mobility limitations

| | | |
|----------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.06 [1.01;1.11] | 0.99 [0.94;1.04] |
| Major limitations | 1.08 [1.01;1.15] | 1.01 [0.94;1.08] |

Sensory limitations

| | | |
|----------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate limitations | 1.01 [0.97;1.05] | 1.00 [0.95;1.04] |
| Major limitations | 0.95 [0.87;1.05] | 0.95 [0.86;1.05] |

Cognitive limitations

| | | |
|----------------------------|------------------|------------------|
| No limitation | 1.00 (ref.) | 1.00 (ref.) |
| Moderate/major limitations | 0.98 [0.93;1.04] | 0.98 [0.93;1.04] |

Needs family assistance for performing daily living activities

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 1.04 [1.00;1.09] | 1.01 [0.97;1.07] |

The respondent answered him/herself

| | | |
|-----|------------------|------------------|
| No | 1.00 (ref.) | 1.00 (ref.) |
| Yes | 0.99 [0.94;1.05] | 0.99 [0.94;1.05] |

Number of visits to a general practitioner in the last 12 months

| | | |
|------------|---|------------------|
| 1-4 | - | 1.00 (ref.) |
| ≥5 | - | 1.18 [1.13;1.23] |
| Don't know | - | 0.53 [0.44;0.64] |

Chronic disease

| | | |
|-----|---|------------------|
| No | - | 1.00 (ref.) |
| Yes | - | 1.07 [1.01;1.12] |

^a Weighted data. Missing data were excluded from the analysis.

^b Adjusted risk ratio (aRR) and 95% confidence intervals (95% CI).

Accepted Version