

Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion deficiency type III

Paul Saultier, Marie-Christine Alessi, Gérard Michel, Hervé Chambost

▶ To cite this version:

Paul Saultier, Marie-Christine Alessi, Gérard Michel, Hervé Chambost. Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion deficiency type III. Pediatrics & Neonatology, 2017, 58 (6), pp.560 - 561. 10.1016/j.pedneo.2017.07.006 . hal-01753440

HAL Id: hal-01753440 https://amu.hal.science/hal-01753440

Submitted on 29 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion deficiency type III

Paul Saultier, Marie-Christine Alessi, Gérard Michel, Hervé Chambost

PII: \$1875-9572(17)30240-1

DOI: 10.1016/j.pedneo.2017.07.006

Reference: PEDN 693

To appear in: Pediatrics & Neonatology


Received Date: 18 April 2017

Revised Date: 1875-9572 1875-9572

Accepted Date: 18 July 2017

Please cite this article as: Saultier P, Alessi M-C, Michel G, Chambost H, Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion deficiency type III, *Pediatrics and Neonatology* (2017), doi: 10.1016/j.pedneo.2017.07.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



TITLE PAGE

Article category: pediatric hematology

Title: Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion deficiency type III

Running title: Treatment of LAD III

Authors:

Paul Saultier, 1,2 Marie-Christine Alessi, 1,3 Gérard Michel and Hervé Chambost 1,2

¹Aix Marseille Univ, INSERM, INRA, NORT, Marseille, France

²Department of Pediatric Hematology and Oncology, Timone Enfants Hospital, APHM and Aix Marseille Univ, Marseille, France

³APHM, CHU Timone, French Reference Center on Inherited Platelet Disorders, Marseille, France

Corresponding author:

Paul Saultier, Department of Pediatric Hematology, Timone Enfants Hospital, 264 rue St Pierre, 13385 Marseille, France; E-mail: paul.saultier@gmail.com; phone: +33491386778; fax: +33491384989.

TITLE PAGE

Article category: pediatric hematology

Title: Hematopoietic stem cell transplantation for the treatment of leukocyte adhesion

deficiency type III

Running title: treatment of leukocyte adhesion deficiency type III

TEXT

To the Editor,

We read with great interest the article published in *Pediatrics and Neonatology* by Aygun and colleagues.¹ The authors described the cases of two children of Turkish origin who had leukocyte adhesion deficiency type III (LAD III). To date, fewer than 40 cases have been reported in the literature. In this article, the authors provided clinical data and conducted a biological investigation, including genetic analysis, that led to the diagnosis. They also provided interesting information regarding the treatment.

The patients described by Aygun and colleagues harbored a Glanzmann-like bleeding phenotype associated with recurrent infections, which are reported in almost all patients with LAD III. The authors stated that hematopoietic stem cell transplantation (HSCT) was the only proven curative therapy for the disease. Indeed, several other groups reported the cure of LAD III using HSCT. One of the two patients reported by Aygun and colleagues died a few months after HSCT because of bacteremia and graft-versus-host disease. They assumed that the delayed diagnosis and HSCT accounted for the outcome. They stated, like other authors, that performing HSCT in early infancy will improve the outcome of LAD III. However, an early HSCT would preclude the precise evaluation of disease severity, which can vary among patients with LAD III. Furthermore, HSCT is associated with a high treatment-related mortality. Indeed, several authors reported HSCT-related deaths in patients with LAD III.^{2,3} Other groups reported graft failure. As proposed by several investigators, the recovery of recipient hematopoiesis after HSCT may be related to the osteopetrosis-like increased bone density observed in several patients with LAD III. Overall, no clear evidence supports the systematic and early use of HSCT in LAD III. Optimizing the treatment of bleeding and infections without performing HSCT may be an appropriate approach for the management of this disease, at least for the patients with the mildest symptoms. Indeed, several authors

reported that patients treated for LAD III survived even without HSCT.² However, these investigators provided limited data regarding the treatment.

The optimal management of LAD III remains to be delineated. Owing to the rarity of the disease, evaluation of the HSCT benefit-to-risk ratio is not straightforward. Reporting of LAD III cases is pivotal for a precise description of the disease phenotype. Data regarding long-term management of patients with LAD III treated without HSCT are scarce but would be greatly useful to evaluate the different therapeutic options.

CONFLICT OF INTEREST STATEMENT

The authors have no conflict of interest to declare.



ACKNOWLEDGMENTS

This work was supported by the "Fondation pour la recherche médicale FRM" (grant to P. Saultier FDM20150633607).



REFERENCES

- 1. Aygun D, Nepesov S, Gershoni R, Camcioglu Y. Leukocyte Adhesion Deficiency III: Report of Two Siblings. *Pediatr Neonatol* 2017;**58**:99–100.
- 2. Kuijpers TW, van de Vijver E, Weterman MAJ, de Boer M, Tool ATJ, van den Berg TK, et al. LAD-1/variant syndrome is caused by mutations in FERMT3. *Blood* 2009;**113**:4740–6.
- 3. Stepensky PY, Wolach B, Gavrieli R, Rousso S, Ben Ami T, Goldman V, et al. Leukocyte adhesion deficiency type III: clinical features and treatment with stem cell transplantation. *J Pediatr Hematol Oncol* 2015;**37**:264–8.