



HAL
open science

Screening for intrathoracic goiters before thyroid surgery: does chest X-Rays have a place?

Fares Benmiloud, Michel Grino, Charles Oliver, Anne Denizot

► To cite this version:

Fares Benmiloud, Michel Grino, Charles Oliver, Anne Denizot. Screening for intrathoracic goiters before thyroid surgery: does chest X-Rays have a place?. *Langenbeck's Archives of Surgery*, 2017, 402 (2), pp.251-255. 10.1007/s00423-016-1534-9 . hal-01753792v1

HAL Id: hal-01753792

<https://amu.hal.science/hal-01753792v1>

Submitted on 3 Apr 2018 (v1), last revised 3 Apr 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fares Benmiloud¹. Michel Grino^{2,3}. Charles Oliver^{3,4}. Anne Denizot¹

Screening for intrathoracic goiters before thyroid surgery: does chest X-Rays have a place?

¹European Hospital. Endocrine surgery department, 6 rue Désirée Clary, 13003 Marseille, France.

²Inserm UMR 1062, Faculté de Médecine secteur Timone, 27 boulevard Jean Moulin, 13385 Marseille, France.

³Aix-Marseille University, Faculté de Médecine secteur Timone, 27 boulevard Jean Moulin, 13385 Marseille, France.

⁴Division of Geriatric Medicine, State Geriatric Centre, 176 Avenue de Montolivet, 13012 Marseille, France.

Address all correspondence to: Dr A. Denizot, European Hospital. Endocrine surgery department. 6 rue Désirée Clary, 13003 Marseille, France. anne.denizot@dbmail.com, Phone: (33) 413 427 338, Fax: (33) 413 427 724

Abstract

Purpose Since intrathoracic goiters (IG), either cervico-mediastinal goiters (CMG) or mediastinal nodules (MN), can lead to sternotomies and/or evitable reoperations, their detection is mandatory before thyroid surgery. A systematic screening by CTscan or MRI is not conceivable ~~(because too~~ expensive of their expensiveness). We tested if ~~c~~Chest XRay (CXr) could remain a good screening tool for IG before thyroid surgery.

Methods In this prospective study (2,554 patients), CXr usefulness was evaluated in relation with patients complaints, clinical examination, neck US, and anatomical and surgical findings.

Results CMG (n = 67) and MN (n = 42) were symptomatic in 10 and 5 patients, respectively. Clinical examination or neck US suspected their existence in 25 and 13, and 45 and 17 patients, respectively. Among the 50 IG detected by CXr (42 CMG and 8 MN), 5 CMG and 2 MN were missed by clinical examination or neck US. CXr failed to detect IG in 59 patients (54%): 25 CMG (37 %) and 34 MN (80%). 28 IG (9 CMG and 19 NM) were discovered during surgery. CXr resulted in false positive in 88 out of 2,445 patients (3.5%). CXr potentially avoided reoperation in 2 patients (a maximum saving of 6160 €, whereas the total cost of CXr was 54,895 €).

Conclusions CXr should not be used routinely for the preoperative detection of IG. Surgeons should preferably use clinical examination or neck US and directly perform CT scan when a mediastinal extension is suspected.

Key words Chest X-rays; thyroid surgery; intrathoracic goiters; cervico-mediastinal goiters; mediastinal nodules.

Author's contributions

Study conception and design: FB, CO, AD

Acquisition of data: FB, CO, AD

Analysis and interpretation of data: FB, CO, AD

Drafting of manuscript: FB, CO, AD, MG

Critical revision of manuscript: FB, CO, AD, MG

Introduction

The presence of thyroid tissue in the mediastinum occurs in 2 to 19% of patients who undergo thyroidectomy [1]. These intrathoracic goiters (IG), also called retrosternal, substernal, cervico-mediastinal or mediastinal goiter, can be connected to the thyroid, like in cervicomedial goiters (CMG) or disconnected from the thyroid body, resulting in independent thyroid mediastinal nodules (MN).

There are two caveats in the surgery of IG: (a) surgical difficulty during the cervical approach which may need sternotomy (b) risk of missing a MN with a subsequent need for reoperation [2]. Therefore, it is mandatory to detect IG preoperatively. This can be challenging since these lesions are asymptomatic in 20 to 30% of the cases [3, 4] and develop in an area not easily accessible to clinical examination or neck ultrasonography (US).

In the past, chest X-ray (CXr) was the reference for the preoperative diagnosis of IG. Nowadays, computerized tomography (CT) scan or magnetic resonance imaging (MRI), are the accurate tools for the characterization of IG [5, 6]. However, CT and MRI are not used to screen for IG because they are expensive [7].

We hypothesized that CXr, because of its inexpensiveness, could remain a good screening test to detect IG unsuspected by clinical examination and neckUS and to select patients who will benefit from further assessment (by CT or MRI) for an appropriate surgical strategy. To the best of our knowledge there is no prospective study which confirmed (or invalidated) our hypothesis.

In this report we describe the results of a prospective study aimed at evaluating the usefulness of a systematic screening by CXr (as a complement of clinical examination and neck US) for the detection of IG before thyroidectomy.

Materials and Methods

This prospective study included 2,570 patients referred to the Endocrine Surgery Department of Paul Desbieff Hospital, Marseilles, France during a 7-year period (2005-2011) for surgery of thyroid goiters.

Preoperative investigations included anamnesis, clinical examination searching for mediastinal extension and/or compression with specific attention to patients' complaints (dyspnea, dysphagia, hoarseness), physical signs (lower pole of the gland not palpable, tracheal deviation, signs of venous thoracic compression), neck US (access or not to the lower pole of the gland) and CXr (posterior-anterior CXr performed with both arms over the head). The CXr abnormalities revealing mediastinal thyroid extension were classified as anterior, lateral or latero-tracheal deviation and mediastinal opacities.

When a mediastinal extension was found (lower thyroid lobe margins extending below the sternal manubrium), ~~they were it was described-recorded~~ in the ~~anatomical charts~~ operative report. IG were separated into CMG (with parenchymal continuum between the cervical and mediastinal thyroid gland) and MN (lined by its own capsule and disconnected or attached with only a thin fibrous tissue to the cervical thyroid gland). The size of the thoracic portion of the IG, and the distance between MN and the thyroid were recorded. Clinical, imaging and surgical characteristics were compared between CMG and MN. At the end of the operations, visual examination and palpation of the superior mediastinum were routinely performed, searching for occult MN. The majority of thyroidectomies were performed by the same surgeon (AD).

Statistical analysis was performed with the Graph Pad Prism 5 software (La Jolla, CA). The χ^2 test was used for categorical variables and the Mann-Whitney U test was used for continuous variables. A *P* value less than 0.05 was considered significant.

Results

2,554 patients were included in the study since CXr was not interpretable in 16 of them.

Populations characteristics (table 1)

For the entire group of patients, the age ranged between 13 and 86 years with a mean (\pm SD) of 50.4 ± 14.0 years. The female/male ratio was 4:1 (2,074/496).

Patients with CMG were significantly ($P = 0.0126$) older than those with NM (61.3 ± 12.5 vs. 54.3 ± 16.4 years, respectively). There was no significant difference in the female/male ratio between the two groups (2.52 and 5.0, $P = 0.1457$, for patients with CMG or NM, respectively)

Anatomical characteristics of CMG and MN (table 2)

IG were found in 109/2,554 patients (4.3 %). As above mentioned, CMG ($n = 67$) showed a parenchymal continuum between the cervical and mediastinal portion of the thyroid gland. MN ($n = 42$) were independent ($n = 35$) or connected to the cervical thyroid by a thin fibrous tissue ($n = 7$). The distance between MN and main thyroid varied from 0 cm (contiguous MN) to 3 cm (complete intrathoracic nodule, $n = 7$). No difference was observed in the size of the thoracic portion of CMG and MN. Partial sternotomy (manubriotomy) was necessary in 2 cases of NM. No total sternotomy was performed.

Preoperative findings

As shown in table 3, CMG and MN were symptomatic in 10/67 (14.9 %) and 5/42 (11.9 %) patients, respectively. Physical examination suspected their presence in 25/67 (37.3 %) and 13/42 (30.9 %) patients, and neck US suspected them in 45/67 (67.2 %) and 17/42 (40.5%) patients, respectively. Overall, clinical examination combined to neck US allowed suspicion of 54/67 CMG (80.6 %) and 21/42 MN (50.0 %).

Performance of CXr

2,554/2,570 CXr were interpretable. CXr detected the mediastinal thyroid extensions in 50/109 patients (45.9 % of true positives): 42/67 CMG (62.7 %) and 8/42 MN (19.0 %). CXr was the only tool that allowed the detection of IG in 6/109 (6.4 %) patients: 4 CMG and 2 MN, which have not

been detected by clinical examination and neck US. The 4 CMG had a mediastinal portion of 7, 6, 2 and 2 cm. One MN detected only by CXr measured 6 cm; it was located 1 cm from the inferior pole of the thyroid and was connected to the thyroid by a fibrous bridge. The other MN measured 3 cm and was disconnected, separated from the thyroid by a 3 cm distance. No manubriotomy was necessary in these 6 patients.

CXr failed to detect IG in 59/109 patients (54.1 % of false negatives) separated into 25/67 (37.3 %) CMG and 34/42 (80.9 %) MN. A total of 28 IG (9 CMG and 19 NM) were undetectable by clinical examination, neck US or CXr and discovered only during surgery. Among the 9 CMG, the mediastinal extension measured less than 3 cm in 8 cases and 4 cm in one case. None of them was difficult to remove surgically. Three out of the 19 NM discovered during the operation have been difficult to visualize. In this study, 2 patients were reoperated for MN since it had not been detected during the first operation. Clinical examination and neck US were positive in 31 (16 CMG and 15 MN) of the 59 patients (52.5 %) negative for CXr.

False positive images were observed by CXr in 88 out of 2445 patients without IG (3.5% false positives). The false positive for CMG (n = 88) were separated into curved tracheal deviations (n = 50), straight deviations of the whole trachea (n = 24) and “bayonet-like” deviations (n = 4). The false positives for MN included mediastinal opacities (n = 10). All 88 patients had strict cervical goiter; in 11 cases, the inferior limit of the goiter reached the superior limit of the thoracic inlet without crossing it.

In the present study, CxR specificity was good (96.8 and 95.3 % for for CMG and MN, respectively) but its sensitivity was low (62.7 and 19.0 % for CMG and MN, respectively).

Since CXR revealed in 2 cases a MN that could have remained undetected, we assume that CXr saved a maximum of 2 patients from second surgery, which means a maximum saving of 6160 € (3080 € being the cost of an operation for mediastinal lesions resection, as it is evaluated by French Social Insurance). The total cost of CXr in this study was 54,895 €.

In this series, the cumulated irradiation dose was very low (55 mSv for 2570 CxR).

Discussion

Among 2570 patients referred for thyroid surgery, our series include 109 IG separated into 67 CMG and 42 MN, based on their anatomical characteristics, in the continuum of the cervical thyroid for CMG and independent or attached only by a thin fibrous tissue to the cervical thyroid for MN. Among these 109 IG, 75 (54 CMG and 21 MN) have been suspected after clinical examination and neck US; 6 further cases (4 CMG and 2 MN) have been detected after CXr. In 29 cases, clinical examination and imaging failed to detect IG which was discovered during the first surgery in 27 patients (9 CMG and 19 MN). 2 MN were missed during first surgery and removed after reoperation. Although the diagnosis performance of CXr was low (56% of false negative), this procedure was useful in 6 cases (6% of the IG but 0,02% of the thyroidectomies).

In France and in many other countries, routine preoperative CXr is no longer recommended for all non-cardiac non-thoracic surgeries [8]. It is even not systematically reimbursed by the French health insurance since 2012, unless if the indication for CXr (like “search for Ig”, for instance) is clearly noticed [8]. However, CXr is still described as a diagnostic tool in reference textbooks [4] and in recent studies on the management of large descending goiters [9-12]. The usefulness of CXr in thyroid surgery, as in many other types of surgery, has been questioned in several studies. In two studies, tracheal deviation detected by CXr was not correlated with any difficulty of intubation [13, 14]. In our series, although its performance in the screening for IG was mediocre, CXr added some useful information to the preoperative investigations, especially for the CMG. Indeed, CXr allowed the diagnosis of 4 cases of CMG and 2 cases of MN unsuspected by clinical examination and US (both MN would have probably been missed in the absence of CXr).

In this study, CXr was useless to predict the need for sternotomy or thoracotomy. Classically, predictive factors for sternotomy are the extension below the carina or the aortic arch posterior mediastinum extension and the presence of mediastinal nodule or conical shape of the goiter in which the thoracic inlet becomes a ring of constriction [[14](#),[15](#), -16]. Our series include 2 manubriotomies: one for a “snowman-shaped” MN with two contiguous parts separated by a tight, quasi-non-existent collar,

the largest part being intrathoracic and one for a CMG descending no more than 1 cm below the thoracic inlet but with strong adherence to the trachea.

Regarding the false negative of the preoperative procedures, all of the 9 CMG that were found during operation were small and were not difficult to remove. However, this was not the case with the 27 MN. Three MN were discovered by the surgeon during systematic inspection and palpation of the superior mediastinum following thyroidectomy. In the present series, 2 MN were missed and needed reoperation. These 2 cases were revealed by ¹³¹I fixation during radio-therapy for thyroid cancer. One cannot exclude that more MN were missed.

False positive CXr signs of tracheal deviations may be artefactual and the consequence of slight deviations of X-rays during imaging. Some mediastinal opacities, especially those situated in the upper part of the mediastinum, may also be explained by the different positions of the patients when CXr and surgery were performed.

In our series, we have also attempted to classify IG on the basis of anatomical characteristics. The IG are usually separated into primary or secondary, depending on their site of origin [17, 18]. Primary IG are rare (less than 1% of all IG); they arise from “ectopic” mediastinal thyroid without connection with the cervical thyroid and they receive their blood supply exclusively from mediastinal vessels. Secondary IG which account for most cases develop through the downward growth of the cervical thyroid. Based on the literature and our experience, we have separated IG into two subgroups: CMG with a continuum between the cervical and mediastinal portions and MN separated or connected only by a thin fibrous tissue to the cervical thyroid. MN may develop through the extension of a nodule arising in the lower pole of a thyroid lobe. The nodule follows a path of least resistance into the mediastinum and its attachment to the cervical thyroid is broken or reduced to a thin fibrous tissue. Alternatively, they may correspond to the rare primary IG with their blood supply originating exclusively from mediastinal vessels. Our results do not allow ~~the distinction to distinguish~~ between both origins for MN. MN have been described under different names: aberrant thyroid adenoma found in 32/88 IG [19], ectopic nodules in 4/108 IG [20], grades III and IV thyrothymic rests [1] or ~~the~~ complete, as opposed to partial, intrathoracic goiter IG which were more frequent before than after 65

~~years (17/58 vs. 12/94, respectively) as opposed to the partial intrathoracic goiter [22]. For these authors, the percentage of complete IG was higher before than after 65 years (17/58 vs 12/94).~~ As reported by Cohen and Cho [23], the MN did not receive sufficient recognition although they may be the source of unsuccessful thyroidectomy reported by several authors under the name of forgotten thyroid [24-29].

In our series, preoperative CXr helped to alert the surgeon from potential surgical difficulty in 4 cases (and in one of the cases, CXr itself allowed the detection of the thyroid problem). In 2 other patients, CXr revealed a MN that could have remained undetected. Therefore, CXr saved 2 patients from second surgery, which means a maximum saving of 6160 €. The total cost of CXr in the study being 54,895 €, CXr screening was clearly not cost effective, although the extra cost represents less than 1% of the minimal cost of thyroidectomy (21€ vs. 2356 €). Furthermore, if a CT scan (which costs 266 €) or an MRI (406 €) would have been performed in each of the false positive patients, this would have led to an additional unnecessary cost of 88 x 266 € or 88 x 406 € respectively (that is 23,408 or 35,728 €, respectively).

Mis en forme : Police :Italique

In conclusion, systematic screening by CXr in order to detect IG had mediocre performance (50% true positive) and, although it allowed to suspect 6% of the IG and potentially avoided reoperation in 2 patients, it was not cost effective. CXr should therefore reasonably be abandoned in routine use for the preoperative screening for IG before thyroid surgery. Surgeons should preferably rely on clinical examination or neck US and perform directly CT scan when a mediastinal extension is suspected. However, they should be aware that undiagnosed IG may be found during surgery in about 1% of the patients.

Conflict of Interest The authors declare that they have no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with

the 1964 Helsinki declaration and its later amendments or comparable ethical standards. For this type of study formal consent is not required.

References

1. White ML, Doherty GM, Gauger P (2008) Evidence-based surgical management of substernal goiter. *World J Surg* 32:1285-1300. Code de champ modifié
2. Hegedüs L, Bonnema SJ (2010) Approach to management of the patient with primary or secondary intrathoracic goiter. *J Clin Endocrinol Metab* 95:5155-5162. Code de champ modifié
3. Hedayati N, McHenry CR (2002) The clinical presentation and operative management of nodular and diffuse substernal thyroid disease. *Am J Surg* 68:245-251. Code de champ modifié
4. Textbook of Endocrine Surgery (2nd Ed, 2005) Elsevier Saunders, Philadelphia.
5. Bashist B, Ellis K, Gold RP (1983) Computed tomography of intrathoracic goiters. *AJR Am J Roentgenol* 140:455-460. Code de champ modifié
6. Buckley JA, Stark P (1999) Intrathoracic mediastinal thyroid goiter: imaging manifestations. *AJR Am J Roentgenol* 173:471-475. Code de champ modifié
7. AACE/AME Guidelines Gharib H, Papini E, Paschke R (2008) Thyroid nodules: a review of current guidelines, practices, and prospects. *Eur J Endocrinol* 159:493-505. Code de champ modifié
8. Haute Autorité de la Santé Principales non-indications de la radiographie du thorax (2009). Available at [http:// www. has.fr](http://www.has.fr)
9. Ben Nun A, Soudack M, Best LA (2006) Retrosternal thyroid goiter: 15 years experience. *Isr Med Assoc J* 8:106-109. Code de champ modifié
10. Shen WT, Kebebew E, Duh QY, Clark OH (2004) Predictors of airway complications after thyroidectomy for substernal goiter. *Arch Surg* 139:656-659. Code de champ modifié
11. Bizakis J, Karatzanis A, Hajjioannou J, Bourolias C, Maganas E, Spanakis E, Bizaki A, Velegarakis G (2008) Diagnosis and management of substernal goiter at the University of Crete. *Surg Today* 38:99-103. Code de champ modifié

12. Rugiu MG, Piemonte M (2009) Surgical approach to retrosternal goitre: do we still need sternotomy? Acta Otorhinolaryngol Ital 29:331-338. Code de champ modifié
Code de champ modifié
13. Voyagis GS, Kyriakos KP (1997) The effect of goiter on endotracheal intubation. Anesth Analg 84:611-612. Code de champ modifié
14. Hong BW, Mazeh H, Chen H, Sippel RS (2012) Routine chest X-ray prior to thyroid surgery: is it always necessary? World J Surg 36:2584-2589. Code de champ modifié
15. Makeieff M, Marlier F, Khudjadze M, Garrel R, Crampette L, Guerrier B (2000) Substernal goiter. Report of 212 cases. Ann Chir 125:18-25. Code de champ modifié
16. Riffat F, Del Pero MM, Fish B, Jani P (2013) Radiologically predicting when a sternotomy may be required in the management of retrosternal goiters. Ann Otol Rhinol Laryngol 122:15-19. Code de champ modifié
17. Huins CT, Georgalas C, Mehrzad H, Tolley NS (2008) A new classification system for retrosternal goitre based on a systematic review of its complications and management. Int J Surg 6:71-76. Code de champ modifié
18. Hashmi SM, Premachandra DJ, Bennett AM, Parry W (2006) Management of retrosternal goitres: results of early surgical intervention to prevent airway morbidity, and a review of the English literature. J Laryngol Otol 120:644-649. Code de champ modifié
19. Cichoń S, Anielski R, Konturek A, Baczyński M, Cichoń W, Orlicki P (2008) Surgical management of mediastinal goiter: risk factors for sternotomy. Langenbecks Arch Surg 393:751-757. Code de champ modifié
20. Cohen JP (2009) Substernal goiters and sternotomy. Laryngoscope 119:683-688. Code de champ modifié
21. Sackett WR, Reeve TS, Barraclough B, Delbridge L (2002) Thyrothymic thyroid rests: incidence and relationship to the thyroid gland. J Am Coll Surg 195:635-640. Code de champ modifié
Code de champ modifié
Code de champ modifié
Code de champ modifié
22. Huang W-C, Huang C-H, Hsu H-S, Hsien C-C, Hsu W-H, Tsai T-H, Chen YJ (2013) Intrathoracic goiter in elderly patients Int J Gerontol 7:8-12.
23. Cohen JP, Cho HT (1994) Surgery for substernal goiter. In: Feldman M (ed) Operative techniques in otolaryngology and head and neck surgery. WB Saunders Philadelphia, PA, pp 118-125.

24. Uludag M, Isgor A, Yetkin G, Citgez B (2009) Ectopic mediastinal thyroid tissue: cervical or mediastinum originated? *BMJ Case Rep*. doi: 10.1136/bcr.09.2008.1004 **Code de champ modifié**
25. Sahbaz A, Aksakal N, Ozcinar B, Onuray F, Caglayan K, Erbil Y (2013) The "forgotten" goiter after total thyroidectomy. *Int J Surg Case Rep* 4:269-271. **Code de champ modifié**
26. Casadei R, Perenze B, Calculli L, Minni F, Conti A, Marrano D (2002) "Forgotten" goiter: clinical case and review of the literature. *Chir Ital* 54:855-860. **Code de champ modifié**
27. Basaria S, Cooper DS (1999) Graves' disease and recurrent ectopic thyroid tissue. *Thyroid* 9:1261-1264. **Code de champ modifié**
28. Calò PG, Tatti A, Medas F, Petruzzo P, Pisano G, Nicolosi A (2012) Forgotten goiter. Our experience and a review of the literature. *Ann Ital Chir* 83:487-490. **Code de champ modifié**
29. Massard G, Wihlm JM, Jeung MY, Roeslin N, Dumont P, Witz JP, Morand G (1992) Forgotten mediastinal goiter: seven cases. *Ann Chir* 46:770-773. **Code de champ modifié**

Table 1 Description of the population (n = 2570)

Age (years, mean \pm SD)	50.4 \pm 14.0
Female/male ratio	4.42 (2,074/469)
	n
Begnin disease	2029
Multinodular disease	1138
Solitary thyroid nodule	484
Solitary toxic nodule	35
Toxic multinodular disease	112
Grave's disease	218
Thyroiditis	28
Amiodarone-induced hyperthyroidism	1
C cell hyperplasia	13
Malignant disease	524
Papillary	363
Follicular variant papillary	99
Follicular	22
Hurtle cell	19
Medullary	12
Poorly differentiated/anaplastic	3
Lymphoma	1 Hodgkin/1 thyroid lymphoma
Metastasis	2 kidney/1 breast
Parathyroid cancer	1
Tumor of undetermined prognosis	17

Table 2 Anatomopathological characteristics of cervico-mediastinal goiters (CMG) and mediastinal nodules (MN).

Anatomopathological parameters	n (%) [*]		<i>P</i> value [†]
	CMG (n = 67)	MN (n = 42)	
Localization			
anterior	11 (16.4)	21 (50)	0.0002
antero-lateral	17 (25.4)	4 (9.5)	0.0412
lateral	39 (58.2)	17 (40.5)	0.0056
Intrathoracic size (cm, mean ± SD)	2.76 ± 2.18	1.79 ± 1.23	0.0251 ‡
Distance from cervical thyroid (cm, mean ± SD)		1.28 ± 1.41	
Differentiated cancer	6 (8.9)	1 (2.4)	0.173

^{*}All data are presented as numbers (%) except when indicated otherwise

[†]*P* values for differences between CMG and MN were calculated using the χ^2 test except when indicated otherwise

[‡]*P* value for differences between CMG and MN was calculated using the Mann-Whitney U test.

Table 3 Efficiency of the various diagnostic procedures in the evaluation of cervico-mediastinal goiters (CMG) and mediastinal nodules (MN)

Diagnostic procedures	n (%) [*]		P value [†]
	CMG n = 67	MN n = 42	
Patients complaints	10 ‡ (14.9)	5 ‡‡ (11.9)	0.6559
Physical examination	25 (37.3)	13 (30.9)	0.4976
Neck US	45 (67.2)	17 (40.5)	0.0062
Physical examination + Neck US	54 (80.6)	21 (50.0)	0.0008
CXr	42 (62.7)	8 (19.0)	< 0.0001
Physical examination + Neck US + CXr	58 (86.6)	23 (54.8)	0.0002
Intraoperative finding	9 (13.4)	19 (45.2)	< 0.0001
Postoperative finding	0	2 (4.8)	0.0714

Neck US: Neck ultrasonography, CXr: chest X-ray.

^{*}All data are presented as numbers (%)

[†]P value for differences between CMG and MN was calculated using the χ^2 test.

[‡]: dyspnea, n = 7; dysphagia, n = 3; venous compression, n = 2; syncope, n = 1.

^{‡‡}: dyspnea, n = 4; dysphagia, n = 1; dysphonia, n = 1.