



**HAL**  
open science

## Mahorones, Highly Brominated Cyclopentenones from the Red Alga *Asparagopsis taxiformis*

Stephane Greff, Mayalen Zubia, Grégory Genta-Jouve, Lionel Massi, Thierry  
Perez, Olivier P. Thomas

► **To cite this version:**

Stephane Greff, Mayalen Zubia, Grégory Genta-Jouve, Lionel Massi, Thierry Perez, et al.. Mahorones, Highly Brominated Cyclopentenones from the Red Alga *Asparagopsis taxiformis*. *Journal of Natural Products*, 2014. hal-01760562

**HAL Id: hal-01760562**

**<https://amu.hal.science/hal-01760562>**

Submitted on 6 Apr 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


This document is confidential and is proprietary to the American Chemical Society and its authors. Do not copy or disclose without written permission. If you have received this item in error, notify the sender and delete all copies.

**Mahorones, highly brominated cyclopentenones from the red alga *Asparagopsis taxiformis***

Journal:	<i>Journal of Natural Products</i>
Manuscript ID:	np-2013-01094h.R3
Manuscript Type:	Full Paper
Date Submitted by the Author:	09-Apr-2014
Complete List of Authors:	GREFF, stephane; Aix-Marseille Universite, DFME IMBE UMR 7263 Zubia, Mayalen; UMR EIO, Genta-Jouve, Gregory; UMR 8638 CNRS, Massi, Lionel; 2Institut de chimie de Nice (ICN),, Perez, Thierry; IMBE - AMU, Thomas, Olivier; University of Nice-Sophia Antipolis, Chemistry

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



708x271mm (72 x 72 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Mahorones, Highly Brominated Cyclopentenones

## from the Red Alga *Asparagopsis taxiformis*

*Stéphane Greff,*<sup>†</sup> *Mayalen Zubia,*<sup>‡</sup> *Grégory Genta-Jouve,*<sup>§</sup> *Lionel Massi,*<sup>⊥</sup> *Thierry Perez,*<sup>†</sup> *Olivier P. Thomas*<sup>⊥\*</sup>

<sup>†</sup> Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, IMBE UMR 7263 CNRS / IRD / Aix-Marseille Université / Université d'Avignon et des Pays du Vaucluse, Station Marine d'Endoume, rue de la Batterie des Lions, 13007 Marseille, France.

<sup>‡</sup> Agence pour la Recherche et la Valorisation Marine, ARVAM, 97490 Sainte-Clotilde, La Réunion, France.

<sup>§</sup> Laboratoire de Pharmacognosie, UMR 8638 CNRS, Université Paris Descartes, Sorbonne Paris Cité, Faculté des Sciences Pharmaceutiques et Biologiques, 4 avenue de l'Observatoire, 75006 Paris, France

<sup>⊥</sup> Nice Institute of Chemistry-PCRE, UMR 7272 CNRS, Université de Nice Sophia-Antipolis, Faculté des Sciences, Parc Valrose 06108, Nice, France.



**ABSTRACT**

The red alga *Asparagopsis taxiformis* (Rhodophyta, Bonnemaisoniaceae) has been shown to produce a large diversity of halogenated volatile organic compounds, with one to four carbons. As the distribution of this alga may expand worldwide, we implemented a research program that aims to understand the functions of its specialized metabolome in marine ecosystems. Phytochemical investigations performed on *A. taxiformis* gametophyte stages from the Indian Ocean revealed two new highly brominated cyclopentenones named mahorone (**1**) and 5-bromomahorone (**2**). They are the first examples of natural 2,3-dibromocyclopentenone derivatives. Their structure elucidation was achieved using spectrometric methods including NMR and MS. A standardized ecotoxicological assay was used as an assessment of their role in the environment revealing high toxicities for both compounds ( $EC_{50}$  0.16  $\mu$ M for **1** and **2**). Additionally, both compounds were evaluated in antibacterial, antifungal and cytotoxicity assays. Compounds **1** and **2** exhibit mild antibacterial activities against the human pathogen *Acinetobacter baumannii*.

1  
2  
3 The presence of non-indigenous species in marine ecosystems has strongly increased during  
4 the last decades due to globalisation of trade exchanges and transports.<sup>1-3</sup> In this context,  
5 seaweeds have attracted much attention,<sup>4-9</sup> because of the negative impacts they can have on the  
6 environment counterbalanced by the benefits they may provide to human societies.<sup>10-11</sup> While a  
7 general consensus has arisen on the need to minimise these impacts and to limit them in the  
8 future, little is known about the ecological and chemical processes which trigger the proliferation  
9 of an introduced species and the impact that chemical cues may have on native species.<sup>12</sup> Widely  
10 distributed from tropical to temperate waters, species of the genus *Asparagopsis* (Rhodophyta,  
11 Bonnemaisoniaceae) are spreading worldwide, affecting several marine ecosystems.<sup>5,13</sup> The  
12 genus is composed of cryptogenic populations, and it contains only two species to date: *A.*  
13 *armata* Harvey and *A. taxiformis* (Delile) Trevisan de Saint-Léon. From a chemical perspective,  
14 these algae are particularly interesting due the production of a high diversity of halogenated  
15 metabolites.<sup>14</sup> Thoroughly studied in the 1970s, *A. taxiformis* produces low molecular weight  
16 halogenated compounds with one to four linear carbons, including methanes, ethenes, acetic  
17 acids, acetamides, propanols, propanones, propenes, acrylic acids, propylene oxide, propyl and  
18 propenyl acetates, butenols and butenones,<sup>15-21</sup> which exhibit an array of biological activities  
19 (Supporting Information).<sup>22</sup>

20  
21  
22 To better understand the mechanisms associated with the possible proliferation of these algae,  
23 we initiated a chemical ecological study that explores the role of secondary metabolites in their  
24 interaction with native species. In the present part of the study our objectives were: i) to isolate  
25 and characterize the secondary metabolites produced by this alga; ii) to assess the  
26 ecotoxicological activities of fractions and pure compounds from this alga; iii) and to evaluate  
27 their potential as therapeutic agents due to the large biomass of algae available. Unexpectedly,  
28 the phytochemical study of specimens of *A. taxiformis* collected in the Indian Ocean led to the  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

isolation of two new highly brominated cyclopentenones, named mahorone (**1**) and 5-bromomahorone (**2**). We report herein the isolation, the structure elucidation, and also the ecotoxicological as well as antiinfective and antitumor activities of these compounds.

mahorone (**1**)5-bromomahorone (**2**)

## RESULTS AND DISCUSSION


Specimens of *A. taxiformis* were collected off the coast of Mayotte (Indian Ocean). Samples were kept frozen, freeze-dried and ground before extraction by MeOH/CH<sub>2</sub>Cl<sub>2</sub> (1:1) and sonication. The resulting extract was then fractionated by reversed-phase (C<sub>18</sub>) vacuum liquid chromatography using solvents of decreasing polarities (H<sub>2</sub>O, MeOH, CH<sub>2</sub>Cl<sub>2</sub>). The chemical profile of the methanolic fraction showed a very promising UHPLC-DAD-ELSD profile, evidencing major compounds with UV profiles unknown for this alga. A succession of HPLC purification steps on this fraction resulted in the isolation of both new compounds **1** and **2**.

The isotopic pattern in the (-)-HRESIMS spectrum of **1** was indicative of a pentabromo-derivative with the most intense peak at  $m/z$  506.6090, which was in accordance with a molecular formula C<sub>6</sub>H<sub>5</sub>Br<sub>5</sub>O<sub>2</sub> if associated with the most common [M-H]<sup>-</sup> ion observed in (-)-ESIMS. The very low H/C ratio and the presence of a large number of heteroatoms made the structure elucidation highly challenging by NMR. Indeed, the <sup>1</sup>H NMR spectrum of **1**, first performed in CD<sub>3</sub>OD, indicated three signals for a total integration of three, two of them corresponding to an AB system at  $\delta_H$  3.33 (d,  $J$  = 18.5 Hz, 1H, H-5a) and 2.77 (d,  $J$  = 18.5 Hz, 1H, H-5b) and a singlet at  $\delta_H$  6.06 (1H, H-6) (Table 1). The <sup>13</sup>C NMR and HSQC spectra confirmed the presence


1  
2 of six carbons corresponding to the following signals:  $\delta_C$  194.9 (C, C-1) assigned to a ketone  
3  
4 function;  $\delta_C$  131.8 (C, C-2) and 161.2 (C, C-3) assigned to a tetrasubstituted double bond; one  
5  
6 saturated methylene, one saturated methine, and one deshielded and saturated quaternary carbon.  
7  
8 The lack of vicinal and longer range proton-proton coupling information led us to rely on HMBC  
9  
10 correlations to assess the structure of **1** (Figure 1). The proton signals corresponding to the C-5  
11  
12 AB methylene system were HMBC correlated to the other five carbons of the molecule while the  
13  
14 proton signal of the C-6 methine was only correlated to C-3, C-4, and C-5. These data suggested  
15  
16 that the C-5 methylene, the C-1 ketone and the C-2/C-3 unsaturation were placed within a cyclic  
17  
18 system while the C-6 methine was expected to be exocyclic. In this case, the C-6 methine should  
19  
20 be substituted on the resulting five-membered ring. An acyclic system would not lead to H-6/C-3  
21  
22 coupling while, in the case of a six-membered ring system, H-5 and C-3 would be  $^4J$  coupled and  
23  
24 would not exhibit an intense HMBC correlation as observed. Additionally, H-6 would be  
25  
26 expected to couple to four carbons and, in our case, only three couplings are observed. Among  
27  
28 the several possibilities still existing for the arrangement around this ring, the H-5a/C-6 and H-  
29  
30 5b/C-6 together with the H-6/C-5 HMBC correlations were only consistent with a substitution of  
31  
32 the C-6 methine at the C-4 quaternary carbon, one of the positions  $\alpha$  to the C-5 methylene.  
33  
34 Finally, the H-6/C-3 HMBC correlation allowed us to rule out a last possibility for the endocyclic  
35  
36  $\alpha,\beta$ -unsaturated ketone, and the ketone was unambiguously placed at C-1, the second position  $\alpha$   
37  
38 to the C-5 methylene. Among the five bromine atoms identified by HRMS, two of them were  
39  
40 placed on the C-2/C-3 tetrasubstituted double bond, which was in agreement with the detected  
41  
42 chemical shifts of the corresponding  $^{13}\text{C}$  atoms.<sup>23</sup>  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1.**  $^1\text{H}$  (500 MHz) and  $^{13}\text{C}$  (125 MHz) NMR Data for **1** and **2** in  $\text{CD}_3\text{OD}$ .

position	mahorone ( <b>1</b> )			5-bromomahorone ( <b>2</b> )	
	$\delta_{\text{C}}$ , mult.	$\delta_{\text{H}}$ , mult. ( $J$ in Hz)	HMBC (H $\rightarrow$ C)	$\delta_{\text{C}}$ , mult.	$\delta_{\text{H}}$ , mult.
1	194.9, C			192.1, C	
2	131.8, C			130.4, C	
3	161.2, C			159.9, C	
4	84.0, C			82.8, CH	
5a	46.3, $\text{CH}_2$	3.33, d (18.5)	1, 2, 3, 4, 6	50.9, CH	5.29, s
5b		2.77, d (18.5)			
6	50.7, CH	6.06, s	3, 4, 5	49.3, CH	6.20, s

**Figure 1.** a. Key HMBC (H $\rightarrow$ C) correlations for **1**; b. key NOESY (double headed arrow) correlation for **2**.


In order to determine whether the resulting bromines and alcohol were placed on the C-4 quaternary carbon or the C-6 methine, we first relied on a possible metabolic pathway leading to these compounds (Scheme 1). Indeed, chemical shifts modeling would not lead to any definitive conclusion on the position of the substituents at C-4 and C-6 due to close electronic effects of these substituents.



**Scheme 1.** Biogenetic hypothesis for mahorone (**1**).

The six carbons of **1** suggested that the construction of the skeleton could originate from the condensation of two brominated acetones, as these derivatives have already been reported from this alga.<sup>15,18,24</sup> The enol reactivity of the ketone would be involved, first through the connection of the second ketone partner by nucleophilic substitution, and secondly through the cyclization by an aldol-type reaction. Consequently, this hypothesis allowed us to propose the presence of the alcohol at C-4 and two bromines at C-6. In order to ascertain the position of the substituents on the cycle, we performed <sup>13</sup>C NMR modeling for the three possible isomers a, b and c (Table 2). Data obtained for **1** fit best with theoretical values of a, even more when considering values obtained for a compound already reported in the literature with bromines at C-2 and C-3.<sup>23</sup>

**Table 2.** Comparison between Experimental and Theoretical  $^{13}\text{C}$  NMR Chemical Shifts for isomers **a**, **b** and **c** of **1**.



position	a			b		c		Best prediction
	Exp. $\delta_{\text{C}}$	$\delta_{\text{C}}$	$\Delta$ (ppm)	$\delta_{\text{C}}$	$\Delta$ (ppm)	$\delta_{\text{C}}$	$\Delta$ (ppm)	
1	194.9	199.8	4.9	209.6	14.7	201.8	6.9	a
2	131.8	117.7	14.1	102.3	29.5	130.1	1.7	c
3	161.2	157.7	3.5	181.2	20	154	7.2	a
4	84	76.7	7.3	58.2	25.8	56.6	27.4	a
5	46.3	42	4.3	41.2	5.1	41.3	5	a
6	50.7	57.9	7.2	55.7	5	55.3	4.6	c

The fifth bromine atom of the molecular formula obtained by (-)-HRESIMS was consequently suspected to be an additional bromide ion obtained through an ion-molecule reaction during ionization leading to  $[\text{M}+\text{Br}]^-$  and then a molecular formula of  $\text{C}_6\text{H}_4\text{Br}_4\text{O}_2$  for **1**. The EIMS spectrum of **1** was fully consistent with this assumption. The ion cluster with four bromine atoms was observed at  $m/z$  428 (1:4:6:4:1) which are odd values and that should then correspond to the molecular peak. The base peak at  $m/z$  255 (1:2:1,  $[\text{M}-\text{CHBr}_2]$ ) confirmed the five-membered ring and the exocyclic  $-\text{CHBr}_2$  substituent. Bromide adducts are not frequent in (-)-ESIMS and they have usually been observed when halide salts are added to the solvents used.<sup>25-26</sup> In our case the bromide may be present in the methanolic solution just like for sodium adducts or by exchange with another molecule during the ionization process. This reactivity may originate from the absence of acidic protons in the mahorones but also because of the high polarisability of the numerous bromine atoms of the molecules.<sup>27</sup> In order to test this hypothesis, we decided to add salts to our samples to induce the formation of ion adducts. Positive ions were not observed by

1  
2 (+)-ESIMS (adding LiCl or NH<sub>4</sub>Cl) but when adding NH<sub>4</sub>Cl we observed an isotopic ion at *m/z*  
3  
4 459 in the (-)-ESIMS spectrum. The presence of this chloride adduct then confirms our  
5  
6 assumption (Supporting Information).  
7

8  
9 Due to the low level of flexibility for mahorone (**1**), we anticipated that the use of electronic  
10  
11 circular dichroism could easily lead to the determination of the absolute configuration. Even if  
12  
13 the helicity rule proposed by Snatzke is well adapted for *s-trans*  $\alpha,\beta$ -unsaturated ketones,<sup>28</sup> some  
14  
15 recent computational studies applied to 4-substituted 2-cyclohexenones demonstrated that  
16  
17 molecular modeling was necessary to determine the absolute configuration in these cases.<sup>29</sup> We  
18  
19 consequently decided to first perform a conformational analysis of **1** in order to assess the  
20  
21 theoretical ECD spectrum of each calculated conformer (Supporting Information). While the  
22  
23 conclusion was not clear for the  $\pi_{C=C}-\pi^*_{C=O}$  transition at 288 nm, all of the calculated conformers  
24  
25 obtained for the *S* configuration show a positive Cotton effect at 372 nm for the less intense  $n_{C=O}-$   
26  
27  $\pi^*_{C=O}$  electronic transition. The experimental data showing a negative Cotton effect for this  
28  
29 transition were consistent with an *R* configuration for the asymmetric carbon at C-4.  
30  
31  
32  
33  
34

35 The molecular formula of **2** was deduced as C<sub>6</sub>H<sub>3</sub>Br<sub>5</sub>O<sub>2</sub> on the basis of the (-)-HRESIMS  
36  
37 spectrum exhibiting an isotopic cluster centered at *m/z* 586.5183 assuming the same bromide  
38  
39 adduct seen for **1**. In the <sup>1</sup>H NMR spectrum of **2** performed in CD<sub>3</sub>OD, only two singlets were  
40  
41 observed at  $\delta_H$  5.29 (H-5) and 6.20 (H-6). Due to the additional bromine of the molecular  
42  
43 formula and the disappearance of one proton from the methylene group of **1**, we suspected the  
44  
45 substitution of one proton of the methylene at C-5 by a fifth bromine atom. Support was given by  
46  
47 EIMS through the isotopic cluster centered at *m/z* 506 (1:5:10:10:5:1; 5 Br) and a base peak at  
48  
49 *m/z* 333 (1:3:3:1; [M-CHBr<sub>2</sub>]). The relative configuration around the five-membered ring was  
50  
51 assessed using NOESY. Indeed, a clear nOe was observed between H-6 and H-5 which is  
52  
53 consistent with a most likely *cis*-configuration between the hydroxy at C-4 and the bromine at C-  
54  
55  
56  
57  
58  
59  
60

1  
2 5 (Figure 1). Because the ECD spectrum of **2** presented the same Cotton effects as those  
3  
4 encountered for **1**, the 4*R*,5*S* absolute configuration was assigned for this compound.  
5  
6

7 Ecotoxicological activities of chromatographic fractions and pure compounds **1** and **2** were  
8  
9 evaluated against a marine bioluminescent bacteria (*Vibrio fischeri*) using a standardized  
10  
11 Microtox assay in order to target bioactive metabolites and hypothesize their role in the  
12  
13 environment. Bioactivity was quantified by measuring the direct effect on the metabolism of the  
14  
15 bacteria indicated by a decrease in light emitted after 15 min exposure. Toxicity was expressed  
16  
17 as: i) EC<sub>50</sub>, that represents the concentration (in  $\mu\text{g.mL}^{-1}$ ) of compounds/extracts that reduces  
18  
19 bacteria light to a half the initial light; and ii) as gamma units relative to one mg dry mass.mL<sup>-1</sup>  
20  
21 as this index appears to be relevant for ecological comparisons.<sup>30</sup> Interestingly, the MeOH  
22  
23 fraction containing **1** and **2** displays the lowest EC<sub>50</sub> (2.8  $\mu\text{g.mL}^{-1}$ ) and the maximum gamma  
24  
25 (6.452) (Table 3). According to Martì et al., who set the threshold between toxic and non-toxic  
26  
27 samples at 0.5 gamma units, this fraction appears as the most bioactive. As a reference, the  
28  
29 extract of *Falkenbergia rufolanosa*, the tetrasporophytic stage of *A. armata*, presented a gamma  
30  
31 value of 5.204, and was consequently considered as highly toxic.<sup>30</sup> Pure compounds were also  
32  
33 evaluated following the same methodology and revealed high toxicities with EC<sub>50</sub> values of 0.07  
34  
35  $\mu\text{g.mL}^{-1}$  (0.16  $\mu\text{M}$ ) and 0.08  $\mu\text{g.mL}^{-1}$  (0.16  $\mu\text{M}$ ) for **1** and **2**, respectively.<sup>31</sup>  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 3.** EC<sub>50</sub> ( $\mu\text{g}\cdot\text{mL}^{-1}$ ) and Gamma Units of Fractions of *Asparagopsis taxiformis* Gametophyte Stage (Microtox Bioassay after 15 min Exposure). EC<sub>50</sub> values are means  $\pm$  SD (n = 3).

Fraction	Solvent fractionating	EC <sub>50</sub> ( $\mu\text{g}\cdot\text{mL}^{-1}$ )		Gamma
		Mean	SD	
A1	H <sub>2</sub> O	<1000	0.0	0.000
A2	H <sub>2</sub> O/MeOH (1:1)	6.0	0.6	0.135
A3	MeOH	2.8	0.3	<b>6.452</b>
A4	MeOH/CH <sub>2</sub> Cl <sub>2</sub> (1:1)	11.6	1.4	0.012
A5	CH <sub>2</sub> Cl <sub>2</sub>	10.2	0.9	0.000
B1	Hexane	2.9	0.2	0.028
B2	Hexane/EtOAc (1:1)	5.8	0.9	0.081
B3	EtOAc	48.0	3.4	0.001
B4	EtOAc/MeOH	170.5	1.6	0.003
B5	MeOH	391.8	1.2	0.001

Several studies already reported the antimicrobial/antifungal activities of *A. taxiformis* extracts from medium to low polarity fractions, evidencing a large range of biological activities.<sup>32-37</sup> However, biological assays using single compounds from the genus remain scarce.<sup>22,38-39</sup> The release of brominated metabolites involved in the control of epiphytic bacterial communities has been reported from *A. armata* and *Bonnemaisonia asparagoides*.<sup>39-40</sup> Antibacterial and quorum sensing inhibition activities of MeOH extracts of *A. taxiformis* have also been demonstrated.<sup>41</sup> The high toxicities exhibited by the methanolic fraction and compounds **1** and **2** of *A. taxiformis* against a bacterial marine pathogen may be considered as a real advantage for the species. Although their release in the environment has not been demonstrated, we might anticipate their potential role in the interaction of this species with other micro- and/or macro-organisms.

At the same time, we decided to assess the pharmaceutical potential of these compounds, keeping in mind a possible biotechnological use of this widespread macroalga. Isolated metabolites **1** and **2** were thus tested against human bacterial and fungal pathogens: Gram negative bacteria (*Acinetobacter baumannii*, two strains of *Escherichia coli* 2884 and 5746,

*Pseudomonas aeruginosa*), Gram positive bacteria (*Staphylococcus aureus* MRSA/MSSA - methicillin resistant/sensitive strains) and fungi (*Aspergillus fumigatus* and *Candida albicans*). Mahorones **1** and **2** exhibited the strongest effect against *A. baumannii* (MIC<sub>80</sub> of 8 and 16  $\mu\text{g.mL}^{-1}$  respectively) (Table 4). Compound **1** exhibited a weak effect on *E. coli* 5746, while both compounds inhibited the cell growth of Gram positive *S. aureus* MRSA in the same range.

**Table 4.** Antibacterial and Antifungal MIC<sub>80</sub> values of **1** and **2**.

		<b>1</b>	<b>2</b>	<b>Control</b>
		MIC <sub>80</sub> ( $\mu\text{g.mL}^{-1}$ )		
Gram-negative bacteria	<i>A. baumannii</i>	8	16	Rifampicin 1
	<i>E. coli</i> (2884)	> 32	> 32	Novobiocin 0.3
	<i>E. coli</i> (5746)	16	32	Novobiocin 0.1
	<i>P. aeruginosa</i>	> 10	> 10	Ciprofloxacin 1.5
Gram-positive bacteria	<i>S. aureus</i> MRSA	16	16	Imipenem 8
	<i>S. aureus</i> MSSA	> 32	> 32	Penicillin 0.01
Filamentous fungi	<i>A. fumigatus</i>	> 32	> 32	
	<i>A. fumigatus</i> + caspofungin enhancer	> 32	> 32	
	caspofungin enhancer	> 32	> 32	
Yeast	<i>C. albicans</i>	> 32	> 32	

Cytotoxicities of both compounds were also evaluated on an immortalized hepatocyte (Fa2N4) and several human tumor cell lines including lung (A549), liver (HepG2), colon (HT29) and breast (MCF7). Neither compound showed any inhibition of tumor cells, while compound **1** evidenced a cytotoxic effect on healthy liver cells (53 % inhibition of cell growth at 5  $\mu\text{M}$ ). No other study has reported cytotoxic activities of *A. taxiformis* compounds. In addition, Genovese et al. did not find any effect when studying the toxicity of extracts of *A. taxiformis* on digestive glands of mussels, *Mytilus galloprovincialis*,<sup>36</sup> while Zubia et al. highlighted significant cytotoxicities of MeOH/CH<sub>2</sub>Cl<sub>2</sub> extracts of *A. armata* against two tumor cell lines.<sup>42</sup>

## EXPERIMENTAL SECTION

**General Experimental Procedures.** Optical rotations were determined using a Perkin Elmer 343 polarimeter. UV and ECD spectra were measured at 20 °C on a J-810 spectropolarimeter (Jasco), and the IR spectrum was measured on a Bruker Tensor 27 spectrometer. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker Avance 500 MHz spectrometer; chemical shifts were referenced to residual solvent signals (CD<sub>3</sub>OD,  $\delta_{\text{H}} = 3.31$ ,  $\delta_{\text{C}} = 49.0$ ). EIMS spectra were recorded on an Agilent 6890 gas chromatograph coupled to a mass spectrometer 5973N at 70 eV. HRESIMS was performed on a LTQ Orbitrap mass spectrometer (Thermo Finnigan) in negative mode. UHPLC data were acquired on a Dionex Ultimate 3000 equipped with Ultimate 3000 RS pump, 3000 Diode Array Detector and Agilent Technologies 380-ELSD. HPLC purifications of the fractions were achieved on a Jasco LC-2000 system equipped with a PU-2087 Plus preparative pumping system and a UV-2075 Plus detector.

**Collection and Identification of the Alga.** *Asparagopsis taxiformis* gametophyte stages were sampled off the coast of Mayotte (Airport, 12.8213° S, 45.2901° E) in April 2011. Our chemical investigations were focused on the gametophyte stage of the algal life cycle because it represents the highest biomass in native ecosystems and this stage interacts frequently with autochthonous species such as corals and native macrophytes. Voucher specimens were deposited at the MARS Herbarium (MARS07731/Saint-Charles center at Aix-Marseille University). Fresh material was conveyed in ice to the laboratory, frozen at -20 °C and freeze-dried before grinding. Ground material was protected from moisture by adding silica gel packed into paper bags, and stored at -70 °C before extractions.

**Extraction and Isolation.** Dry material (185 g) was extracted three times with 500 mL MeOH/CH<sub>2</sub>Cl<sub>2</sub> (1:1) and sonication (5 min). After filtration, the solutions were pooled and concentrated to dryness at 40 °C to give 17.7 g of a brown residue. The extract was first

1  
2 solubilized in CH<sub>3</sub>CN and then in CH<sub>2</sub>Cl<sub>2</sub> to give fractions A and B (3.7 g and 1.9 g,  
3  
4 respectively). Fraction A was subjected to reversed-phase vacuum liquid chromatography (VLC,  
5  
6 non end-capped C<sub>18</sub> Polygoprep 60-50, Macherey-Nagel) eluting with H<sub>2</sub>O, H<sub>2</sub>O/MeOH (1:1),  
7  
8 MeOH, MeOH/CH<sub>2</sub>Cl<sub>2</sub> (1:1) and CH<sub>2</sub>Cl<sub>2</sub>, giving fractions A1 to A5. Fraction B was subjected to  
9  
10 normal phase VLC (diol, LiChroprep DIOL 40-63 μm, Macherey-Nagel) eluting with hexane,  
11  
12 hexane/EtOAc (1:1), EtOAc, EtOAc/MeOH (1:1) and MeOH, giving fractions B1 to B5.  
13  
14 Chemical analyses of A1-A5, B3-B5 were performed by UHPLC-DAD-ELSD. The purification  
15  
16 was performed on a preparative XSELECT CSH Phenylhexyl column (19 mm x 250 mm, 5 μm,  
17  
18 Waters). Elution rate was set at 12 mL.min<sup>-1</sup>. Initial conditions, maintained during 5 min, were  
19  
20 followed by a linear gradient of CH<sub>3</sub>CN in H<sub>2</sub>O (35 to 90 % for A3) over 22 min. The first  
21  
22 purification afforded pure compound **1** (14.3 mg). Compound **2** was further purified on a semi-  
23  
24 preparative column Synergi Fusion-RP 80A (10 mm x 250 mm, 4 μm, Phenomenex). Elution  
25  
26 was performed at 5 mL.min<sup>-1</sup> in isocratic mode at 55:45 (CH<sub>3</sub>CN/H<sub>2</sub>O), and this second  
27  
28 purification afforded pure compound **2** (0.8 mg).  
29  
30  
31  
32  
33

34  
35 **Mahorone (1)**: yellow amorphous solid;  $[\alpha]_D^{20} +3$  (*c* 0.45, MeOH); UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ )  
36  
37 254 (3.2) nm; ECD (*c* 9.75x10<sup>-5</sup> M, MeOH)  $\lambda_{\max}$  ( $\Delta\epsilon$  M<sup>-1</sup>.m<sup>-1</sup>) 372 (-0.25), 288 (-0.37) nm; IR  
38  
39 (neat)  $\nu_{\max}$  3200 (br), 1728, 1582, 1212, 1045, 1022, 1000, 735 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR data see  
40  
41 Table 1; EIMS (70 eV) *m/z* (%) 428 [M<sup>+</sup>, <sup>79</sup>Br<sup>2</sup><sup>81</sup>Br<sub>2</sub>] (<1), 349 (4), 347 (4), 258 (3), 257 (50),  
42  
43 256 (6), 255 (100), 253 (53), 229 (6), 227 (13), 225 (7), 199 (4), 197 (3), 173 (3), 161 (3), 159  
44  
45 (3), 133 (8), 131 (10), 119 (4), 117 (5), 51 (4), 50 (3), 42 (3); (-)-HRESIMS *m/z* 506.6090  
46  
47 [M+Br]<sup>-</sup> (calcd for C<sub>6</sub>H<sub>4</sub>Br<sub>5</sub>O<sub>2</sub>, 506.6092,  $\Delta$  -0.2 ppm).  
48  
49  
50  
51

52  
53 **5-Bromomahorone (2)**: yellow amorphous solid;  $[\alpha]_D^{20} +2$  (*c* 0.27, MeOH); UV (MeOH)  $\lambda_{\max}$   
54  
55 (log  $\epsilon$ ) 254 (3.2) nm; ECD (*c* 5.64x10<sup>-5</sup> M, MeOH)  $\lambda_{\max}$  ( $\Delta\epsilon$  M<sup>-1</sup>.m<sup>-1</sup>) 370 (-0.59), 288 (-0.70)  
56  
57 nm; <sup>1</sup>H and <sup>13</sup>C NMR data see Table 1; EIMS (70 eV) *m/z* (%) 506 [M<sup>+</sup>, <sup>79</sup>Br<sub>3</sub><sup>81</sup>Br<sub>2</sub>] (<1), 337  
58  
59  
60

1  
2 (32), 335 (95), 334 (11), 333 (100), 332 (16), 331 (35), 330 (14), 268 (14), 266 (13), 255 (17),  
3  
4 253 (15), 251 (14), 225 (19), 223 (19), 197 (11), 187 (9), 185 (9), 175 (10), 173 (14), 133 (19),  
5  
6 131 (24), 79 (11), 62 (11), 61 (10), 50 (10); (-)-HRESIMS  $m/z$  586.5183  $[M+Br]^-$  (calcd for  
7  
8  $C_6H_3Br_6O_2$ , 586.5177,  $\Delta$  +1.02 ppm).  
9

10  
11 **Ecotoxicological Assays.** Microtox (Microbics) is a standardized ecotoxicological bioassay  
12 that measures the toxic effect of compounds/extracts on bioluminescent marine bacteria, *Vibrio*  
13 *fischeri* (NRRL B-11177 strain).<sup>43</sup> Tests were carried out to evaluate the toxicity of A1 to B5  
14 fractions, as well as those of pure compounds **1** and **2**. Fractions were made up to an initial  
15 concentration of 1000 or 2000  $\mu\text{g.mL}^{-1}$  (500  $\mu\text{g.mL}^{-1}$  for pure compounds) using artificial  
16 seawater containing 2% acetone to assist compound dissolution. Concentrations tested were 45,  
17 22.5, 11.25 and 5.625% of the initial concentrations after 15 min of bacteria exposure to toxins.  
18 Samples were diluted when necessary to fit apparatus recommendations. Toxicities, given as  
19 conventional gammas from Microtox, are measured as  $(I_0 / I_t) - 1$  where  $I_0$  and  $I_t$  are the intensity  
20 of the bioluminescences before and after exposure time, respectively. A linear relationship is  
21 obtained when plotting the log of gamma against the log of the tested concentrations, and permit  
22 the direct determination of an  $EC_{50}$  value (equivalent to a gamma of 1) representing the  
23 concentration of fractions (or compounds) that reduces the initial bioluminescence to 50%.  
24 However, because extract concentrations (and consecutive  $EC_{50}$ ) are expressed as  $\mu\text{g}$  of  
25 extracts. $\text{mL}^{-1}$  of solution independent of extraction yields, comparisons between samples/species  
26 were difficult. So, new regression curves were generated replacing original concentrations ( $\mu\text{g}$   
27 extracts. $\text{mL}^{-1}$ ) by modified ones (mg DM alga. $\text{mL}^{-1}$  of solution) as described:  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51  
52 Concentrations (mg DM alga. $\text{mL}^{-1}$ ) = concentration of extract ( $\mu\text{g.mL}^{-1}$ ). dry mass of initial  
53 sample (mg) / dry mass of fraction ( $\mu\text{g}$ ).  
54  
55  
56  
57  
58  
59  
60

1  
2 Using modified concentrations, mathematical relationships were generated using Statgraphics  
3  
4 (four concentrations in triplicate, except for fraction B5 because of insufficient material), and  
5  
6 unconventional gamma units relative to one mg of alga.mL<sup>-1</sup> of solution were calculated,  
7  
8 according to the formula.  
9

10  
11 Gamma = 10<sup>B</sup> where B is the intercept of generated regressions for each fraction (Supporting  
12  
13 Information).  
14

15  
16 **Antimicrobial Assays.** All bioassays were performed by Fundacion Medina (Granada -  
17  
18 Spain). Antimicrobial assays were carried out against six bacteria (*Acinetobacter baumannii*,  
19  
20 *Escherichia coli*, *Pseudomonas aeruginosa* as Gram negative bacterial models; *Staphylococcus*  
21  
22 *aureus* MSSA, *Staphylococcus aureus* MRSA as examples of Gram-positive bacteria). Positive  
23  
24 controls were used against Gram-negative strains (rifampicin, novobiocin, ciprofloxacin for *A.*  
25  
26 *baumannii*, *E. coli*, *P. aeruginosa* respectively) and Gram positive strains (imipenem, penicillin  
27  
28 and tunicamycin for *S. aureus* MSSA, *S. aureus* MRSA respectively). Amphotericin B was used  
29  
30 as negative controls against all strains. The liquid assay procedure was employed to measure  
31  
32 microbial susceptibility of all strains. Antifungal assays were performed on *Candida albicans* as  
33  
34 a yeast model and *Aspergillus fumigatus* as a filamentous fungus, using the liquid assay  
35  
36 procedure.  
37  
38  
39  
40  
41

42 In the liquid assay procedure, compounds were tested in a liquid growth medium dispensed in  
43  
44 96 or 384 well plates (Costar 3370 and 3680 from Corning) inoculated with a bacterial or fungal  
45  
46 suspension. Following overnight incubation at 37 °C, the plates were examined for visible  
47  
48 bacterial/fungal growth as evidenced by turbidity or fluorescence. Compounds were dissolved in  
49  
50 DMSO to a stock solution of 1 mM, and tested at the final concentration of 5 μM in triplicate in  
51  
52 the same experiment. For *A. baumannii*, *P. aeruginosa*, *E. coli* and *S. aureus* liquid assays, 90 μL  
53  
54 of the appropriate diluted inoculums are mixed with 8.4 μL of medium (Luria Broth) and 1.6 μL  
55  
56  
57  
58  
59  
60

1  
2 of a 0.312 mM (DMSO 100%) stock solution of compounds per well. The *A. fumigatus* and *C.*  
3  
4 *albicans* liquid assays were performed in 384-well microtiter plates: 0.5  $\mu\text{L}$  per well of the  
5  
6 compounds from a 0.5 mM solution followed by the inoculum to a final volume of 50  $\mu\text{L}$ . The  
7  
8 medium used was a modified RPMI medium: 10.4  $\text{g}\cdot\text{L}^{-1}$  of RPMI-1640 medium (R8755 from  
9  
10 Sigma), 6.7  $\text{g}\cdot\text{L}^{-1}$  of Yeast Nitrogen Base (YNB) (BD – Becton, Dickinson and Company), 1.8 %  
11  
12 (w/v) glucose and 40 mM HEPES (pH 7.1). An amphotericin B curve (0.5 to 4  $\mu\text{g}\cdot\text{mL}^{-1}$ ) was  
13  
14 used as a control. Cellular viability was scored using resazurin at the final concentration of 0.002  
15  
16 %.<sup>45</sup>  
17  
18  
19

20  
21 **Cytotoxicity Assays.** Five human cancer cell lines (A549, HepG2, HT29, MCF7, MiaPACA2)  
22  
23 and healthy liver cells (Fa2N4) were used to evaluate cytotoxic effects according to the MTT  
24  
25 methodology.<sup>46</sup> Cells were seeded at a concentration of  $3\times 10^4$  cells/well in 200  $\mu\text{L}$  culture  
26  
27 medium and incubated at 37 °C at 5%  $\text{CO}_2$ . Compounds **1** and **2** were prepared at 1 mM in 100%  
28  
29 DMSO. One  $\mu\text{L}$  of this solution was added to 199  $\mu\text{L}$  of culture medium (1/200 dilution), and  
30  
31 left in contact with the cells for 24 h at 37 °C in a 5%  $\text{CO}_2$  incubator. After this time, a MTT  
32  
33 solution was prepared at 5  $\text{mg}\cdot\text{mL}^{-1}$  in PBS 1X and then diluted at 0.5  $\text{mg}\cdot\text{mL}^{-1}$  in MEM without  
34  
35 phenol red. 100  $\mu\text{L}$  of the MTT solution was added to each well. The plates were gently shaken  
36  
37 and incubated for 3 h at 37 °C in 5%  $\text{CO}_2$  incubator. The supernatant was removed and 100  $\mu\text{L}$  of  
38  
39 DMSO 100% was added. The plates were gently shaken to solubilize the formed formazan. The  
40  
41 absorbance was measured using a multireader Victor at a wavelength of 570 nm. Methyl  
42  
43 methanesulfonate MMS (8 mM) was used as positive control, and DMSO 1 % as a negative  
44  
45 control (same concentration as compounds).  
46  
47  
48  
49  
50

51  
52 **Calculations.**  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 NMR prediction was achieved using the PERCH software (PERCH Solutions Ltd). After  
3  
4 importation of the molecular model, geometry optimisation was conducted using Monte Carlo  
5  
6 analysis. The most stable conformer was used to obtain the calculated chemical shifts.  
7  
8

9  
10 ECD calculations were performed at 298 K using the Gaussian03 program package.<sup>47</sup> The  
11  
12 Density Functional Theory (DFT) was used to scan the potential energy surface at the B3LYP/6-  
13  
14 311G\* level to identify the most stable conformers. TDDFT was employed to calculate  
15  
16 excitation energy (in eV) and rotatory strength R in dipole velocity ( $R_{vel}$ ) and dipole length ( $R_{len}$ )  
17  
18 forms. The calculated rotatory strengths were simulated in ECD curve by using the software  
19  
20 package SpecDis.<sup>48</sup>  
21  
22

23  
24 **Statistical Analyses.** Statgraphics Centurion 15.2.11.0 was used to determine linear  
25  
26 regressions.  
27

28  
29 **Supporting Information Available.** All NMR spectra, HRESIMS, EIMS and ECD spectra for  
30  
31 **1** and **2**. This material is available free of charge via the Internet at <http://pubs.acs.org>.  
32  
33

### 34 **Corresponding Author**

35  
36  
37 \* E-mail: [olivier.thomas@unice.fr](mailto:olivier.thomas@unice.fr), Tel: +33(0)49207-6134, Fax: +33(0)49207-6189.  
38  
39

### 40 **Present Addresses**

41  
42 ‡ Ecosystème Insulaires Océaniens, UMR 241 IFREMER/IRD/ILM, Université de la Polynésie  
43  
44 Française, BP 6570, 98702 Faa'a, Tahiti  
45  
46  
47

### 48 **Author Contributions**

49  
50 The manuscript was written through contributions of all authors. All authors have given approval  
51  
52 to the final version of the manuscript.  
53  
54  
55

### 56 **Notes**

1  
2 The authors declare no competing financial interest.  
3  
4

5 **ACKNOWLEDGMENT**  
6  
7

8 This work was supported by the ERA-NET Biome project “SEAPROLIF”. M. Gaysinski  
9 (PFTC Nice) and J.-M. Guignonis (Plateforme Bernard Rossi) are acknowledged for performing  
10 the NMR and HRMS spectra respectively. The ECD spectra were obtained with the help of D.  
11  
12  
13  
14  
15  
16 Bonhomme.  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## REFERENCES

- 1
- 2
- 3
- 4
- 5
- 6 (1) Ruiz, G. M.; Carlton, J. T.; Grosholz, E. D.; Hines, A. H. *Am. Zool.* **1997**, *37*, 621-632.
- 7
- 8 (2) Streftaris, N.; Zenetos, A.; Papathanassiou, E., *Globalisation in marine ecosystems : The*
- 9 *story of non-indigenous marine species across European seas*. CRC Press: Boca Raton, FL,
- 10 USA, 2005; Vol. 43, p 35.
- 11
- 12
- 13
- 14
- 15 (3) Molnar, J. L.; Gamboa, R. L.; Revenga, C.; Spalding, M. D. *Front. Ecol. Environ.* **2008**,
- 16 *6*, 485-492.
- 17
- 18
- 19
- 20 (4) Valentine, J. P.; Magierowski, R., H.; Johnson, C. R. *Bot. Marina* **2007**, *50*, 351.
- 21
- 22 (5) Boudouresque, C. F.; Verlaque, M. *Mar. Pollut. Bull.* **2002**, *44*.
- 23
- 24 (6) Pickering, T. D.; Skelton, P.; Sulu, R. J. *Bot. Marina* **2007**, *50*, 338.
- 25
- 26 (7) Meinesz, A. *Bot. Marina* **2007**, *50*, 373-384.
- 27
- 28 (8) Johnson, C. R.; Chapman, A. R. O. *Bot. Marina* **2007**, *50*, 321-325.
- 29
- 30 (9) Hewitt, C. L.; Campbell, M. L.; Schaffelke, B. *Bot. Marina* **2007**, *50*, 326.
- 31
- 32 (10) Boudouresque, C. F. *Les espèces introduites et invasives en milieu marin*; GIS Posidonie
- 33 Publisher: Marseilles, France, 2008.
- 34
- 35
- 36
- 37
- 38 (11) Schaffelke, B.; Hewitt, C. L. *Bot. Marina* **2007**, *50*, 397.
- 39
- 40 (12) Shine, C.; Williams, N.; Gündling, L., A Guide to Designing Legal and Institutional
- 41 Frameworks on Alien Invasive Species. In *Environmental Policy and Law Paper No. 40*, Union,
- 42 I.-T. W. C., Ed. 2000.
- 43
- 44
- 45
- 46
- 47
- 48 (13) Andreakis, N.; Kooistra, W. H. C. F.; Procaccini, G. *Mol. Ecol.* **2009**, *18*, 212-226.
- 49
- 50 (14) Kladi, M.; Vagias, C.; Roussis, V. *Phytochem. Rev.* **2004**, *3*, 337-366.
- 51
- 52 (15) Fenical, W. *Tetrahedron Lett.* **1974**, *15*, 4463-4466.
- 53
- 54 (16) Burreson, B. J.; Moore, R. E.; Roller, P. *Tetrahedron Lett.* **1975**, 473-476.
- 55
- 56 (17) Woolard, F. X.; Moore, R. E.; Roller, P. P. *Tetrahedron* **1976**, *32*, 2843-2846.
- 57
- 58
- 59
- 60

- 1  
2 (18) McConnell, O.; Fenical, W. *Phytochemistry* **1977**, *16*, 367-374.  
3  
4 (19) Woolard, F. X.; Moore, R. E.; Roller, P. P. *Phytochemistry* **1979**, *18*, 617-620.  
5  
6 (20) Abrahamsson, K.; Ekdahl, A.; Collen, J.; Pedersen, M. *Limnol. Oceanogr.* **1995**, *40*,  
7  
8 1321-1326.  
9  
10 (21) Sugano, M.; Sato, A.; Nagaki, H.; Yoshioka, S.; Shiraki, T.; Horikoshi, H. *Tetrahedron*  
11  
12 *Lett.* **1990**, *31*, 7015-7016.  
13  
14 (22) McConnell, O. J.; Fenical, W. Antimicrobial agents from marine red algae of the family  
15  
16 *Bonnemaisoniaceae*. In *Marine Algae in Pharmaceutical Sciences*, Hoppe, H. A., Levring, T.,  
17  
18 Tanaka, Y., Ed. Water de Gruyter: Berlin, New York, 1979; pp 403-427.  
19  
20 (23) Vadalà, A.; Finzi, Paola V.; Zanoni, G.; Vidari, G. *Eur. J. Org. Chem.* **2003**, 642-648.  
21  
22 (24) Burreson, B. J.; Moore, R. E.; Roller, P. P. *J. Agric. Food Chem.* **1976**, *24*, 856-861.  
23  
24 (25) Kumar, M. R.; Prabhakar, S.; Kumar, M. K.; Reddy, T. J.; Vairamani, M. *Rapid*  
25  
26 *Commun. Mass Spectrom.* **2004**, *18*, 1109-1115.  
27  
28 (26) Cai, Y.; Cole, R. B. *Anal. Chem.* **2002**, *74*, 985-991.  
29  
30 (27) Zhai, H.; Zhang, X. *Water Res.* **2009**, *43*, 2093-2100.  
31  
32 (28) Snatzke, G. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 363-377.  
33  
34 (29) Kwit, M.; Gawronski, J.; Boyd, D. R.; Sharma, N. D.; Kaik, M. *Org. Biomol. Chem.*  
35  
36 **2010**, *8*, 5635-5645.  
37  
38 (30) Marti, R.; Uriz, M. J.; Turon, X. *Mar. Ecol. Prog. Ser.* **2004**, *282*, 73-85.  
39  
40 (31) Botsford, J. L. *ATLA-Altern. Lab. Anim.* **2002**, *30*, 539-550.  
41  
42 (32) Bouhlal, R.; Riadi, H.; Martínez, J.; Bourgougnon, N. *Afr. J. Biotechnol.* **2010**, *9*, 6365-  
43  
44 6372.  
45  
46 (33) Genovese, G.; Leitner, S.; Minicante, S. A.; Lass-Flörl, C. *Mycoses* **2013**, *56*, 516-519.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2 (34) Manilal, A.; Sujith, S.; Kiran, G. S.; Selvin, J.; Shakir, C.; Gandhimathi, R.; Lipton, A. P.  
3  
4 *Ann. Microbiol.* **2009**, *59*, 207-219.  
5  
6 (35) Gonzalez del Val, A.; Platas, G.; Basilio, A.; Cabello, A.; Gorrochategui, J.; Suay, I.;  
7  
8 Vicente, F.; Portillo, E.; Jimenez del Rio, M.; Reina, G. G.; Pelaez, F. *Int. Microbiol.* **2001**, *4*,  
9  
10 35-40.  
11  
12 (36) Genovese, G.; Faggio, C.; Gugliandolo, C.; Torre, A.; Spano, A.; Morabito, M.; Maugeri,  
13  
14 T. L. *Mar. Environ. Res.* **2012**, *73*, 1-6.  
15  
16 (37) Genovese, G.; Tedone, L.; Hamann, M. T.; Morabito, M. *Mar. Drugs* **2009**, *7*, 361-366.  
17  
18 (38) Fenical, W.; McConnell, O. J.; Stone, A. In *Antibiotics and antiseptic compounds from*  
19  
20 *the family Bonnemaisoniaceae (Florideophyceae)*, Proceedings of the Ninth International  
21  
22 Seaweed Symposium, Santa Barbara - California, 20-27 August 1977, 1979; Jensen, A.; Stein, J.  
23  
24 R., Eds. Science Press, Princeton: Santa Barbara - California, 1979; pp 387-400.  
25  
26 (39) Paul, N. A.; de Nys, R.; Steinberg, P. D. *Mar. Ecol. Prog. Ser.* **2006**, *306*, 87-101.  
27  
28 (40) Nylund, G. M.; Persson, F.; Lindegarth, M.; Cervin, G.; Hermansson, M.; Pavia, H.  
29  
30 *FEMS Microbiol. Ecol.* **2010**, *71*, 84-93.  
31  
32 (41) Jha, B.; Kavita, K.; Westphal, J.; Hartmann, A.; Schmitt-Kopplin, P. *Mar. Drugs* **2013**,  
33  
34 *11*, 253-265.  
35  
36 (42) Zubia, M.; Fabre, M.-S.; Kerjean, V.; Deslandes, E. *Bot. Marina* **2009**, *52*, 268-277.  
37  
38 (43) Johnson, B. T. Microtox® Acute Toxicity Test. In *Small-scale Freshwater Toxicity*  
39  
40 *Investigations*, Blaise, C.; Férard, J.-F., Eds. Springer Netherlands: 2005; pp 69-105.  
41  
42 (44) Barry, A. L. Procedure for Testing Antimicrobial Agents in Agar Media: Theoretical  
43  
44 Considerations. In *Antibiotics in Laboratory Medicine*, Lorian, V., Ed. Williams & Wilkins:  
45  
46 Baltimore, Maryland, 1986; pp 1-26.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 (45) Suay, I.; Arenal, F.; Asensio, F.; Basilio, A.; Angeles Cabello, M.; Teresa Díez, M.;  
3  
4 García, J.; González del Val, A.; Gorrochategui, J.; Hernández, P.; Peláez, F.; Francisca Vicente,  
5  
6 M. *Antonie Van Leeuwenhoek* **2000**, *78*, 129-140.  
7

8  
9 (46) Supino, R. MTT Assays. In *In Vitro Toxicity Testing Protocols*, O'Hare, S.; Atterwill, C.  
10  
11 K., Eds. Humana Press: Hatfield UK, 1995; Vol. 43, pp 137-149.  
12

13 (47) *Gaussian, Inc.: Wallingford, CT, USA* **2003**.  
14

15  
16 (48) Bruhn, T.; Schaumlöffel, A.; Hemberger, Y.; Bringmann, G. *Chirality* **2013**, *25*, 243–  
17  
18 249.  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Table of Contents

