

Enterococcus hirae, an unusual pathogen in humans causing urinary tract infection in a patient with benign prostatic hyperplasia: first case report in Algeria

Nadjette Bourafa, L. Loucif, N Boutefnouchet, Jean-Marc Rolain

▶ To cite this version:

Nadjette Bourafa, L. Loucif, N Boutefnouchet, Jean-Marc Rolain. Enterococcus hirae, an unusual pathogen in humans causing urinary tract infection in a patient with benign prostatic hyperplasia: first case report in Algeria. New Microbes and New Infections, 2015, 8, pp.7-9. 10.1016/j.nmni.2015.08.003. hal-01769615

HAL Id: hal-01769615 https://amu.hal.science/hal-01769615

Submitted on 18 Apr 2018 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Enterococcus hirae, an unusual pathogen in humans causing urinary tract infection in a patient with benign prostatic hyperplasia: first case report in Algeria

N. Bourafa^{1,2,3}, L. Loucif^{1,4}, N. Boutefnouchet² and J.-M. Rolain¹

 Unité de recherche sur les maladies infectieuses et tropicales émergentes (URMITE), UM 63, CNRS 7278, IRD 198, INSERM 1095, IHU Méditérranée Infection, Faculté de Médecine et de Pharmacie, Aix-Marseille-Université, Marseille, France, 2) Laboratoire de microbiologie et biochimie appliquée, Département de biochimie, Université Badji mokhtar Annaba, 3) Département de biologie, Faculté des sciences de la nature et de la vie, Université Mohamed Cherif Messaadia-Souk-ahras and 4) Laboratoire de Biotechnologie des Molécules Bioactives et de la Physiopathologie Cellulaire (LBMBPC), Université El Hadj Lakhdar, Batna, Algeria

Abstract

Enterococcus hirae is a zoonotic pathogen rarely isolated from human infections. This case is the first description of *E. hirae* causing urinary tract infection in a diabetic man with benign prostatic hyperplasia from Algeria. The clinical isolate was identified by MALDI-TOF MS and displayed a multisensitivity antibiotic profile.

New Microbes and New Infections © 2015 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

Keywords: Algeria, *Enterococcus hirae*, MALDI-TOF MS, prostate enlargement, urinary tract infection

Original Submission: 8 July 2015; Accepted: 10 August 2015 Article published online: 17 August 2015

Corresponding author: J.-M. Rolain, Unité de recherche sur les maladies infectieuses et tropicales émergentes (URMITE), UM 63, CNRS 7278, IRD 198, INSERM 1095, IHU Méditérranée Infection, Faculté de Médecine et de Pharmacie, Aix-Marseille-Université, Marseille, France

E-mail: jean-marc.rolain@univ-amu.fr

Introduction

Enterococci are important opportunistic pathogens and have become increasingly known as a significant cause of nosocomial and community-acquired infections; *Enterococcus faecalis* and *E. faecium* are the most common species implicated [1]. *E. hirae* is known to cause infections in animals but is rarely isolated from human clinical samples [2]. In the present report, we describe a case of *E. hirae* causing urinary tract infection in a diabetic man with benign prostatic hyperplasia. This case represents the first report of infection by this organism in Algeria.

Case report

A 50-year-old man presented with a medical history of diabetes mellitus type 2 and had prostate enlargement for approximately I year. He was previously hospitalized twice as a result of an inability to fully empty his bladder. He underwent urinary catheterization; however, no antibacterial therapy was administered. The last hospitalization was 2 months before this present admission.

Clinically, the patient presented with symptoms shared between benign prostatic hyperplasia, including straining to urinate, weak urine stream and inability to fully empty the bladder, and symptomatic lower urinary tract infection, including dysuria with cloudy urine, suprapubic pain, urinary frequency and urgency. The patient had a negative blood culture; however, the microscopic analysis of urine indicated the presence of white blood cells (>10 leukocytes per high-power field). The urine culture was positive, and the isolate was identified at the genus level using phenotypic methods including Gram staining, catalase test and growth on bile esculin azide agar and 6.5% NaCl media. The primary bacteriologic diagnosis indicated that the isolated strain belonged to the genus Enterococcus. The biochemical identification using an API 20 Strep system (bio-Mérieux, Marcy l'Etoile, France) could not identify the species involved. Characterization of the clinical isolate to the species level was achieved by MALDI-TOF MS (matrix-assisted laser desorption/ionization time-of-flight mass spectrometry) (Microflex; Bruker Daltonics, Bremen, Germany) using Flex Control and Biotyper 3.0 software (Bruker Daltonics) as previously described [3]. The identification of the strain by MALDI-TOF MS revealed the E. hirae species, with a correct identification score of 2.263. The isolate was multisensitive against nearly all antibiotics tested, including high-level (HL) aminoglycosides (HL gentamicin and HL kanamycin), ampicillin, linezolid, ciprofloxacin, nitrofuran and vancomycin. However, the

New Microbe and New Infect 2015; 8: 7-9

strain presented resistance only to trimethoprim/sulfamethoxazole. The patient received oral treatment with antimicrobial combinations of ampicillin and gentamicin for 10 days. A urethral catheter was inserted for 5 days to empty the bladder. Urine samples were analyzed during treatment until the culture result was negative. The clinical status of the patient improved after 5 days of hospitalization, and the patient was discharged and continued antibiotic therapy at home.

Discussion

Urinary tract infection (UTI) caused by enterococci is very common and is primarily due to *E. faecalis* and *E. faecium* species [4]. Most cases of urinary tract infections occur in women and are uncommon in men [5]. Prostate enlargement, also called benign prostatic hyperplasia, represents an important risk factor for UTI and bacterial prostatitis in men [5,6]. This structural abnormality is mainly associated with aging and most often affects men who are 60 years of age and older [7]. This chronic condition can prevent the bladder from emptying completely, which increases the likelihood that bacteria will grow and trigger an infection [6].

In our case, we report the occurrence of symptomatic lower UTI in a diabetic man with an enlarged prostate caused by an unusual pathogen in humans, *E. hirae*. Clinical diagnosis of our patient showed that the infection was limited to the lower urinary tract based on the absence of clinical signs related to upper UTI, such as flank pain, vomiting and nausea [4], and to prostatitis, such as prostate pain, fever, chills, body aches and perineal pain [5,8]. Notably, in addition to prostate enlargement, urinary tract instrumentation, infection with HIV, interventions of the male urogenital tract and underlying illnesses, such as diabetes, are the main predisposing risk factors for UTI in men [8,9]. In the current case, benign prostatic hyperplasia was the main factor for the occurrence of UTI. Nonetheless, diabetes and urinary catheter indwelling performed for this patient during the previous hospitalizations represent also two other factors that may increase the risk of UTI.

The causative agent isolated in this case was identified as *E. hirae*. This species causes infections in animals, but reported cases in humans are rare [2]. This species was identified for the first time in young chickens [4]. The first report of a human infection caused by *E. hirae* was described in 1998 by Gilad *et al.* [10] in a case of septicemia in a 49-year-old man with renal insufficiency treated with hemodialysis. Since then, few cases have been described in humans [2,4] (Table 1). Notably, among all enterococcal infections in humans, the low incidence of *E. hirae* species has previously been described [1,4]. The bacterium may be underdiagnosed or misdiagnosed by standard identification approaches [4]. Our case demonstrates that MALDI-TOF MS is an important tool useful for rapidly and

TARI	ΕI	•	'aco ro	norte	~f	Enterococcus	hiraa i	in l	human	infocti	inne
IADL			ase re	puius	U.	Lincelococcus	ini ue i		nunnan	mecu	0113

Patient no.	Age (years)/sex, country	Diagnosis	Predisposing factors	Method of E. hirae identification	Source of sample	Antibiotic treatment	Reference
I	49/M, Israel	Septicemia	Hemodialysis catheter	Rapid ID 32 Strep system (bioMérieux, Marcy l'Etoile, France)	Blood	VAN	[10]
2	72/M, France	Native valve Endocarditis	Coronary artery disease	(sodA _{int} gene) ^b sequencing	Blood	AMP, GEN, RIF, VAN	[15]
3	55/M, Spain	Spondylodiscitis	Diabetes mellitus	VITEK 2 automated system (bioMérieux); 16S rRNA gene sequencing	Blood	AMP, GEN, LEV, SXT	[13]
4	78/F, France	Infective endocarditis	Diabetes mellitus, bioprosthetic valve	16S rRNA sequencing;	Blood	AMP, GEN, RIF	[16]
5	62/F, Taiwan	Acute pyelonephritis	Unidentified	BD Phoenix ID/AST Panel Inoculation System (Becton Dickinson, Franklin Lakes, NJ, USA); mur-2 gene ^a sequencing	Blood Urine	AMX	[4]
6	86/F, Taiwan	Acute cholangitis	Unidentified	BD Phoenix ID/AST Panel Inoculation System (Becton Dickinson); mur-2 gene ^a sequencing	Blood	CFM	[4]
7	61/M, Korea	Bacterial peritonitis	Liver cirrhosis	Automated MicroScan WalkAway system; sugar fermentation tests	Blood Ascetic fluid	AMP	[17]
8	44/M, France	Bacteremia Pyonephrosis	Alcoholic liver disease	Undetermined	Blood Urine Kidney biopsy	AMX, CFT, AMK	[14]
9	56/M, Brazil	Native Valve Endocarditis	Diabetes, cardiac arrhythmia with surgical ablation	Undetermined	Blood	AMP, RIF, AMX	[2]
10	50/M, Algeria	Symptomatic Iower UTI	BPH, diabetes mellitus, urinary catheterization	MALDI-TOF MS	Urine	AMP, GEN	This study

AMK, amikacin; AMP, ampicillin; AMX, amoxicillin; BPH, benign prostatic hyperplasia; CFM, cefmetazole; CFT, ceftriaxone; CIP, ciprofloxacin; GEN, gentamicin; LEV, levofloxacin; MALDI-TOF MS, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; RIF, rifampin; SXT, trimethoprim/sulfamethoxazole; VAN, vancomycin. ^aMuramidase gene (*mur-2*) of *E. hirae*.

^bGene encoding manganese-dependent superoxide dismutase.

New Microbes and New Infections © 2015 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases, NMNI, 8, 7–9 This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)

correctly identifying the pathogen at the species level without any complementary tests. The ability of MALDI-TOF MS to identify bacterial colonies from agar was evaluated for a broad range of clinically relevant bacterial strains. The proper identification to the species level is achieved in 80-95% of bacterial isolates [11]. Although the strain isolated in our patient showed a sensitive profile against the majority of antibiotics tested, the study of Robredo *et al.* [12] showed that this species may harbour vancomycin resistance genes and may participate in their transmission to other species of enterococci. The resistance of *E. hirae* to ampicillin and high-level gentamicin has also been reported [13]. More importantly, enterococci long-term urinary tract colonization can worsen and can lead to subsequent invasive infections, such as bacteraemia [4,14].

In conclusion, we report the first case of symptomatic UTI involving *E. hirae* in a patient with benign prostatic hyperplasia from Algeria; this is the first report describing *E. hirae* in this country. This case emphasizes that MALDI-TOF MS is a powerful diagnostic tool for fast and accurate identification of this unusual pathogen at the species level.

Conflict of interest

None declared.

Acknowledgements

We are grateful to L. Hadjadj for technical assistance. We thank American Journal Experts for English-language editing. This work was partly funded by CNRS (CNRS 7278) and IHU Méditerranée Infection.

References

[I] Chan TS, Wu MS, Suk FM, Chen CN, Chen YF, Hou YH, et al. Enterococcus hirae-related acute pyelonephritis and cholangitis with bacteremia: an unusual infection in humans. Kaohsiung J Med Sci 2012;28:111-4.

- [2] Salem-Bekhit MM, Moussa IM, Muharram MM, Alanazy FK, Hefni HM. Prevalence and antimicrobial resistance pattern of multidrug-resistant enterococci isolated from clinical specimens. Indian J Med Microbiol 2012;30:44–51.
- [3] Anghinah R, Watanabe RG, Simabukuro MM, Guariglia C, Pinto LF, Goncalves DC. Native valve endocarditis due to *Enterococcus hirae* presenting as a neurological deficit. Case Rep Neurol Med 2013;2013: 636070.
- [4] Benagli C, Rossi V, Dolina M, Tonolla M, Petrini O. Matrix-assisted laser desorption ionization-time of flight mass spectrometry for the identification of clinically relevant bacteria. PLoS One 2011;6:e16424.
- [5] Wagenlehner FM, Weidner W, Pilatz A, Naber KG. Urinary tract infections and bacterial prostatitis in men. Curr Opin Infect Dis 2014;27: 97–101.
- [6] Bradway C, Bixby MB, Hirschman KB, McCauley K, Naylor MD. Case study: transitional care for a patient with benign prostatic hyperplasia and recurrent urinary tract infections. Urol Nurs 2013;33. 177–179, 200.
- [7] Zhang SJ, Qian HN, Zhao Y, Sun K, Wang HQ, Liang GQ, et al. Relationship between age and prostate size. Asian J Androl 2013;15: 116-20.
- [8] Seminerio JL, Aggarwal G, Sweetser S. 26-year-old man with recurrent urinary tract infections. Mayo Clin Proc 2011;86:557–60.
- [9] Arrellano-Valdez F, Urrutia-Osorio M, Arroyo C, Soto-Vega E. A comprehensive review of urologic complications in patients with diabetes. Springerplus 2014;3:549.
- [10] Gilad J, Borer A, Riesenberg K, Peled N, Shnaider A, Schlaeffer F. Enterococcus hirae septicemia in a patient with end-stage renal disease undergoing hemodialysis. Eur J Clin Microbiol Infect Dis 1998;17: 576–7.
- [11] Vlek AL, Bonten MJ, Boel CH. Direct matrix-assisted laser desorption ionization time-of-flight mass spectrometry improves appropriateness of antibiotic treatment of bacteremia. PLoS One 2012;7:e32589.
- [12] Robredo B, Singh KV, Torres C, Murray BE. Streptogramin resistance and shared pulsed-field gel electrophoresis patterns in vanA-containing *Enterococcus faecium* and *Enterococcus hirae* isolated from humans and animals in Spain. Microb Drug Resist 2000;6:305–11.
- [13] Canalejo E, Ballesteros R, Cabezudo J, Garcia-Arata MI, Moreno J. Bacteraemic spondylodiscitis caused by *Enterococcus hirae*. Eur J Clin Microbiol Infect Dis 2008;27:613–5.
- [14] Brule N, Corvec S, Villers D, Guitton C, Bretonniere C. Life-threatening bacteremia and pyonephrosis caused by *Enterococcus hirae*. Med Mal Infect 2013;43:401–2.
- [15] Poyart C, Lambert T, Morand P, Abassade P, Quesne G, Baudouy Y, et al. Native valve endocarditis due to *Enterococcus hirae*. J Clin Microbiol 2002;40:2689–90.
- [16] Talarmin JP, Pineau S, Guillouzouic A, Boutoille D, Giraudeau C, Reynaud A, et al. Relapse of *Enterococcus hirae* prosthetic valve endocarditis. J Clin Microbiol 2011;49:1182–4.
- [17] Sim JS, Kim HS, Oh KJ, Park MS, Jung EJ, Jung YJ, et al. Spontaneous bacterial peritonitis with sepsis caused by *Enterococcus hirae*. J Korean Med Sci 2012;27:1598–600.