



HAL
open science

Global change effects on land management in the Mediterranean region

Žiga Malek, Peter H. Verburg, Ilse R. Geizendorffer, Alberte Bondeau,
Wolfgang Cramer

► **To cite this version:**

Žiga Malek, Peter H. Verburg, Ilse R. Geizendorffer, Alberte Bondeau, Wolfgang Cramer. Global change effects on land management in the Mediterranean region. *Global Environmental Change*, 2018, 50, pp.238-254. 10.1016/j.gloenvcha.2018.04.007 . hal-01773448

HAL Id: hal-01773448

<https://amu.hal.science/hal-01773448v1>

Submitted on 8 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Global change effects on land management in the Mediterranean region

Žiga Malek¹, Peter H Verburg^{1,2}, Ilse R Geijzendorffer³, Alberte Bondeau⁴, Wolfgang Cramer⁴

¹ Institute for Environmental studies, Vrije Universiteit Amsterdam, De Boelelaan 1087, 1081 HV Amsterdam, the Netherlands

² Swiss Federal Institute for Forest Snow and Landscape Research, WSL Zürcherstrasse 111, CH-8903, Birmensdorf, Switzerland

³ Tour du Valat Research Institute for the Conservation of Mediterranean wetlands, Le Sambuc, 13200 Arles, France

⁴ Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale (IMBE), Aix Marseille Université, CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée Bât. Villemin – BP 80, F-13545 Aix-en-Provence cedex 04, France

Corresponding author: Žiga Malek, z.malek@vu.nl. Telephone: +31 20 59 88710

Acknowledgements

The research in this paper has been supported by the European Research Council under the European Union's Seventh Framework Programme project LUC4C (Grant No. 603542), OPERAs (Grant No. 308393) and ERC grant GLOLAND (no. 311819). This paper contributes to the objectives of the Global Land Project (<http://glp.earth/>) and to the objectives of the Labex OTMed (no. ANR-11-LABX-0061) funded by the French Government Investissements d'Avenir program of the French National Research Agency (ANR) through the A*MIDEX project (no. ANR-11-IDEX-0001-02).

1. Introduction

Most of the Earth's land surface has been changed as a result of human use, with large environmental consequences and both positive and negative impacts on human well-being (Ellis *et al.*, 2010; Schmitz *et al.*, 2012). With limited resources of land and water, a large societal challenge consist of meeting the increasing demand for food and living space for growing populations in the context of climate change. These challenges are especially significant for the Mediterranean, a dynamic and densely populated region with severe constraints on land and water resources (Giorgi & Lionello, 2008; Giannakopoulos *et al.*, 2009; García-Ruiz *et al.*, 2011; Fader *et al.*, 2016). The Mediterranean has a long history of land use, resulting in valuable cultural landscapes created throughout centuries (Blondel *et al.*, 2010; Tieskens *et al.*, 2017), and is one of the most rich areas in terms of biodiversity (Cuttelod *et al.*, 2009). On the other side, human activities in the region have resulted in significant degradation of soil and water resources (García-Ruiz *et al.*, 2011; Karamesouti *et al.*, 2015).

Resulting from its cultural and environmental characteristics and its long land use history, the Mediterranean Basin hosts a diversity of land systems of varying intensities and levels of (multi)functionality. Intensive systems have higher yields and produce most of the crops in the region, a large part of them being exported. These systems however also have high water demands (Daccache *et al.*, 2014) and can negatively affect the quality of soil contributing to land degradation (Karamesouti *et al.*, 2015). Traditional mosaic systems represent landscapes, where human activities and environmental conditions are intricately linked. An example is the *dehesa / montado* system of Spain and Portugal, where different activities such as livestock grazing, cereal production, and forestry occur simultaneously (Joffre *et al.*, 1999). Although these areas have lower yields, they contribute significantly to total regional food production (Blondel, 2006; McAdam *et al.*, 2008). Many of the traditional mosaic systems are associated with high biodiversity values (Médail & Quézel, 1999). These landscapes are particularly vulnerable to global change, threatening their supply of not only food, but a number of ecosystem services (Zamora *et al.*, 2007; Guiot & Cramer, 2016).

The Middle Eastern and North African part of the Mediterranean region is characterized with high population pressures and increasing dependence on food imports (Wright & Cafiero, 2011). Depending heavily on food imports makes the region more vulnerable to fluctuations in food supply and prices (Sowers *et al.*, 2010). The region hosts a considerable portion of cropland with relatively low yields, meaning that future cropland expansion and intensification will play a crucial role in satisfying the demand for food (Mueller *et al.*, 2012). This can however exacerbate soil and water degradation, and appropriate land management will be needed to reduce these consequences, or restore soil and water resources (Cerdan *et al.*, 2010; García-Ruiz, *et al.*, 2011). Moreover, water and land grabbing are also significant issues in the region, leading to conflicts (GRAIN, 2012; Houdret, 2012). The European Mediterranean area hosts high-input intensive agricultural systems significant for regional food production and global

37 commodity markets. However, recent socio-economic development, such as the Greek financial crisis,
38 have influenced the steadiness of supply of agricultural products (Pfeiffer & Koutantou, 2015). Other
39 global change effects are the abandonment of traditional livestock grazing systems due to low economic
40 competitiveness and reduction of livestock productivity (de Rancourt *et al.*, 2006; Bernués *et al.*, 2011). In
41 summary, future global change, particularly changes to climate and population, could significantly impact
42 the potential food supply of the Mediterranean region (Evans, 2008; Sowers *et al.*, 2010).

43 Published global land change scenarios suggest significant intensification of crop production and grazing,
44 together with urban expansion in the Mediterranean region (Hurtt *et al.*, 2011; Letourneau *et al.*, 2012;
45 Souty *et al.*, 2012; van Asselen & Verburg, 2013). These global studies often do not consider specific
46 regional characteristics that could affect these processes, such as the existence of a large share of
47 permanent crops and traditional mosaic systems, and severe water limitations. Water limitation also
48 affects intensification and cropland expansion, feedback loops which are currently not possible to study
49 with land use models in which water availability is represented by a proxy, such as precipitation (NRC,
50 2014). Simplified proxies can only influence the spatial distribution of intensive systems and do not limit
51 cropland expansion or intensification based on available water resources. Land use modeling studies that
52 do take into account water scarcity are often unable to generate land use patterns with the spatial detail
53 of most biophysical models (see for example Lotze-Campen *et al.*, 2010).

54 In this study, we determine the impact of two potential future scenarios on land management in the
55 Mediterranean region to study environmental consequences of increasing food production in the
56 Mediterranean for the year 2050. We advance from the existing knowledge by combining global outlooks
57 of socio-economic and climate change in a land system change model with regional spatial
58 characteristics and configuration of land use. We are particularly interested in how global change might
59 affect traditional Mediterranean landscapes and water resources. We also demonstrate how water
60 resources limitations can be represented in land system models.

61 **2. Future challenges for land management in the Mediterranean region**

62 2.1 The Mediterranean region

63 We focused on the Mediterranean ecoregion defined by the approximate extent of representative
64 Mediterranean natural communities from a biogeographical study (Olson *et al.*, 2001). We expanded the
65 ecoregion by also including the Nile Delta, the Po floodplain and numerous “islands” of similar ecoregions
66 within the Mediterranean ecoregion (Fig. 1). Thematically, we divided the region into two parts, North and
67 South, which, based on land use characteristics and biodiversity trends (Galewski *et al.*, 2011), were
68 subdivided into two sub-regions each (Fig. 1). In addition, this subdivision accounted for more uniform
69 markets that need to fulfil their own demands for food and living space and took into account socio-
70 economic disparities between the Northwest and South. In total, the study area covers 2.3 million km² in

71 27 countries with around 420 million inhabitants in 2015 (CAPMAS, 2015; EUROSTAT, 2016c; IIASA,
72 2016).

73 We identified four major challenging trends for the region based on various documents on the future of
74 the Mediterranean (Appendix A): 1) increasing population and continuous urban sprawl; 2) agriculture and
75 food production; 3) threatened biodiversity, and 4) significant climate change impacts and increasing
76 water scarcity.

77 2.2 Population and urban expansion

78 The total population in the southern (Fig. 1) Mediterranean countries is expected to increase by 43% until
79 2050, and by 16% in the North (Fig. 1, including Western Balkans and Turkey) following the Shared
80 Socioeconomic Pathways 2 (SSP2) scenario (Kc & Lutz, 2014; IIASA, 2016). Urban population in the
81 South is expected to increase even more by 82% (Jiang & O'Neill, 2015). Urban expansion in the
82 Mediterranean region is defined by dispersed and poorly managed urban sprawl (Benoit & Comeau,
83 2012; Salvati *et al.*, 2012). Significant portions of Mediterranean coasts are being transformed to built-up
84 coastal landscapes (Munoz, 2003; Parcerisas *et al.*, 2012). Agriculture is pushed further into wetlands
85 where surface water is available (MWO, 2012). As fertile areas in the region are limited to river plains and
86 coastlines, a large extent of cropland is lost due to soil sealing (Mediterranean 2030 Consortium, 2011).
87 Future population increase will undoubtedly result in further pressure on coastal urban areas, especially
88 near existing large urban agglomerations (European Commission, 2011).

89 2.3 Agriculture and food production

90 Agriculture in the region reflects its socio-economic disparities. The northern part hosts high-input
91 intensive agricultural systems, with most of the crops being exported. In contrast, the Southern
92 Mediterranean hosts a considerable portion of cropland with low yields and less efficient agricultural
93 management (Mueller *et al.*, 2012). Trade regulations, protection of the European market and food safety
94 requirements pose serious constraints to the Southern Mediterranean countries to export commodities
95 such as fruit and vegetables (Larson *et al.*, 2002; Cioffi & dell'Aquila, 2004; García Martínez & Poole,
96 2004). Access to agricultural technology, subsidies and loans is unequally distributed in the region and is
97 more accessible to farmers in the north-western part. Pasture based systems are becoming less
98 competitive, due to high labour costs – on-farm resources are being substituted with external inputs,
99 promoting intensive livestock systems near urban areas (Steinfeld *et al.*, 2006; Bernués *et al.*, 2011). This
100 has resulted in the increase in landless livestock systems in the Mediterranean region (Steinfeld *et al.*,
101 2006).

102 2.4 Biodiversity

103 The Mediterranean ecoregion was identified as one of the world's biodiversity hotspots, as it hosts a large
104 number of plant and animals species, numerous among them endemic (Cuttelod *et al.*, 2009). The
105 presence of these species is closely related to extensive Mediterranean landscapes, particularly agro-
106 silvo-pastoral mosaic systems (Médail & Quézel, 1999) and wetlands (Cuttelod *et al.*, 2009). The
107 transformation of these systems into intensive, single function cropland systems, or their abandonment
108 into woodland can have a significant impact on the biodiversity of the region, including changes in
109 landscape and plant community structure (Médail & Quézel, 1999). Only 5.5% of the region's area is
110 protected, with 90% of protected areas in the Mediterranean North (Benoit & Comeau, 2012; FAO, 2013).
111 Among the main threats to the region's biodiversity are also urban concentration and expansion in coastal
112 areas (FAO, 2013). Human activities have also affected Mediterranean wetlands, through increased
113 water extraction or livestock grazing (Houérou, 1993; Ayache *et al.*, 2009).

114 2.5 Climate and water constraints

115 The region is projected to experience warming exceeding global trends, with most climate change
116 scenarios also resulting in reduced water availability (Chenoweth *et al.*, 2011; Keenan *et al.*, 2011; Guiot
117 & Cramer, 2016). Climate change and water scarcity have even been proposed as potential contributors
118 to conflicts in the region (e.g. Gleick, 2014; Kelley *et al.* 2015). Water scarcity is likely to pose severe
119 limitations to the agricultural sector in the future, as numerous countries risk not being able to meet
120 irrigation requirements (Fader *et al.*, 2016). Already today, freshwater resources in the region are being
121 extracted at unsustainable rates, not allowing for natural replenishment (FAO, 2016). However, improved
122 irrigation efficiency and a shift to crops with lower irrigation demands could considerably lower the
123 requirement for irrigation water withdrawal in the region (Daccache *et al.*, 2014; Fader *et al.*, 2016).

124 **3. Methods**

125 3.1 Land system change simulation

126 Land systems characterize human-environment interactions in landscapes and are defined as
127 combinations of land use and land cover, livestock and management type and intensity (van Asselen &
128 Verburg, 2012; Turner *et al.*, 2013). The use of land systems is a particularly suitable approach for the
129 Mediterranean region with its diverse mosaic landscapes that may not be easily disentangled into
130 separate land cover classes. Moreover, land systems allow a clear quantification of goods and services
131 provided by each land systems unit, necessary to simulate future change. To simulate future land system
132 change until 2050, we applied the CLUMondo model (Fig. 2). CLUMondo simulates land system changes
133 driven by defined demands for specific goods or services provided by land systems, taking into account
134 the local spatial characteristics (van Asselen & Verburg, 2013). As a baseline, we used the Mediterranean
135 land systems map for 2010 (Fig. 3, Malek & Verburg, 2017a). Each land system is characterized by an
136 average cropland and urban extent (% of the land system unit area of 4 km²), and livestock density
137 (livestock units in nr. per unit). In the land systems map, irrigated cropland systems are present on areas

138 equipped with irrigation. Intensive rain-fed cropland was identified on areas where fertilizer application,
139 field size or yields indicate intensive agriculture. The remaining rain-fed cropland was identified as
140 extensive (for details on the method see Malek & Verburg, 2017a). Every land system provides annual
141 and/or permanent crops, livestock and consists of a certain fraction of built-up area (Appendix B). The
142 land systems' output of crops (annual and permanent) is a value specific to each land system. These
143 values were calculated based on crop production statistics to land systems and are described in
144 Appendix B.

145 The actual output of annual and permanent crops was based on agricultural production statistics for 2010
146 (EUROSTAT, 2013; 2016a, 2016b). We used subnational statistics to exclude non-Mediterranean parts of
147 countries (France, Spain, Italy, Turkey), where a significant share of crops is produced. For countries in
148 the Southern Mediterranean, we looked at national statistics on cropland areas and production in
149 subnational units within the Mediterranean ecoregion. In most countries, virtually all crops were produced
150 in the Mediterranean part (for example, 99% in Algeria and Morocco). The only exception was Egypt,
151 where 65% of the crops are produced in the Nile delta, which we used to adjust national crop production
152 statistics (FAO, 2016; Mohamed, 2016). Crop production was aggregated for each sub-region, and all
153 land system output values present a mean for a specific sub-region. We used the SPAM database to
154 identify the shares of crops produced in irrigated, intensive rain-fed and extensive rain-fed systems (You
155 et al., 2014). Total crop production was then scaled according to these shares and calculated for the
156 appropriate land system type (e.g. irrigated systems were associated with crops produced on irrigated
157 cropland, intensive rain-fed with crops produced in intensive cropland). We used agricultural statistics for
158 2010 for two reasons. First, the year 2010 presents the most recent year where crop production statistics
159 were consistently reported for subnational units in our study area. Secondly, the reported national crop
160 production for 2010 deviates the least from the average national crop production of the last 20 years (less
161 than 2%).

162 Annual crops consist of cereals (wheat, maize, barley and rice), and vegetables (fresh vegetables,
163 potatoes and tomatoes). In the Mediterranean, wheat, maize and barley present the majority of cereals,
164 with rice having a significant share in Egypt and Italy (EUROSTAT, 2013, 2016a, 2016b). Permanent
165 crops consist of fruit, olives and dates. Olives, grapes and citrus alone amount to over 20% of total crop
166 production in the Mediterranean region (Daccache *et al.*, 2014). Livestock output is based on the values
167 derived from a global livestock database, with values for bovines, goats and sheep calculated to livestock
168 units (Robinson *et al.*, 2014).

169 CLUMondo allocates changes to land systems based on spatial preference, spatial restrictions and
170 competition between land systems (van Asselen & Verburg, 2013). Spatial preference or local suitability
171 describes the probability of each location (grid cell) to host a specific land system, based on its
172 biophysical and socioeconomic conditions. CLUMondo allocates future land change in locations with
173 highest preference for a defined land system. To calculate spatial preferences the relationships between

174 the spatial occurrence of a specific land system and location factors (explanatory biophysical and
175 socioeconomic variables presented in Appendix C) are investigated using logistic regression. This results
176 in maps presenting the likelihood of occurrence of different land systems as a function of local
177 environmental and socio-economic conditions (Appendix D). Spatial preference maps are updated
178 annually to account for population and climate change, as described in later sections. Spatial restrictions
179 are constraints for changing specific land systems, such as protected areas. Spatial restrictions can either
180 completely constrain land change, or allow predefined changes to land systems (e.g. woodlands can
181 change into forests in a protected area). To achieve a solution, CLUMondo iterates different land system
182 allocation combinations until it fulfils all demands for a specific year for a specific scenario. Although the
183 model can handle the provision of numerous crops and livestock units by land systems, it will promote the
184 most competitive (while accounting for both the occurrence likelihood and competitiveness in terms of the
185 services delivered towards the demand) system for that demand. The model is described in more detail
186 by (van Asselen & Verburg, 2013) and is available (as open-source) at
187 <http://www.environmentalgeography.nl/site/data-models/data/clumondo-model/>.

188 3.2 Limiting water resources

189 Besides achieving the most likely land system allocation under the defined demands, CLUMondo can
190 take into account the limitations of the allocated land systems in terms of resource use. In this study, use
191 of water resources was constrained, by applying a threshold on the total irrigation water withdrawal. This
192 situation is implemented using physical limitations as a maximum level of resource use which cannot be
193 overshoot. If the demand cannot be satisfied fully with irrigated land systems under the given constraints,
194 CLUMondo has to fulfill it with rain-fed land systems. This limits the allocation of irrigated land systems
195 with high water requirements, although they have the highest crop output (Appendix B) and are therefore
196 promoted by the model if irrigation is not limited. To satisfy the demand for annual and permanent crops,
197 CLUMondo needs to consider other land systems without irrigation demands, but also with lower output
198 for these crops, having a tradeoff on the conversion of (semi-)natural areas.

199 To associate irrigated land systems with irrigation water withdrawal, we used spatially explicit irrigation
200 data on areas equipped with irrigation (Siebert *et al.*, 2005, 2013). Irrigation demand values for irrigated
201 land systems were based on the extent of irrigated areas, which we linked to national and subnational
202 irrigation water withdrawal statistics (EUROSTAT, 2013, 2016a, 2016b; FAO, 2016). This resulted in
203 mean values of irrigation water withdrawal per cell of irrigated land system for each region (Appendix B).
204 The available water resources, that served as a limit for irrigation water withdrawal, are based on water
205 resource statistics on the national (Mediterranean South: FAO, 2016) and subnational level
206 (Mediterranean North: EUROSTAT, 2013, 2016a, 2016b; Egypt: FAO, 2016; Mohamed, 2016).

207 3.3 Scenarios

208 3.3.1 Scenario introduction

209 To study potential pathways to fulfill the growing food production and the demand for living area in the
210 Mediterranean region, we have developed two contrasting scenarios: 'growth' and 'sustainability'. The
211 scenarios follow different assumptions in terms of production of annual and permanent crops, and
212 livestock, and the demand for built-up area up to 2050 (Table 1). Both scenarios are based on global
213 SSP2 projections for the region in terms of regionally produced food, however modified to fit the storyline
214 of each scenario as described in Table 2 and Appendix E. The production values for annual and
215 permanent crops, and livestock are based on the SSP2 marker scenario projections for food production
216 (Fricko *et al.*, 2016; Riahi *et al.*, 2016; Popp *et al.*, 2017). These projections are a result of integrated
217 assessment models, where the remaining consumption is satisfied with imports. In the projections, the
218 Mediterranean North maintains producing more than it consumes in 2050, whereas the dependency of
219 the Mediterranean South on food imports increases despite significant crop and livestock production
220 growth (Alexandratos & Bruinsma, 2012). The Mediterranean South imported 56% of its total crop
221 consumption in 2007 (Wright & Cafiero, 2011), which is projected to increase to 73% in 2050 (World
222 Bank, 2009; Alexandratos & Bruinsma, 2012). The demand for built-up areas is based on the population
223 growth rate of the SSP2 scenario (Kc & Lutz, 2014). The scenarios differ from each other significantly
224 regarding the handling of major challenges the Mediterranean region will be facing in the future. In
225 particular, in the sustainability scenario, freshwater is explicitly regarded as a limited resource for
226 irrigation, while in the growth scenario, it is not. Technical details of translating the scenario storylines to
227 the CLUMondo land system model are described in Appendix E.

228 3.3.2 Growth scenario – Production maximization through growth

229 Under the growth scenario the current economic disparities between the sub-regions are assumed to
230 remain unchanged. Each sub-region follows its own goals, aiming to satisfy its own demands.
231 Environmental issues, such as freshwater resources depletion are not considered.

232 *Urban expansion.* Urban expansion has priority to any other land process. Low density peri-urban
233 systems are promoted, continuing urban sprawl. Rural abandonment continues in the North of the
234 Mediterranean Basin. In the South, rural population increases until 2020, followed by a gradual decrease
235 (Jiang & O'Neill, 2015).

236 *Agriculture and food production.* Agricultural intensification has priority – extensive areas are promoted to
237 convert to intensive ones, achieved by higher use of fertilizers, irrigation and herbicides. Yield
238 improvement technology remains unequally distributed in the region. Yield gaps are not closed, however
239 achieve 75% of the attainable yield for irrigated systems. Landless livestock systems are promoted –
240 indoor breeding systems on intensive cropland or near urban areas are subject to 15% intensification. No
241 food waste reduction efforts are made.

242 *Biodiversity.* The extent of protected areas in the Mediterranean does not increase, but existing protected
243 areas are conserved. Cropland and livestock activities with low intensity (extensive cropland and mosaic
244 systems) are allowed in protected areas. Wetlands continue to be used for cropland and used for
245 intensive livestock grazing. Grazing in arid areas is uncontrolled.

246 *Climate and water constraints.* Same crops are being grown as today which, given the reduction in
247 precipitation and increases in evaporation, results in lower suitability for rain-fed intensive systems
248 occurring in semi-arid areas due to climate change. Water resources can be exploited without limitations.

249 3.3.3 Sustainability scenario – Fulfilling the Mediterranean sustainable development goals

250 The sustainability scenario describes a future, where the future Mediterranean region is characterized by
251 prosperity, solidarity and stability. Throughout the region, national governments and other actors agree to
252 follow shared environmental and development goals.

253 *Urban expansion.* Urban sprawl is limited and well planned. A new type of a Mediterranean compact city
254 emerges, denser urban areas are promoted. Rural population remains stable.

255 *Agriculture and food production.* A common Mediterranean market involving all countries around the
256 Mediterranean sea (Fig. 1) is established. It results in a liberalization of agricultural trade, with improved
257 access to agricultural technology, loans and subsidies for everybody. The entire region can easier satisfy
258 its demand through additional food with imports, and can export even more commodities such as
259 permanent crops. Yield gaps on irrigated and rain-fed intensive cropland are closed in the North, and
260 reach 90% (irrigated) and 75% (rain-fed intensive) of attainable yield in the South. Land based livestock
261 systems are promoted. The efficiency of livestock production systems in terms of output is improved, for
262 example by improving breeds or fattening of herds (Bernués *et al.*, 2011). A moderate reduction in food
263 waste due to changed dietary habits (North) and improvement in the food supply chain (South) results in
264 a lower increase in total regional demand for agricultural production.

265 *Biodiversity.* The entire region applied coherent and consistent environmental policies and conservation
266 tools. The extent of protected areas in the region reaches the 17% of terrestrial ecosystems as specified
267 by the “Aichi target” of the UN Convention for Biodiversity (<http://www.cbd.int/sp/targets>), being a
268 considerable improvement for the Mediterranean South (Pouzols *et al.*, 2014) (Appendix F). Low intensity
269 cropland and livestock grazing (extensive cropland and mosaic systems) are allowed in newly established
270 protected areas. Wetlands use by intensive crop production system and grazing of livestock is decreased.
271 Grazing intensification in arid areas is limited in order to combat desertification. Ecological focus areas
272 are assigned as 5% of area set aside in all rain-fed systems. This way, landscape elements with higher
273 biodiversity values are being protected or established.

274 *Climate and water constraints.* Due to improvement in cultivars and crop changes (for example with crops
275 with lower water demands), intensive rain-fed areas can expand more than they do in the growth scenario
276 (Daccache *et al.*, 2014). Improvements in irrigation infrastructure and changes to irrigation type (e.g. drip
277 irrigation lead to a 35% improvement in irrigation efficiency . Irrigation water withdrawal is limited to 75%
278 of current total withdrawal, to allow more water for biodiversity and ecosystems.

279 3.4 Climate and population change

280 In both scenarios, global change was included either through a change in the location factors which
281 underpin the spatial preference map for land system transitions, or as a land system change constraint.
282 We used results from downscaled global climate models from CMIP5 (Hijmans *et al.*, 2005; Taylor *et al.*,
283 2012) forced by the RCP4.5 greenhouse gas radiative forcing representative concentration pathway.
284 RCP4.5 is a cost-minimizing mitigation scenario, presenting a “median” and probable pathway compared
285 to other scenarios (Thomson *et al.* 2011). We calculated the mean of 19 CMIP5 simulation outputs
286 (Appendix G) for both temperature and precipitation for the years 2041-2060, referred to as year 2050.
287 Based on these projections, we generated annual temperature and precipitation maps, which were then
288 used to derive annual potential evapotranspiration (PET) and aridity index (AI) maps (Trabucco *et al.*,
289 2008; Zomer *et al.*, 2008) (Appendix H). Temperature, precipitation and PET served as location factors in
290 the logistic regression, and AI was used to limit particular processes (Appendix E). For example, forest
291 expansion was possible only in areas with $AI > 0.65$ (Zomer *et al.*, 2008).

292 To account for population change, we updated the 2010 population density map using the SSP2
293 projection growth rates (CIESIN, 2015; Jiang & O’Neill, 2015). We used urban population growth rates for
294 areas with higher population density (>250 inhabitants/km²) for both scenarios (Appendix I). This
295 corresponds well with population change projections for the Mediterranean region (Benoit & Comeau,
296 2012). Changes to rural population were applied to the rural population density map (Appendix C), and
297 were also based on global SSP2 projections (CIESIN *et al.*, 2011; Jiang & O’Neill, 2015).

298 3.5 Changes to productivity of land systems

299 Changes to productivity were implemented by changing the output of each land system in time (Appendix
300 B). We first studied the current average yield gap of land systems using yield gap data for major crops
301 (Foley *et al.*, 2011). Then, we calculated annual growth rates of land system output changes to achieve
302 crop yields assumed attainable under the scenario conditions (Tables 2 and 3). The attainable yields
303 were based on plausible improvements to nutrient management and irrigation as proposed by Mueller *et al.*
304 *et al.* (2012). Mueller *et al.* (2012) did not consider climate change impacts on crop yields, although the
305 potential to close yield gaps in different regions was constrained by climate conditions.

306 Because the used productivity changes do not explicitly account for climate change impacts on
307 productivity, we compared the annual productivity increases in both scenarios (Table 3), with other

308 scenario studies that focus on future yield change in the Mediterranean region. Our assumed productivity
309 increases are lower than in other studies, except in both climate change studies (Parry et al., 2004;
310 Giannakopoulos et al., 2009). In these two studies, technological improvements and adaptation were
311 however limited or not considered. In contrast to the studies mentioned in Table 3, we assumed cropland
312 productivity will only increase in intensive rain-fed and irrigated systems, and not overall in the cropland
313 sector. We did not improve the productivity of existing extensive cropland or traditional multifunctional
314 mosaic systems. Moreover, we did not consider potential increases to water demands of different crops.
315 Studies have demonstrated that improving the irrigation efficiency can help maintaining cropland
316 productivity and allow for expansion of irrigated areas in the Mediterranean (Fader et al., 2016; Malek &
317 Verburg 2017b; Saadi et al., 2015).

318 **4 Results**

319 4.1 Changes to management of Mediterranean land systems

320 In both scenarios, extensive cropland systems decreased most (Table 4). This is either through
321 intensification of management, or due to abandonment and subsequent conversion to woodlands. There
322 is also a drastic increase in urban systems in the Middle Eastern region, particularly Syria, Israel,
323 Palestine and Jordan. The largest expansions of irrigated areas can be observed in Turkey, Tunisia and
324 Morocco. The inclusion of water limitation for crop irrigation in the sustainability scenario impacts the way
325 in which the demands for food are satisfied in comparison to the growth scenario (Fig. 4). In some parts
326 of North-West Africa and Turkey, limited freshwater availability for irrigation results in more rain-fed
327 intensive areas compared to the growth scenario (Fig. 5). In Algeria, the Western Balkans and Turkey,
328 limited water availability does not necessarily result in more rain-fed intensive area, as a significant part of
329 food demands is fulfilled by intensifying mosaics, higher efficiency improvement, while simultaneously the
330 overall food demand is lower due to a decrease in food waste. Land system results for both scenarios are
331 depicted in Fig. 4, with more detailed land system conversions shown in Table 4, Fig. 5 and Fig. 6,
332 Appendix J and downloadable in GIS format from www.environmentalgeography.nl. Fig. 7 presents the
333 two future scenarios in three focus areas, the Iberian peninsula, Middle East and Turkey, and in Tunisia.

334 4.2 Changes to traditional Mediterranean landscapes

335 More mosaic systems are preserved in the sustainability scenario: 60% more compared to the growth
336 scenario. Whereas a significant share of mosaic systems remain the same, still around 58% and 32% of
337 mosaics change in the growth and sustainability scenario respectively. Around 36% of mosaics in the
338 growth and 21% in the sustainability scenario were changed to other mosaic systems (Table 5). Based on
339 the level of multifunctionality, the mosaics either changed to another mosaic with the same level, or to a
340 mosaic with a higher level. The level of multifunctionality was identified based on the number of activities
341 defining the system. A change from a mosaic land system with 2 activities (e.g. woodland-rangeland
342 mosaic) to another mosaic with 2 activities (woodland-cropland mosaic) was identified as a change in

343 management on a same level of multifunctionality. The extent of mosaics, that change from another type
344 of mosaic while keeping the same level of multifunctionality is significant in both scenarios (Table 5).
345 Such processes were projected in southern Spain, Portugal, Western Balkans and south-western Turkey
346 (Fig. 6). An interesting process is the increase in functionality, occurring in both scenarios. This process
347 represents additional activities on existing land systems, such as introducing livestock to woodlands or
348 woodland-cropland mosaics. The model transformed land systems with lower output in terms of crops and
349 livestock such as woodlands, to land systems with higher output occurring in similar locations (similar
350 spatial preference) such as woodland-cropland or woodland-rangeland mosaics. These processes are
351 projected to occur on extensive areas in Algeria, Western Balkans and western Syria (Fig. 6). Differences
352 in the extent of losses of Mediterranean agro-silvo-pastoral mosaic systems between the two scenarios
353 are mostly a consequence of cropland expansion and intensification – more than twice as many mosaics
354 are transformed to intensive or irrigated cropland in the growth scenario compared to the sustainability
355 scenario (Table 5). In some areas, agro-silvo-pastoral mosaic systems expand. This is mostly caused by
356 an increase in livestock density in extensive cropland systems, or through the introduction of livestock to
357 cropland that was abandoned during the simulated time period. Increases in livestock density and grazing
358 on abandoned cropland were projected mostly in NW Africa, but also in southern Spain (Fig. 6 and 7).

359 4.3 Water resources

360 As the use of water resources was constrained only in the sustainability scenario, the overall lower water
361 extraction values are considerably lower than in the growth scenario. Clearly, the constraints on water
362 use have influenced the model's choice of land systems. In the growth scenario, all regions demonstrate
363 a significant increase of irrigation water withdrawal (Table 6). The highest expansions of irrigated
364 cropland systems and consequent increases in irrigation water withdrawal are projected in the sub-
365 regions Western Balkans and Turkey and NW Africa. We used the pressure on freshwater resources
366 index (PFR), to describe the water stress of the sub-regions in the two scenarios (Table 6). It is defined as
367 the share of total irrigation water withdrawal in available freshwater resources (FAO, 2016). There are
368 considerable differences between the regions in the baseline year (FAO, 2016) (Table 6). Both sub-
369 regions of the Northern Mediterranean have a PFR index that is below 20%, which still enables biological
370 functioning of freshwater bodies and does not result in water stress as a limiting development factor
371 (Arnell, 1999). This is mostly due to the more abundant freshwater resources in these regions and a lower
372 dependence of their agriculture on irrigation (Appendix B). The Middle East and NE Africa sub-region
373 already has unsustainable freshwater extraction rates, as some countries (e.g. Egypt and Libya) are
374 already extracting more resources compared to their renewable water resources (Table 6).

375 5 Discussion

376 5.1 Future changes of Mediterranean land systems

377 Modelling the future of land use and management intensity is a central part in integrated assessment
378 models (Wise *et al.*, 2009; Stehfest *et al.*, 2014). However, given the global character of these
379 assessments, and the strongly simplified land cover representations, little insight into the potential
380 changes in the specific Mediterranean land systems is acquired from these studies. While single case
381 studies have documented the (potential) response to global change in the Mediterranean (Bugalho *et al.*,
382 2011; Keenan *et al.*, 2011; Nainggolan *et al.*, 2012; Parcerisas *et al.*, 2012), no earlier study specifically
383 focused on the impacts of global change across the entire region and took into account the impacts of the
384 limited water availability. Land systems provide highly appreciated benefits for the regional population: the
385 region is a significant producer of highly demanded commodities, such as olives and grapes, and the
386 region hosts vast areas of traditional multifunctional mosaic landscapes, that are well studied (Blondel,
387 2006; Daccache *et al.*, 2014). The region is projected to witness significant climate changes, mostly in the
388 form of temperature increases, decreases of precipitation and more frequent climate extremes. At the
389 same time, demographic changes and fast urbanization will pose additional challenges to land systems.

390 The two fundamentally different scenarios we developed, demonstrate two potential pathways of how
391 land systems may respond to a growing population in the Mediterranean. While the future of land
392 management of the region is more likely to be between these two scenarios, the scenarios sketch how
393 differences in the level of technological development and the implementation of policies concerning rural
394 development, water management and biodiversity lead to strongly different land system outcomes. Both
395 scenarios represent a future, where the southern Mediterranean countries continue to depend on food
396 imports for a significant part of their food demand and are based on global integrated assessment
397 calculations of trade, demand and supply between regions. In all global level scenarios it is acknowledged
398 that to feed the growing population of the region, significant food imports will still be needed (Wright &
399 Cafiero, 2011). Our results indicate that already under those conditions enormous changes in land
400 systems are needed to meet such demands for production in the region. Moreover, when looking at the
401 changes between the two scenarios, we can identify adaptation opportunities in land management to
402 avoid changes to mosaic land systems and cropland expansion (Fig. 8). Significantly more areas are
403 subject to irrigation, intensification and changes to mosaics in the growth scenario compared to the
404 sustainability scenario, suggesting the potential outcomes of successfully following common sustainable
405 development goals. Technological improvements and nutrient management, improved irrigation
406 efficiency, and protection of traditional land use systems were thus recognized as successful measures to
407 increase the resilience of traditional Mediterranean landscapes. Nevertheless, the sustainability scenario
408 also projected changes to intensity and irrigation differently from the growth scenario. These locations
409 mostly present the tradeoffs of expanding the protected areas network (Fig. 8).

410 Despite a lower demand for agricultural products in the sustainability scenario, more land system
411 transformations to intensive rain-fed cropland were projected for some areas, notably in NW Africa. In this
412 region rain-fed intensive areas provide much less output compared to other regions and are characterized

413 by high yield gaps (Mueller *et al.*, 2012). The lower use of irrigated cropland to meet agricultural demands
414 therefore comes at the cost of a larger expansion of rain-fed croplands in this scenario. The intensification
415 of rain-fed cropland was mostly projected in areas that will maintain more favorable climatic conditions in
416 the future, such as northeast Spain. Other locations of intensifying rain-fed cropland (e.g. north of the
417 Atlas mountains) correspond well with other research (van Asselen & Verburg, 2012; Mueller *et al.*,
418 2012).

419 The model projected significant increases in irrigated cropland in both scenarios, because irrigated
420 systems in most sub-regions have the highest output of crops, and were thus promoted by the model to
421 fulfil agricultural demands. Equipping rain-fed cropland with irrigation, particularly in semi-arid regions as
422 the Mediterranean, is among the most implemented adaptation options to reduce risks to climate change
423 (Smit & Skinner, 2002). Due to projected climate change, the extent of areas most suitable for rain-fed
424 intensive cropland systems decreased. A major limiting factor for rain-fed agriculture is high aridity, and
425 arid and hyper-arid areas are projected to expand in the NW Africa sub-region (Appendix H). Although
426 necessary improvements in irrigation efficiency to maintain current water withdrawal with projected
427 expansion of irrigated areas are possible (Fader *et al.*, 2016), the two southern Mediterranean sub-
428 regions already today have unsustainable water withdrawal levels. Furthermore, the growth scenario
429 does not consider projected decreases in water resources as may be expected due to depletion of
430 aquifers and climate change (Vörösmarty *et al.*, 2010; Chenoweth *et al.*, 2011). Despite the efforts on
431 finding new water resources, water reuse and desalination, the growth scenario is strongly overestimating
432 the water availability.

433 The drastic expansion of irrigated areas projected in the sub-regions Western Balkans and Turkey, and
434 NW Africa is expected under the 'growth' scenario. Particularly in NW Africa, irrigation is needed in order
435 to increase the yields, as studies suggest that only improving nutrient management and mechanization
436 will not be enough to improve cropland productivity in this area due to local climatic conditions (Mueller *et al.*
437 *et al.*, 2012). Generally, the sustainability scenario resulted in less intensive rain-fed and irrigated cropland
438 (Fig. 8), which can also be attributed to a 5% lower demand due to a decrease in food waste.
439 Nevertheless, our reduction in demand for agricultural products is conservative, as studies suggest higher
440 potentials to reduce agricultural demands in case of drastic improvements in the supply chain or diet
441 change (Parfitt *et al.*, 2010; Garrone *et al.*, 2014).

442 Expansion of multifunctional systems, projected by the model in both scenarios can be explained two-
443 fold. First, these areas are subject to expansion of protected areas in the sustainability scenario, which
444 prevents more intensive land systems to be established, but allows for the conversions into other (more)
445 traditional extensive systems. Secondly, climatic and soil characteristics of these areas constrain rain-fed
446 intensive agriculture. Similar transitions have already been observed at the local scale in areas such as
447 south-eastern Spain, as a result of environmental conditions, policies favoring woodland expansion and
448 less profitable rain-fed agriculture (Nainggolan *et al.*, 2012). This increase of multifunctional areas can be

449 defined as sustainable intensification, where satisfying future demands for crops and livestock occurs
450 simultaneously with meeting sustainability objectives (e.g. biodiversity protection and strengthening rural
451 communities) (Garnett *et al.*, 2013). Multifunctional land systems have been acknowledged as a
452 significant adaptation option to climate change, particularly for smallholder farmers (Verchot *et al.*, 2007).
453 Although multifunctional areas contribute less to satisfying food demand as compared to intensive
454 cropland, they can also be recognized as an effort to rehabilitate and conserve land and water resources.
455 Such increased land productivity (in terms of crop and livestock production) with simultaneous
456 sustainable land use follows the objectives of the United Nations Convention to Combat Desertification
457 (UNCCD, 1994).

458 This study took into account competing demands for food production and living space. Traditional
459 Mediterranean land systems are, however, providing other significant ecosystem services, such as non-
460 timber forest products like mushrooms or cork, fire prevention, carbon storage, soil erosion prevention
461 and landscape aesthetics (Bugalho *et al.*, 2011; Almagro *et al.*, 2013; Guerra *et al.*, 2016). These
462 services could also act as additional demands especially if covered by environmental, rural development
463 and tourism policies (Eitelberg *et al.*, 2016). Future studies should therefore study the effects of
464 maintaining different benefits provided by traditional Mediterranean mosaic systems besides food. This
465 way, the extent of mosaic systems needed to provide a certain extent of important ecosystem services
466 can be determined, as well as potential tradeoffs in irrigation and intensification in other areas.

467 5.2 Water limitation as a contribution to land change modelling

468 Significant improvements in the modelling of future land use have been made in the recent decades,
469 including more precise coverage of spatial, temporal and thematic resolution and moving beyond an
470 approach based only on dominant land cover (Hurtt *et al.*, 2011; Letourneau *et al.*, 2012; Souty *et al.*,
471 2012; Bryan *et al.*, 2016). Land management which used to be represented in a simplified manner as a
472 class of regional management factor (Bouwman *et al.*, 2006; Bondeau *et al.*, 2007), can now be
473 described using different management intensity metrics, such as livestock numbers, fertilizer inputs or
474 yield gaps (Souty *et al.*, 2012; van Asselen & Verburg, 2013). This is necessary, as often socio-economic
475 changes are not limited to direct land cover changes, but predominantly lead to changes in management
476 intensity or irrigation. In this study, we managed to quantify the relative changes required for both land
477 management and land cover to fulfill future food demands. Our results show that for the growth and
478 sustainability scenario respectively 61% and 51% of the increase of agricultural demand is met by new
479 irrigation, 23% and 36% by cropland intensification and only 12% each by cropland expansion. Our study
480 therefore confirms that the inclusion of changes in land management (irrigation and intensification) may
481 be more important than land cover changes in modeling of future scenarios. This way we contribute to
482 better understanding of land system processes leading to increased pressure on land and water
483 resources, and consequent land degradation (UNCCD, 1994).

484 The growth scenario presented a continuation of worldwide trends, where the demands for food and living
485 space are fulfilled by increasing more productive, monofunctional, land systems at the expense of
486 traditional systems, as is also suggested by global scale studies (van Asselen & Verburg, 2013; Eitelberg
487 *et al.*, 2016). In this scenario, the likelihood of spatial variables and location preferences for where
488 irrigated systems were considered in the allocation of land systems by the model. However, the
489 expansion of irrigated agriculture was not limited by water availability. Consequently, the growth scenario
490 resulted in more than twice as much irrigation water withdrawal in the sub-regions NW Africa and Western
491 Balkans and Turkey. An assumption of unlimited water availability can therefore lead to an overestimation
492 of expanding irrigated areas. Such increases of water extraction are unlikely to happen in the
493 Mediterranean basin, partly because of the regional expected impacts of climate change (Elliott *et al.*,
494 2014). On the other side, not constraining water availability is a useful scenario exercise as it
495 demonstrates the land systems distribution that might otherwise be possible (Fig. 8). Our results show the
496 necessity of including the reality of limited water availability when simulating future changes to land
497 systems, particularly in (semi)arid regions. Many global studies disregard this issue and suggest
498 significant cropland expansion and intensification in other semi-arid areas (van Asselen & Verburg, 2013;
499 Eitelberg *et al.*, 2016), which undoubtedly will have an impact on future water resources. The approach
500 presented in this paper can, therefore, be applied in areas that face similar resource constraints, and
501 improve the understanding of how future cropland expansion and intensification are responding to
502 situations of water stress.

503 In this study we used spatially explicit irrigation data, linked to irrigation water withdrawal and freshwater
504 resources statistics. This resulted in mean values per cell of irrigated land system, not considering areas
505 where water withdrawal values can be considerably higher due to higher potential evapotranspiration.
506 Incorporating water cycle processes is needed to improve the availability and constraints of water
507 resources. One example could be to use spatially explicit data on water availability and water extraction, if
508 such data was available on a more detailed resolution (Brauman *et al.*, 2016). A higher resolution of water
509 withdrawal of land systems might be achieved, by operating on a catchment scale. Nevertheless, data on
510 crop production is not available on the same scale, which would result in a mismatch of management
511 levels - water management on catchment scale vs. agricultural management on a regional or national
512 scale. Moreover, using spatially explicit water withdrawal data would mean additional uncertainties to our
513 approach, related to the models used to derive such data. Using regional or national scale irrigation
514 withdrawal data furthermore ensures a higher transferability of our approach to similar (semi)arid areas
515 with increasing food demand and high water stress (e.g. other areas in the Middle East, south Asia,
516 China, North America...). Another challenge would be to consider groundwater reserves, as these are
517 often non-renewable, or their exploitation exceeds groundwater recharge rates. Although there is data on
518 irrigation from groundwater resources, for the Southern Mediterranean it is based on national statistics
519 (Siebert *et al.*, 2010), which makes it difficult to limit their spatial extent and occurrence. Irrigation using

520 groundwater might also occur in the hyper-arid (desert) part in the Middle East and North Africa, outside
521 our study area (Mediterranean ecoregion).

522 We only focused on water withdrawals for agriculture and did not account for the competing demands for
523 municipal and industrial water use. Whereas in the European Union irrigation amounted for around 40%
524 of total water withdrawals, it is the main source of water withdrawals in other sub-regions (Appendix K).
525 Future socio-economic development particularly in the southern Mediterranean will however also likely
526 result in increased demands for non-agricultural water use (Flörke *et al.*, 2013). Livestock water use was
527 also not considered, mostly as the statistics in all sub-regions except the European Union equal irrigation
528 with agricultural water withdrawals (FAO, 2016). Nevertheless, livestock also has significant water
529 demands influencing freshwater resources (Mekonnen & Hoekstra, 2012). Further research on improving
530 the data on water use and how different sector compete for water resources is therefore needed.

531 5.3 Storyline to modeling translation and uncertainties

532 In this study, we present a transparent and clear methodology on translating storylines to modeling, as
533 demonstrated by detailed supplement information. Individual steps of our study are presented, ranging
534 from preparing dynamic input files for changing temperature and precipitation, to defining model
535 parameters reflecting the storyline. This way, we aimed at improving the presentation of the translation of
536 specific policy assumptions to model parameters.

537 In analyzing global change effects on local scale land management, we went beyond applying global
538 demand projections only. We developed two storylines that describe regional challenges on a higher
539 detail – global SSP storylines are more broad and general (Riahi *et al.*, 2016). For example, in our study
540 water management is one of the most crucial challenges for the Mediterranean, influencing the
541 development of scenarios significantly.

542 Model studies are sensitive to uncertainties and errors in the input data. We aimed to reduce the
543 uncertainties in our approach by relying mostly on crop production and irrigation water withdrawal
544 statistics. Nevertheless, combining numerous spatial data can result in an aggregation of uncertainties.
545 For example, our baseline land systems map is heavily dependent on inputs such as land cover, with the
546 southern Mediterranean having higher observed inaccuracies compared to the northern part (Malek &
547 Verburg, 2017a). Moreover, in this study we focused on the Mediterranean ecoregion. Irrigated systems
548 in deserts, such as oasis like date plantations in North Africa, are also contributing to total crop production
549 and also irrigation water withdrawals.

550 One of the biggest uncertainties for land system modelling are assumptions on improvements in
551 technology and efficiency. Agreement in efficiency improvements used in different models is usually
552 higher in developed regions, such as the European Union (Paillard *et al.*, 2014), whereas we observed
553 relatively large variations in other regions, such as the whole Southern Mediterranean and Turkey. This

554 could be linked to the fact that yield gaps are larger in developing countries than in developed countries
555 (Ramankutty *et al.*, 2008; Mueller *et al.*, 2012). Yield improvement scenarios are often optimistic, not
556 considering the expected impacts of climate change (Long, 2006). Our yield improvements were rather
557 conservative, considering the region's socio-economic and environmental characteristics (Mueller *et al.*,
558 2012). We did not apply efficiency improvements to all land systems as it is the case in some similar
559 studies (van Asselen & Verburg, 2013; Eitelberg *et al.*, 2016). For example, in NW Africa, improvements
560 in nutrient management are needed to achieve higher yields. Improved nutrient management on
561 extensive cropland could however also mean a transition to a more intensive cropland, resulting in higher
562 yields. Finally, we did not consider potential new crop production systems in the future. One example of
563 such system are greenhouses with significantly higher agricultural output. Such systems could occur on a
564 wider spatial extent, as they are less dependent on local environmental characteristics. Assumptions on
565 technological improvement, particularly increases of yield are significantly influencing the extent of
566 simulated cropland expansion, intensification and new irrigation and need to be considered carefully. To
567 reduce the uncertainties related to future cropland productivity, future research should focus on
568 differences between projected yields or cropland efficiency.

569 **6 Conclusions: what are the consequences of global change for the** 570 **Mediterranean?**

571 In this article, we assessed how global change might influence future land systems in the Mediterranean.
572 Similarly to global scale studies, we projected significant increases in intensively managed and irrigated
573 cropland for the Mediterranean basin. Our study shows, that to a certain extent, it is possible to preserve
574 traditional Mediterranean mosaic systems. The growth scenario depicts a future, where more mosaic
575 systems will be abandoned or transformed to more intensive systems. Rural development policies
576 focusing on improving the socio-economic conditions of rural areas, and increasing yields within
577 traditional mosaic systems, as shown in the sustainability scenario, can be an alternative to further
578 cropland expansion or conversion to monoculture intensive cropland systems. We have also shown, that
579 an expansion of protected areas in the region is possible without compromising the region's abilities to
580 produce food. The same goes for reducing the intensity of cropland and grazing activities in wetlands.

581 The two scenarios represent different pathways on managing Mediterranean freshwater resources and
582 dealing with water shortages. In the growth scenario, water resources are continued to being depleted at
583 unsustainable rates in the future with significant investments into alternative water resources. Some of
584 them are already taking place today: water reuse, desalination, large infrastructural projects such as
585 dams or channels (Hochstrat *et al.*, 2010; Pedrero *et al.*, 2010; El Gammal & Ali, 2011). Improving the
586 state of freshwater resources, as projected by the sustainability scenario, is possible by increasing the
587 efficiency of rain-fed cropland and strengthening the role of multifunctional mosaic systems.

588 Our results indicate, that increased food production in the Mediterranean can be accompanied by
589 preserving landscapes with higher cultural and biodiversity values, and decreasing the pressure on
590 freshwater resources. However, we also show that such a future is only possible under the
591 implementation of common Mediterranean sustainable development goals and orchestrated agricultural
592 and environmental management strategies.

593

594 **References**

595

596 Alexandratos N, Bruinsma J (2012) World agriculture towards 2030/2050: The 2012 revision. Rome: Food
597 and Agriculture Organization of the United Nations.

598 Almagro M, Vente J de, Boix-Fayos C et al. (2013) Sustainable land management practices as providers
599 of several ecosystem services under rainfed Mediterranean agroecosystems. *Mitigation and*
600 *Adaptation Strategies for Global Change*, 1–15.

601 Arnell N (1999) Climate change and global water resources. *Global Environmental Change*, **9**, S31–S49.

602 van Asselen S, Verburg PH (2012) A Land System representation for global assessments and land-use
603 modeling. *Global Change Biology*, **18**, 3125–3148.

604 van Asselen S, Verburg PH (2013) Land cover change or land-use intensification: simulating land system
605 change with a global-scale land change model. *Global Change Biology*, **19**, 3648–3667.

606 Ayache F, Thompson JR, Flower RJ, Boujarra A, Rouatbi F, Makina H (2009) Environmental
607 characteristics, landscape history and pressures on three coastal lagoons in the Southern
608 Mediterranean Region: Merja Zerga (Morocco), Ghar El Melh (Tunisia) and Lake Manzala
609 (Egypt). *Hydrobiologia*, **622**, 15–43.

610 Benoit G, Comeau A (2012) *A Sustainable Future for the Mediterranean: The Blue Plan's Environment*
611 *and Development Outlook*. Routledge, 465 pp.

612 Bernués A, Ruiz R, Olaizola A, Villalba D, Casasús I (2011) Sustainability of pasture-based livestock
613 farming systems in the European Mediterranean context: Synergies and trade-offs. *Livestock*
614 *Science*, **139**, 44–57.

615 Blondel J, Aronson J, Bodiou J-Y, Boeuf G (eds.) (2010) *The Mediterranean region: biological diversity in*
616 *space and time*, 2nd ed edn. Oxford University Press, Oxford ; New York, 376 pp.

617 Blondel J (2006) The “Design” of Mediterranean Landscapes: A Millennial Story of Humans and
618 Ecological Systems during the Historic Period. *Human Ecology*, **34**, 713–729.

619 Bondeau A, Smith PC, Zaehle S et al. (2007) Modelling the role of agriculture for the 20th century global
620 terrestrial carbon balance. *Global Change Biology*, **13**, 679–706.

621 Bouwman A, Kram T, Klein Goldewijk K (2006) *Integrated Modelling of Global Environmental Change. An*
622 *Overview of IMAGE 2.4*. Netherlands Environmental Assessment Agency, Bilthoven.

623 Brauman KA, Richter BD, Postel S, Malsy M, Flörke M (2016) Water depletion: An improved metric for
624 incorporating seasonal and dry-year water scarcity into water risk assessments. *Elementa:*
625 *Science of the Anthropocene*, **4**, 000083.

626 Bryan BA, Nolan M, McKellar L et al. (2016) Land-use and sustainability under intersecting global change
627 and domestic policy scenarios: Trajectories for Australia to 2050. *Global Environmental Change*,
628 **38**, 130–152.

629 Bugalho MN, Caldeira MC, Pereira JS, Aronson J, Pausas JG (2011) Mediterranean cork oak savannas
630 require human use to sustain biodiversity and ecosystem services. *Frontiers in Ecology and the*
631 *Environment*, **9**, 278–286.

632 CAPMAS (2015) *Egypt in figures*. Central Agency for Public Mobilization and Statistics. Available from:
633 [http://www.msrintranet.capmas.gov.eg/pdf/EgyptinFigures2015/EgyptinFigures/Tables/PDF/1-](http://www.msrintranet.capmas.gov.eg/pdf/EgyptinFigures2015/EgyptinFigures/Tables/PDF/1-%20%D8%A7%D9%84%D8%B3%D9%83%D8%A7%D9%86/pop.pdf)
634 [%20%D8%A7%D9%84%D8%B3%D9%83%D8%A7%D9%86/pop.pdf](http://www.msrintranet.capmas.gov.eg/pdf/EgyptinFigures2015/EgyptinFigures/Tables/PDF/1-%20%D8%A7%D9%84%D8%B3%D9%83%D8%A7%D9%86/pop.pdf) (Accessed on: June 12,
635 2017)

636 Cerdan O, Govers G, Le Bissonnais Y, Van Oost K, Poesen J, Saby N, et al. (2010) Rates and spatial
637 variations of soil erosion in Europe: A study based on erosion plot data. *Geomorphology*, **122**(1–
638 2), 167–177.

639 Chenoweth J, Hadjinicolaou P, Bruggeman A et al. (2011) Impact of climate change on the water
640 resources of the eastern Mediterranean and Middle East region: Modeled 21st century changes
641 and implications. *Water Resources Research*, **47**, W06506.

642 CIESIN (2015) Gridded Population of the World, Version 4. Center for International Earth Science
643 Information Network - CIESIN - Columbia University. NASA Socioeconomic Data and
644 Applications Center (SEDAC). Available from:
645 <http://sedac.ciesin.columbia.edu/data/collection/gpw-v4> (Accessed on: December 12, 2016)

646 CIESIN, IFPRI, CIAT (2011) Global Rural-Urban Mapping Project, Version 1: Population Density Grid.
647 Center for International Earth Science Information Network - CIESIN - Columbia University,
648 International Food Policy Research Institute - IFPRI, The World Bank, and Centro Internacional
649 de Agricultura Tropical - CIAT. Available from:
650 <http://sedac.ciesin.columbia.edu/data/collection/grump-v1> (Accessed on: December 12, 2016)

651 Cioffi A, dell’Aquila C (2004) The effects of trade policies for fresh fruit and vegetables of the European
652 Union. *Food Policy*, **29**, 169–185.

653 Cuttelod A, Garcia N, Malak DA, Temple HJ, Katariya V (2009) The Mediterranean: a biodiversity hotspot
654 under threat. In: *Wildlife in a Changing World: An Analysis of the 2008 IUCN Red List of*
655 *Threatened Species* (eds Vie J-C, Hilton-Taylor C, Stuart SN). International Union for
656 Conservation of Nature, Gland, Switzerland.

657 Daccache A, Ciurana JS, Diaz JAR, Knox JW (2014) Water and energy footprint of irrigated agriculture in
658 the Mediterranean region. *Environmental Research Letters*, **9**, 124014.

659 Eitelberg DA, van Vliet J, Doelman JC, Stehfest E, Verburg PH (2016) Demand for biodiversity protection
660 and carbon storage as drivers of global land change scenarios. *Global Environmental Change*,
661 **40**, 101–111.

662 El Gammal HAA, Ali HMM (2011) Commissioning of abandoned drainage water reuse systems in Egypt:
663 A case study of upgrading the Umoum project, Nile Delta. *Irrigation and Drainage*, **60**, 115–122.

664 Elliott J, Deryng D, Müller C et al. (2014) Constraints and potentials of future irrigation water availability on
665 agricultural production under climate change. *Proceedings of the National Academy of Sciences*,
666 **111**, 3239–3244.

667 Ellis EC, Klein Goldewijk K, Siebert S, Lightman D, Ramankutty N (2010) Anthropogenic transformation of
668 the biomes, 1700 to 2000: Anthropogenic transformation of the biomes. *Global Ecology and*
669 *Biogeography*, no-no.

670 European Commission (ed.) (2011) *EuroMed-2030: Long term challenges for the Mediterranean area.*
671 *Report of an expert group.* Publications Office of the European Union, Luxembourg, 139 pp.

672 EUROSTAT (2013) *Pocketbook on Euro-Mediterranean statistics.* Publications office of the European
673 Union, Luxembourg.

674 EUROSTAT (2016a) *Agriculture statistics - North Africa and Eastern Mediterranean.* Available from:
675 [http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_-](http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_-_North_Africa_and_Eastern_Mediterranean)
676 [_North_Africa_and_Eastern_Mediterranean](http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_-_North_Africa_and_Eastern_Mediterranean) (Accessed on: June 12, 2017)

677 EUROSTAT (2016b) Agriculture statistics at regional level. March 2016. Available from:
678 http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_at_regional_level
679 (Accessed on: June 12, 2017)

680 EUROSTAT (2016c) *Population statistics at regional level.* March 2016. Available from:
681 http://ec.europa.eu/eurostat/statistics-explained/index.php/Population_statistics_at_regional_level
682 (Accessed on: June 12, 2017)

683 Evans JP (2008) 21st century climate change in the Middle East. *Climatic Change*, **92**, 417–432.

684 Fader M, Shi S, von Bloh W, Bondeau A, Cramer W (2016) Mediterranean irrigation under climate
685 change: more efficient irrigation needed to compensate increases in irrigation water
686 requirements. *Hydrology and Earth System Sciences*, **20**, 953–973.

687 FAO (2013) *State of Mediterranean forests 2013.* FAO, Rome.

688 FAO (2016) AQUASTAT website. FAO's Information System on Water and Agriculture. Food and
689 Agriculture Organization of the United Nations. Available from:
690 <http://www.fao.org/nr/water/aquastat/> (Accessed on: June 12, 2017)

691 Flörke M, Kynast E, Bärlund I, Eisner S, Wimmer F, Alcamo J (2013) Domestic and industrial water uses
692 of the past 60 years as a mirror of socio-economic development: A global simulation study.
693 *Global Environmental Change*, **23**, 144–156.

694 Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, et al. (2011) Solutions for a
695 cultivated planet. *Nature*, **478**(7369), 337–342.

696 Fricko O, Havlik P, Rogelj J et al. (2017) The marker quantification of the Shared Socioeconomic Pathway
697 2: A middle-of-the-road scenario for the 21st century. *Global Environmental Change*, **42**, 251-267.

698 Galewski T, Collen B, McRae L, Loh J, Grillas P, Gauthier-Clerc M, Devictor V (2011) Long-term trends in
699 the abundance of Mediterranean wetland vertebrates: From global recovery to localized declines.
700 *Biological Conservation*, **144**, 1392–1399.

701 García Martínez M, Poole N (2004) The development of private fresh produce safety standards:
702 implications for developing Mediterranean exporting countries. *Food Policy*, **29**, 229–255.

703 García-Ruiz JM, López-Moreno JI, Vicente-Serrano SM, Lasanta-Martínez T, Beguería S (2011)
704 Mediterranean water resources in a global change scenario. *Earth-Science Reviews*, **105**, 121–
705 139.

706 Garnett T, Appleby MC, Balmford A et al. (2013) Sustainable Intensification in Agriculture: Premises and
707 Policies. *Science*, **341**, 33–34.

708 Garrone P, Melacini M, Perego A (2014) Opening the black box of food waste reduction. *Food Policy*, **46**,
709 129–139.

710 Giannakopoulos C, Le Sager P, Bindi M, Moriondo M, Kostopoulou E, Goodess CM (2009) Climatic
711 changes and associated impacts in the Mediterranean resulting from a 2 °C global warming.
712 *Global and Planetary Change*, **68**, 209–224.

713 Giorgi F, Lionello P (2008) Climate change projections for the Mediterranean region. *Global and*
714 *Planetary Change*, **63**, 90–104.

715 Gleick PH (2014) Water, Drought, Climate Change, and Conflict in Syria. *Weather, Climate, and Society*,
716 **6**, 331–340.

717 GRAIN (2012) Land Grab Deals Dataset. Available from: [www.grain.org/article/entries/4479-grain-](http://www.grain.org/article/entries/4479-grain-releases-data-set-with-over-400-global-land-grabs)
718 [releases-data-set-with-over-400-global-land-grabs](http://www.grain.org/article/entries/4479-grain-releases-data-set-with-over-400-global-land-grabs) (Accessed on: June 12, 2017)

719 Guerra CA, Maes J, Geijzendorffer I, Metzger MJ (2016) An assessment of soil erosion prevention by
720 vegetation in Mediterranean Europe: Current trends of ecosystem service provision. *Ecological*
721 *Indicators*, **60**, 213–222.

722 Guiot J, Cramer W (2016) Climate change: The 2015 Paris Agreement thresholds and Mediterranean
723 basin ecosystems. *Science*, **354**, 465–468.

724 Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate
725 surfaces for global land areas. *International Journal of Climatology*, **25**, 1965–1978.

726 Hochstrat R, Wintgens T, Kazner C, Melin T, Gebel J (2010) Options for water scarcity and drought
727 management—the role of desalination. *Desalination and Water Treatment*, **18**, 96–102.

728 Houdret A (2012) The Water Connection: Irrigation, Water Grabbing and Politics in Southern Morocco.
729 *Water Alternatives*, **5**, 284–303.

730 Houérou HNL (1993) Salt-tolerant plants for the arid regions of the Mediterranean isoclimatic zone. In:
731 *Towards the rational use of high salinity tolerant plants* (eds Lieth H, Masoom AAA), pp. 403–422.
732 Springer Netherlands.

733 Hurtt GC, Chini LP, Frohking S et al. (2011) Harmonization of land-use scenarios for the period 1500–
734 2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting
735 secondary lands. *Climatic Change*, **109**, 117–161.

736 IAASTD (2009). Agriculture at a crossroads. Global report of the International Assessment of Agricultural
737 Knowledge, Science, and Technology for Development. Island Press, Washington, DC.

738 IIASA (2016) SSP Database (Shared Socioeconomic Pathways) - Version 1.1. Available from:
739 <https://tntcat.iiasa.ac.at/SspDb> (Accessed on: August 24, 2016)

740 Jiang L, O'Neill BC (2015) Global urbanization projections for the Shared Socioeconomic Pathways.
741 *Global Environmental Change*.

742 Joffre R, Rambal S, Ratte JP (1999) The dehesa system of southern Spain and Portugal as a natural
743 ecosystem mimic. *Agroforestry Systems*, **45**, 57–79.

744 Karamesouti M, Detsis V, Kounalaki A, Vasiliou P, Salvati L, Kosmas C (2015) Land-use and land
745 degradation processes affecting soil resources: Evidence from a traditional Mediterranean
746 cropland (Greece). *CATENA*, **132**, 45–55.

747 Kc S, Lutz W (2014) The human core of the shared socioeconomic pathways: Population scenarios by
748 age, sex and level of education for all countries to 2100. *Global Environmental Change*.

749 Keenan T, Maria Serra J, Lloret F, Ninyerola M, Sabate S (2011) Predicting the future of forests in the
750 Mediterranean under climate change, with niche- and process-based models: CO2 matters!
751 *Global Change Biology*, **17**, 565–579.

752 Kelley CP, Mohtadi S, Cane MA, Seager R, Kushnir Y (2015) Climate change in the Fertile Crescent and
753 implications of the recent Syrian drought. *Proceedings of the National Academy of Sciences*,
754 **112**(11): 3241–6.

755 Larson BA, Nicolaides E, Al Zu'bi B et al. (2002) The Impact of Environmental Regulations on Exports:
756 Case Study Results from Cyprus, Jordan, Morocco, Syria, Tunisia, and Turkey. *World*
757 *Development*, **30**, 1057–1072.

758 Letourneau A, Verburg PH, Stehfest E (2012) A land-use systems approach to represent land-use
759 dynamics at continental and global scales. *Environmental Modelling & Software*, **33**, 61–79.

760 Long SP (2006) Food for Thought: Lower-Than-Expected Crop Yield Stimulation with Rising CO2
761 Concentrations. *Science*, **312**, 1918–1921.

762 Lotze-Campen H, Popp A, Beringer T, Müller C, Bondeau A, Rost S, Lucht W (2010) Scenarios of global
763 bioenergy production: The trade-offs between agricultural expansion, intensification and trade.
764 *Ecological Modelling*, **221**, 2188–2196.

765 Malek Ž, Verburg PH (2017a) Mediterranean land systems: representing diversity and intensity of
766 complex land systems in a dynamic region. *Landscape and Urban Planning*, **165**, 102–116.

767 Malek Ž, Verburg PH (2017b) Adaptation of land management in the Mediterranean under scenarios of
768 irrigation water use and availability. *Mitigation and Adaptation Strategies for Global Change*.

769 McAdam JH, Burgess PJ, Graves AR, Riguero-Rodriguez A, Mosquera-Losada MR (2008) Classifications
770 and Functions of Agroforestry Systems in Europe. In: *Agroforestry in Europe: Current Status and*
771 *Future Prospects* (eds Rodríguez AR, McAdam J, Mosquera-Losada MR). Springer Science &
772 Business Media.

773 MEA (2005). Millennium ecosystem assessment. Ecosystems and human wellbeing: a framework for
774 assessment Washington, DC: Island Press.

775 Médail F, Quézel P (1999) Biodiversity Hotspots in the Mediterranean Basin: Setting Global Conservation
776 Priorities. *Conservation Biology*, **13**, 1510–1513.

777 Mediterranean 2030 Consortium (2011) *Tomorrow, the Mediterranean. Scenarios and projections for*
778 *2030*. IPEMED, Paris.

779 Mekonnen MM, Hoekstra AY (2012) A Global Assessment of the Water Footprint of Farm Animal
780 Products. *Ecosystems*, **15**, 401–415.

781 Mohamed NN (2016) Land Degradation in the Nile Delta. In Negm AM (Ed.), *The Nile Delta* (Vol. 55, pp.
782 235–264). Cham: Springer International Publishing.

783 Mueller ND, Gerber JS, Johnston M, Ray DK, Ramankutty N, Foley JA (2012) Closing yield gaps through
784 nutrient and water management. *Nature*, **490**, 254–257.

785 Munoz F (2003) Lock living: Urban sprawl in Mediterranean cities. *Cities*, **20**, 381–385.

786 MWO (2012) *Mediterranean Wetlands: Outlook. First Mediterranean Wetlands Observatory report*.
787 Mediterranean Wetlands Observatory, Tour du Valat, France.

788 Nainggolan D, de Vente J, Boix-Fayos C, Termansen M, Hubacek K, Reed MS (2012) Afforestation,
789 agricultural abandonment and intensification: Competing trajectories in semi-arid Mediterranean
790 agro-ecosystems. *Agriculture, Ecosystems & Environment*, **159**, 90–104.

791 NRC (2014) *Advancing Land Change Modeling: Opportunities and Research Requirements. National*
792 *Research Council. Committee on Needs and Research Requirements for Land Change Modeling*.
793 National Academies Press, Washington, D.C.

794 Olson DM, Dinerstein E, Wikramanayake ED et al. (2001) Terrestrial Ecoregions of the World: A New
795 Map of Life on Earth A new global map of terrestrial ecoregions provides an innovative tool for
796 conserving biodiversity. *BioScience*, **51**, 933–938.

797 Paillard S, Treyer S, Dorin B (eds.) (2014) *Agrimonde - scenarios and challenges for feeding the world in*
798 *2050*. Springer, Dordrecht ; New York, 250 pp.

799 Parcerisas L, Marull J, Pino J, Tello E, Coll F, Basnou C (2012) Land use changes, landscape ecology
800 and their socioeconomic driving forces in the Spanish Mediterranean coast (El Maresme County,
801 1850–2005). *Environmental Science & Policy*, **23**, 120–132.

802 Parfitt J, Barthel M, Macnaughton S (2010) Food waste within food supply chains: quantification and
803 potential for change to 2050. *Philosophical Transactions of the Royal Society B: Biological*
804 *Sciences*, **365**, 3065–3081.

805 Pedrero F, Kalavrouziotis I, Alarcón JJ, Koukoulakis P, Asano T (2010) Use of treated municipal
806 wastewater in irrigated agriculture—Review of some practices in Spain and Greece. *Agricultural*
807 *Water Management*, **97**, 1233–1241.

808 Pfeiffer T, Koutantou A (2015) Greece debt crisis: Olive oil supplies threatened as banking system grinds
809 to a halt. *The Independent*.

810 Pouzols FM, Toivonen T, Di Minin E et al. (2014) Global protected area expansion is compromised by
811 projected land-use and parochialism. *Nature*, **516**, 383–386.

812 Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., et al. (2017) Land-use futures
813 in the shared socio-economic pathways. *Global Environmental Change*, **42**, 331-345.

814 Ramankutty N, Evan AT, Monfreda C, Foley JA (2008) Farming the planet: 1. Geographic distribution of
815 global agricultural lands in the year 2000: Global Agricultural Lands in 2000. *Global*
816 *Biogeochemical Cycles*, **22**.

817 de Rancourt M, Fois N, Lavín MP, Tchakerian E, Vallerand F (2006) Mediterranean sheep and goats
818 production: An uncertain future. *Small Ruminant Research*, **62**, 167–179.

819 Riahi K, van Vuuren DP, Kriegler E et al. (2016) The Shared Socioeconomic Pathways and their energy,
820 land use, and greenhouse gas emissions implications: An overview. *Global Environmental*
821 *Change*.

822 Robinson TP, Wint GRW, Conchedda G et al. (2014) Mapping the Global Distribution of Livestock. *PLoS*
823 *ONE*, **9**, e96084.

824 Ronzon, T (2014). Food Crop Yields in 2050, in: Paillard, S., Treyer, S., Dorin, B. (Eds.), *Agrimonde –*
825 *Scenarios and Challenges for Feeding the World in 2050*. Springer Netherlands, pp. 101–122.

826 Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizzigalli C, Lionello P (2015). Climate change and
827 Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation
828 requirements and yield. *Agric. Water Manag.* **147**, 103–115.

829 Salvati L, Munafo M, Morelli VG, Sabbi A (2012) Low-density settlements and land use changes in a
830 Mediterranean urban region. *Landscape and Urban Planning*, **105**, 43–52.

831 Schmitz C, Biewald A, Lotze-Campen H et al. (2012) Trading more food: Implications for land use,
832 greenhouse gas emissions, and the food system. *Global Environmental Change*, **22**, 189–209.

833 Siebert S, Döll P, Hoogeveen J, Faures J-M, Frenken K, Feick S (2005) Development and validation of
834 the global map of irrigation areas. *Hydrology and Earth System Sciences*, **9**, 535–547.

835 Siebert S, Burke J, Faures JM, Frenken K, Hoogeveen J, Döll P, Portmann FT (2010) Groundwater use
836 for irrigation – a global inventory. *Hydrology and Earth System Sciences*, **14**, 1863–1880.

837 Siebert S, Henrich V, Frenken K, Burke J (2013) Update of the digital global map of irrigation areas to
838 version 5. Rheinische Friedrich-Wilhelms-Universität, Bonn, Germany and FAO, Rome, Italy,
839 170pp.

840 Smit B, Skinner, MW (2002) Adaptation options in agriculture to climate change: a typology. *Mitigation*
841 *and Adaptation Strategies for Global Change*, **7**(1), 85–114.

842 Souty F, Brunelle T, Dumas P et al. (2012) The Nexus Land-Use model version 1.0, an approach
843 articulating biophysical potentials and economic dynamics to model competition for land-use.
844 *Geoscientific Model Development*, **5**, 1297–1322.

845 Sowers J, Vengosh A, Weinthal E (2010) Climate change, water resources, and the politics of adaptation
846 in the Middle East and North Africa. *Climatic Change*, **104**, 599–627.

847 Stehfest E, Vuuren DP van, Kram T et al. (2014) *Integrated Assessment of Global Environmental Change*
848 *with IMAGE 3.0 – Model Description and Policy Applications*. PBL Netherlands Environmental
849 Assessment Agency, the Hague.

850 Steinfeld H, Wassenaar T, Jutzi S (2006) Livestock production systems in developing countries: status,
851 drivers, trends. *Revue scientifique et technique*, **25**, 505–516.

852 Taylor KE, Stouffer RJ, Meehl GA (2012) An Overview of CMIP5 and the Experiment Design. *Bulletin of*
853 *the American Meteorological Society*, **93**, 485–498.

854 Thomson AM, Calvin KV, Smith SJ, Kyle GP, Volke A, Patel P, et al. (2011) RCP4.5: a pathway for
855 stabilization of radiative forcing by 2100. *Climatic Change*, **109**(1):77.

856 Tieskens KF, Schulp CJE, Levers C, Lieskovský J, Kuemmerle T, Plieninger T, Verburg PH (2017)
857 Characterizing European cultural landscapes: Accounting for structure, management intensity
858 and value of agricultural and forest landscapes. *Land Use Policy*, **62**, 29–39.

859 Trabucco A, Zomer RJ, Bossio DA, van Straaten O, Verchot LV (2008) Climate change mitigation through
860 afforestation/reforestation: A global analysis of hydrologic impacts with four case studies.
861 *Agriculture, Ecosystems & Environment*, **126**, 81–97.

862 Turner BL, Janetos AC, Verburg PH, Murray AT (2013) Land system architecture: Using land systems to
863 adapt and mitigate global environmental change. *Global Environmental Change*, **23**, 395–397.

864 UNCCD (1994) United Nations Convention to Combat Desertification. Convention text. Bonn. Available
865 from [http://www2.unccd.int/sites/default/files/relevant-links/2017-](http://www2.unccd.int/sites/default/files/relevant-links/2017-01/UNCCD_Convention_ENG_0.pdf)
866 [01/UNCCD_Convention_ENG_0.pdf](http://www2.unccd.int/sites/default/files/relevant-links/2017-01/UNCCD_Convention_ENG_0.pdf) (Accessed on: June 12, 2017)

867 Verchot LV, Van Noordwijk M, Kandji S, Tomich T, Ong C, Albrecht A, et al. (2007) Climate change:
868 linking adaptation and mitigation through agroforestry. *Mitigation and Adaptation Strategies for*
869 *Global Change*, **12**(5), 901–918.

870 Vörösmarty CJ, McIntyre PB, Gessner MO et al. (2010) Global threats to human water security and river
871 biodiversity. *Nature*, **467**, 555–561.

872 Wise M, Calvin K, Thomson A et al. (2009) Implications of Limiting CO₂ Concentrations for Land Use and
873 Energy. *Science*, **324**, 1183–1186.

874 Wright B, Cafiero C (2011) Grain reserves and food security in the Middle East and North Africa. *Food*
875 *Security*, **3**, 61–76.

876 World Bank (2009) Improving Food Security in Arab Countries. Washington, DC: The World Bank,
877 International Fund for Agricultural Development (IFAD), Food and Agriculture Organization
878 (FAO).

879 You L, Wood-Sichra U, Fritz S, Guo Z, See L, Koo L (2014) Spatial Production Allocation Model (SPAM)
880 2005 v2.0. Available from <http://mapspam.info> (Accessed on: June 12, 2017)

881 Zamora J, Verdú JR, Galante E (2007) Species richness in Mediterranean agroecosystems: Spatial and
882 temporal analysis for biodiversity conservation. *Biological Conservation*, **134**, 113–121.

883 Zomer RJ, Trabucco A, Bossio DA, Verchot LV (2008) Climate change mitigation: A spatial analysis of
884 global land suitability for clean development mechanism afforestation and reforestation.
885 *Agriculture, Ecosystems & Environment*, **126**, 67–80.
886

887 **List of tables**

888 **Table 1.** Demands for annual and permanent crops, livestock and built-up areas for the Mediterranean
889 region for 2010 (EUROSTAT, 2013; EUROSTAT, 2016a, 2016b), and 2050 (modified from Fricko *et al.*,
890 2016; Riahi *et al.*, 2016; Popp *et al.*, 2017).

891 **Table 2.** Summary of main storyline elements of the two scenarios

892 **Table 3.** Future changes to cropland productivity (in % per year), together with considered climate
893 change effects and future technological improvements in this study and comparable studies. EU:
894 European Union, WBTU: Western Balkans and Turkey, MENA: Middle East and North Africa, NWA:
895 Northwest Africa

896 **Table 4.** Changes to spatial extent of Mediterranean land systems in %

897 **Table 5.** Changes to agro-silvo-pastoral mosaic system until 2050

898 **Table 6.** Irrigation water withdrawals and pressure on freshwater resources (PFR) in the Mediterranean in
899 2010 and 2050. Irrigation water withdrawals and freshwater resources and are based on national and
900 subnational statistics (EUROSTAT, 2013, 2016a, 2016b; FAO, 2016).

901

902 **Table 1.** Demands for annual and permanent crops, livestock and built-up areas for the Mediterranean
 903 region for 2010 (EUROSTAT, 2013; EUROSTAT, 2016a, 2016b), and 2050 (modified from Fricko *et al.*,
 904 2016; Riahi *et al.*, 2016; Popp *et al.*, 2017).

Region	2010		Growth 2050		Sustainability 2050	
	North	South	North	South	North	South
Annual crops (10 ⁶ t)	138.23	78.62	163.53	110.61	153.19	101.89
Permanent crops (10 ⁶ t)	68.40	21.73	80.88	30.56	78.33	30.12
Livestock (10 ⁶ nr)	22.44	13.66	28.23	16.92	26.33	15.99
Built-up areas (10 ³ km ²)	21.52	12.63	25.07	18.71	23.72	16.23

905

906

907 **Table 2.** Summary of main storyline elements of the two scenarios

	Growth	Sustainability
Population change and urban expansion		
Population in 2050	19.5% increase in the North, 47.4% increase in the South (SSP2)	
Demand for built up areas	10% higher than annual population growth rate	30% lower than annual population growth rate
Spatial pattern	Urban sprawl allowed, urban land has priority over all other uses. No expansion in protected areas	Compact and denser urban areas promoted, no expansion in protected areas
Agriculture and food production		
Food demand	Projected SSP2 marker scenario food production	10% lower annual crops demand growth. 10% higher permanent crops growth due to easier exports. Reduction in food waste resulting in total 5% lower food demand.
Yields	Irrigated systems achieve 75% of the attainable yield	Northern Mediterranean: closed yield gap for rain-fed intensive and irrigated systems Southern Mediterranean: extensive systems reach 50%, rain-fed intensive 75% and irrigated 90% of the attainable yield.
Livestock	15% efficiency improvement to landless livestock systems	5% livestock efficiency improvement to all systems
Access to irrigation	Same as baseline, 5% lower spatial preference for areas with low market accessibility	Improved accessibility
Biodiversity		
Protected areas	Only existing PAs	17% of national priority areas designated as PAs, transformations to high-intensity land systems in such areas are not possible
Wetland management	No changes	Intensity in cropland and livestock

Grazing in arid areas	No limitations	reduced by 30%
<hr/>		
Climate change and water		
<hr/>		
Climate change scenario	RCP 4.5	RCP 4.5
Location specific deduction to rain-fed systems	reduced spatial preference in areas with aridity index < 0.5 by 0.1	reduced spatial preference in areas with aridity index < 0.5 by 0.05
Water resources	No limitations and changes to water withdrawal	Limited water withdrawal. Withdrawal reduced by 25%
Irrigation efficiency	N.A.	35% more efficient

908

909 **Table 3.** Future changes to cropland productivity (in % per year), together with considered climate change effects and future technological
 910 improvements in this study and comparable studies. EU: European Union, WBTU: Western Balkans and Turkey, MENA: Middle East and North
 911 Africa, NWA: Northwest Africa

	EU	WBTU	MENA	NWA	Climate change	Technological improvements
Observed yields change (1961-2000)	1.92	0.34		2.49		
International Assessment of Agricultural Knowledge, Science and Technology for Development (IAASTD, 2009)						
High estimate	1.33	1.33		1.75	Crop responses to temperature and precipitation change, water stress (relatively small climate impacts in 2050)	Depend on investments in agricultural science and technology, and water productivity
Low estimate	0.71	0.71		0.79		
Millenium Ecosystem Assessment (MEA, 2005)						
High estimate	0.87	0.75		1.05	Crop responses to temperature and precipitation change, water stress	Increased fertilization and irrigation efficiency, major investments in agricultural research, GMOs, high mechanization level Insufficient investments in irrigation and cropland productivity, difficulties to maintain fertility of land
Low estimate	0.35	0.42		0.63		
Agrimonde (Ronzon, 2014)						
High estimate	0.81	2.22		0.67	Water stress, slower yield increase, increased variability of precipitation	Rapid technological improvements enable to overcome the impacts of climate change Rural development and ecological intensification to increase cropland productivity, irrigation techniques, water preservation
Low estimate	0	1.33		0.24		
Parry et al. (2004)						

High emissions	-0.59 to +0.04	0.08		-0.08	Crop responses to temperature and precipitation under current agricultural management	Limited: changes in planting dates, additional fertilization and irrigation on existing cropland
Low emissions	-0.17 to +0.17	0.04		-0.04		
Giannakopoulos et al. (2009)	-0.01 to +0.11	0.11 to 0.29	-0.27 to - 0.13	-0.1	Crop responses to temperature and precipitation, and seasonality	No improvements. If adaptation is implemented, cropland productivity can stay the same or increase with changing sowing dates and cycles and irrigation.
This study						
Sustainability – irrigated / int. rain-fed	0.11 / 0.05	0.33 / 0.16	0.56 / 0.43	0.94 / 0.43	Productivity of rain-fed cropland is limited by current climate (Mueller et al. 2012), future climate limits the spatial extent of rain-fed cropland	Investments both in rain-fed and irrigated systems, resulting in moderate productivity increase
Growth – irrigated / int. rain-fed	0 / 0	0.16 / 0	0.45 / 0	0.78 / 0.26		Investments focus on high output systems (irrigated only), low productivity increase

913 **Table 4.** Changes to spatial extent of Mediterranean land systems in %

Land system		Growth	Sustainability
Forest systems	medium intensity forest	-6.0	-12.4
	(semi)natural forest	24.8	22.0
	high intensity forest	-79.9	-43.1
Arid grazing systems	extensive arid system	-71.9	-18.2
	intensive arid system	84.1	16.6
Agro-silvo-pastoral mosaics	closed wooded rangeland	24.4	23.7
	open woodland	-92.1	-61.6
	open wooded rangeland	57.8	78.0
	cropland/wooded rangeland	-11.5	-34.9
	cropland/rangeland	29.0	63.1
Extensive rain-fed cropland	extensive annual	-82.9	-64.1
	extensive permanent	32.3	-53.3
	extensive mosaic	-70.8	-64.3
Intensive rain-fed cropland	rain-fed intensive annual	8.2	15.4
	rain-fed intensive permanent	41.6	-14.7
	rain-fed intensive mosaic	44.5	44.1
Irrigated cropland	irrigated annual	79.1	12.6
	irrigated permanent	26.0	38.5
	irrigated mosaic	-49.3	31.2
Settlements	peri-urban	10.2	2.1
	urban	72.2	44.7

914

915

916 **Table 5.** Changes to agro-silvo-pastoral mosaic system until 2050

Change process (%)	Growth	Sustainability
Persistent multifunctional systems	42.43	67.98
Multifunctionality loss towards monoculture	21.57	10.66
Similar level of multifunctionality, different management system	23.48	15.13
Increase of functionality within mosaic systems	12.52	5.86
Extensive cropland transformed to multifunctional systems	44.87	53.40

917

918

919 **Table 6.** Irrigation water withdrawals and pressure on freshwater resources (PFR) in the Mediterranean in
 920 2010 and 2050. Irrigation water withdrawals and freshwater resources and are based on national and
 921 subnational statistics (EUROSTAT, 2013, 2016a, 2016b; FAO, 2016).

	Sub-regions			
	W. Balkans and Turkey	EU	Middle East and NE Africa	NW Africa
Changes in irrigation water withdrawal compared to baseline levels (%)				
2050 – growth	+55.9	+37.0	+21.4	+59.9
2050 – sustainability	-18.0	-21.5	-27.9	-33.6
PFR (%)				
2010	11.8	10.2	94.4	30.2
2050 – growth	18.4	13.9	114.7	49.6
2050 – sustainability	9.7	8.0	68.1	20.0
Irrigation efficiency improvement to maintain current water extraction in growth scenario (%)	35.8	27.0	17.6	37.4

922

923

924 **Figure captions**

925 **Fig. 1** The studied Mediterranean ecoregion with its 4 sub-regions

926 **Fig. 2** CLUMondo land system change concept

927 **Fig. 3** Mediterranean land systems map (based on: Malek & Verburg, 2017a)

928 **Fig. 4** Future land systems in 2050 as simulated for the two scenarios: (a) growth, (b) sustainability. High
929 resolution version of the map is available in Appendix J.

930 **Fig. 5** Changes in land management intensity in the (a) growth and (b) sustainability scenario. High
931 resolution version of the map is available in Appendix J.


932 **Fig. 6** Land system change affecting mosaic land systems in the growth (a) and sustainability (b)
933 scenario. High resolution version of the map is available in Appendix J.

934 **Fig. 7** Future 2050 land systems scenarios in focus areas, (a) Spain and Portugal, (b) Middle East and
935 Turkey, (c) Tunisia

936 **Fig. 8** Future land management opportunities for the Mediterranean region, defined as spatially explicit
937 changes between the two scenarios, with the sustainability scenario as a reference. The two maps
938 present the consequences of implementing the policies of the sustainability scenario, described as
939 avoided and consequent changes to a) mosaic land systems and b) irrigation, intensification, cropland
940 expansion and abandonment. Values in brackets are in 1000 km².

941

942	List of appendices
943	Appendix A: List of consulted documents on Mediterranean future
944	Appendix B: Mediterranean land system characteristics per 4 km ² land system unit
945	Appendix C: Location factors used in calculating spatial preference maps using logistic regression
946	Appendix D: example of spatial preference maps
947	Appendix E: technical details of both scenarios
948	Appendix F: protected areas in the Mediterranean ecoregion
949	Appendix G: CMIP5 simulations of the RCP4.5 scenario used to update precipitation, temperature, PET
950	and AI maps for the year 2050 (average of 2041-2060)
951	Appendix H: PET and AI calculation and map example
952	Appendix I: Changes to urban population
953	Appendix J: Scenarios: high resolution figures
954	Appendix K: Freshwater resources and irrigation water withdrawal
955	
956	
957	




Demand scenarios	Annual crops	Built-up areas
	Permanent crops	Livestock

Crops	Living space	Supply
Irrigation demands	Livestock	

Limited water resources

Efficiency increase





Policy

Global change

Conversion rules	Conversion restrictions	Neighborhood effect
	Conversion resistance	Competitive advantage

Population	Market access	Spatial preference
Climate	Soil and terrain	



Arid systems

- ext. arid system
- int. arid system

Forests

- medium intensity forest
- (semi)natural forest
- high intensity forest

Agro-silvo-pastoral mosaics

- closed wooded rangeland
- open woodland
- open wooded rangeland
- cropland/wooded rangeland
- cropland/rangeland

Cropland systems


- ext. annual cropland
- ext. perm. cropland
- ext. ann-perm mosaic
- rain. int. annual
- rain. int. perm.
- rain int. ann-perm
- irr. annual
- irr. permanent
- irr. ann-perm

Settlements


- peri-urban
- urban


Other

- wetlands






0 250 500 750 1,000 km






0 250 500 750 1,000
km





0 250 500 750 1,000
km










■ mosaic persistence ■ multifunctionality increase
■ multifunctionality reduction ■ kept multifunct. different system




Growth oriented


Sustainability



a



b



Appendix A – List of consulted documents on Mediterranean future

Benoit G, Comeau A (2012) A Sustainable Future for the Mediterranean: The Blue Plan's Environment and Development Outlook. Routledge, 465 pp.

European Commission (ed.) (2011) EuroMed-2030: Long term challenges for the Mediterranean area. Report of an expert group. Publications Office of the European Union, Luxembourg, 139 pp.

FAO (2013) State of Mediterranean forests 2013. FAO, Rome.

Mediterranean 2030 Consortium (2011) Tomorrow, the Mediterranean. Scenarios and projections for 2030. IPEMED, Paris.

Paillard S, Treyer S, Dorin B (eds.) (2014) Agrimonde - scenarios and challenges for feeding the world in 2050. Springer, Dordrecht ; New York, 250 pp.

Sanna S, Le Tellier J (2013) Building on the Mediterranean scenario experiences. Cross-cutting approaches between regional foresight analysis and participatory prospective. Plan Bleu, Paris.

Appendix B – Mediterranean land system output calculation and characteristics

Land systems present an approach, where different data on land cover, management intensity, irrigation and livestock are combined. The initial land systems map of 2010 (Malek and Verburg, 2017) provides values for each land system as a combination of annual and permanent crops, livestock density and urban areas (see tables on next pages). Usually, land cover based simulations focus only on crop productions on cropland (as defined in a land cover product) and neglect agricultural activities in other areas, for example in peri-urban systems, and traditional mosaics. These areas contribute significantly to crop production in the Mediterranean ecoregion.

In this study, each land system in the region provides four services: annual and permanent crops, livestock, and urban areas. Their provision is based on the average values of each land system for cropland extent (%), permanent cropland extent (%), livestock density (livestock units in nr.) and urban extent (%). All four services are described as the average output of a land system unit (4 km²). The numbers deviate from reported statistics on yields, due to the fact, that every land system is a combination of different land use and land cover. In the Mediterranean, an average irrigated cropland system has between 42 (Turkey) to 68% (North-west Africa) of cropland. Irrigated land systems, present landscapes dominated (but not completely covered) with irrigated cropland. The rest of the landscape can either be covered with woodland, arid areas, or infrastructure (roads, ditches, etc.). All these values are based on global data on land cover, cropland extent, irrigation and land management, and are described in more detail in Malek and Verburg (2017).

To associate crop production to land systems in a sub-region, we first aggregated subnational and national crop production statistics for a specific sub-region. Then, we divided the crop production based on the share of crops produced in irrigated, intensive and extensive rainfed cropland, where we used data from SPAM (You et al. 2014). For example, total values of crops produced on irrigated cropland were assigned to irrigated land systems, based on their average cropland extent (mean % of land system unit) and their regional coverage. Values on the regional coverage and mean cropland extent are described in the next pages per sub-region.

The same is valid for irrigation values – reported irrigation water withdrawals were assigned to irrigated systems. Crop production in irrigated system is therefore also associated with a cost in terms of water use. Irrigation values present mean values for each unit of land system and not the actual demand of crops per ha.

Some values of land system output deviate from common crop production statistics. These seemingly surprising values can be explained threefold. First, we work with reported statistics on crop production. If statistics on a specific crop had errors, were inconsistent or unavailable, this was reflected in the lower land system output. Secondly, in several parts of the Mediterranean, the climate or irrigation enables multicropping. This could lead to higher average values of crop output per land system unit. Finally, the initial map of 2010 was generated using global data, where some of intensive cropland was possibly not captured. This could also result in higher output values per land system unit, particularly in the Southern Mediterranean.

Table B.1a Mediterranean North – European Union (% or land system unit or output per km²)

Land System	Regional coverage (% of region)	Mean cropland extent (% of land systems unit)	Mean permanent cropland extent (% of land systems unit)	Annual crops (t)	Permanent crops (t)	Livestock (nr)	Built up (ha)	Demand for water (m³)
wetlands	0.2	16.5	3.7	56.39	30.52	6.80	1.17	
medium inten. forest	10.6	14.37	3.1	6.91	7.50	7.30	0.57	
(semi)natural forest	5.9	10.9	2.3	5.28	5.55	7.78	0.25	
high inten. forest	2.4	9.8	1.6	4.99	3.93	8.04	0.29	
ext. arid system	0.5	12.0	3.3	5.27	8.14	7.19	1.34	
int. arid system	0.7	17.9	6.9	6.75	16.76	14.48	0.76	
closed wooded rangel.	2.8	17.3	2.7	8.97	6.46	20.82	0.24	
open woodland	5.7	14.7	4.7	6.11	11.41	6.18	0.97	
open wooded rangel.	7.3	19.3	4.3	9.15	10.60	16.56	0.43	
cropl./wooded rangel.	8.8	36.6	5.1	19.28	12.45	16.78	0.66	
cropland/rangel.	5.5	44.7	5.5	24.01	13.44	30.12	0.56	
exten. annual	3.3	46.5	5.6	25.07	13.52	11.81	0.84	
exten. permanent	2.3	37.5	31.1	3.90	75.82	9.03	1.26	
exten. mosaic	3.7	43.3	14.4	17.69	35.07	12.18	0.90	
rainfed inten. annual	8.0	49.7	3.3	331.35	27.19	17.29	0.70	
rainfed inten. perm.	3.3	46.1	34.0	86.32	281.03	12.11	1.07	
rainfed inten. mosaic	2.3	50.2	12.5	269.09	103.37	13.62	0.86	
irrigated annual	10.0	44.4	4.4	194.17	36.24	22.08	1.33	232.03
irrigated permanent	4.1	44.3	36.1	18.68	295.11	9.75	1.56	213.32
irrigated mosaic	4.5	43.9	14.8	128.42	121.25	14.95	1.50	199.43
peri-urban	6.3	38.3	13.7	108.59	112.17	13.93	9.34	120.47
urban	1.7	29.0	10.1	83.51	82.63	7.48	28.03	89.56

Table B.1b Mediterranean North – European Union – total crop production per land system group

Total crop production in 2010 (EUROSTAT, 2013, 2016a, 2016b). Crop production category "Irrigated" relates to irrigated and urban land systems, "Rainfed high" to rainfed intensive land systems, and "Rainfed low" to all remaining land systems, and is based on the shares provided by You et al. (2014).

	Share (%)	Production (t)
Annual crops		
Irrigated	41.6	33104235
Rainfed high	48.7	35277415
Rainfed low	9.7	7036980
Total annual		75418630
Permanent crops		
Irrigated	56.7	28885123
Rainfed high	27.2	13826362
Rainfed low	16.1	8197011
Total permanent		50908496

Table B.2a Mediterranean North – Western Balkans and Turkey (% or land system unit or output per km²)

Land System	Regional coverage (% of region)	Mean cropland extent (% of land systems unit)	Mean permanent cropland extent (% of land systems unit)	Annual crops (t)	Permanent crops (t)	Livestock (nr)	Built up (ha)	Demand for water (m³)
wetlands	1.1	12.9	0.3	51.21	7.91	4.52	0.22	
medium inten. forest	6.4	13.4	0.6	36.28	8.62	11.03	0.18	
(semi)natural forest	0.1	15.3	0.0	43.41	0.27	7.58	0.32	
high inten. forest	1.3	18.1	1.5	47.05	20.47	14.60	0.34	
ext. arid system	4.2	21.4	0.4	59.51	5.96	5.26	0.08	
int. arid system	5.9	24.6	0.6	68.22	8.07	12.97	0.17	
closed wooded rangel.	1.4	18.3	0.9	49.46	12.18	33.53	0.11	
open woodland	6.5	17.0	0.8	45.90	11.26	7.90	0.29	
open wooded rangel.	6.8	21.4	1.6	56.16	22.01	19.97	0.30	
cropl./wooded rangel.	6.0	32.8	1.2	89.92	15.74	17.40	0.29	
cropland/rangel.	5.8	41.2	2.8	109.10	38.29	24.80	0.48	
exten. annual	20.1	40.6	0.7	113.36	9.58	11.28	0.31	
exten. permanent	0.6	32.8	23.9	25.32	326.86	19.58	0.75	
exten. mosaic	0.8	40.3	8.0	91.94	108.77	13.80	0.48	
rainfed inten. annual	11.0	44.7	1.2	240.28	21.38	16.36	0.44	
rainfed inten. perm.	1.1	40.0	34.7	29.29	608.17	16.54	0.75	
rainfed inten. mosaic	0.8	43.7	11.1	180.43	193.67	17.08	0.50	
irrigated annual	16.4	45.1	1.0	179.85	23.02	20.31	0.63	306.17
irrigated permanent	0.4	42.2	16.3	64.83	611.44	25.76	1.45	279.89
irrigated mosaic	0.8	43.4	8.7	141.65	201.33	23.46	0.84	283.18
peri-urban	1.5	40.3	4.8	196.24	83.95	21.27	10.87	121.12
urban	0.8	29.7	4.1	141.31	72.39	18.14	33.70	77.05

Table B.2b Mediterranean North – Western Balkans and Turkey – total crop production per land system group

Total crop production in 2010 (EUROSTAT, 2013, 2016a, 2016b). Crop production category "Irrigated" relates to irrigated and urban land systems, "Rainfed high" to rainfed intensive land systems, and "Rainfed low" to all remaining land systems, and is based on the shares provided by You et al. (2014).

	Share (%)	Production (t)
Annual crops		
Irrigated	26.1	16645751
Rainfed high	27.9	17473003
Rainfed low	46.0	28688356
Total annual		62807110
Permanent crops		
Irrigated	24.2	4232387
Rainfed high	38.9	6800665
Rainfed low	36.9	6456238
Total permanent		17489290

Table B.3a Mediterranean South – Middle East and North-East Africa (% or land system unit or output per km²)

Land System	Regional coverage (% of region)	Mean cropland extent (% of land systems unit)	Mean permanent cropland extent (% of land systems unit)	Annual crops (t)	Permanent crops (t)	Livestock (nr)	Built up (ha)	Demand for water (m³)
wetlands	1.6	5.4	1.3	39.25	15.60	89.34	0.67	
medium inten. forest	0.4	33.0	12.2	46.02	98.27	17.36	0.36	
(semi)natural forest	0.0	5.8	3.9	4.22	31.44	17.34	0.00	
high inten. forest	0.1	38.8	10.7	40.03	86.23	192.62	1.72	
ext. arid system	16.2	2.4	0.6	3.97	4.73	1.41	0.23	
int. arid system	13.6	7.4	1.8	12.41	14.39	15.40	0.53	
closed wooded rangel.	0.1	28.5	11.5	37.58	92.77	53.82	1.00	
open woodland	2.4	7.8	1.9	13.01	15.37	38.50	0.60	
open wooded rangel.	4.1	9.0	2.3	14.72	18.63	23.77	0.67	
cropl./wooded rangel.	0.7	43.2	7.9	77.85	64.17	29.03	1.10	
cropland/rangel.	11.9	59.6	9.3	111.06	75.39	21.22	1.38	
exten. annual	4.2	54.7	3.1	113.96	24.97	10.21	0.93	
exten. permanent	0.6	66.6	25.8	68.07	208.25	19.14	0.42	
exten. mosaic	2.9	57.4	12.9	98.37	103.84	16.90	1.16	
rainfed inten. annual	0.7	56.3	4.6	501.43	113.87	35.42	1.05	
rainfed inten. perm.	0.0	43.0	19.1	231.42	721.35	18.74	0.25	
rainfed inten. mosaic	1.3	52.0	10.5	402.31	260.42	81.33	1.15	
irrigated annual	10.6	63.2	2.4	587.39	28.41	33.90	1.96	877.67
irrigated permanent	1.1	62.8	28.7	232.89	334.23	20.96	1.43	738.03
irrigated mosaic	10.5	66.4	10.7	538.19	124.91	49.04	1.99	902.86
peri-urban	12.9	60.1	7.3	510.05	85.45	47.83	9.59	734.42
urban	4.0	35.0	5.7	283.35	66.24	25.04	45.79	292.28

Table B.3b Mediterranean South – Middle East and North-East Africa – total crop production per land system group

Total crop production in 2010 (EUROSTAT, 2013, 2016a, 2016b). Crop production category "Irrigated" relates to irrigated and urban land systems, "Rainfed high" to rainfed intensive land systems, and "Rainfed low" to all remaining land systems, and is based on the shares provided by You et al. (2014).

	Share (%)	Production (t)
Annual crops		
Irrigated	82.9	38857839
Rainfed high	4.4	1759748
Rainfed low	12.7	5027599
Total annual		45645186
Permanent crops		
Irrigated	58.2	6560864
Rainfed high	8.0	900796
Rainfed low	33.8	3809004
Total permanent		11270664

Table B.4a Mediterranean South – North-West Africa (% or land system unit or output per km²)

Land System	Regional coverage (% of region)	Mean cropland extent (% of land systems unit)	Mean permanent cropland extent (% of land systems unit)	Annual crops (t)	Permanent crops (t)	Livestock (nr)	Built up (ha)	Demand for water (m³)
wetlands	0.9	6.6	1.5	29.89	12.15	288.60	0.12	
medium inten. forest	1.2	30.8	5.3	14.11	14.23	16.08	0.33	
(semi)natural forest	0.3	25.2	0.5	13.66	1.39	14.14	0.07	
high inten. forest	0.3	44.5	5.4	21.67	14.46	18.83	0.48	
ext. arid system	29.8	2.2	0.2	1.12	0.52	1.57	0.09	
int. arid system	20.1	6.2	1.0	2.90	2.57	8.49	0.10	
closed wooded rangel.	0.6	33.6	4.7	16.00	12.59	27.38	0.40	
open woodland	0.4	30.5	3.1	9.60	8.42	9.30	0.57	
open wooded rangel.	1.8	24.8	2.5	12.34	6.70	15.07	0.14	
cropl./wooded rangel.	4.4	52.9	2.2	28.08	5.84	15.73	0.28	
cropland/rangel.	13.2	64.9	6.0	32.63	16.05	18.47	0.31	
exten. annual	13.0	58.2	1.6	31.33	4.29	7.38	0.33	
exten. permanent	0.8	69.4	32.9	14.66	88.20	13.80	0.48	
exten. mosaic	2.4	57.6	11.1	25.73	29.82	13.22	0.37	
rainfed inten. annual	0.4	69.3	3.6	586.14	12.31	12.93	0.72	
rainfed inten. perm.	1.9	65.8	32.5	296.55	112.55	22.45	0.23	
rainfed inten. mosaic	0.4	61.8	11.7	447.05	40.33	18.46	0.42	
irrigated annual	3.0	48.7	2.3	388.31	18.91	12.43	0.76	338.74
irrigated permanent	1.2	68.8	33.7	205.62	274.28	17.11	0.93	439.42
irrigated mosaic	1.6	61.4	14.1	276.70	114.93	14.83	0.94	357.07
peri-urban	1.6	55.1	8.2	274.41	66.76	22.56	11.54	106.19
urban	0.8	37.2	6.5	179.82	52.66	10.05	39.80	54.17

Table B.4b Mediterranean South – North-West Africa – total crop production per land system group

Total crop production in 2010 (EUROSTAT, 2013, 2016a, 2016b). Crop production category "Irrigated" relates to irrigated and urban land systems, "Rainfed high" to rainfed intensive land systems, and "Rainfed low" to all remaining land systems, and is based on the shares provided by You et al. (2014).

	Share (%)	Production (t)
Annual crops		
Irrigated	52.8	17419629
Rainfed high	22.0	7242574
Rainfed low	25.2	8310512
Total annual		32972716
Permanent crops		
Irrigated	49.0	5117591
Rainfed high	16.6	1733763
Rainfed low	34.4	3604275
Total permanent		10455630

Appendix C – Location factors used in calculating spatial preference maps using logistic regression

Location Factor	Unit/description	Resolution	Date	Source
Socio-economic				
Population density*	People/km ²	1 km	2010	CIESIN (2015)
Rural population*	Rural population/km ²	1 km	2000	CIESIN et al. (2011)
Market accessibility	Index (0-1)	1 km	2000-2010	Verburg et al. (2011b)
Market influence	USD/person (ppp)	1 km	2000-2010	Verburg et al. (2011b)
Accessibility**	Distance to roads (m)	vector	1999	NGIA (2015)
Soil				
Drainage	Drainage class	1 km	2010	Hengl et al. (2014)
Sand content	Sand mass in %	1 km	2010	Stoorvogel et al. (2016)
Clay content	Clay mass in %	1 km	2013	Stoorvogel et al. (2016)
Cation Exchange Capacity (CEC)	cmol/kg	1 km	2010	Hengl et al. (2014)
pH	log(h+)	1 km	2010	Hengl et al. (2014)
Organic carbon content	g/kg in the top 50 cm	1 km	2013	Stoorvogel et al. (2016)
Soil depth	cm	1 km	2013	Stoorvogel et al. (2016)
Terrain				
Altitude	m above sea level	1 km	2005	Hijmans et al. (2005)
Slope	Slope degrees	1 km	2005	derived from Hijmans et al. (2005)
Climate				
Precipitation*	annual precipitation (sum of monthly means) in mm	1 km	1960-1990	Hijmans et al. (2005)
Temperature*	Temperature (mean of monthly means) Celsius degree	1 km	1960-1990	Hijmans et al. (2005)
Solar radiation	Horizontal surface irradiation (kWh/m ²), 1998-2011 mean	1.5 arc minute	2012	Huld et al. (2012)
Other				
Potential Evapotranspiration (PET)*	annual PET in mm	1 km	1960-1990	Zomer et al. (2008)
Potential vegetation	Pot. vegetation classes	10 km	2000	Ellis & Ramankutty (2008)

* dynamic factor, updated annually

** stays the same in growth scenario, improved accessibility in the sustainable scenario

Appendix D – example of spatial preference maps




Fig. D.1 Spatial preference map for the urban system




Fig. D.2 Spatial preference map for the cropland/wooded rangeland mosaic system




Fig. D.3 Spatial preference map for the irrigated annual cropland system

Appendix E – technical details of both scenarios

Table continues on several pages

Model setting	Sustainable scenario	Growth oriented
Dynamic driving factors		
Population density	Growth rates based on SSP2 projections of Jiang and O'Neill (2015) applied to high population areas only (> 250 per km ² , S9)	Growth rates based on SSP2 projections of Jiang and O'Neill (2015) applied to high population areas only (> 250 per km ² , S9)
Rural population density	Rural population stabile	Change rates (growth/decrease) based on SSP2 projections of Jiang and O'Neill (2015) applied to rural population density map
Temperature, climate, potential evapotranspiration	Based on RCP4.5 climate change maps (mean of all model runs for RCP4.5). PET calculated using PET equations.	Based on RCP4.5 climate change maps (mean of all model runs for RCP4.5). PET calculated using PET equations.
Aridity index	Calculated using the AI equation, dynamic for every year (RCP4.5)	Calculated using the AI equation, dynamic for every year (RCP4.5)
Static driving factors		
Market index and accessibility	All inaccessibility above average reduced by 40 %.	Stays as it is (differences)
Demands		
Annual crops	<ol style="list-style-type: none"> 1. Due to the common Mediterranean market, MENA can easier satisfy its food demand by imports, the demands for MENA are so lower (10 % lower growth rate compared to growth scenario). 2. 5 % decrease in total demand due to a reduction of food waste. 	Annual crops follow the SSP2 marker scenario for food production.
Permanent crops	<ol style="list-style-type: none"> 1. Due to the common Mediterranean market, exports of permanent crops are easier and thus higher (10 % higher growth rate compared to the growth scenario). 2. 5 % decrease in total demand due to a reduction of food waste. 	Permanent crops follow the SSP2 marker scenario for food production.
Livestock	<ol style="list-style-type: none"> 1. Due to the common Mediterranean market, MENA can easier satisfy its livestock by imports, and thus the demands for MENA are lower (10 % lower growth rate compared to growth scenario). 2. 5 % decrease in total livestock demand due to a reduction of food waste. 	Livestock numbers follow the SSP2 scenario for livestock numbers.

Built up areas	Demand linked to population change (SSP2), with 30 % lower growth rate due to increased density and compactness of new urban areas.	Demand linked to population change (SSP2), with a 10 % higher growth rate than population growth.
Land system specific settings		
<i>Wetlands</i>		
Supply	Wetlands still contribute fulfilling the demand, but the output is reduced: 30 % decrease in crops and livestock output of wetlands.	Same supply as baseline (overgrazing of wetland areas still possible).
Spatial pattern/change process	Protected – no change possible	Protected – no change possible
Other wetland conservation	Sustainable water management implementation on the regional scale. Irrigated land systems are defined with a water demand value (demand table, S2). Each region has limited water resources, cannot allocate more irrigated areas.	No other wetland conservation measures.
<i>Settlement systems</i>		
Supply	<ol style="list-style-type: none"> 1. Remains the same as in the baseline for built up areas 2. Efficiency improvement for livestock (5 % increase) 3. Efficiency improvement for crop production (closing yield gaps in EU, 90 % attainable yield achieved in other regions) 	<ol style="list-style-type: none"> 1. Remains the same as in the baseline for built up areas 2. Efficiency improvement for livestock (15 % increase due to hosting indoor livestock breeding facilities) 3. Efficiency improvement for crop production (75 % attainable yield achieved)
Spatial pattern/change process	<ol style="list-style-type: none"> 1. Lower conversion order for peri-urban systems (2), higher density urban systems preferred. 2. Neighborhood of land system allocation more compact (only 2 neighboring cells). 3. Conversion to urban only possible in non-protected areas. 4. More difficult urban expansion on the account of best cropland (higher conversion resistance for intensive cropland systems). 	<ol style="list-style-type: none"> 1. Higher conversion order for peri-urban systems (3), higher conversion order for livestock provision 2. Larger neighborhood of urban and peri-urban land system allocation (3 cell neighborhood). 3. Urban land has absolute priority. Any system (except natural forests and wetlands) can be converted to a settlement.
<i>Forest systems</i>		
Supply	N.A.	N.A.
Spatial pattern/change process	<ol style="list-style-type: none"> 1. All forest systems (including closed wooded rangeland) can change to open woodland in areas with an aridity index < 0.65. 2. Forest expansion (open 	<ol style="list-style-type: none"> 1. All forest systems (including closed wooded rangeland) can change to open woodland in areas with an aridity index < 0.65. 2. Forest expansion (open

	woodlands to forests) only possible in areas with an AI > 0.65	woodlands to forests) only possible in areas with an AI > 0.65
<i>Rainfed intensive</i>		
Supply	<ol style="list-style-type: none"> 1. Yields reach 75 % of attainable yield - higher yields in the region often not possible without irrigation (Mueller <i>et al.</i>, 2012) 2. 5 % decrease in total cropland output due to ecological focus areas (EFA) – set aside policy. 	<ol style="list-style-type: none"> 1. Rainfed intensive systems have to be at 50% of attainable yield (all regions already achieve that, except the Maghreb region). 2. Increased livestock efficiency by 15 %.
Spatial pattern/change process	<ol style="list-style-type: none"> 1. Possible only in areas with AI > 0.2. In areas with $0.2 < AI < 0.65$, there is a decrease in probability of rainfed intensive areas (-0.05) to account for extreme climate events. This value is lower than in the growth scenario, as it considers crop change or cultivar improvements. 2. Not possible in new protected areas (unless existing). 3. Can be transformed to less intensive systems after 10 years (annual crops) or 15 years (permanent crops) 	<ol style="list-style-type: none"> 1. Possible only in areas with AI > 0.2. In areas with $0.2 < AI < 0.65$, there is a decrease in probability of rainfed intensive areas (-0.1) to account for extreme climate events. This value is higher than in the sustainable scenario, as there are no changes in crops. 2. Can be transformed to less intensive systems after 10 years (annual crops) or 15 years (permanent crops)
<i>Extensive cropland</i>		
Supply	<ol style="list-style-type: none"> 1. 50 % of attainable yield achieved due to crop change or labor intensification (due to rural development policies and small farm promotion). 2. 5 % decrease in annual cropland output due to ecological focus areas (EFA) – set aside policy. 3. Increase in livestock output by 5 % due to improvements in breeds and herd fattening. 	No efficiency increase, considered as economically unattractive areas.
Spatial pattern/change process	<ol style="list-style-type: none"> 1. No spatial limitations. 2. Conversions to woodlands after being inactive for 10 years (abandonment). 	<ol style="list-style-type: none"> 1. No spatial limitations. 2. Conversions to woodlands after being inactive for 10 years (abandonment).
<i>Irrigated cropland</i>		
Supply	<ol style="list-style-type: none"> 1. Yield gaps closed in EU, in other regions yields reach 90 % of attainable yield. 2. Improved irrigation efficiency by 35 % 3. Increase in livestock output by 5 % due to improvements in breed and herd fattening 	<ol style="list-style-type: none"> 1. Yields achieve 75 % of attainable yield. 2. Increased livestock efficiency by 15 %
Spatial	Irrigated areas limited by amount of	Irrigated areas have a reduced spatial

pattern/change process	available water resources (by region)	probability (locspect -0.1) in areas with a market index below 0.3 (areas with extremely low investment potential).
Mosaic systems		
Supply	<ol style="list-style-type: none"> 1. Increase in livestock output by 5 % due to improvements in breeds and herd fattening 	<ol style="list-style-type: none"> 1. No increase in output of traditional systems 2. Lower conversion resistance of traditional systems as they experience further marginalisation
Spatial pattern/change process	<ol style="list-style-type: none"> 1. Higher conversion resistance of mosaic systems (less likely to be changed to more intensive, or abandoned). 2. Higher conversion order of mosaic systems to satisfy the demand for livestock and crops (we tell the model it needs to promote those systems) 	More intensive rangeland mosaics preferred (so a 2 function open woodland preferred to a 3 function medium tree cover woodland)
Arid systems		
	<ol style="list-style-type: none"> 1. Arid systems have reduced potential to fulfill demand for livestock. 2. Grazing intensification in arid systems not allowed. 	<ol style="list-style-type: none"> 1. Arid systems' potential to fulfill demand for livestock remains the same. 2. Grazing intensification in arid systems allowed.
Other		
Biodiversity/nature protection	Aichi targets reached – 17 % of terrestrial areas defined as protected areas (Pouzols <i>et al.</i> , 2014), S6. Conversions to less intensive systems, and systems with low intensity is possible in the protected areas.	Aichi targets not reached. National parks and similar areas are protected and excluded from changes, no expansion of PAs in MENA.
Ecological Focus Areas (EFA)	A Mediterranean Common Mediterranean Agricultural Policy (CAP) results in a set aside policy – farmers have to ensure that 5 % of their land is set aside to receive payments from the CAP. EFAs are either buffer strips, set aside land or similar. Land systems in question will experience a 5 % decrease in the annual cropland output of their systems. This decrease in output will occur gradually from year 0 to year 5.	No such measure applied.

Appendix F – protected areas in the Mediterranean ecoregion




Fig. F Existing and recommended protected area (PA) extent. Recommended areas used in the sustainability scenario are based on the national priorities for protected areas (Pouzols *et al.*, 2014) and are defined as 17 % of national terrestrial areas

Appendix G – CMIP5 simulations of the RCP4.5 scenario used to update precipitation, temperature, PET and AI maps for the years 2041-2060 (representing 2050)

Model	Institution
ACCESS1-0	Commonwealth Scientific and Industrial Research Organisation (CSIRO) and Bureau of Meteorology, Australia
bcc-csm1-1	Beijing Climate Center, China Meteorological Administration
CCSM4	National Center for Atmospheric Research, USA
CESM1(CAM5.1, FV2)	National Science Foundation, Department of Energy, National Center for Atmospheric Research, USA
CNRM-CM5	Centre National de Recherches Meteorologiques / Centre Europeen de Recherche et Formation Avancees en Calcul Scientifique, France
GFDL-CM3	NOAA, Geophysical Fluid Dynamics Laboratory, USA
GFDL-ESM2G	NOAA, Geophysical Fluid Dynamics Laboratory, USA
GISS-E2-R	NASA Goddard Institute for Space Studies, USA
HadGEM2-AO	National Institute of Meteorological Research / Korea Meteorological Administration
HadGEM2-CC	Met Office Hadley Centre, UK
HadGEM2-ES	Met Office Hadley Centre / Instituto Nacional de Pesquisas Espaciais)
INM-CM4	Institute for Numerical Mathematics, Russia
IPSL-CM5A-LR	Institut Pierre-Simon Laplace, France
MIROC-ESM-CHEM	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies
MIROC-ESM	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies
MIROC5	Atmosphere and Ocean Research Institute (The University of Tokyo), National Institute for Environmental Studies, and Japan Agency for Marine-Earth Science and Technology
MPI-ESM-LR	Max Planck Institute for Meteorology (MPI-M), Germany
MRI-CGCM3	Meteorological Research Institute, Japan
NorESM1-M	Norwegian Climate Centre




Fig. G.1 Changes in precipitation until 2050 (compared to current climate, represented by the 1960-1990 average) based on the mean of 19 CMIP5 simulations of the RCP4.5 for the Mediterranean ecoregion



Fig. G.2 Changes in temperature in °C until 2050 (compared to current climate, represented by the 1960-1990 average) based on the mean of 19 CMIP5 simulations of the RCP4.5 for the Mediterranean ecoregion

Appendix H – PET and AI calculation and map example

Potential evapotranspiration (PET) represents the ability of the atmosphere to remove water through evapotranspiration processes, and was introduced by the FAO (Allen & FAO, 1998; Trabucco *et al.*, 2008). We used the Hargreaves model (Hargreaves & Allen, 2003) to calculate future PET in this study, as it was also used to calculate the current PET in Zomer *et al.* (2008):

$$PET = 0.0023 \times RA \times (T_{mean} + 17.8) \times TD^{0.5}$$

PET – monthly average PET (mm/year)

RA – annual extra-terrestrial radiation, radiation on top of atmosphere expressed (mm/year)

T_{mean} – annual mean temperature (°C)

TD – annual mean daily temperature range (°C)

We used spatial distributions on RA, T_{mean} and TD from the Worldclim dataset (Hijmans *et al.*, 2005). Future temperature spatial distributions were acquired by calculating the mean from 19 CMIP5 simulations of the RCP4.5 scenario (S7).




Fig. H.1 Changes in annual PET until 2050 based on the 19 CMIP5 simulations of the RCP4.5 for the Mediterranean ecoregion. The future decrease in the temperature range is balancing the increase in the mean temperature, leading to a decrease in PET in some areas (marked as green on the figure).

The Aridity index serves as an indicator to quantify precipitation deficits over atmospheric water demand (UNEP, 1997; Zomer *et al.*, 2008). It is defined as a function of precipitation and potential evapotranspiration (PET):

$$AI = \frac{MAP}{MAE}$$

AI – aridity index

MAP – mean annual precipitation (mm/year)

MAE – mean annual potential evapotranspiration (PET, mm/year)




Fig. H.2 Current aridity index (Zomer et al. 2008)




Fig. H.3 Changes to aridity until 2050 based on the 19 CMIP5 simulations of the RCP4.5 for the Mediterranean ecoregion

Appendix I – Changes to urban population




Fig. I Areas with high population density where urban population change trends based on the SSP2 scenario were applied (Kc & Lutz, 2014; Jiang & O'Neill, 2015)

Appendix J – Scenarios: high resolution figures




Fig. J.1 Future land systems in 2050 as simulated for the two scenarios : (a) growth, (b) sustainability




Fig. J.2 Changes in land management intensity in the (a) growth and (b) sustainability scenario




Fig. J.3 Land system change affecting mosaic land systems in the growth (a) and sustainability (b) scenario

Appendix K – Freshwater resources and irrigation water withdrawal (EUROSTAT, 2013, 2016; FAO, 2016)

	North		South	
	W Balkans and Turkey	European Union	Middle East and NE Africa	NW Africa
Renewable freshwater resources (km ³)	255.3	478.5	62.8	55.3
% of irrigation water withdrawal in total water withdrawal	78.9	39.3	93.8	78.2
Irrigation water withdrawal (km ³ /yr)				
2010	30.1	48.6	59.3	16.7
Growth scenario	46.9	66.6	72.0	27.4
Sustainability scenario	24.7	38.2	42.8	11.1

References

- Allen RG, FAO (eds.) (1998) *Crop evapotranspiration: guidelines for computing crop water requirements*. Food and Agriculture Organization of the United Nations, Rome, 300 pp.
- CIESIN (2015) Gridded Population of the World, Version 4. Center for International Earth Science Information Network - CIESIN - Columbia University. NASA Socioeconomic Data and Applications Center (SEDAC).
- CIESIN, IFPRI, CIAT (2011) Global Rural-Urban Mapping Project, Version 1: Population Density Grid. Center for International Earth Science Information Network - CIESIN - Columbia University, International Food Policy Research Institute - IFPRI, The World Bank, and Centro Internacional de Agricultura Tropical - CIAT.
- Ellis EC, Ramankutty N (2008) Putting people in the map: anthropogenic biomes of the world. *Frontiers in Ecology and the Environment*, **6**, 439–447.
- EUROSTAT (2013) *Pocketbook on Euro-Mediterranean statistics*. Publications office of the European Union, Luxembourg.
- EUROSTAT (2016a) Agriculture statistics - North Africa and Eastern Mediterranean. Available from: http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_-_North_Africa_and_Eastern_Mediterranean (Accessed on: June 12, 2017)
- EUROSTAT (2016b) Agriculture statistics at regional level. March 2016. Available from: http://ec.europa.eu/eurostat/statistics-explained/index.php/Agriculture_statistics_at_regional_level (Accessed on: June 12, 2017)
- FAO (2016) AQUASTAT website. FAO's Information System on Water and Agriculture. Food and Agriculture Organization of the United Nations.
- Hargreaves G, Allen R (2003) History and Evaluation of Hargreaves Evapotranspiration Equation. *Journal of Irrigation and Drainage Engineering*, **129**, 53–63.
- Hengl T, de Jesus JM, MacMillan RA et al. (2014) SoilGrids1km — Global Soil Information Based on Automated Mapping. *PLoS ONE*, **9**, e105992.
- Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**, 1965–1978.
- Huld T, Müller R, Gambardella A (2012) A new solar radiation database for estimating PV performance in Europe and Africa. *Solar Energy*, **86**, 1803–1815.
- Jiang L, O'Neill BC (2015) Global urbanization projections for the Shared Socioeconomic Pathways. *Global Environmental Change*.
- Kc S, Lutz W (2014) The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100. *Global Environmental Change*.
- Malek Ž, Verburg PH (2017) Mediterranean land systems: representing diversity and intensity of complex land systems in a dynamic region. *Landscape and Urban Planning*, **165**, 102–116.
- Mueller ND, Gerber JS, Johnston M, Ray DK, Ramankutty N, Foley JA (2012) Closing yield gaps through nutrient and water management. *Nature*, **490**, 254–257.
- NGIA (2015) VMap0 data. National Geospatial Intelligence Agency. <http://gis-lab.info/qa/vmap0-eng.html>.
- Pouzols FM, Toivonen T, Di Minin E et al. (2014) Global protected area expansion is compromised by projected land-use and parochialism. *Nature*, **516**, 383–386.
- Stoorvogel JJ, Bakkenes M, Temme AJAM, Batjes NH, ten Brink B (2016) S-World: a Global Soil Map for Environmental Modelling: S-World: a Global Soil Map for Environmental Modelling. *Land Degradation & Development*.

- Trabucco A, Zomer RJ, Bossio DA, van Straaten O, Verchot LV (2008) Climate change mitigation through afforestation/reforestation: A global analysis of hydrologic impacts with four case studies. *Agriculture, Ecosystems & Environment*, **126**, 81–97.
- UNEP (1997) *World atlas of desertification*. United Nations Environment Programme, London ; New York : New York.
- Verburg PH, Ellis EC, Letourneau A (2011) A global assessment of market accessibility and market influence for global environmental change studies. *Environmental Research Letters*, **6**, 034019.
- You L, Wood-Sichra U, Fritz S, Guo Z, See L, Koo L (2014) Spatial Production Allocation Model (SPAM) 2005 v2.0.
- Zomer RJ, Trabucco A, Bossio DA, Verchot LV (2008) Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, Ecosystems & Environment*, **126**, 67–80.