



'Bacteroides cutis,' a new bacterial species isolated from human skin.

S. Belkacemi, N. Cassir, J. Delerce, F. Cadoret, B. La Scola

► To cite this version:

S. Belkacemi, N. Cassir, J. Delerce, F. Cadoret, B. La Scola. 'Bacteroides cutis,' a new bacterial species isolated from human skin.. New Microbes and New Infections, 2018, 22, pp.17-18. 10.1016/j.nmni.2017.12.007 . hal-01795913

HAL Id: hal-01795913

<https://amu.hal.science/hal-01795913>

Submitted on 22 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

'*Bacteroides cutis*,' a new bacterial species isolated from human skin

S. Belkacemi, N. Cassir, J. Delerce, F. Cadoret and B. La Scola

Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, Aix-Marseille Université, Marseille, France

Abstract

We report the main characteristics of '*Bacteroides cutis*' sp. nov., strain Marseille-P4118^T (= CSUR P4118), a new species within the genus *Bacteroides*. This strain was isolated from a skin sample of a 75-year-old man from Marseille.

© 2017 Published by Elsevier Ltd.

Keywords: *Bacteroides cutis*, culturomics, intensive care unit patient, skin microbiota, taxonogenomics

Original Submission: 29 November 2017; **Accepted:** 21 December 2017

Article published online: 29 December 2017

Corresponding author: N. Cassir, URMITE, CNRS (UMR 7278), IRD (198), INSERM (U1095), AMU (UM63), Faculté de Médecine, Aix-Marseille Université, 27 Boulevard Jean Moulin, 13385 Marseille Cedex 5, France
E-mail: cassirnadim@gmail.com

In April 2017, as part of the culturomics study to assess the microbial diversity of the human skin [1,2], a new bacterial species was isolated from a skin sample of a 75-year-old man hospitalized in the Nord hospital, Marseille, France. Although the patient was dermatologically asymptomatic, we isolated from his neck a bacterial strain that could not be identified by our systematic matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) screening on a MicroFlex spectrometer (Bruker Daltonics, Bremen, Germany) [3]. The study was approved by the Institut Fédératif de Recherche 48 (agreement number 09-022, Marseille, France), and the patient's consent was obtained.

Strain Marseille-P4118^T was first isolated on 5% sheep's blood-enriched Columbia agar (bioMérieux, Marcy l'Etoile, France) at 37°C in anaerobic atmosphere (AnaeroGen Compact; Oxoid, Thermo Scientific, Dardilly, France) after a 3-day preincubation in an anaerobic bottle of blood culture (Becton Dickinson, Le Pont-de-Claix, France) containing 5% sheep's blood supplemented with 5% rumen fluid previously filter sterilized through a 0.2 µm pore filter (Thermo Fisher Scientific, Villebon sur Yvette, France). Colonies were

colourless, with a diameter ranging from 0.5 to 1.5 mm. Bacterial cells were Gram-negative, motile and rod-shaped bacilli with 0.6 µm wide by 2.8 µm long.

The strain Marseille-P4118^T grew after 48 hours of incubation under anaerobic conditions at a temperature ranging from 28°C to 40°C, but optimally at 37°C. Growth was impossible in microaerophilic (campyGEN; Oxoid) conditions. The bacterial cells tolerated a pH of 6.5 to 7, and a NaCl concentration less than 50 mg/L. After 20 minutes of thermal shock at 80°C, this bacterium did not grow at 37°C on Columbia agar enriched with 5% sheep's blood, confirming the spore search result by microscopy, which was negative. The catalase and oxidase tests for the strain Marseille-P4118^T were negative.

After three failed identifications by MALDI-TOF MS screening on a MicroFlex spectrometer (Bruker Daltonics) [3], the 16S rRNA gene was sequenced using universal primers FD1 and RP2 (Eurogentec, Angers, France) as previously described [4] and a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France). Strain Marseille-P4118^T exhibited a 97.4% sequence identity with the type strain *Bacteroides uniformis*, the phylogenetically closest species with standing in nomenclature (Fig. 1). Therefore, strain Marseille-P4118 was classified as a member of the genus *Bacteroides* in the order of *Bacteroidales* within the *Bacteroidetes* phylum. Because the sequence identity with the phylogenetically closest validated species was <98.7%, which is the threshold recommended to define a species according to the nomenclature [5,6], we suggest the creation of a new species named '*Bacteroides cutis*' sp. nov. strain Marseille-P4118^T




FIG. 1. Phylogenetic tree showing position of '*Bacteroides cutis*' strain Marseille-P4118^T (red) relative to other phylogenetically close neighbours. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using maximum-likelihood method within MEGA software. Numbers at nodes are percentages of bootstrap values obtained by repeating analysis 1000 times to generate majority consensus tree. Only bootstraps scores of at least 95% were retained. Scale bar indicates 2% nucleotide sequence divergence.

(ku.tis, L. fem., from *cutis*, 'skin,' the site from which strain Marseille-P4118^T was first isolated).

MALDI-TOF MS spectrum

The MALDI-TOF MS spectrum of '*Bacteroides cutis*' Marseille-P4118^T is available online (<http://www.mediterraneeinfection.com/>).

Nucleotide sequence accession number

The 16S rRNA gene sequence was deposited in GenBank under accession number LT908453.

Deposit in a culture collection

Strain Marseille-P4118^T was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR) under number P4118.

Acknowledgement

This research was funded by the Agence Nationale de la Recherche as part of the Méditerranée Infection I0-IAHU-03 project.

Conflict of interest

None declared.

References

- [1] Lagier JC, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, et al. Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat Microbiol* 2016;1:16203.
- [2] Cassir N, Papazian L, Fournier PE, Raoult D, La Scola B. Insights into bacterial colonization of intensive care patients' skin: the effect of chlorhexidine daily bathing. *Eur J Clin Microbiol Infect Dis* 2015;34:999–1004.
- [3] Seng P, Abat C, Rolain JM, Colson P, Lagier JC, Gourié F, et al. Identification of rare pathogenic bacteria in a clinical microbiology laboratory: impact of matrix-assisted laser desorption ionization–time of flight mass spectrometry. *J Clin Microbiol* 2013;51:2182–94.
- [4] Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol* 2000;38:3623–30.
- [5] Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* 2006;33:152.
- [6] Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi M, Sentausa E, et al. A polyphasic strategy incorporating genomic data for the taxonomic description of novel bacterial species. *Int J Syst Evol Microbiol* 2014;64:384–91.