



HAL
open science

**'Massilimalia timonensis' sp. nov., a new bacterium
species isolated from human faeces after
decontamination with alcohol.**

P. Afouda, G. Dubourg, F. Cadoret, Pierre-Edouard Fournier, Didier Raoult

► **To cite this version:**

P. Afouda, G. Dubourg, F. Cadoret, Pierre-Edouard Fournier, Didier Raoult. 'Massilimalia timonensis' sp. nov., a new bacterium species isolated from human faeces after decontamination with alcohol.. New Microbes and New Infections, 2018, 21, pp.86-87. 10.1016/j.nmni.2017.10.011 . hal-01795915

HAL Id: hal-01795915

<https://amu.hal.science/hal-01795915>

Submitted on 22 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

'*Massilimalia timonensis*' sp. nov., a new bacterium species isolated from human faeces after decontamination with alcohol

P. Afouda, G. Dubourg, F. Cadoret, P. E. Fournier and D. Raoult

Aix Marseille Université, Marseille, France

Abstract

We describe the main characteristics of a new bacterial species '*Massilimalia timonensis*' sp. nov. The type strain Marseille-P3753^T (=CSUR P3753) was isolated from the stool of a healthy Senegalese man.

© 2017 Published by Elsevier Ltd.

Keywords: Culturomics, Gut microbiota, Human microbiota, Massilimalia, Taxonogenomics

Original Submission: 4 August 2017; **Revised Submission:** 14 September 2017; **Accepted:** 26 October 2017

Article published online: 26 November 2017

Corresponding author: D. Raoult, Aix-Marseille Université, Unité de Recherche sur les Maladies Infectieuses et Tropicales Emergentes (URMITE), CNRS 7278, IRD 198, INSERM 1095, UM63, Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, 27 Boulevard Jean Moulin, 13385, Marseille cedex 5, France
E-mail: didier.raoult@gmail.com

Thanks to the culturomics approach [1], a new bacterial species was isolated from the stool sample of a healthy 32-year-old Senegalese man in January 2017. Patient consent was obtained, and the study approved by the Institut Fédératif de Recherche 48 (Faculty of Medicine, Marseille, France), under agreement 09-022. For eliminating vegetative forms, stools were decontaminated with 70% alcohol (v/v) as previously described [2]. Strain Marseille-P3753 was isolated for the first time at 37°C in anaerobic conditions on 5% sheep's blood-enriched Columbia agar (bioMérieux, Marcy l'Etoile, France) after 5 days' pre-incubation in an anaerobic bottle of blood culture containing 5% sheep's blood. Strain Marseille-P3753 grows correctly after 72 hours' incubation in microaerophilic and anaerobic conditions at a temperature ranging from 28°C to 45°C, with an optimal growth temperature at 37°C. On sheep's blood-enriched Columbia agar medium, strain Marseille-P3753 formed transparent microcolonies with diameters ranging from 0.2 to 0.7 mm. '*Massilimalia timonensis*' is nonmotile, rod shaped and non-spore forming. Bacterial cells observed under electron

microscopy have a length ranging from 0.7 to 2.9 µm, and their growth is observed at a NaCl level below 5 g/L. The catalase and oxidase tests for strain Marseille-P3753 showed no activity.

The 16S rRNA gene was sequenced using universal primers FDI and RP2 (Eurogentec, Angers, France) as previously described [3] using a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France). Strain Marseille-P3753 showed a 98.05% sequence identity with the strain '*Massilimalia massiliensis*' Marseille-P2963 (GenBank accession no. LT576408), the phylogenetically closest species. Therefore, this bacterium was classified as a new species of the genus *Massilimalia*. Other close species belonging to the same *Firmicutes* phylum are shown in Fig. 1.

Strain Marseille-P3753 shows a 16S rRNA gene sequence divergence of >1.3% with its phylogenetically closest species validated species [4,5]. Accordingly, a new species was proposed, named '*Massilimalia timonensis*' (ti.mo.nen'sis, L. adj. fem., from 'Timone,' the name of the main hospital of Marseille, France, where the strain was first isolated). Strain Marseille-P3753^T is the type strain of the new species '*Massilimalia timonensis*'.

MALDI-TOF MS spectrum

The matrix-assisted desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) spectrum of '*Massilimalia timonensis*' Marseille-P3753^T is available online (<http://www>.




FIG. 1. Phylogenetic tree showing position of 'Massilimalia timonensis' strain Marseille-P3753 relative to other phylogenetically close neighbours. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using maximum-likelihood method within MEGA software. Numbers at nodes are percentages of bootstrap values obtained by repeating analysis 1000 times to generate majority consensus tree. Only bootstrap scores of at least 90% were retained.

mediterranean-infection.com/article.php?laref=256&titre=urms-database).

Nucleotide sequence accession number

The 16S rRNA gene sequence was deposited in GenBank under accession number LT854295.

Deposit in a culture collection

Strain Marseille-P3753^T was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR, WDCM 875) under number P3753.

Acknowledgement

This study was funded by the Fondation Méditerranée Infection.

Conflict of interest

None declared.

References

- [1] Lagier JC, Hugon P, Khelaifa S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev* 2015;28:237–64.
- [2] Browne HP, Forster SC, Anonye BO, Kumar N, Neville BA, Stares MD, Goulding D, et al. Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation. *Nature* 2016;533:543–6.
- [3] Morel AS, Dubourg G, Prudent E, Edouard S, Gouriet F, Casalta JP, et al. Complementarity between targeted real-time specific PCR and conventional broad-range 16S rDNA PCR in the syndrome-driven diagnosis of infectious diseases. *Eur J Clin Microbiol Infect Dis* 2015;34:561–70.
- [4] Huson DH, Auch AF, Qi J, Schuster SC. MEGAN analysis of metagenomic data. *Genome Res* 2010;17:377–86.
- [5] Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microbiol Today* 2006;33:152.