



**HAL**  
open science

**'Colidextribacter massiliensis' gen. nov., sp. nov.,  
isolated from human right colon**

D. Ricaboni, M. Mailhe, F. Cadoret, V. Vitton, Pierre-Edouard Fournier,  
Didier Raoult

► **To cite this version:**

D. Ricaboni, M. Mailhe, F. Cadoret, V. Vitton, Pierre-Edouard Fournier, et al.. 'Colidextribacter massiliensis' gen. nov., sp. nov., isolated from human right colon. *New Microbes and New Infections*, 2017, 17 (C), pp.27-29. 10.1016/j.nmni.2016.11.023 . hal-01795965

**HAL Id: hal-01795965**

**<https://amu.hal.science/hal-01795965v1>**

Submitted on 22 May 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0  
International License

# '*Colidextribacter massiliensis*' gen. nov., sp. nov., isolated from human right colon

D. Ricaboni<sup>1,3</sup>, M. Mailhe<sup>1</sup>, F. Cadoret<sup>1</sup>, V. Vitton<sup>2</sup>, P.-E. Fournier<sup>1</sup> and D. Raoult<sup>1</sup>

1) Aix-Marseille Université, URMITE, UM63, CNRS7278, IRD198, Inserm 1095, Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, 2) Service de Gastroenterologie, Hôpital Nord, Assistance Publique-Hôpitaux de Marseille, Marseille, France and 3) Department of Biomedical and Clinical Sciences, 3rd Division of Clinical Infectious Disease, University of Milan, Luigi Sacco Hospital, Milan, Italy

## Abstract

We report here the main characteristics of '*Colidextribacter massiliensis*' strain Marseille-P3083<sup>T</sup> (CSURP3083), which was isolated from a human right colon lavage sample.

© 2016 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

**Keywords:** *Colidextribacter massiliensis*, Culturomics, human gut, microbiota, taxonogenomics

**Original Submission:** 25 October 2016; **Revised Submission:** 20 November 2016; **Accepted:** 23 November 2016


**Article published online:** 28 November 2016

**Corresponding author:** D. Raoult, Aix-Marseille Université, URMITE, UM63, CNRS7278, IRD198, Inserm 1095, Institut Hospitalo-Universitaire Méditerranée-Infection, Faculté de médecine, 27 Boulevard Jean Moulin, 13385, Marseille cedex 05, France  
**E-mail:** [didier.raoult@gmail.com](mailto:didier.raoult@gmail.com)

In 2016, as a part of culturomics study of the human microbiome [1], we isolated from the right colon of a 27-year-old obese patient a bacterial strain that could not be identified by our systematic matrix-assisted desorption ionization–time of flight mass spectrometry (MALDI-TOF MS) screening on a Microflex spectrometer (Bruker Daltonics, Bremen, Germany) [2]. The patient provided signed informed consent, and the study was validated by the ethics committee of the Institut Fédératif de Recherche IFR48 under number 2016-010.

Strain Marseille-P3083<sup>T</sup> growth was obtained on 5% sheep's blood–enriched Columbia agar medium (bioMérieux, Marcy l'Étoile, France) in anaerobic atmosphere (anaeroGEN, Oxoid, Dardilly, France) after a 30-day enrichment of the fresh right colon sample in an anaerobic haemoculture bottle (Becton Dickinson, Pont de Claix, France) added with 5 mL of sheep's blood (bioMérieux) and 5 mL of 0.2 µm filtered (Thermo Fisher Scientific, Villebon-sur-Yvette, France) rumen at 37°C.

After 96 hours of anaerobic incubation on 5% sheep's blood–enriched agar (bioMérieux) colonies were punctiform, approximately round and translucent, with a mean diameter of 0.1 to 0.3 mm. Bacterial cells were small Gram-negative cocci ranging in length from 0.4 to 0.6 µm. Strain Marseille-P3083<sup>T</sup> tested catalase and oxidase negative. Sporulation test (20 minutes at 80°C) was negative, and no growth was achieved under aerobic or microaerophilic (campyGEN, Oxoid) conditions. The 16S rRNA gene was sequenced using fDI-rP2 primers as previously described [3] using a 3130-XL sequencer (Applied Biosciences, Saint Aubin, France). Strain Marseille-P3083<sup>T</sup> exhibited a 93.57% sequence identity with *Pseudoflavonifractor capillosus* strain ATCC 29799<sup>T</sup> (GenBank accession no. AY136666), the phylogenetically closest species with standing in nomenclature (Fig. 1). This value putatively classifies strain Marseille-P3083<sup>T</sup> as a member of a new genus within the *Clostridiales* cluster IV in the phylum *Firmicutes*. *Clostridiales* cluster IV was created in 1994 and collects microorganisms that exhibit *Clostridium* and non-*Clostridium*-like characteristics [4], including Gram-negative microorganisms [5]. *Intestinimonas butyriciproducens* DSM 26588<sup>T</sup> (16S rRNA similarity of 93.42%) is another phylogenetically very close species (Fig. 1) that exhibits the peculiar characteristic of appearing as a



**FIG. 1.** Phylogenetic tree showing position of '*Colidextribacter massiliensis*' strain Marseille-P3083<sup>T</sup> relative to other phylogenetically close neighbours. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using maximum-likelihood method within MEGA software. Numbers at nodes are percentages of bootstrap values obtained by repeating analysis 1000 times to generate majority consensus tree. Only bootstrap scores of at least 90% were retained.

Gram-negative rod while electron microscopy demonstrated Gram-positive cell wall characteristics [6].

On the basis of the morphologic differences (morphology and Gram stain) and the 16S rRNA sequence divergence of strain Marseille-P3083<sup>T</sup> with the phylogenetically closest species with standing in nomenclature [7,8] we propose here the creation of the new genus *Colidextribacter* (Co.li.dex.tri.bac.ter, L. neut. n. *colon* or *colum*, 'colon'; L. adj. *dexter*, 'right'; N.L. masc. n. *bacter*, 'a rod'; N.L. masc. n. *Colidextribacter*, 'a rod from the right colon'); and a new species *Colidextribacter massiliensis* for which Marseille-P3083<sup>T</sup> (= CSUR P3083) is the type strain.

Strain Marseille-P3083<sup>T</sup> is the type strain of *Colidextribacter massiliensis* gen. nov., sp. nov. (mas.si.li.en'sis, L., masc. adj., *massiliensis*, for Massilia, the Roman name of Marseille).

### MALDI-TOF MS spectrum

The MALDI-TOF MS spectrum of '*Colidextribacter massiliensis*' strain Marseille-P23083<sup>T</sup> is available online (<http://www.mediterranee-infection.com/article.php?laref=256&titre=urms-database>).

[mediterranee-infection.com/article.php?laref=256&titre=urms-database](http://www.mediterranee-infection.com/article.php?laref=256&titre=urms-database).

### Nucleotide sequence accession number

The 16S r RNA gene sequence was deposited in GenBank under accession number LT598546.

### Deposit in a culture collection

Strain Marseille-P3083<sup>T</sup> was deposited in the Collection de Souches de l'Unité des Rickettsies (CSUR, WDCM 875) under number P3083.

### Acknowledgement

This study was funded by the Fondation Méditerranée Infection.

## Conflict of Interest

---

None declared.

## References

---

- [1] Lagier JC, Hugon P, Khelaifia S, Fournier PE, La Scola B, Raoult D. The rebirth of culture in microbiology through the example of culturomics to study human gut microbiota. *Clin Microbiol Rev* 2015;28:237–64.
- [2] Seng P, Abat C, Rolain JM, Colson P, Lagier JC, Gouriet F, et al. Identification of rare pathogenic bacteria in a clinical microbiology laboratory: impact of matrix-assisted laser desorption ionization–time of flight mass spectrometry. *J Clin Microbiol* 2013;51:2182–94.
- [3] Drancourt M, Bollet C, Carlouz A, Martelin R, Gayral JP, Raoult D. 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J Clin Microbiol* 2000;38:3623–30.
- [4] Collins MD, Lawson PA, Willems A, Cordoba JJ, Fernandez-Garayzabal J, Garcia P, et al. The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *Int J Syst Bacteriol* 1994;44:812–26.
- [5] Carlier JP, Bedora-Faure M, K'ouas G, Alauzet C, Mory F. Proposal to unify *Clostridium orbiscindens* Winter et al. 1991 and *Eubacterium plautii* (Séguin 1928) Hofstad and Aasjord 1982, with description of *Flavonifactor plautii* gen. nov., comb. nov., and reassignment of *Bacteroides capillosus* to *Pseudoflavonifactor capillosus* gen. nov., comb. nov. *Int J Syst Evol Microbiol* 2010;60:585–90.
- [6] Kläring K, Hanske L, Bui N, Charrier C, Blaut M, Haller D, et al. *Intestinimonas butyriciproducens* gen. nov., sp. nov., a butyrate-producing bacterium from the mouse intestine. *Int J Syst Evol Microbiol* 2013;63:4606–12.
- [7] Huson DH, Auch AF, Qi J, Schuster SC. MEGAN analysis of metagenomic data. *Genome Res* 2007;17:377–86.
- [8] Kim M, Oh HS, Park SC, Chun J. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* 2014;64:346–51.