



HAL
open science

Adult spawning and early larval development of the endangered bivalve *Pinna nobilis*

Sergio Trigos, Nardo Vicente, Patricia Prado, Francisco J. Espinós

► **To cite this version:**

Sergio Trigos, Nardo Vicente, Patricia Prado, Francisco J. Espinós. Adult spawning and early larval development of the endangered bivalve *Pinna nobilis*. *Aquaculture*, 2018, 483, pp.102-110. 10.1016/j.aquaculture.2017.10.015 . hal-01828955

HAL Id: hal-01828955

<https://amu.hal.science/hal-01828955>

Submitted on 15 Mar 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Adult spawning and early larval development of the endangered bivalve *Pinna nobilis*

Sergio Trigos^{a,*}, Nardo Vicente^{b,c}, Patricia Prado^d, Francisco J. Espinós^e

^a Innovation Network in Aquaculture Industries of the Valencian Community (RIIA-CV), Avda. Naranjos s/n. Edificio Colegio Mayor Galileo Galilei, Local 15, 46022, Valencia, Spain

^b Institut Océanographique Paul Ricard, Ile des Embiez, 83140 Six Fours les Plages, France

^c Institut Méditerranéen de la Biodiversité et de l'Ecologie marine et continentale (IMBE), Aix-Marseille Université, Avignon Université, CNRS, IRD, France

^d IRTA-Aquatic Ecosystems, Ctra. Poble Nou Km 5.5, 43540 Sant Carles de la Ràpita (Tarragona), Spain

^e ACUMA Research Center (Aquaculture and Environmental R.C.), Polytechnic University of Valencia, Camino de Vera s/n, 46022, Valencia, Spain

* Corresponding author at: Innovation Network in Aquaculture Industries of the Valencian Community (RIIA-CV), Avda. Naranjos s/n. Edificio Colegio Mayor Galileo Galilei, Local 15, 46022, Valencia, Spain. E-mail address: strigos@riia.es (S. Trigos).

Abstract: The development of aquaculture activities has posed an alternative solution for the preservation of some overexploited shellfish fisheries worldwide. In the same way, endemic Mediterranean bivalves such as *Pinna nobilis*, highly threatened by habitat loss and coastal pollution, could find in aquaculture a solution for pre-serving the continuity of the species. Given the endangered status of the species, the biological and ecological processes regulating natural populations have been well studied, but there are still important knowledge gaps preventing the development of viable artificial cultures. This study describes for the first time the larval development of *P. nobilis* (from fertilization until pediveliger larval stages) in captivity conditions. Moreover, different rearing tanks of 5, 16 and 80 L, larvae density from 1 to 600 larvae mL⁻¹, light conditions, food doses, were tested in order to establish the bases for the optimal rearing of the species and provide a source of individuals for restoring field populations. Results showed that 16 L tanks with a concentration of 2 larvae mL⁻¹, constant temperature of 21 °C, 12/12 h photoperiod and fed with an “optimal” mixture of 25 cells per µL of *Chaetoceros calcitrans* + 33.3 cells per µL of *Pavlova lutheri* + 100 cells per µL of *Isochrysis galbana* appear to be the best conditions to rear larvae of *P. nobilis*. Different captivity conditions such as lower or higher tank volume, larvae density, or food doses; light privation did not report better results for larval development.

Keywords: *Pinna nobilis*, Conservation, Larvae development, Rearing, Captivity

1. Introduction

In the last decades the rearing of endangered bivalve species for both commercial and ecological purposes has received considerable attention (Ellis, 2000; Knop, 2009). In spite of the economic and scientific interest, the rearing of bivalve species has been proved to be rather difficult due to the great larval mortalities of the group (Rumrill, 1990). Field studies show that larval mortality in natural conditions ranges from 25 to 80% during first days of life (Talmage and Gobler, 2009; Gazeau et al., 2010). Both natural and/or anthropogenic-driven mortality in bivalve larvae –i.e., from egg to postlarval recruit– are often difficult to measure and are considered as a “constant” (Philippart et al., 2003). Possible causes may include failure of fertilization, which is often related to sea acidification and may produce shape abnormalities in embryos or avoid development in over 50% of the larvae (Kurihara et al., 2007); coastal development (Gómez et al., 2000; Guest

et al., 2008); shortage of food resources (Rico-Villa et al., 2006), absence of a suitable substrate for benthic settlement (Su et al., 2007); combined effects of short planktonic durations and local patterns of marine circulation (Shanks and Brink, 2005); and/or lethal environmental temperatures (Philippart et al., 2003). Any of these scenarios usually leads to a reduction of recruitment rates and consequently, to the disappearance of populations.

Historically, mortality problems have been observed in the marine bivalve industry for decades (Samain and McCombie, 2008; Barton et al., 2015). The main goal in hatchery production is to improve larval and post-larval survival by achieving larval growth and metamorphosis success. In this term, the proper formulation of larval diets has been considered as the most critical aspect in hatchery operations focusing on the type of microalgae used as feed (Knuckey et al., 2002; Ponis et al., 2006; Rico-Villa et al., 2006; Pettersen et al., 2010; Ragg et al., 2010; Gui et al., 2016) or even in artificial substitutes to phytoplankton

diet (Coutteau and Sorgeloos, 1992; Knauer and Southgate, 1999; Nevejan et al., 2007; Gui et al., 2016).

In other cases, recurrent mortality episodes are described as a result of virus or *Vibrio*-like bacteria infecting bivalve hatcheries and nurseries (Renault and Arzul, 2001; Dubert et al., 2015; Rojas et al., 2015) and reducing commercial production by approximately 40% as reported for French farming facilities of Pacific oyster (*Crassostrea gigas*) since 2008 (Pernet et al., 2014).

From an ecological perspective, the rearing of endangered bivalves in captivity conditions may pose a potential solution for the rehabilitation of seriously damaged populations (Ronquillo and Mckinley, 2006; Vincie, 2008; Thomas et al., 2010; Theodorou et al., 2015) and face the same early stages' bottlenecks than those reported for bivalve commercial farming (Arnold, 2008). One of these endangered species is the Mediterranean fan mussel, *Pinna nobilis*. The populations of this bivalve have significantly declined during the last decades (De Gaulejac and Vicente, 1990; Vicente and Moreteau, 1991; Garcia-March, 2005; Basso et al., 2015; Rouanet et al., 2015) as a result of different anthropogenic stressors such as coastal development, fishing pressure, and/or accidental harvesting by trawling and shell breakage by anchoring (Katsanevakis, 2005; Acarli et al., 2011; Hendriks et al., 2011). For these reasons, the European Union included it as an endangered species in the ANNEX IV of the Council Directive 92/43/EEC (EC Habitats Directive), and strictly forbids any kind of culling. Although these measures have helped certain recovery of populations in some Mediterranean regions (Pérez-Vallazza et al., 2008; Theodorou et al., 2017), the species is still at risk by unknown factors such as those causing mass mortality event along the Spanish Mediterranean coast and the Balearic Islands during the summer of 2016 (Darriba, 2017; Vázquez-Luis et al., 2017). Temperature effects due to climate change, coastal pollution, and pathogenic mechanisms are among potential explanatory factors, all of them mediated by human influence (Rodrigues et al., 2015). In this scenario, the rearing of *P. nobilis* in captivity appears as an alter-native solution to obtain healthy stocks of recruits that could be re-introduced in suitable areas with declining population densities (Trigos, 2017). Yet, the duration of the different larval stages from fertilized eggs, as well as the main factors influencing survival rates are still largely unknown, although they are proposed to follow similar patterns to other bivalves. Following this hypothesis, Vicente (1986) described the larval stages of trochophore, veliger, and pediveliger followed by metamorphosis, and juvenile development of *P. nobilis*. De Gaulejac (1989), observed larvae shells with electronic microscope and pointed the possibility of a time gap of 5 to 10 days between gametes expulsion and substrate settlement. Peharda and Vilibić (2008) suggested that veliger fan mussel larvae could have a negative phototactism that allow vertical migration to deeper waters during the day to return to upper bathymetric ranges at night, as also indicated for other bivalves (Gosling, 2003), presumably to avoid UV light or predator activity (Manuel et al., 1996). However, there is still no available information on the behavioural responses of *P. nobilis* larvae to different environmental factors under captivity conditions that could provide a solid basis for designing and implementing a viable larval hatchery. In addition, factors determining patterns of benthic recruitment are virtually unknown, although in other species is often linked to the availability of preferential substrates for settlement (Prado et al., 2012), which highlights the importance of simulating optimal settlement conditions to rear *P. nobilis* in captivity. Other biological and physical aspects such as the possible correlation between adult size and numbers of oocytes expelled also require further investigation.

In this context, the main objective of this work was to describe, for the first time, the larval development of the ecologically important Mediterranean fan mussel (*P. nobilis*) with the aim of developing cultivation techniques and providing a practical guideline for problem identification during larval rearing in this species. More specifically we conducted experiments aimed (1) at different larval densities (2) with different tank volumes, (3) fed with three doses of microalgae mixture

and (4) reared in light/dark conditions.

2. Materials and methods

2.1. Field sampling

A total of 40 individuals of *P. nobilis* were collected around the Embiez archipelago, South East of France from April 2012 to September 2014. The collected specimens were carefully transported to laboratory facilities within portable tanks and their total height (Ht) registered and labelled. Shells were brushed to remove epibiontic organisms, including other bivalves (Rabaoui et al., 2015) that could affect the fertilization process. All individuals were reintroduced to the field at the end of the experiments.

2.2. Gamete release and fertilization

Small groups of 4–6 individuals were placed in two 120 L tanks within controlled temperature facilities (20–21 °C) with filtered (1 µm) and treated (UV) seawater for a maximum of 24 h. A ≈ 10 °C gradient is necessary to induce a thermal shock causing gametes release in bi-valves (Helm et al., 2006) which takes place at various temperature thresholds, depending on the species (Drent, 2004). Herein, in order to keep this temperature gradient two induction tanks were prepared; one tank was maintained at 15 °C while a second tank was heated to 25 °C thus respecting the conformability range of temperatures described for *P. nobilis* (Trigos et al., 2015).

Individuals were transferred to the induction tanks and individuals were translocated between both of them every 50 min for a maximum of 6 times. If no response was achieved, individuals were introduced in 2600 L tanks with constant aeration and water renovation, and the process repeated on the following days. When only males expelled gametes, small volumes of sperm were poured in the tanks on the following thermal shocks, with the aim to stimulate the females (if any) while the thermal shock was occurring. When male and female spawning occurred, individuals were introduced individually in 60 L tanks in order to avoid possible polyspermia. Both oocytes and spermatozooids were recovered from individual tanks using a sterilized 60 mL micropipette and then filtered through a 30 µm sieve in order to remove any possible fecal waste that could be attached (mainly to oocytes). Volumes of 4 mL of a homogenized sample of oocytes were counted ($N = 3$) in order to determine the possible correlation between adult size and numbers of oocytes expelled. After that, gametes were introduced together in a 15 L tank favouring the mixture, thus enhancing fertilization.

2.3. Larval cultures

Preliminary essays were conducted to determine the optimal rearing conditions according to survival time observed. To this aim and after fertilization, larvae were maintained in the same tank with constant aeration and temperature (21 °C) until they reached the trochophore stage. Then, they were transferred to different small rearing tanks of 5, 16 and 80 L ($N = 3$) in order to better control the larvae development and cultures were also adjusted at different larval concentrations of 1, 2 and 600 larvae mL⁻¹ respectively, according to Helm et al. (2006) with water renewal fluxes (600 mL·h⁻¹) through a 35 µm strainer and constant aeration. In addition to exhaustive filtration of seawater circuits periodic, bacterial cultures were carried out to monitor the presence of pathogens within tanks. For each concentration and rearing tank, survival time was estimated daily by sampling ($N = 1$) of 4 mL and counting of living larvae.

Larvae diet was established using a mixture of three microalgae (*Isochrysis galbana*, *Pavlova lutheri* and *Chaetoceros calcitrans*), as suggested in the literature (Pernet et al., 2005; Milke et al., 2004, 2006). The final concentration mix of the three phytoplankton species was

adjusted according to three different doses named as “low”, “optimal” and “high” using the formula proposed by Helm et al. (2006) for bivalves feeding in breeding facilities:

$$V_{dose}(L) = \frac{\text{cell density needed} [\mu\text{L}] \cdot V_{tank}}{\text{cell density available} [\mu\text{L}]}$$

where:

V_{dose} = supplied dose in liters.

Cell density needed [μL] = cell concentration according to low, optimal or high dose.

$V_{tank}(L)$ = tank volume.

Cell density available [μL] = cell concentration in laboratory cultures.

For each concentration mix, the equation included the cell density suggested by the author: "15 *Chaetoceros* cells per μL + 25 *Pavlova* cells per μL + 50 cells per μL of *Isochrysis*" for the “low” dose, "25 *Chaetoceros* cells per μL + 33.3 *Pavlova* cells per μL + 100 cells per μL of *Isochrysis*" for the “optimal” dose, and "30 *Chaetoceros* cells per μL + 50 *Pavlova* cells per μL + 150 cells per μL of *Isochrysis*" for the “high” dose.

2.4. Effect of rearing conditions on larval development and settlement

Once previous essays allowed determination of the maximum survival time of larvae under the different parameters established, cultures were also used for testing the influence of light on larvae development due to negative phototactism, as observed for certain bivalve larvae (Raby et al., 1994). To this end, six 16 L tanks were set with open circulation at densities of 2 larvae·mL⁻¹. Three of those tanks were kept at 12/12 h photoperiod whereas the other three tanks were covered with opaque plastic (darkness conditions). Microalgae doses were identically established as “low”, “optimal” and “high” to elucidate if the use of any of them could affect directly the larvae growth. The latter was determined by registering larval length with a “Leica DM2500” microscope. A total of 60 fan mussel larvae were daily placed on a dig dish, and their length measured for average estimations (\pm SD). The process was carried out by triplicate using larvae coming from different spawns.

For all tanks, abiotic parameters were kept as stable as possible. Temperature: 19–21 °C; salinity: 32–37; pH: 7.7–8.4; and dissolved oxygen: 5.9–7.0 mg O₂. Larvae concentrations were daily monitored ($N = 1$ per tank) to determine the conditions with lower mortality rates. Moreover, an artificial substrate made of 250 μm PVC mesh was deployed in all tanks in order to enhance larvae settlement. The substrate was autoclaved (120 °C, 30 min) to prevent the introduction of pathogens within rearing tanks.

2.5. Statistical analyses

All statistical analyses were conducted with SPSS® Statistics 21 program. The possible association between the size of broodstocks and the number of oocytes expelled was studied using the Pearson correlation factor.

The best survival rate was determined studying the effect of the tank volume (fixed factor, three levels) and the larvae density (fixed factor, three levels) and calculated by a two-way factorial ANOVA.

The effect of photoperiod (fixed factor, two levels), food dose (fixed factor, three levels) and rearing day on larvae survival, was investigated using a three-way factorial ANOVA followed by a *post-hoc* analysis (DHS-Tukey) to establish significant groupings. All data were tested for ANOVA assumptions of normality (Levene's test) and homogeneity of variances (Cochran's test).




Fig. 1. Correlation between broodstock size (Ht) and number of oocytes expelled per L. $N = 26$ adults.

3. Results

3.1. Gamete release and fertilization

A total of 31 individuals (47.10 ± 10.61 cm Ht), of the 40 subjected to thermal shock, released gametes for periods of 40 min and, in some instances for up to 3 h. Among individuals that responded positively, 5 of them (16.1%) were strictly males and 14 (45.2%) expelled only oocytes. The remaining 12 specimens (38.7%) released almost simultaneously male and female gametes. The release of female gametes in *P. nobilis* was estimated in $1.9 \cdot 10^6$ oocytes L⁻¹ (averaged for the 26 individuals releasing female gametes) with a mean Ht of 51.8 ± 9.98 cm (Fig. 1).

There was a significant a positive association between the size of individuals and the number of expelled oocytes ($F = 0.765$, $p < 0.01$) as reported for other bivalve species (Helm et al., 2006). The smallest spawning female was 37.7 cm Ht, and 34.6 cm Ht in males. In those instances in which there was only male spawning, sperm was stored in a temperature controlled room at 4 °C and visual observations revealed that sperm remained alive for a maximum of 3 days.

3.2. Larval cultures

Viable oocytes were spherical with an average diameter of $\emptyset = 5.5 \pm 1 \mu\text{m}$ whereas spermatocytes hardly exceed $1 \mu\text{m}$ length. Embryonic development of *P. nobilis* started with the rapid fertilization of oocytes by surrounding spermatocytes at a temperature of 21 °C (Table 2). After 15–30 min the appearance of the first polar body and the formation of a perivitelline membrane confirmed successful fertilization (Fig. 2). The first zygote inclusions were observed after 40 min and gradually increase the number of blastomeres in a successive formation of inclusions until they attained a ciliated blastula stage 5 h later. Herein, the phase in which the motility of larvae begins (Fig. 3) and last for 24 h until the trocophore stage at an average size of $65 \pm 5 \mu\text{m}$. This was followed by a period of frenetic activity where larvae can reach speeds of 0.5 to $1 \text{ cm} \cdot \text{s}^{-1}$ and displayed a helical swimming pattern as observed in other bivalves (Troost et al., 2008). This speed was considerably reduced after the first 48 h when the larvae started to generate their own shell (Prodissoconch I) and become a D-larva or early veliger stage ($85 \pm 3 \mu\text{m}$). Here, the appearance of a ciliary structure or “vellum” produced an incessant movement that generates a current of attraction that allows the capture of phyto-plankton cells. Later, the carbonate shell that protects the visceral

Table 1

Different established doses depending on volume and cell density.

	Low dose (L)	Optimal dose (L)	High dose (L)
Tank 5 L			
<i>Chaetoceros calcitrans</i>	–	0.04	–
<i>Isochrysis galbana</i>	–	0.14	–
<i>Pavlova lutheri</i>	–	0.05	–
Total (L)	–	0.23	–
Tank 16 L			
<i>Chaetoceros calcitrans</i>	0.07	0.11	0.14
<i>Isochrysis galbana</i>	0.23	0.46	0.69
<i>Pavlova lutheri</i>	0.11	0.15	0.23
Total (L)	0.41	0.72	1.05
Tank 80 L			
<i>Chaetoceros calcitrans</i>	–	0.57	–
<i>Isochrysis galbana</i>	–	2.29	–
<i>Pavlova lutheri</i>	–	0.76	–
Total (L)	–	3.62	–

Table 2Time of larvae development in *Pinna nobilis* at 21 °C. pH 8.50. O₂ 6.5 mgO₂L⁻¹ and salinity 38.0 psu.

Stage	Cumulated time (h:min)	Size (µm)
Spawning	0:00	50
Sperm attachment	0:00	50
1st polar body	0:15	50
Double membrane	0:30	50
1st inclusion	0:40	55
Blastule	5:00	55
Gastrule	8:00	55
Early trocophore	22:00	65
Late trocophore	30:00	70
Early veliger	48:00	85
Late veliger	72:00	90
Early umbonate	144:00	100
Pediveliger	168:00	110

cavity of the larvae and the food become more important, reducing cilia movement to an intermittent rotation. Progressively, the larvae secrete more calcium carbonate causing the thickening of shell layers and the development of the first growth rings (Prodissoconch II). Herein, the characteristic straight hinge that gives name to the “D” larva tends to bend thus reaching the umbonate phase. From this moment the larvae stopped swimming completely but few developed the foot that allows benthic settlement. (See Table 1.)

From the spawning to pediveliger stage the average growth of larvae was estimated at 8.57 µm·day⁻¹. However, at some point of the experiments (day 6 in 80 L tanks and day 7 in 16 L tanks) all growth was stopped and no more larval development was observed regardless of the rearing conditions and coinciding with the period when larvae stop swimming (red line) (Fig. 4). Thus, larvae remained alive for a maximum of 22 days and statistical analysis showed that there is a significant difference between survival of larvae and the parameters selected for the rearing activity occurring best survival results in 16 L tanks with an initial larvae density of 2 larvae·mL⁻¹ ($F = 13.542, p < 0.05$).

3.3. Effect of rearing conditions on larval development and settlement

Results from ANOVA showed higher survival of larvae due to the presence of light in the tanks ($F = 4.597, p < 0.05$) and evidenced important differences according to the dose of food provided ($F = 3.434, p < 0.05$) while interaction between “Tank + Dose” was not significant ($F = 2.910, p = 0.58$). Post-hoc analysis showed that the “optimal” dose significantly improved ($p < 0.05$) larval survival compared to “low” and “high” doses where > 80% of larvae had died after 48 h and 96 h, respectively.

The relative daily mortality registered in tanks with photoperiod 12/12 showed mean values of 54.2% when the dose was “low”, 18.3% when the dose was “optimal” and 34.4% when the dose was “high”. In addition, the highest values of mortality were also recorded during mainly the first three days, during the trocophore and veliger phases (Fig. 5).

3.4. Pathologies observed

In most cases typical symptoms of bacterial infection (Fig. 6) are suspected to have caused the observed mortalities over 80% during the first 2–9 days. Herein, the loss of the larval “vellum” was observed during the veliger phase. Despite larvae continue alive, the absence of the ciliary structure prevents them to feed properly and after some days die. Another disease observed in dead larvae is easily identifiable by the continuous movement of the bacteria around and inside larvae shells referred as “swarming”. In some cases, bacterial infection seems to affect larval motility, generating large clusters of larvae referred as “spotting” at the bottom of the tanks due to the secretion of mucous filaments.

4. Discussion

This study reports for the first time the successful spawning and larval rearing to pediveliger stage in captivity conditions of the endangered bivalve *P. nobilis*. This is also the first daily graphic documentation of the early development on this species.

Recent episodes of mass mortality occurred in southwestern Mediterranean coasts for *P. nobilis* accentuates the population regression registered in the last decades and consequently points captivity cultures as a potential solution to restore damaged populations. As described in Trigos (2017) the maintenance of large number of adults in captivity conditions for prolonged periods of time is necessary to obtain enough gametes of both sexes in a species in which the absence of external sexual dimorphism difficult the development of a hatchery protocol.

This study shows how 12 specimens (38.7%) from the total studied, released almost simultaneously male and female gametes, an event so far unknown in this species, which is described as a successive hermaphroditism, as a mechanism to prevent self-fertilization (De Gaulejac et al., 1995a,b). Interestingly, some expelled oocytes were still in a division process, suggesting that internal fertilization may occur as described for other bivalves such as *Ostrea edulis* (Peteiro et al., 2007). This observation coupled with the fact that oocytes present higher density than seawater and tended to sink after being released to the media could be indicate of internal fertilization as a common mechanism enhancing the survival of larvae, but further research is needed to confirm this hypothesis.

Spawned oocytes stayed at the tank bottom until the ciliated blastula stage (5 h post-spawning) when they became motile and swimming activity was observed. Life stages from late embryos to the veliger phase are considered as planktonic and potentially disperse by currents. Once the pediveliger stage was reached, larvae ceased to be planktonic and searched for a suitable substrate to start benthic development. Our study observed up to 22 days to achieve the pediveliger stage then to acquire the capability to get attached. In this context, the hypothesis proposed by De Gaulejac (1989) with a period between 5 and 10 days to settlement phase, contrast with our longer period which might be affected by captivity conditions. This could indicate that artificial conditions can be improved in order to obtain better results.

The massive mortality rates observed (up to 100% at day 4 and day 22, respectively in dark and light cultures), prevented observing more development phases for *P. nobilis* but as indicated by Hernández-Hernández (2000) and Robles-Mungaray (2004) for other species of Pinnidae, simultaneously to foot development, the progresses of the pediveliger phase (110 ± 10 µm size) is characterized by a gradual loss of the “vellum” that gives way to the formation of gills.




Fig. 2. Early developmental stages of *Pinna nobilis*: (A) first polar body (00:15); (B) fertilized oocytes with double membrane (00:30); (C) first inclusion (00:40); (D) first complete division (01:00); (E) 3th and 4th division (03:00); (F) end of cell division phase (04:30). cp. polar body; ep. perivitelline membrane; ma. macromere; mi. micromere.

Subsequently the process of metamorphosis begins with the secretion of new shell from the edge of the Prodissoconch II. Based on the type of growth of other specimens of the same family, the Prodissoconch II should continue its growth in a transverse direction to that previously recorded (Robles-Mungaray, 2004). The appearance of this new structure called dissoconch represents the turning point at which individuals reach the juvenile phase and acquire all adult characteristics such as the typical “pen” shape of the Pinnidae.

In our case, the mortality rates observed at this pediveliger phase were possibly associated to bacterial pathologies which are widespread in hatcheries of commercial species, as well as in regular experimental activities in the laboratory (Andersen et al., 2000; Prado et al., 2016). The infection of larval cultures may be caused by external or horizontal factors such as bacteria escaping the mechanisms of water filtration or other broodstock individuals (Fontanez and Cavanaugh, 2014). The other type of transmission can be vertical when bacterial contamination occurs in gonads and intestinal tracts of broodstock and passes to offspring (Beninger et al., 2003; Prado et al., 2013). These pathologies are thought to mainly affect larvae because they are much more susceptible to bacterial infections than adults (Lambert and Nicolas, 1998). In particular, *Vibrio* species are regarded as central pathogens in larval bivalve cultures (Gómez-León et al., 2008; Elston et al., 2008; Ksaracodi-Watson, 2009) with new species described in the last years (Prado et al., 2005; Dubert et al., 2015). The most common problem arising from *Vibrio* action is the necrosis of soft tissues and ciliary structures (Sugumar et al., 1998; Neo et al., 2011), thus preventing filtration and feeding mechanisms that cause the death of the larvae (Dubert et al., 2016). This disease is easily identifiable by the

continuous movement of the bacteria around and inside larvae shells referred as “swarming” (Beaz-Hidalgo et al., 2010). The loss of the larval “vellum” during the veliger phase was a typical symptom of bacterial infection. Hence, *Vibrio* is suspected to have caused the observed mortalities over 80% during the first 2–9 days, depending on light conditions and food dose. A low dose of phytoplankton could weaken the larvae being consequently more susceptible to infection. By the contrary, the addition of high doses of the microalgae mixture could be responsible of an excess of non-profit food in the tanks, thus enhancing the proliferation of bacteria. The tanks exposed to the darkness registered mortalities higher than 80% on the second day and survival tended to improve with the “high” dose. Therefore, the photoperiod is presented as a limiting factor for the development of *P. nobilis* larvae and the high mortality observed could be explained by the absence of light which move the larvae away from natural conditions. This fact is supposed to stress considerably the larvae being consequently more susceptible to infection. According to our results, *P. nobilis* needs light to complete its larvae cycle in contrast with described by Peharda and Vilibić (2008) who suggested that *P. nobilis* veliger larvae have a negative phototaxis and migrates vertically to deeper waters during daylight and returns to superficial areas at night, as also indicated for other bivalves (Gosling, 2003).

The loss of vellum tissue also affected larval motility, generating large clusters of larvae referred as “spotting” at the bottom of the tanks due to the secretion of mucous filaments by the larval foot that may measure up to one meter in length (Gérard et al., 1989; Bachelet et al., 1992; Rojas et al., 2009). Further studies need be conducted to determine whether these larvae clusters following the secretion of mucous




Fig. 3. Early larval stages in *Pinna nobilis*: (A) early trocophore (22:00); (B) late trocophore (30: 00); (C and D) isometric view of late veliger (72:00); (E and F) early umbonate (144:00); (G) pediveliger (168:00) (H) lateral view of attached larva. ci. cili; fa. Apical flagella; v. vellum; ma. posterior adductor muscle; gd. digestive glandule; u. umbo; p. foot; pl. Prodissoconch I; plI. Prodissoconch II.

filaments are a side effect of the veil loss or a mechanism to improve larval buoyancy and facilitate dispersal as proposed by Beninger et al. (2003) thus, discerning if it is a natural process or a negative consequence of bacterial activity.

In other Pinnidae such as *Atrina maura*, success in larvae rearing has not been achieved despite its commercial interest has prompted the study of aquaculture conditions for more than a decade. Coupled with pathologic problems, the larvae of *A. maura* appear to show a high hydrophobicity which causes the larvae adhesion to the water surface and consequently the death from desiccation and/or starvation (Maeda-Martínez, 2008). According to González-Corona (2003) and Robles-Mungaray (2004) there are still some biological and technical aspects such as the adjustment of larval density or cleaning protocol, that need be optimized in order to reduce mortality and allow the sustainable commercial production of this bivalve. Therefore, there is a lack of empirical knowledge regarding the mechanisms that trigger disease transmission (Arechavala-Lopez et al., 2013), thus, the optimisation in

the hatchery process is accordingly necessary and involves a better understanding of bivalve physiological requirements.

To conclude, this work presents the first detailed information on the biological cycle of *Pinna nobilis* and provides information concerning important variables determining larval mortality and settlement success, such as light conditions or food dose, establishing the bases for the rearing of the endangered fan mussel *P. nobilis* in captivity. Yet, the closure of its biological cycle in captivity appears to be rather difficult since large mortality rates are observed during first days of life (4 to 22 depending on light treatment). Given that our results are conclusive on the suitability of light conditions and “optimal” food doses, the experimental activity should be intended from a pathological approach, considering bacterial infection as one of the main bottlenecks in the rearing of *P. nobilis* larvae and preventing the development of this species at pediveliger stages.




Fig. 4. Larval growth at the different experimental volumes (5, 16, and 80 L). The red line indicates the moment when larvae stop swimming, concurring with the end of growth within 16 and 80 L tanks (arrows). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 5. Daily evolution of larval survival according to photoperiod and phytoplankton dose.





Fig. 6. Common diseases observed in *Pinna nobilis*: (A) loss of vellum structure; (B) bacterial movement inside and around the larvae "swarming"; (C) larvae clusters attached by mucus "spotting".

Acknowledgements

The present study was financed by the Caisse d'Épargne Côte d'Azur. We are also grateful to the research crew of the Institut Océanographique Paul Ricard and the Catholic University of Valencia for their technical support and help collecting and maintaining fan mussels. Special thanks to the reviewers for their constructive and necessary suggestions.

References

- Acarli, S., Loek, A., Acarli, D., 2011. Preliminary spat settlement of fan mussel *Pinna nobilis* Linnaeus, 1758 on a mesh bag collector in Karantina island (eastern Aegean Sea, Turkey). *Presenius Environ. Bull.* 10, 6.
- Andersen, S., Burnell, G., Bergh, Ø., 2000. Flow-through systems for culturing great scallop larvae. *Aquac. Int.* 8 (2–3), 249–257.
- Archavala-Lopez, P., Sanchez-Jerez, P., Bayle-Sempere, J.T., Uglem, I., Mladineo, I., 2013. Reared fish, farmed escapees and wild fish stocks — a triangle of pathogen transmission of concern to Mediterranean aquaculture management. *Aquac. Environ. Interact.* 3, 153–161.
- Arnold, W.S., 2008. Application of Larval Release for Restocking and Stock Enhancement of Coastal Marine Bivalve Populations. *Rev. Fish. Sci.* 16 (1–3), 65–71.
- Bachelet, G., Guillou, J., Labourg, P.J., 1992. Adult larval and juvenile interactions in the suspension-feeding bivalve. *Cerastoderma edule* (L.): field observations and experiments. *Ann. Inst. Oceanogr. Paris* 68, 75–87.
- Barton, A., Waldbusser, G.G., Feely, R.A., Weisberg, S.B., Newton, J.A., Hales, B., Cudd, S., Eudeline, B., Langdon, C.J., Jefferds, I., King, T., Suhrbier, A., McLaughlin, K., 2015. Impacts of coastal acidification on the Pacific northwest shellfish industry and adaptation strategies implemented in response. *Oceanography* 28 (2), 146–159. <http://dx.doi.org/10.5670/oceanog.2015.38>.
- Basso, L., Vázquez-Luis, M., García-march, J.R., Deudero, S., Alvarez, E., Vicente, N., Duarte, C.M., Hendriks, I.E., 2015. The pen shell: a review of population status and recommended research priorities in the Mediterranean Sea. *Adv. Mar. Biol.* (71), 109–160.
- Beaz-Hidalgo, R., Balboa, S., Romalde, J.L., Figueras, M.J., 2010. Diversity and pathogenicity of *Vibrio* species in cultured bivalve molluscs. *Environ. Microbiol. Rep.* 2 (1), 34–43.
- Beninger, P.G., Le Penneç, G., Le Penneç, M., 2003. Demonstration of nutrient pathway from the digestive system to oocytes in the gonad intestinal loop of the scallop *Pecten maximus* L. *Biol. Bull.* 205, 83–92.
- Coutteau, P., Sorgeloos, P., 1992. The use of algal substitutes and the requirement for live algae in the hatchery and nursery rearing of bivalve molluscs: an international survey. *J. Shellfish Res.* 11, 467–476.
- Darriba, S., 2017. First haplosporidan parasite reported infecting a member of the superfamily Pinnoidae (*Pinna nobilis*) during a mortality event in Alicante (Spain, western Mediterranean). *J. Invertebr. Pathol.* 148, 14–19. <http://dx.doi.org/10.1016/j.jip.2017.05.006>.
- De Gaulejac, B., 1989. Ecologie de *Pinna nobilis* (L.). Mollusque eulamellibranche en baie de Calvi. Étude de la coquille larvaire. Étude des possibilités de réimplantation de l'espèce. DEA "Environnement Marin". Université d'Aix-Marseille III (220 pp).
- De Gaulejac, B., Vicente, N., 1990. Ecologie de *Pinna nobilis* (L.) mollusque bivalve sur les côtes de Corse. Essais de transplantation et expériences en milieu contrôlé. *Haliotis* 10, 17.
- De Gaulejac, B., Henry, M., Vicente, N., 1995a. An ultrastructural study of the gametogenesis of the marine bivalve *Pinna nobilis* (Linnaeus 1758) I. Oogenesis. *J. Moll. Stud.* 61, 375–392.
- De Gaulejac, B., Henry, M., Vicente, N., 1995b. An ultrastructural study of the gametogenesis of the marine bivalve *Pinna nobilis* (Linnaeus 1758) II. Spermatogenesis. *J. Moll. Stud.* 61, 393–403.
- Drent, J., 2004. Life history variation of a marine bivalve (*Macoma balthica*) in a changing world (PhD thesis). Rijksuniversiteit Groningen, pp. 95–106.
- Dubert, J., Romalde, J.L., Prado, S., Barja, J.L., 2015. *Vibrio bivalvicida* sp. nov. a novel larval pathogen for bivalve molluscs reared in a hatchery. *Syst. Appl. Microbiol.* <http://dx.doi.org/10.1016/j.syapm.2015.10.006>.
- Dubert, J., Nelson, D.R., Spinard, E.J., Kessler, L., Gomez-chiarri, M., Da Costa, F.D., Prado, S., Barja, J.L., 2016. Following the infection process of Vibriosis in Manila clam (*Ruditapes philippinarum*) larvae through GFP-tagged pathogenic *Vibrio* species. *J. Invertebr. Pathol.* 133, 27–33.
- Ellis, S., 2000. Nursery and grow-out techniques for giant clams (Bivalvia: *Tridacnidae*). Center for Tropical and Subtropical Aquaculture Publication CTSA #143 (99 p). Elston, R., Hasegawa, H., Humphrey, K.L., Polyak, I.K., Häse, C.C., 2008. Re-emergence of *Vibrio tubiashii* in bivalve shellfish aquaculture: severity, environmental drivers, geographic extent and management. *Dis. Aquat. Org.* 82, 119–134. Fontanez, K.M., Cavanaugh, C.M., 2014. Evidence for horizontal transmission from multilocus phylogeny of deep-sea mussel (*Mytilidae*) symbionts. *Environ. Microbiol.* 16 (12), 3608–3621.
- García-March, J.R., 2005. Aportaciones al conocimiento de la biología de *Pinna nobilis* Linneo, 1758 (Mollusca Bivalvia) en el litoral mediterráneo ibérico. Universitat de València. Servei de Publicacions, Valencia. http://www.tesisenxarxa.net/TDX/TDX_UV/TESIS/AVAILABLE/TDX-0628106-132411/garcia.pdf.
- Gazeau, F., Gattuso, J.P., Dawber, C., Pronker, A.E., Peene, F., Peene, J., Heip, C.H.R., Middelburg, J.J., 2010. Effect of ocean acidification on the early life stages of the blue mussel *Mytilus edulis*. *Biogeosciences* 7, 2051–2060.
- Gérard, A., Salaün, M., Tritar, S., 1989. Criteres de compétence des larves à la métamorphose chez *Pecten maximus*. *Haliotis* 19, 373–380.
- Gómez, E.D., Mingo-Licuanan, S.S., Roa-Quiaioit, H.A., 2000. The culture of true giant clam *Tridacna gigas* for conservation in the Philippines. *Moll. Res. Asia* 159–163.
- Gómez-León, J., Villamil, L., Salger, S., Sallum, R., Remacha-Triviño, A., Leavitt, D., Gómez-Chiarri, M., 2008. Survival of eastern oyster *Crassostrea virginica* from three lines following experimental challenge with bacterial pathogens. *Dis. Aquat. Org.* 79, 95–105.
- González-Corona, J., 2003. Estudio de la fisiología reproductiva y gametogénesis del callo de hacha *Atrina maura* (Sowerby, 1835). (Tesis de Maestría. Guaymas. Sonora. México, 74 pp).
- Gosling, E., 2003. Bivalve Molluscs: Biology. Ecology and Culture. Fishing News Books, Oxford (443 pp).
- Guest, J.R., Todd, P.A., Goh, E., Siva, B.S., Reddy, K.P., 2008. Can giant clam (*Tridacna squamosa*) populations be restored on Singapore's heavily impacted coral reefs? *Aquat. Conserv. Mar. Freshwat. Ecosyst.* 18, 570–579.
- Gui, Y., Zamora, L., Dunphy, B.J., Jeffs, A.G., 2016. Evaluation of the formulated diet MySpat for feeding hatchery-reared spat of the green-lipped mussel, *Perna canaliculus* (Gmelin, 1791). *Aquac. Res.* 2016 (47), 3907–3912. <http://dx.doi.org/10.1111/are.12841>.
- Helm, M.M., Bourne, N., Lovatelli, A., 2006. Cultivo de bivalvos en criadero. In: Un manual práctico. vol. 471 FAO, Roma. c. e. F. N. R. (184 pp).
- Hendriks, I.E., Cabanellas-Reboredo, M., Bouma, T.J., Deudero, S., Duarte, C.M., 2011. Seagrass meadows modify drag forces on the shell of the fan mussel *Pinna nobilis*. *Estuar. Coasts* 34, 60–67.
- Hernández-Hernández, O., 2000. Distribución y abundancia de larvas de callo de hacha (Bivalvia: Pinnidae) en el sistema lagunar Corralero-Alotengo (Tesis de Licenciatura en Biología Marina). Universidad del Mar, Puerto Angel Oaxaca México (41 p).
- Katsanevakis, S., 2005. Population ecology of the endangered fan mussel *Pinna nobilis* in a marine lake. *Endanger. Species Res.* 1, 1–9.
- Knauer, J., Southgate, P.C., 1999. A review of the nutritional requirements of bivalves and the development of alternative and artificial diets for bivalve aquaculture. *Rev. Fish. Sci.* 7, 241–280.
- Knop, D., 2009. Riesenmuscheln: Arten und Pflege im Aquarium. Dähne Verlag, Ettlingen (220 pp).
- Knuckey, R.M., Brown, M.R., Barrett, S.M., Hallegraef, G.M., 2002. Isolation of new nanoplanktonic diatom strains and their evaluation as diets for juvenile Pacific oysters (*Crassostrea gigas*). *Aquaculture* 211 (1), 253–274.
- Kurihara, H., Kato, S., Ishimatsu, A., 2007. Effects of increased seawater pCO₂ on early development of the oyster *Crassostrea gigas*. *Aquat. Biol.* 1, 91–98.
- Lambert, C., Nicolas, J.L., 1998. Specific inhibition of chemiluminescent activity by pathogenic *Vibrios* in Hemocytes of two marine bivalves: *Pecten maximus* and *Crassostrea gigas*. *J. Invertebr. Pathol.* 71 (1), 53–63.
- Maeda-Martínez, A.N., 2008. Estado actual del cultivo de bivalvos en México. En A. Lovatelli, A. Farías I. Uriarte (eds). Estado actual del cultivo y manejo de moluscos bivalvos y su proyección futura: factores que afectan su sustentabilidad en América Latina. Taller Técnico Regional de la FAO, 20–24 de agosto de 2007. Puerto Montt, Chile, FAO Actas de Pesca y Acuicultura, No 12. Roma. FAO. pp. 91–100.
- Manuel, J.L., Gallager, S.M., Pearce, C.M., Manning, D.A., O'dor, R.K., 1996. Veligers from different populations of sea scallop *Placopecten magellanicus* have different vertical migration patterns. *Mar. Ecol. Prog. Ser.* 147–163.
- Milke, L.M., Bricelj, V.M., Parrish, C.C., 2004. Growth of postlarval sea scallops, *Placopecten magellanicus*, on microalgal diets, with emphasis on the nutritional role of lipids and fatty acids. *Aquac. Res.* 234, 20.
- Milke, L.M., Bricelj, V.M., Parrish, C.C., 2006. Comparison of early life history stages of the bay scallop, *Argopecten irradians*: effects of microalgal diets on growth and biochemical composition. *Aquac. Res.* 260, 17.
- Neo, M.L., Todd, P.A., Chou, L.M., Teo, S.L.M., 2011. Spawning induction and larval development in the fluted giant clam. *Tridacna squamosa* (Bivalvia: Tridacnidae). *Nat. Sing.* 4, 157–161.
- Nevejan, N., Davis, J., Little, K., Kilona, A., 2007. Use of a formulated diet for mussel spat *Mytilus galloprovincialis* (Lamarck 1819) in a commercial hatchery. *J. Shellfish Res.* 26, 357–363.
- Peharda, M., Vilibić, I., 2008. Modelling the recruitment effect in a small marine protected area: the example of saltwater lakes on the island of Mljet (Adriatic Sea). *Acta Adriat.* 49, 25–35.
- Pérez-Vallazza, C., Álvarez-Vázquez, R., Cardona, L., Pintado, C., Hernández-Brito, J., 2008. Cetacean diversity at the west coast of La Palma Island (Canary Islands). *J. Mar. Biol. Assoc. UK* 88 (06), 1289–1296.
- Pernet, F., Bricelj, V.M., Parrish, C.C., 2005. Effect of varying dietary levels of ω6 polyunsaturated fatty acids during the early ontogeny of the sea scallop, *Placopecten magellanicus*. *J. Exp. Mar. Biol. Ecol.* 327, 18.
- Pernet, F., Lagarde, F., Le Gall, P., Roque D'orbcastel, E., 2014. Associations between farming practices and disease mortality of Pacific oyster *Crassostrea gigas* in a Mediterranean lagoon. *Aquac. Environ. Interact.* 5 (2), 99–106.
- Peteiro, L.G., Filgueira, R., Ayala, A.M., Fernández-Reiriz, M.J., 2007. Ciclo reproductivo de moluscos bivalvos. (IIM.CSIC. NIPO: 251-07-134-3).
- Petersen, A.K., Turchini, G.M., Jahangard, S., Ingram, B.A., Sherman, C.D., 2010. Effects of different dietary microalgae on survival, growth, settlement and fatty acid composition of blue mussel (*Mytilus galloprovincialis*) larvae. *Aquaculture* 309, 115–124.
- Philippart, C.J., Van Aken, H.M., Beukema, J.J., Bos, O.G., Cadee, G.C., Dekker, R., 2003. Climate-related changes in recruitment of the bivalve *Macoma balthica*. *Limnol. Oceanogr.* 48 (6), 2171–2185.
- Ponis, E., Robert, I., Véron, B., Mathieu, M., Robert, R., 2006. New microalgae for the Pacific oyster *Crassostrea gigas* larvae. *Aquaculture* 253, 618–627.

- Prado, S., Romalde, J.L., Montes, J., Barja, J.L., 2005. Pathogenic bacteria isolated from disease outbreaks in shellfish hatcheries. First description of *Vibrio neptunius* as an oyster pathogen. *Dis. Aquat. Org.* 67 (3), 209–215.
- Prado, P., Tomas, F., Pinna, S., Farina, S., Roca, G., Ceccherelli, G., Romero, J., Alcoverro, T., 2012. Habitat and scale shape the demographic fate of the keystone sea urchin *Paracentrotus lividus* in Mediterranean macrophyte communities. *PLoS One* 7, e35170.
- Prado, S., Dubert, J., Da Costa, F., Martínez-Patiño, D., Barja, J.L., 2013. *Vibrios* in hatchery cultures of the razor clam. *Solen marginatus* (Pulteney). *J. Fish Dis.* 37, 209–217.
- Prado, P., Roque, A., Pérez, J., Ibáñez, C., Alcaraz, C., Casals, F., Caiola, N., 2016. Warming and acidification-mediated resilience to bacterial infection determine mortality of early *Ostrea edulis* life stages. *Mar. Ecol. Prog. Ser.* 545, 189–202.
- Rabaoui, L., Belgacem, W., Ismail, D.B., Mansour, L., Tlig-Zouari, S., 2015. Engineering effect of *Pinna nobilis* shells on benthic communities. *Oceanologia* 57 (3), 271–279.
- Raby, D., Lagadeuc, Y., Dodson, J.J., Mingelbier, M., 1994. Relationship between feeding and vertical distribution of bivalve larvae in stratified and mixed waters. *Mar. Ecol. Prog. Ser.* 103, 275–284.
- Ragg, N.L.C., King, N., Watts, E., Morrish, J., 2010. Optimising the delivery of the key dietary diatom *Chaetoceros calcitrans* to intensively cultured Greenshell™ mussel larvae, *Perna canaliculus*. *Aquaculture* 306, 270–280.
- Renault, T., Arzul, I., 2001. Herpes-like virus infections in hatchery-reared bivalve larvae in Europe: specific viral DNA detection by PCR. *J. Fish Dis.* 24 (3), 161–168.
- Rico-Villa, B., Le Coz, J.R., Mingant, C., Robert, R., 2006. Influence of phytoplankton diet mixtures on microalgae consumption, larval development and settlement of the Pacific oyster *Crassostrea gigas* (Thunberg). *Aquaculture* 256 (1), 377–388.
- Robles-Mungaray, M., 2004. Desarrollo de la biotecnología para la producción de semilla en laboratorio. diploide y triploide. de callo de hacha *Atrina maura* (Sowerby. 1835). Universidad Autónoma de Baja California Sur. Baja California Sur. México, pp. 66.
- Rodrigues, L.C., van den Bergh, J.C.J.M., Massa, F., Theodorou, J.A., Ziveri, P., Gazeau, F., 2015. Sensitivity of Mediterranean bivalve Mollusc aquaculture to climate change and ocean acidification: results from a producers' survey. *J. Shellfish Res.* 34 (3), 1161–1176.
- Rojas, R., Miranda, C.D., Amaro, A.M., 2009. Pathogenicity of a highly exopolysaccharide-producing Halomonas strain causing epizootics in larval cultures of the Chilean scallop *Argopecten purpuratus* (Lamarck. 1819). *Microb. Ecol.* 57 (1), 129–139.
- Rojas, R., Miranda, C.D., Romero, J., Asenjo, F., Valderrama, K., Segovia, C., Ugalde, J.A., Santander, J., 2015. Genome sequence of *Vibrio* VPAP30, isolated from an episode of massive mortality of reared larvae of the scallop *Argopecten purpuratus*. *Genome Announc.* 3 (4), e00745–15.
- Ronquillo, J.D., Mckinley, R.S., 2006. Developmental stages and potential mariculture for coastal rehabilitation of endangered Pacific angelwing clam, *Pholas orientalis*. *Aquaculture* 256 (1), 180–191.
- Rouanet, E., Trigos, S., Vicente, N., 2015. From youth to death of old age: the 50-year story of a *Pinna nobilis* fan mussel population at Port-Cros Island Port-Cros National Park, Provence, Mediterranean Sea. *Sci. Rep. Port-Cros Natl. Park* 29, 209–222.
- Rumrill, S.S., 1990. Natural mortality of marine invertebrate larvae. *Ophelia* 21, 163–198.
- Samain, J.F., McCombie, H. (Eds.), 2008. Summer Mortality of Pacific Oyster *Crassostrea gigas*, the Morest Project. Ifremer/Quae Éditions, Versailles.
- Shanks, A.L., Brink, L., 2005. Upwelling, downwelling, and cross-shelf transport of bivalve larvae: test of a hypothesis. *Mar. Ecol. Prog. Ser.* 302, 1–12.
- Su, Z., Huang, L., Yan, Y., Li, H., 2007. The effect of different substrates on pearl oyster *Pinctada martensii* (dunker) larvae settlement. *Aquaculture* 271 (1), 377–383.
- Sugumar, G., Nakai, T., Hirata, Y., Matsubara, D., Muroga, K., 1998. *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis. Aquac. Org.* 33 (2), 111–118.
- Talmage, S.C., Gobler, C.J., 2009. The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and eastern oysters (*Crassostrea virginica*). *Limnol. Oceanogr.* 54, 2072–2080.
- Theodorou, J.A., James, R., Tzovenis, I., Hellio, C., 2015. The recruitment of the endangered fan mussel (*Pinna nobilis*, Linnaeus 1758) on the ropes of a Mediterranean mussel long line farm. *J. Shellfish Res.* 34 (2), 409–414.
- Theodorou, J.A., James, R., Tagalis, D., Tzovenis, I., Hellio, C., Katselis, G., 2017. Density and size structure of the endangered bivalve *Pinna nobilis*, (fan mussel) in the shallow water zone of Maliakos Gulf, Greece. *Acta Adriat.* 58 (581), 63–76.
- Thomas, G.R., Taylor, J., De Leaniz, C.G., 2010. Captive breeding of the endangered freshwater pearl mussel *Margaritifera margaritifera*. *Endanger. Species Res.* 12 (1), 1–9.
- Trigos, S., 2017. Estudio de la ecofisiología y ensayo de cultivo de la nacra *Pinna nobilis* Linnaeus, 1758 (Tesis doctoral). Universidad Católica de Valencia, Valencia, España.
- Trigos, S., Garcia-March, J.R., Vicente, N., Tena, J., Torres, J., 2015. Respiration rates of the fan mussel *Pinna nobilis* at different temperatures. *J. Molluscan Stud.* 81, 217–222. <http://dx.doi.org/10.1093/mollus/eyu07r5>.
- Troost, K., Veldhuizen, R., Stamhuis, E.J., Wolff, W.J., 2008. Can bivalve veligers escape feeding currents of adult bivalves? *J. Exp. Mar. Biol. Ecol.* 358, 185–196.
- Vázquez-Luis, M., Álvarez, E., Barraón, A., García-March, J.R., Grau, A., Hendriks, I.E., Jiménez, S., Kersting, D., Moreno, D., Pérez, M., Ruiz, J.M., Sánchez, J., Villalba, A., Deudero, S., 2017. S.O.S. *Pinna nobilis*: a mass mortality event in western Mediterranean Sea. *Front. Mar. Sci.* 4, 220. <http://dx.doi.org/10.3389/fmars.2017.00220>.
- Vicente, N., 1986. La grande Nacre de Méditerranée. Parc national de Port-Cros. ([plaque de présentation]. 4 volets).
- Vicente, N., Moreteau, V., 1991. Statut de *Pinna nobilis* L. en Méditerranée (Mollusque Eulamellibranche). In: Boudouresque, C.F., Avon, M., Gravez, V. (Eds.), *Les Espèces Marines à Protéger en Méditerranée*. GIS Posidonie, Marseille, pp. 159–168.
- Vincie, M.E., 2008. Development of a Suitable Diet for Endangered Juvenile Oyster Mussels, *Epioblasma capsaeformis* (Bivalvia: Unionidae), Reared in a Captive Environment. Doctoral dissertation, Virginia Tech.