



**HAL**  
open science

## From which level of competition in clubs are adolescents at greater risk of injury compared with outside-of-clubs athletes? A school-based study

Maxime Luiggi, Victoria Rindler, Jean Griffet

### ► To cite this version:

Maxime Luiggi, Victoria Rindler, Jean Griffet. From which level of competition in clubs are adolescents at greater risk of injury compared with outside-of-clubs athletes? A school-based study. *Physician and Sports Medicine*, 2017, 46 (1), pp.66-72. 10.1080/00913847.2018.1402665 . hal-01950366

**HAL Id: hal-01950366**

**<https://amu.hal.science/hal-01950366>**

Submitted on 14 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

This is an Accepted Manuscript of an article published by Taylor & Francis in The Physician and Sportsmedicine on November, 15, 2017, available online:

<http://dx.doi.org/10.1080/00913847.2018.1402665>

Manuscript title: From which level of competition in clubs are adolescents at greater risk of injury compared with outside-of-clubs athletes? A school-based study.

Authors: Maxime Luigi<sup>a</sup>, Victoria Rindler<sup>a</sup>, Jean Griffet<sup>a</sup>

<sup>a</sup>Institute of Movement Sciences, Aix Marseille Univ, CNRS, ISM, Marseille, France

To cite this article: Maxime Luigi, Victoria Rindler & Jean Griffet (2018) From which level of competition in clubs are adolescents at greater risk of injury compared with outside-of-clubs athletes? A school-based study, The Physician and Sportsmedicine, 46(1):66-72, DOI:

<http://dx.doi.org/10.1080/00913847.2018.1402665>

1 **From which level of competition in clubs are adolescents at**  
2 **greater risk of injury compared with outside-of-clubs athletes? A**  
3 **school-based study.**

---

4  
5 **ABSTRACT**

6 **Objectives:** Sport practice is a key factor in a person's physical and mental health but, for  
7 adolescent athletes, some injuries lead to health problems in the long term. The literature  
8 provides multiple factors for understanding injury but does not give information about  
9 injury risk related to each level of play in a large sample of multisport athletes. This study  
10 investigates this relationship in 14- to 19-year-old adolescents.

11 **Methods:** The survey on adolescents and health was conducted in classrooms of France,  
12 from February to March 2015. Only sports players were included in the analyses (n =  
13 986). The levels of play were divided into five categories: Outside of a club/no  
14 competition, club player/no competition, club player/local level, club player/state level  
15 and club player/national and higher level. A three-step binary logistic regression analysis  
16 with age, sex, type of sport, weekly hours of exposure, and level of play was used.

17 **Results:** During the past year, 48.1% of the adolescents were injured. Age and sex were  
18 not risk factors. The injury risk associated with the increases in level of play is higher than  
19 those related to the hours of exposure per week or the type of sport. In clubs, adolescents  
20 who do not compete or play at a local level showed no evidence of greater injury risk  
21 whereas state-level and national- and higher-level athletes were at greater risk than  
22 outside-of-club players (OR= 2.18, 95%CI= 1.13-3.94 and OR= 3.89, 95%CI= 2.07-7.31,  
23 respectively).

24 **Conclusion:** Adolescents who play sports in clubs are clearly more exposed to injury than  
25 those who play outside of a club, mainly from state level. Age and sex are not related to

1 injury. Future epidemiological studies should control adolescents' level of play. Special  
2 attention should be accorded to the injury risk of athletes playing at these levels of  
3 competition.

4 **Keywords:** athletic injuries; sports injury; risk factors; injury prevention; public health;  
5 sport level

---

## 6 **Introduction**

7 Sport practice is regarded as a key factor in a person's physical [1] and mental health [2,3]  
8 and as a means to reduce the risk of such diseases as cancer, diabetes type II or osteoporosis  
9 [4]. However, sport practice also involves significant risks. For adolescent athletes, besides  
10 immediate effects of injuries that cause pain, immobilization, and absences from school, some  
11 injuries lead to health problems in the long term [5]. For instance, knee injuries significantly  
12 increase the risk of osteoarthritis in adulthood [6]. To inform public health policies, coaches,  
13 and parents, and to guide future prevention initiatives, students most vulnerable to sports  
14 injuries need to be identified. Multiple factors are related to the occurrence of injuries (social,  
15 psychological, biomechanical, physiological) [7,8]. We can act on some of them (hours of  
16 exposure, muscles, body mass index (BMI), attitude or equipment) to reduce their influence,  
17 while others are non-modifiable because they are specific either to the sport and performance  
18 goals (type of sport, context of practice, level of competition) or to the athlete's biological  
19 characteristics (age, sex). Our study focuses mainly on non-modifiable factors that can be  
20 controlled without medical supervision or additional measurement tools (electromyography,  
21 etc.) and helps to identify groups of adolescents at high risk of injury.

22 Age, sex, and hours of exposure [9] have been found to increase the risk of injury in sports  
23 [10]. Regarding body mass index, longitudinal and retrospective studies have demonstrated  
24 that BMI status does not modify the relation between sport participation and injury risk

1 among adolescents [11,12], except in overweight and obese populations [13-15]. Next to the  
2 greater injury risk in collective sports compared with individual sports [16], retrospective  
3 studies show that the risk of injuries is greater for those playing sports in clubs than outside of  
4 clubs [13, 17, 18]. This difference could be explained by the fact that the level of play and the  
5 related intensity of practice may be higher in clubs. These studies use the club/outside of a  
6 club context variable as an indicator of the athlete's level. However, in clubs, there are also  
7 various contexts (practice and competition) and competition levels (i.e. levels of play) which  
8 have different influences on the risk of injury. For example, in retrospective studies, the  
9 competitive context has been shown to be a risk factor [19]. Similarly, cohort studies of young  
10 high-level athletes and high-school athletes demonstrate that the risk of injuries is greater in  
11 the competitive context than in the practice context. Injury rates are higher [20, 21] and  
12 injuries are more severe [20, 22]. Regarding the influence of competition levels in clubs,  
13 existing studies on gymnastics [23] and soccer [24, 25] have shown that injury risk is higher  
14 for professional and international athletes than for amateur or state and national athletes.

15 Recent studies have shown that adolescents playing sports in clubs were more exposed to  
16 injury than those playing outside clubs [13, 17, 18]. However, to our knowledge, at present no  
17 study has measured, in general adolescent populations, the adolescents' odds of being injured  
18 at each level of competition in clubs compared with adolescents playing sport outside of  
19 clubs. The objective of our study was to determine the risk of injury of adolescent students  
20 related to each level of competition in clubs compared with outside of clubs while considering  
21 potential confounders (age, sex, BMI, type of sport, hours of exposure). Injury prevention is  
22 an important public health issue that should be considered to all adolescents' athletes. In  
23 addition, our findings will help to determine at which level of play a continued attention of the  
24 injury risk should be done.

25

## 1 **Materials and Methods**

### 2 *Participants*

3 Data on French adolescents and their health were collected between February and March 2015  
4 (n = 1343). The study was conducted on junior high-school, senior high-school, and first-year  
5 university students. All the participants responded to the survey under the supervision of a  
6 physical education teacher and a researcher. The study was authorized by the Superintendent  
7 of Schools and approved by the Aix-Marseille University Ethics Committee. Only adolescents  
8 who agreed to participate and returned signed parental informed consent forms, or those aged  
9 18 years and over, answered the survey (response rate: 98.5%). This study included only  
10 sports players (n = 986). To define sport, we use a definition adopted by the Council of  
11 Europe (1992) [26]: ‘Sport means all forms of physical activity which, through casual or  
12 organized participation, aim at expressing or improving physical fitness and mental well-  
13 being, forming social relationships or obtaining results in competition at all levels.’ Following  
14 the questionnaire used by Lamprecht et al. (2015) [27] students were invited to give their  
15 sport participation outside of physical education classes at school. Thus, students were invited  
16 to answer the question: ‘Apart from physical education classes at school, do you practice  
17 sports?’ Those who said ‘yes’ were asked to report the sport they preferred to practice outside  
18 of school and their usual amount of practice per week. Preferred activities were listed in a  
19 separate table and a committee of physical education teachers and researchers in sport  
20 sciences was formed to decide whether or not these activities could be considered sports in  
21 terms of the definition of the Council of Europe (1992) [26]. Students playing their preferred  
22 activity – considered a sport by the committee – and practicing it for one hour or more per  
23 week were considered sport players.

24

## 1 *Instruments/Procedure*

2 Sociodemographic, sport participation, and injury variables were used. Students reported their  
3 age (Mean = 16.83, SD = 1.81), weight, height and sex. BMI was calculated by dividing  
4 weight (kg) by height squared (m<sup>2</sup>). Sports activities preferred by adolescents were classified  
5 under collective sports or individual sports by the same committee of physical education  
6 teachers and researchers in sport sciences. The ‘weekly hours of exposure’ continuous  
7 variable was divided into five equal groups (<= 2 hours, <= 3 hours, <= 5 hours, <= 7 hours,  
8 > 7 hours). Adolescents reported the sport participation context (club/outside of a club) and  
9 the level of competition (no competition, local level, state level, national and higher level). A  
10 ‘level of play’ variable was created by combining the sport participation context and the level  
11 of competition. Five levels of play were developed (Outside of a club, no competition; Club  
12 player, no competition; Club player at local level; Club player at state level; Club player at  
13 national and higher level). Athletes classified under Outside of a club at competitive level  
14 were excluded from the level of play analysis because there were too few of them (n = 33).

15 Injury was defined according to consensus statements on injury definition in soccer [28] and  
16 rugby [29]. Injury was defined as: ‘Any physical complaint which is caused by a transfer of  
17 energy that exceeds the body's ability to maintain its structural and/or functional integrity, that  
18 is sustained by a player during a match or training, irrespective of the need for medical  
19 attention or time-loss from his sport activity. An injury that results in a player receiving  
20 medical attention is referred to as a “medical-attention” injury and an injury that results in a  
21 player being unable to take a full part in future training or match play as a “time-loss” injury.’

22 As in other studies [13], we investigate risk factors for (i) all sport injury (ASI), irrespective  
23 of the need for medical attention or time-loss from adolescents’ activity; (ii) injury treated by  
24 a medical practitioner – medically treated injury (MTI). As did by Backx [19] and many  
25 others studies, information about injuries were self-reported by adolescents. They were asked

1 to report whether they had been injured in the past 12 months during their preferred sport  
2 activity, the numbers of injuries sustained, the nature of the injuries (sprain, overuse, tendon  
3 strain, muscle strain, contusion, fracture, dislocation) and the severity of the most serious one.  
4 For the nature and severity of injury, we used the same questions as the Western Australian  
5 Sports Injury Study (WASIS) [30]. Severity was graded into three levels. A minor injury is an  
6 injury declared as self-treated, or treated by a sports trainer, coach, parent, or similar. A  
7 moderate injury is an injury declared as treated by a medical practitioner. A severe injury is an  
8 injury requiring hospitalization. The validity of nature and severity of injury (minor and  
9 moderate) reported by the WASIS method have been previously assessed on a four-week self-  
10 report sports injury recall against external sources (hospital records, health practitioners'  
11 records, and third parties) [31]. Contrary with initial WASIS method, in this study we used a  
12 one-year self-report recall for injury. Information about the adolescents' sports experiences,  
13 mechanisms of injuries, localization, and context of the emergence (match and training) was  
14 also asked for. Results from the latter data are not reported in this article but will be used in  
15 subsequent papers on injury mechanisms.

### 16 *Data Analysis*

17 Statistical analyses were performed using SPSS 18.0.0. First, we presented in a table sample  
18 distribution by sex, type of sport, hours of exposure per week, and level of play. For each  
19 variable we report the proportions (%) of athletes who have sustained an injury in the past 12  
20 months. Chi<sup>2</sup> statistics were used to measure the effect of each variable on injury. Second, we  
21 reported in a table the nature of the injuries sustained by athletes at each level of competition.  
22 Odds ratios (OR) and their 95% confidence interval were calculated. The reference category  
23 for odds ratio was injury risk for athletes playing sport outside clubs. Third, a three-step  
24 binary logistic regression was developed. Only significant variables at 0.05 during multiple  
25 regressions were included in the final model. The first step of the logistic regression model



1 included the sociodemographic and anthropomorphic variables (sex, age, BMI). The second  
2 step concerned sport-related variables already known to be linked to injury (type of sport and  
3 weekly hours of exposure). The third step dealt with the main objective of this study: to  
4 investigate the link between the athletes' level of play and injury. This model was performed  
5 twice: for ASI, then for MTI. Adjusted OR estimated in these analyses are presented in Table  
6 3 with confidence intervals and p-values and were used to approximate injury risk.

## 7 **Results**

8 The descriptive statistics and injury rates are provided in Table 1. Soccer had the highest  
9 participation rates (15.7%), followed by dance (14.5%), basketball (6.0%), tennis (5.3%),  
10 strength training (4.1%), boxing (4.1%), rugby (3.9%), swimming (3.8%), handball (3.8%),  
11 and horse riding (3.2%). 78 activities were identified. Distribution of the adolescent athletes  
12 by sport are in Additional File 1, Table 1. For information, in Additional File 1, Table 2, we  
13 present sport activities declared by the adolescents which were not considered sports by the  
14 committee.

15 A total of 474 (48.1%) students had suffered at least one sport injury within the past 12  
16 months. In all, 758 injuries had been sustained. The severity was minor for 61.4%, moderate  
17 for 32.3% and severe for 6.3% of the injuries. Table 2 presents the nature of injuries sustained  
18 at each level of play and the OR for each kind of injury compared with athletes playing  
19 outside clubs. Adolescents playing sport in clubs at local level, state level, and national and  
20 higher level had a lower overuse injury risk compared with outside-of-club players (OR = 0.6,  
21 95%CI: 0.44-0.84, OR = 0.6, 95%CI: 0.44-0.80 and OR = 0.9, 95%CI: 0.71-1.25,  
22 respectively). Adolescents playing sport in clubs at state level and national and higher level  
23 had a greater risk of tendon strain and fracture. At state level adolescents were 1.9 times more  
24 likely (95%CI: 1.36-2.66) to report a tendon strain and 13.2 times more likely (95%CI: 1.56-  
25 111.27) to report a fracture compared with outside-of-club players. At national and higher

1 level, adolescents were 2.5 times more likely (95%CI: 1.76-3.50) to report a tendon strain and  
2 14.2 times more likely (95%CI: 1.64-122.71) to report a fracture compared with outside-of-  
3 club players.

4 According to the multiple logistic regressions, BMI was not associated with injury (OR =  
5 0.97, 95%CI = 0.93-1.02). Thus, it was not included in the further models. Table 3  
6 summarizes the results from the three-step binary logistic regression model. For all injuries, in  
7 the first step there was an effect of sex (OR = 1.32, 95%CI: 1.00-1.71). In the second step,  
8 there was no evidence of an effect of sex (OR = 1.02, 95%CI: 0.77-1.36), but there was an  
9 effect of age (OR = 0.91, 95%CI: 0.84-0.98). Regarding the type of sport, collective sports  
10 players are more likely to have sustained an injury than individual sports players (OR = 1.40,  
11 95%CI: 1.03-1.89). The risk increased as the hours of exposure grew. Athletes who practiced  
12 from 3 to more than 7 hours per week were significantly more exposed to the risk of injury  
13 than the lowest-participation group. The highest-participation group had 3.18 times the odds  
14 of sustaining an ASI than the lowest-participation group (95%CI: 2.07-4.91). In the third step,  
15 the risk of injury in collective sports compared with individual sports is the same as in the  
16 second step and the risk of injury related to the hours of exposure decreased compared with  
17 the second step. The athletes practicing between 5 hours to 7 hours per week and more than 7  
18 hours per week have almost the same risk of injury compared with the lowest-participation  
19 group (OR = 2.27, 95%CI: 1.45-3.59 and OR = 2.38, 95%CI: 1.48-3.84, respectively).  
20 Regarding the level of play, the risk increased as this rose. In-club players who do not  
21 compete and local-level competitors had no significant greater risk of ASI than outside-of-  
22 club players (OR = 1.42, 95%CI: 0.96-2.09 and OR = 1.36, 95%CI: 0.88-2.10, respectively).  
23 In contrast, state-level and national- or higher-level players were more likely to have  
24 sustained an injury in the past year (OR = 1.64, 95%CI: 1.05-2.57 and OR = 2.39, 95%CI:  
25 1.44-3.99, respectively).

1 Regarding MTIs, in the first step, there was an effect of age (OR = 1.14, 95%CI: 1.04-1.25)  
2 and sex (OR = 1.49, 95%CI: 1.06-2.09). In the second step, these effects disappeared.  
3 Collective sport players were more likely to have sustained an injury than individual sports  
4 players (OR = 2.07, 95%CI: 1.43-2.98). As the hours of exposure increased, so did the risk.  
5 From 3 hours to more than 7 hours of exposure per week, athletes had 1.9 to 4.0 times the  
6 odds of sustaining an MTI, compared with the lowest-participation group. In the third step,  
7 age and sex did not affect the results, and the risk associated with the increase of hours of  
8 exposure decreased compared with the second step. Adolescents playing less than 7 hours per  
9 week were at no greater risk of MTIs than the lowest-participation group. The risk of injury  
10 was greater for athletes who practiced sports more than 7 hours per week (OR = 2.12, 95%CI:  
11 1.13-3.94). Regarding the level of play, club players at a non-competitive level and at a local-  
12 level had no greater risk of MTIs than outside-of-club players, whereas the odds were 2.18  
13 times greater for state-level players (95%CI: 1.19-3.99) and 3.89 times greater for national-  
14 and higher-level players (95%CI: 2.07-7.31).

15 For the reader's interest, additional logistic regression models with severity of injury are in  
16 Additional Files 2.

## 17 **Discussion**

18 We observed no increased risk with BMI for ASI and MTI. Much of this result is certainly  
19 attributed to the small number of overweight and obese adolescents in our sample (less than  
20 5.0% and consistent with previous French studies, e.g. [32]). Effects of BMI for injury have  
21 been shown only for overweight and obese adolescents [11-15]. There was a minor effect of  
22 age for MTI but not for ASI. Regarding the sex effect, the first step of the logistic regression  
23 demonstrated that boys are more prone to injury than girls. In the second and third step, we  
24 observed an increased risk of ASI and MTI for collective sports compared with individual  
25 sports and with increasing hours of exposure per week and level of play. The sex effect then

1 disappeared. Therefore, the explanatory factors were neither the age nor the athletes' sex but  
2 rather the type of sport, the weekly hours of exposure and the level of play. These results were  
3 consistent with those from scientific observations indicating no sex effect on injuries [9,13].

4 It has already been shown that collective sports put adolescents at a greater risk of injury than  
5 individual sports [16]. Hours of exposure have also long been recognized as a risk factor [9,  
6 10]. In large populations of young athletes, it has also been shown that athletes in clubs are  
7 more exposed to injury than those outside clubs [13, 17, 18]. In addition to previous findings,  
8 according to our results, in a multisport population, the risk of injury in clubs varies across  
9 competitive levels. In clubs, non-competitive and local-level players showed no evidence of  
10 greater risk of injury than outside-of-club athletes. In contrast, exposure to injury in clubs  
11 compared with outside of a club increased from state level. To our knowledge, this is the first  
12 time that, in a large sample of adolescent athletes, injury risk is reported by level of play and  
13 that below state level, playing sport in clubs does not increase injury risk in adolescents  
14 compared with the outside-of-club context. Also, we observed that the introduction of the  
15 level of play in the third steps of the models (ASI and MTI) reduced the intensity of the  
16 relation between hours of exposure and injury observed in the second steps.

17 For ASIs, in the second step, from 3 hours to more than 7 hours of exposure per week,  
18 athletes were at greater risk of injury ranging from 1.9 to 3.2 times more than athletes  
19 practicing sports less than 2 hours per week. In the third step, this effect decreased  
20 substantially. For each group of athletes, the injury risk was lower than in the second step and  
21 athletes practicing more than 7 hours per week were not at great at greater risk compared with  
22 those playing 5 hours per week (approximately 2.3 times more risk of injury than athletes  
23 practicing less than 2 hours per week). Regarding the level of play, adolescents practicing  
24 sports at state or national- and higher-level were more likely to have sustained an injury  
25 (respectively 1.64 and 2.39 times more). As ASIs, for MTIs, in the second step, from 3 hours

1 to more than 7 hours of exposure per week, athletes were at greater risk of injury, ranging  
2 from 1.9 to 4.0 times more than athletes practicing sports less than 2 hours per week. In the  
3 third step, only athletes who practiced sport more than 7 hours per week were more likely to  
4 have sustained an injury than the lowest-participation group. In contrast, risk of injury related  
5 to the level of play was notably higher for state-level and national- and higher-level athletes  
6 (respectively, 2.18 and 3.89 times more likely). This suggests that the increasing injury risk  
7 with increasing hours of exposure observed in the second step is directly related to the level of  
8 play.

9 Therefore, these results show that knowing the number of weekly hours of exposure is not  
10 sufficient for understanding injury. Some scientific studies highlight the importance of  
11 considering the intensity of play [13, 33, 34]. The level of competition has been shown to be  
12 positively related to the intensity of internal and external forces acting on athletes' muscular-  
13 skeletal system (e.g. soccer [35], rugby [36]) and to injury incidence and severity [20, 23, 25].  
14 Some authors suggest that the greater risk may be due to an increase in intensity of play and  
15 contact during the game. Our results could be explained by these previous findings. We  
16 hypothesize that, in competition below state level, the increase in intensity does not affect  
17 adolescents' risk of injury whereas, above that level, it would reach a threshold that puts them  
18 at a greater risk of injury. Moreover, our results showed that adolescents playing at state level  
19 and national level had a higher risk of more serious injury than outside-of-club players. The  
20 positive association between level of play and injury was greater for MTI than for ASI. More  
21 precisely, additional analyses (see File 2) showed that the level of play was not associated  
22 with minor injury risk, whereas a high association was observed for moderate and severe  
23 injury. State level and national- and higher-level athletes reported respectively 2.71 and 5.39  
24 times greater risk of moderate or severe injury compared with outside-of-club athletes. They  
25 are especially at greater risk of fractures. This greater risk could be also explained by a

1 positive relation between level of play and intensity of internal and external forces acting on  
2 athletes' muscular-skeletal system [35-36] and is consistent with the greater incidence of  
3 severe injuries reported in competitive contexts [20, 22]. To sum up, in addition to a greater  
4 injury risk at state level and higher, adolescents were also at a greater risk of severe injuries,  
5 such as fractures. To our knowledge this is the first time that this kind of results is observed in  
6 a large sample of adolescent athletes.

7 The threshold effect observed at state level and above needs to be specifically studied.  
8 Researchers, coaches, parents, physical education teachers, and public health officials should  
9 try to understand this phenomenon and develop specific injury prevention programs [37, 38]  
10 toward adolescents playing sport at state level and higher. These interventions would take into  
11 account the sport that adolescents practice most and the injury pattern associated with it. The  
12 literature has shown that injury incidence and pattern vary greatly among sport activities [39,  
13 40]. The highest injury rates are observed in motorcycling, handball, and skating; and contact  
14 sport players are at greater risk of fractures or dislocation. In other sports, such as climbing,  
15 tendinopathy is the most common injury [41]. Obviously, characteristics of sport activity are  
16 linked to the localization of injury. Public health officials, coaches, as well as parents or  
17 physical education teachers, should consider sports playing by adolescents and could develop  
18 appropriate preventive initiatives. Relevant sport and injury specific prevention programs [36,  
19 37] can be found in sports sciences journals.

## 20 ***Limitations***

21 Variation of injury incidence and pattern among sport activities leads to the major limitations  
22 of our work. The sports included in our sample are heterogeneous, and conclusions would  
23 certainly be different if the sport sample changed. Similar analyses conducted in other  
24 samples should be made. In addition to the increased risk for collective sports compared with  
25 individual sports, another classification exists, based on the probability of contact with the

1 environment or the adversary and the related dangerousness. A classification developed by the  
2 Committee on Sports Medicine and Fitness [42] categorizes sport activities according to this  
3 criterion. Future studies could examine the influence of the level of play on injury while  
4 considering this classification. In our opinion, the absence of consensus with respect to the  
5 classification system to be used demands that they all be taken into account so that the most  
6 relevant one for this study object can be determined. A literature review regarding the links  
7 found between each sport classification and the injuries rates could be useful.

8 Secondly, our results did not bring information about the injury risk for athletes who play  
9 sports outside of clubs and who compete. Our sample number was not sufficient to perform  
10 reliable statistical analyses. Thus, it is impossible to know whether the level of competition is  
11 more determinant than the context of sport participation to understand injury occurrence. As  
12 we observed for athletes who do not compete both in clubs and outside of clubs, it is plausible  
13 that athletes playing outside of clubs and who compete have the same risk of injury as athletes  
14 playing in clubs and who compete. However, it is also possible that the context of sport  
15 participation is more determinant, in other words, that athletes playing outside of clubs and  
16 who compete have a lower injury risk than those playing in clubs who do not compete. No  
17 study provides evidence to answer this question. Further research is needed.

18 The retrospective design is another weakness of our study. Although we used a retrospective  
19 report for injury which has shown good reliability, some authors have pointed out that  
20 retrospective reports of injuries in youth and adolescents were not precise enough for back  
21 pain symptoms [43]. Due to our 12-months recall period for injury, it is possible that some  
22 injuries had been forgotten by the adolescents. Further studies should develop longitudinal  
23 analyses to assess the validity of this result over a long time period. For example, further  
24 studies could ask adolescents every month about the injuries they had sustained in the

1 previous four weeks. These studies would help determine more precisely the adolescents'  
2 injury risk for each level of play.

### 3 **Conclusions**

4 Age and sex are not related to injury. However, the 'level of play' variable, created by  
5 combining the context of sport practice and the level of competition, is relevant to  
6 understanding the occurrence of injuries. This variable as well as the hours of exposure per  
7 week help to identify the most vulnerable populations. Moreover, increases in level of play  
8 put adolescents at greater risk of injury more than increases in hours of play per week. This  
9 effect could be due to the positive association between level of play and intensity.  
10 Adolescents who play sports in clubs are clearly more exposed to injury than those who play  
11 outside of a club mainly from state level. At this level and higher, adolescents were also at  
12 greater risk of fracture. Future epidemiological studies should control adolescent's level of  
13 play. For example, recent studies have shown that the younger and more exclusive the sports  
14 specialization is, the greater the injury risk, irrespective of context of participation, sex, BMI,  
15 or hours of play per week [18, 44, 45]. One could argue that the level of play should mediate  
16 this relation. A high level of specialization without a high level of play could not be  
17 associated with a greater injury risk, whereas the opposite should be true. Future studies could  
18 be developed to test this hypothesis. In a public health perspective, a special attention should  
19 be done regarding the injury risk of adolescents playing sport at state level and higher.

### 20 **References**

- 21 1. Janssen I, LeBlanc AG. Systematic review of the health benefits of physical activity  
22 and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act* 2010;7:1–16.
- 23 2. Eime R, Young J, Harvey J, et al. A systematic review of the psychological and social  
24 benefits of participation in sport for adults: informing development of a conceptual  
25 model of health through sport. *Int J Behav Nutr Phys Act* 2013;10:135.



- 1 3. Laborde S, Dosseville F, Allen MS. Emotional intelligence in sport and exercise: A  
2 systematic review. *Scand J Med Sci Sports* 2016;26(8):862-874.  
3 <http://dx.doi.org/10.1111/sms.12510>
- 4 4. US Department of Health and Human Services 2008. *Physical Activity Guidelines*  
5 *Advisory Committee Report 2008*. Washington DC. Available at:  
6 <http://www.health.gov/paguidelines/Report/>. Accessed January 30, 2017.
- 7 5. Maffulli N, Longo UG, Gougoulas N, Caine D, Denaro V. Sport injuries: a review of  
8 outcomes. *Br Med Bull* 2011;97(1):47–80. <http://dx.doi.org/10.1093/bmb/ldq026>
- 9 6. McNicholas MJ, Rowley DI, McGurty D, Adalberth T, Abdon P, Lindstrand A, et al.  
10 Total meniscectomy in adolescence. A thirty-year follow-up. *J Bone Joint Surg Br*  
11 2000;82(2):217–221.
- 12 7. Meeuwisse WH. Assessing causation in sport injury: a multifactorial model. *Clin J*  
13 *Sport Med* 1994;4:166-170. Available at:  
14 [http://journals.lww.com/cjsportsmed/Abstract/1994/07000/Assessing\\_Causation\\_in\\_S](http://journals.lww.com/cjsportsmed/Abstract/1994/07000/Assessing_Causation_in_Sport_Injury_A.4.aspx)  
15 [port\\_Injury\\_A.4.aspx](http://journals.lww.com/cjsportsmed/Abstract/1994/07000/Assessing_Causation_in_Sport_Injury_A.4.aspx). Accessed June 29, 2017.
- 16 8. Bahr R, Holme I. Risk factors for sports injuries—a methodological approach. *Br J*  
17 *Sports Med* 2003;37(5):384–392. <http://doi.org/10.1136/bjism.37.5.384>
- 18 9. Rose MS, Emery CA, Meeuwisse WH. Sociodemographic Predictors of Sport Injury  
19 in Adolescents. *Med Sci Sports Exerc* 2008 Mar;40(3):444-50.  
20 <http://dx.doi.org/10.1249/MSS.0b013e31815ce61a>
- 21 10. Emery CA. Risk factors for injury in child and adolescent sport: a systematic review  
22 of the literature. *Clin J Sport Med* 2003;13(4):256–268.
- 23 11. Warsh J, Pickett W, Janssen I. Are Overweight and Obese Youth at Increased Risk for  
24 Physical Activity Injuries? *Obes Facts* 2010;3(4):225–230.  
25 <http://dx.doi.org/10.1159/000319322>

- 1 12. Kemler E, Vriend I, Paulis WD, Schoots W, van Middelkoop M, Koes B. Is  
2 overweight a risk factor for sports injuries in children, adolescents, and young adults?  
3 Scand J Med Sci Sports 2015;25(2):259–264. <http://dx.doi.org/10.1111/sms.12180>
- 4 13. Richmond SA, Kang J, Emery CA. Is body mass index a risk factor for sport injury in  
5 adolescents? J Sci Med Sport 2013;16(5):401–405.  
6 <http://dx.doi.org/10.1016/j.jsams.2012.11.898>
- 7 14. Jespersen E, Verhagen E, Holst R, Klakk H, Heidemann M, Rexen CT, Franz C,  
8 Wedderkopp N: Total body fat percentage and body mass index and the association  
9 with lower extremity injuries in children: a 2.5-year longitudinal study. Br J Sports  
10 Med 2014;48(20):1497-1502.
- 11 15. Yard E, Comstock D. Injury patterns by body mass index in US high school athletes. J  
12 Phy Act Health 2011;8:182-191.
- 13 16. Theisen D, Frisch A, Malisoux L, Urhausen A, Croisier JL, Seil R. Injury risk is  
14 different in team and individual youth sport. J Sci Med Sport 2013;16(3):200–204.  
15 <http://dx.doi.org/10.1016/j.jsams.2012.07.007>
- 16 17. Mattila VM, Parkkari J, Koivusilta L, Kannus P, Rimpelä A. Participation in sports  
17 clubs is a strong predictor of injury hospitalization: a prospective cohort study. Scand  
18 J Med Sci Sports 2009;19(2):267-273. [http://dx.doi.org/10.1111/j.1600-  
19 0838.2008.00800.x](http://dx.doi.org/10.1111/j.1600-0838.2008.00800.x)
- 20 18. Jayanthi N, LaBella CR, Fischer D, Dugas L. Sports-specialized intensive training and  
21 the risk of injury in young athletes: a clinical case-control study. Am J Sports Med  
22 2015;43(4):794-801.
- 23 19. Backx FJG, Erich WB, Kemper AB, Verbeek AL. Sports injuries in school-aged  
24 children. An epidemiologic study. Am J Sports Med 1989;17(2):234-240.

- 1 20. Rechel JA, Yard EE, Comstock RD. An epidemiologic comparison of high school  
2 sports injuries sustained in practice and competition. *J Athl Train* 2008;43(2):197.  
3 <http://dx.doi.org/10.4085/1062-6050-43.2.197>
- 4 21. Malisoux L, Frisch A, Urhausena A, Seil R, Theisen D. Monitoring of sport  
5 participation and injury risk in young athletes. *J Sci Med Sport* 2013;16(6):504-508.  
6 <http://dx.doi.org/10.1016/j.jsams.2013.01.008>
- 7 22. Darrow CJ, Collins CL, Yard EE, Comstock RD. Epidemiology of Severe Injuries  
8 Among United States High School Athletes: 2005-2007. *Am J Sports Med*  
9 2009;37(9):1798–1805. <http://dx.doi.org/10.1177/0363546509333015>
- 10 23. Kolt GS, Kirkby RJ. Epidemiology of injury in elite and subelite female gymnasts: a  
11 comparison of retrospective and prospective findings. *Br J Sports Med*  
12 1999;33(5):312–318.
- 13 24. Kucera KL, Marshall SW, Kirkendall DT, Marchak PM, Garrett WE. Injury history as  
14 a risk factor for incident injury in youth soccer. *Br J Sports Med* 2005;39(7):462–462.  
15 <http://dx.doi.org/10.1136/bjism.2004.013672>
- 16 25. van Beijsterveldt AMC, Stubbe JH, Schmikli SL, van de Port IG, Backx FJG.  
17 Differences in injury risk and characteristics between Dutch amateur and professional  
18 soccer players. *J Sci Med Sport* 2015;18(2):145-149.  
19 <http://dx.doi.org/10.1016/j.jsams.2014.02.004>
- 20 26. Council of Europe (1992) European Sport Charter (revised 2001). Strasbourg: Council  
21 of Europe. 2001.  
22 [https://search.coe.int/cm/Pages/result\\_details.aspx?ObjectID=09000016804c9dbb](https://search.coe.int/cm/Pages/result_details.aspx?ObjectID=09000016804c9dbb).  
23 [Accessed June 26, 2017.](#)
- 24 27. Lamprecht M, Fischer A, Wiegand D, Stamm HP (2015). Sport Suisse 2014: Rapport  
25 sur les enfants et les adolescents. Macolin: Office fédéral du sport OFSPO.

- 1 28. Fuller CW, Ekstrand J, Junge A, Andersen TE, Bahr R, Dvorak J, et al. Consensus  
2 statement on injury definitions and data collection procedures in studies of football  
3 (soccer) injuries. *Scand J Med Sci Sports* 2006;16(2):83-92.  
4 <http://dx.doi.org/10.1111/j.1600-0838.2006.00528.x>
- 5 29. Fuller CW, Molloy MG, Bagate C, Bahr R, Brooks JHM, Donson H, et al. Consensus  
6 statement on injury definitions and data collection procedures for studies of injuries in  
7 rugby union. *Br J Sports Med* 2007;41(5):328-331.  
8 <http://dx.doi.org/10.1136/bjism.2006.033282>
- 9 30. Stevenson MR, Hamer P, Finch CF, Elliot B, Kresnow M. Sport, age, and sex specific  
10 incidence of sports injuries in Western Australia. *Br J Sports Med* 2000;34:188-194.
- 11 31. Valuri G, Stevenson M, Finch C, Hamer P, Elliott B. The validity of a four week self-  
12 recall of sports injuries. *Injury Prevention* 2005;11(3):135-137.  
13 <http://dx.doi.org/10.1136/ip.2003.004820>
- 14 32. Thibault H, Contrand B, Saubusse E, Baine M, Maurice-Tison S. Risk factors for  
15 overweight and obesity in French adolescents: Physical activity, sedentary behavior  
16 and parental characteristics. *Nutrition* 2010;26(2):192-200.  
17 <http://dx.doi.org/10.1016/j.nut.2009.03.015>
- 18 33. Gabbett JT. Influence of training and match intensity on injuries in rugby league. *J*  
19 *Sports Sci* 2004;22(5):409-417. <http://dx.doi.org/10.1080/02640410310001641638>
- 20 34. Vetter RE, Symonds ML. Correlations between injury, training intensity, and physical  
21 and mental exhaustion among college athletes. *J Strength Cond Res* 2010;24(3):587-  
22 96. <http://dx.doi.org/10.1519/JSC.0b013e3181c7c2eb>
- 23 35. Hölmich P, Larsen K, Krogsgaard K, Gluud C. Exercise program for prevention of  
24 groin pain in football players: a cluster-randomized trial. *Scand J Med Sci Sports*  
25 2010;20(6):814-821. <http://dx.doi.org/10.1111/j.1600-0838.2009.00998.x>

- 1 36. Preatoni E, Stokes KA, England ME, Trewartha G. The influence of playing level on  
2 the biomechanical demands experienced by rugby union forwards during machine  
3 scrummaging. *Scand J Med Sci Sports* 2013;23(3):e178–e184.  
4 <http://dx.doi.org/10.1111/sms.12048>
- 5 37. Abernethy L, Bleakley C. Strategies to prevent injury in adolescent sport: a systematic  
6 review. *Br J Sports Med* 2007;41(10):627–638.  
7 <http://dx.doi.org/10.1136/bjism.2007.035691>
- 8 38. Caine D, Maffulli N, Caine C. Epidemiology of injury in child and adolescent sports:  
9 Injury rates, risk factors, and prevention. *Clinics Sp Med* 2008;27(1):19-50.  
10 <http://dx.doi.org/10.1016/j.csm.2007.10.008>
- 11 39. Åman M, Forssblad M, Henriksson-Larsén K. Incidence and severity of reported acute  
12 sports injuries in 35 sports using insurance registry data: Incidence of acute sports  
13 injuries. *Scand J Med Sci Sports* 2015;26(4):451-462.  
14 <http://dx.doi.org/10.1111/sms.12462>
- 15 40. Hootman JM, Dick R, Agel J. Epidemiology of Collegiate Injuries for 15 Sports:  
16 Summary and Recommendations for Injury Prevention Initiatives. *J Athl Train*  
17 2007;42(2):311–319
- 18 41. Jones G, Asghar A, Llewellyn DJ. The epidemiology of rock-climbing injuries. *Br J*  
19 *Sports Med* 2008;42:773-778. <http://dx.doi.org/10.1136/bjism.2007.037978>
- 20 42. Committee on Sports Medicine and Fitness. Medical Conditions Affecting Sports  
21 Participation. *Pediatrics* 2001;107(5):1205. <http://dx.doi.org/10.1542/peds.107.5.1205>
- 22 43. Johansen B, Wedderkopp N: Comparison between data obtained through real-time  
23 data capture by SMS and a retrospective telephone interview. *Chiropr Osteopat* 2010;  
24 18:10.

- 1 44. Lakomkin N, Sugimoto D, Tepolt F, Stracciolini A. Youth sports specialization and  
2 musculoskeletal injury: a systematic review of the literature. *Phys Sportsmed*  
3 2016;44(3):257-262. <http://dx.doi.org/10.1080/00913847.2016.1177476>
- 4 45. Post E, Trigsted S, Riekena JW, Hetzel S, McGuine TA, Brooks A, et al. The  
5 association of sport specialization and training volume with injury history in youth  
6 athletes. *Am J Sports Med* 2017;45(2):1405-1412.  
7 <http://dx.doi.org/10.1177/0363546517690848>

ACCEPTED MANUSCRIPT

**Table 1** Description variables

Variables	n	%	Sports Injury % (n)	
<i>Sex*</i>				
Girls	448	45.4%	44.2%	(198)
Boys	538	54.6%	51.3%	(276)
<i>Type of sport***</i>				
Individual	668	67.7%	43.9%	(293)
Collective	318	32.3%	56.9%	(181)
<i>Hours of exposure per week***</i>				
<= 2	252	25.6%	34.9%	(88)
<= 3	156	15.8%	40.4%	(63)
<= 5	241	24.4%	49.0%	(118)
<= 7	168	17.0%	59.5%	(100)
> 7	169	17.1%	62.1%	(105)
<i>Level of play***</i>				
Outside of a club, no competition	199	20.2%	35.7%	(71)
Club player, no competition	280	28.4%	42.5%	(119)
Club player at local level	173	17.5%	50.3%	(87)
Club player at state level	176	17.8%	57.4%	(101)
Club player at national and higher level	125	12.7%	66.4%	(83)
Other	33	3.3%		
<i>Had at least one injury in the past 12 months</i>				
No	512	51.9%		
Yes	474	48.1%		

\* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$

1  
2  
3

**Table 2** Nature of injuries sustained according to level during the past 12 months and odds ratio (OR) with 95% confidence interval (95%CI)

All sport	Outside of a club, no competition			Club player, no competition			Club player at local level			Club player at state level			Club player at national and higher level			All
N	199			280			173			176			125			953
N athletes injured	71			119			87			101			83			461
% injured	35.7%			42.5%			50.3%			57.4%			66.4%			48.4%
	% (n)	OR	95%CI	% (n)	OR	95%CI	% (n)	OR	95%CI	% (n)	OR	95%CI	% (n)	OR	95%CI	% (n)
Sprain	30.8% (33)	1.0		25.3% (45)	0.7	(0.58-0.84)	30.1% (37)	0.9	(0.70-1.04)	32.7% (55)	1.4	(1.14-1.66)	26.6% (42)	1.2	(0.96-1.45)	28.9% (212)
Overuse	15.9% (17)	1.0		<b>19.1% (34)</b>	<b>1.3</b>	<b>(1.01-1.6)</b>	<b>11.4% (14)</b>	<b>0.6</b>	<b>(0.44-0.84)</b>	<b>9.5% (16)</b>	<b>0.6</b>	<b>(0.44-0.8)</b>	12.0% (19)	0.9	(0.71-1.25)	13.6% (100)
Tendon strain	9.3% (10)	1.0		8.4% (15)	0.9	(0.6-1.28)	7.3% (9)	0.7	(0.44-1.13)	<b>14.3% (24)</b>	<b>1.9</b>	<b>(1.36-2.66)</b>	<b>15.2% (24)</b>	<b>2.5</b>	<b>(1.76-3.50)</b>	11.2% (82)
Muscle strain	17.8% (19)	1.0		10.7% (19)	0.5	(0.4-0.68)	17.1% (21)	0.9	(0.67-1.13)	14.9% (25)	0.9	(0.70-1.15)	17.1% (27)	1.3	(1.03-1.69)	15.1% (111)
Contusion	22.4% (24)	1.0		29.8% (53)	1.6	(1.3-1.9)	25.2% (31)	1.1	(0.87-1.35)	14.3% (24)	0.6	(0.48-0.77)	16.5% (26)	0.9	(0.71-1.13)	21.5% (158)
Fracture	0.9% (1)	1.0		5.1% (9)	5.7	(0.62-52.92)	7.3% (9)	8.1	(0.87-75.18)	<b>9.5% (16)</b>	<b>13.2</b>	<b>(1.56-111.27)</b>	<b>8.9% (14)</b>	<b>14.2</b>	<b>(1.64-122.71)</b>	6.7% (49)
Dislocation	2.8% (3)	1.0		1.7% (3)	0.6	(0.15-2.27)	1.6% (2)	0.5	(0.10-2.88)	4.8% (8)	1.9	(0.76-5.03)	3.8% (6)	1.8	(0.63-4.97)	3.0% (22)
N injuries	107			178			123			168			158			734



**Table 3** Binary logistic regression model and injury

Step	Type	Predictors	All Sports Injury			Medically Treated Sports Injury		
			OR	95%CI	R <sup>2</sup>	OR	95%CI	R <sup>2</sup>
1	<i>Sociodemographic predictor</i>				0.01			0.02
		Age	0.96	[0.89; 1.03]		1.14**	[1.04; 1.25]	
		Sex <sup>a</sup>	1.32*	[1.02; 1.71]		1.49*	[1.06; 2.09]	
2	<i>Sociodemographic predictor</i>				0.08			0.11
		Age	0.91**	[0.84; 0.98]		1.07	[0.97; 1.17]	
		Sex <sup>a</sup>	1.02	[0.77; 1.36]		0.99	[0.68; 1.43]	
	<i>Type of sport</i>							
		Collective <sup>b</sup>	1.40*	[1.03; 1.89]		2.07***	[1.43; 2.98]	
	<i>Hours of exposure per week<sup>c</sup></i>							
		<= 3	1.20	[0.79; 1.83]		1.10	[0.56; 2.16]	
		<= 5	1.85***	[1.28; 2.69]		1.88*	[1.08; 3.26]	
		<= 7	2.74***	[1.79; 4.19]		2.72***	[1.53; 4.81]	
		> 7	3.18***	[2.07; 4.91]		4.03***	[2.30; 7.06]	
3	<i>Sociodemographic predictor</i>				0.09			0.15
		Age	0.92*	[0.85; 0.99]		1.09	[0.99; 1.20]	
		Sex <sup>a</sup>	1.05	[0.79; 1.40]		1.00	[0.68; 1.46]	
	<i>Type of sport</i>							
		Collective <sup>b</sup>	1.40*	[1.01; 1.92]		1.93***	[1.30; 2.86]	
	<i>Hours of exposure per week<sup>b</sup></i>							
		<= 3	1.15	[0.75; 1.75]		0.97	[0.49; 1.94]	
		<= 5	1.71**	[1.17; 2.51]		1.47	[0.83; 2.61]	
		<= 7	2.27***	[1.45; 3.56]		1.72	[0.93; 3.17]	
		> 7	2.38***	[1.48; 3.84]		2.12*	[1.13; 3.94]	
	<i>Level of play<sup>c</sup></i>							
		Club no competitor	1.42	[0.96; 2.09]		1.01	[0.54; 1.88]	
		Club local level	1.36	[0.88; 2.10]		1.56	[0.84; 2.92]	
		Club state level	1.64*	[1.05; 2.57]		2.18**	[1.19; 3.99]	
		Club national and higher level	2.39***	[1.44; 3.99]		3.89***	[2.07; 7.31]	

Note. OR = odds ratio; 95% CI = 95% Confidence Interval; <sup>a</sup>The reference category is *girls*; <sup>b</sup>The reference category is individual sport; <sup>c</sup>The reference category is *less than 2 hours*; <sup>d</sup>The reference category is *unorganized no competitor*; \* $p < .05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ .

1

2

**Additional file 1. Table 1.**

List of preferred sport activities practiced by adolescents and considered as sport according to the Council of Europe definition

SPORT	N	%	Type of sport
AEROBIC	3	0,3%	INDIVIDUAL
AIKIDO	4	0,4%	INDIVIDUAL
AIRSOFT	1	0,1%	COLLECTIVE
AMERICAN FOOTBALL	3	0,3%	COLLECTIVE
ARTISTIC GYMNASTIC	6	0,6%	INDIVIDUAL
ATHLETICS	25	2,5%	INDIVIDUAL
BADMINTON	31	3,1%	INDIVIDUAL
BALLET DANCE	5	0,5%	INDIVIDUAL
BASEBALL	1	0,1%	COLLECTIVE
BASKET-BALL	59	6,0%	COLLECTIVE
BIKING	6	0,6%	INDIVIDUAL
BMX	5	0,5%	INDIVIDUAL
BMX BICROSS	1	0,1%	INDIVIDUAL
BOXING	40	4,1%	INDIVIDUAL
CLIMBING	15	1,5%	INDIVIDUAL
CROSS-COUNTRY CYCLING	10	1,0%	INDIVIDUAL
CYCLING	5	0,5%	INDIVIDUAL
DANCING	124	12,6%	INDIVIDUAL
FENCING	3	0,3%	INDIVIDUAL
FIGURE SKATING	1	0,1%	INDIVIDUAL
FITNESS	12	1,2%	INDIVIDUAL
FRENCH BOXING	8	0,8%	INDIVIDUAL
GOLF	2	0,2%	INDIVIDUAL
GYMNASTIC	22	2,2%	INDIVIDUAL
HANDBALL	37	3,8%	COLLECTIVE
HIP-HOP DANCE	9	0,9%	INDIVIDUAL
HORSE RIDING	32	3,2%	INDIVIDUAL
ICE HOCKEY	2	0,2%	COLLECTIVE
ICE SKATING	1	0,1%	INDIVIDUAL
JUDO	16	1,6%	INDIVIDUAL
JUJITSU	3	0,3%	INDIVIDUAL
KARATE	13	1,3%	INDIVIDUAL
KAYAK	1	0,1%	INDIVIDUAL
KENDO	1	0,1%	INDIVIDUAL
KRAV MAGA	2	0,2%	INDIVIDUAL
KUNG FU	1	0,1%	INDIVIDUAL
LONGBOARDING	2	0,2%	INDIVIDUAL
MARTIAL ARTS	3	0,3%	INDIVIDUAL
MIXED-MARTIAL ARTS	1	0,1%	INDIVIDUAL
MODERN JAZZ DANCE	3	0,3%	INDIVIDUAL
MOTOCROSS RIDING	4	0,4%	INDIVIDUAL
ORIENTAL DANCE	1	0,1%	INDIVIDUAL
ORIENTEERING RACE	3	0,3%	INDIVIDUAL
PARKOUR	3	0,3%	INDIVIDUAL

RESCUE SWIMMING	1	0,1%	INDIVIDUAL
ROLLER DANCE	1	0,1%	INDIVIDUAL
ROLLER HOCKEY	1	0,1%	COLLECTIVE
ROLLER SKATING	1	0,1%	INDIVIDUAL
ROWING	2	0,2%	INDIVIDUAL
RUGBY FOOTBALL	38	3,9%	COLLECTIVE
RUNNING	29	2,9%	INDIVIDUAL
RYTHMIC GYMNASTIC	3	0,3%	INDIVIDUAL
SAILING	1	0,1%	INDIVIDUAL
SAMBO BOXING	1	0,1%	INDIVIDUAL
SCUBA DIVING	1	0,1%	INDIVIDUAL
SELF DEFENSE	3	0,3%	INDIVIDUAL
SHOOTING SPORT	1	0,1%	INDIVIDUAL
SKATEBOARDING	5	0,5%	INDIVIDUAL
SKIING	9	0,9%	INDIVIDUAL
SOCCER	155	15,7%	COLLECTIVE
SQUASH	5	0,5%	INDIVIDUAL
STRENGTH TRAINING	40	4,1%	INDIVIDUAL
SURFING	9	0,9%	INDIVIDUAL
SWIMMING	37	3,8%	INDIVIDUAL
SYNCHRONIZED SWIMMING	4	0,4%	COLLECTIVE
TABLE TENNIS	2	0,2%	INDIVIDUAL
TAEKWONDO	5	0,5%	INDIVIDUAL
TENNIS	52	5,3%	INDIVIDUAL
THAI BOXING	9	0,9%	INDIVIDUAL
TRAIL RUNNING	1	0,1%	INDIVIDUAL
TRAMPOLINING	2	0,2%	INDIVIDUAL
TRIATHLON	2	0,2%	INDIVIDUAL
VOLLEY-BALL	21	2,1%	COLLECTIVE
WINDSURFING	8	0,8%	INDIVIDUAL
YOGA	2	0,2%	INDIVIDUAL
YOSEIKAN BUDO	1	0,1%	INDIVIDUAL
ZUMBA DANCE	4	0,4%	INDIVIDUAL
<b>All</b>	<b>986</b>	<b>1</b>	

**Additional file 2** Binary logistic regression model and injury severity

Step	Type	Predictors	Minor <sup>w</sup>			Moderate <sup>x</sup>			Severe <sup>y</sup>			Moderate+Severe <sup>z</sup>		
			OR	95% CI	R <sup>2</sup>	OR	95% CI	R <sup>2</sup>	OR	95% CI	R <sup>2</sup>	OR	95% CI	R <sup>2</sup>
1	<i>Sociodemographic predictor</i>				0.01			0.06			0.18			0.09
		Age	1.04	[0.95 ; 1.14]		1.28***	[1.13 ; 1.44]		2.08***	[1.49 ; 2.89]		1.34***	[1.19 ; 1.50]	
		Sex <sup>a</sup>	1.30	[0.96 ; 1.77]		1.68**	[1.13 ; 2.49]		2.25	[0.97 ; 5.23]		1.76**	[1.21 ; 2.56]	
2	<i>Sociodemographic predictor</i>				0.08			0.27			0.37			0.30
		Age	1.02	[0.93 ; 1.12]		1.21**	[1.06 ; 1.37]		1.99***	[1.42 ; 2.78]		1.25***	[1.11 ; 1.42]	
		Sex <sup>a</sup>	1.04	[0.74 ; 1.46]		0.82	[0.51 ; 1.32]		1.09	[0.40 ; 2.94]		0.89	[0.57 ; 1.40]	
	<i>Type of sport</i>													
		Collective <sup>b</sup>	1.17	[0.79 ; 1.72]		2.97***	[1.82 ; 4.83]		1.58	[0.61 ; 4.07]		2.70***	[1.70 ; 4.28]	
	<i>Hours of exposure per week<sup>c</sup></i>													
		<= 3	1.15	[0.72 ; 1.84]		1.03	[0.50 ; 2.13]		1.30	[0.08 ; 21.71]		1.07	[0.53 ; 2.16]	
		<= 5	1.82**	[1.18 ; 2.79]		2.05*	[1.11 ; 3.82]		12.67*	[1.55 ; 103.72]		2.49**	[1.38 ; 4.50]	
		<= 7	2.56***	[1.54 ; 4.23]		3.42***	[1.76 ; 6.63]		18.53**	[2.16 ; 158.68]		4.09***	[2.16 ; 7.73]	
		> 7	4.87***	[2.63 ; 9.01]		11.69***	[5.64 ; 24.21]		61.30***	[6.95 ; 541.01]		13.09***	[6.45 ; 26.56]	
3	<i>Sociodemographic predictor</i>				0.09			0.32			0.41			0.35
		Age	1.03	[0.93 ; 1.13]		1.24**	[1.09 ; 1.42]		2.02***	[1.42 ; 2.88]		1.29***	[1.13 ; 1.46]	
		Sex <sup>a</sup>	1.08	[0.76 ; 1.53]		0.90	[0.55 ; 1.48]		1.24	[0.44 ; 3.53]		0.97	[0.61 ; 1.55]	
	<i>Type of sport</i>													
		Collective <sup>b</sup>	1.17	[0.78 ; 1.76]		2.64***	[1.57 ; 4.44]		1.15	[0.41 ; 3.21]		2.36***	[1.45 ; 3.86]	
	<i>Hours of exposure per week<sup>b</sup></i>													
		<= 3	1.09	[0.68 ; 1.75]		0.89	[0.42 ; 1.88]		1.04	[0.06 ; 17.86]		0.93	[0.45 ; 1.93]	
		<= 5	1.70*	[1.09 ; 2.66]		1.59	[0.83 ; 3.06]		7.76	[0.89 ; 67.85]		1.92*	[1.03 ; 3.58]	
		<= 7	2.22**	[1.30 ; 3.79]		1.94	[0.94 ; 4.02]		9.13	[0.98 ; 85.58]		2.33*	[1.16 ; 4.68]	
		> 7	4.18***	[2.19 ; 7.99]		5.33***	[2.36 ; 12.04]		23.68**	[2.35 ; 238.52]		5.90***	[2.69 ; 12.96]	
	<i>Level of play<sup>c</sup></i>													
		Club no competitor	1.43	[0.91 ; 2.24]		1.18	[0.58 ; 2.39]		0.58	[0.09 ; 3.97]		1.10	[0.56 ; 2.16]	
		Club local level	1.32	[0.79 ; 2.23]		1.71	[0.80 ; 3.63]		2.36	[0.49 ; 11.39]		1.81	[0.90 ; 3.66]	
		Club state level	1.65	[0.94 ; 2.88]		2.71**	[1.29 ; 5.69]		3.51	[0.73 ; 16.78]		2.71**	[1.34 ; 5.46]	
		Club national and higher level	1.81	[0.92 ; 3.55]		5.38***	[2.37 ; 12.19]		3.99	[0.74 ; 21.68]		5.39***	[2.48 ; 11.73]	

---

Note. OR = odds-ratio; 95% CI = 95% Confidence Interval; <sup>a</sup>The reference category is *girls*; <sup>b</sup>The reference category is individual sport; <sup>c</sup>The reference category is *less than 2 hours*; <sup>d</sup>The reference category is *unorganized no competitor*; \* $p < .05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ; <sup>w</sup>At least one minor injury (1)/No injury (0); <sup>x</sup>At least one moderate injury (1)/No injury (0); <sup>y</sup>At least one severe injury (1)/No injury (0); <sup>z</sup>At least one moderate or severe injury (1)/No injury (0)

1

ACCEPTED MANUSCRIPT