



HAL
open science

Test of chlordecone dechlorination by chemical reduction using Fe(II)/Fe(III)-oxides coupling

Jean-Pascal Andraud, S. Criquet, Benjamin Oursel, Stephane Greff, Thierry Woignier, Hervé Macarie

► To cite this version:

Jean-Pascal Andraud, S. Criquet, Benjamin Oursel, Stephane Greff, Thierry Woignier, et al.. Test of chlordecone dechlorination by chemical reduction using Fe(II)/Fe(III)-oxides coupling. The Sixth International Symposium on Environmental Biotechnology and Engineering, Nov 2018, Ciudad Obregón, Mexico. hal-02021158

HAL Id: hal-02021158

<https://amu.hal.science/hal-02021158>

Submitted on 15 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Test of chlordecone dechlorination by chemical reduction using Fe(II)/Fe(III)-oxides coupling

Jean-Pascal ANDRAUD^{1*}, Stéven CRIQUET¹, Benjamin OURSEL², Stéphane GREFF¹, Thierry WOIGNIER^{1,3}, Hervé MACARIE¹

¹ Aix Marseille Univ, Univ Avignon, CNRS, IRD, IMBE, Marseille, France.

² Aix Marseille Univ, Université de Toulon, CNRS, IRD, MIO, Marseille, France.

³ IRD, IMBE, Campus Agro-environnemental Caraïbe, Le Lamentin, Martinique, France.

* Corresponding author: jean-pascal.andraud@imbe.fr

Background & Challenges

Today, **chlordecone (CLD)**, an organochlorine pesticide spread on the soil to control the populations of the banana weevil, is responsible for an unprecedented health, environmental, social and political crisis in the **French West Indies (FWI)**. Despite the health management measures implemented by the authorities to prevent CLD from reaching the consumer's plate, **the population continues to be exposed**. A definitive solution to the problem would be to **destroy the CLD stock in the soil**.



Figure 1: Fe(III)-oxides powders (Goethite, Ferrihydrite, Hematite)

FWI soils are very rich in iron, which accounts for 7 to 11% of their dry weigh. As these soils are oxic, iron is found mainly in the form of **Fe(III)-oxides**, and more precisely in the form of goethite, hematite and ferrihydrite whose proportions vary according to the type of soil (Fig. 1).

The potential (E^0) of the redox couples formed by **Fe(III)-oxides/Fe(II)** and **Fe(II)/Fe⁰** ($E^0 \approx -470$ mV) are similar (Fig. 2 : Redox scale). This indicates that the **Fe(II)** resulting from the **Fe(III)-oxides microbial reduction** could be an electron donor as effective as **Fe⁰** for attacking CLD.

Objectives

Degrade the chlordecone stock of FWI soils by using Fe(III)-oxides present *in situ* via biological iron-reduction.

- Show that chlordecone can be dechlorinated by chemical reduction using Fe(II)/Fe(III)-oxides coupling (Fig.2 : "Abiotic step")
- Couple biological iron reduction with chemical reduction of chlordecone, using Iron-Reducing Bacteria (IRB)
- Test chemical and biological steps on FWI soils incubated under anoxic conditions at lab-scale.

Material & Methods

- Anoxic incubation under agitation at 25°C (room temperature)
- 25 mL of buffer to maintain a pH of the reaction mixture at 7.5-8:
 - Organic buffer: MOPS (25 mM)
 - Inorganic buffer: NaHCO₃ (11.9 mM)
- Fe(III)-oxides: hematite, goethite, ferrihydrite, magnetite
- 1 mM FeSO₄, 7 H₂O
- Redox potentials and pH measurements under anoxic conditions



Results & Conclusions




Figure 3: Reduction potential (E_h) values of goethite suspensions as a function of aqueous Fe^{2+} concentration and pH. Error bars represent the standard deviation from triplicate reactors. No error bars are present for the pH 6.0 and 6.5 reactors because only one reactor was measured for each point. The dashed gray lines represent the model fit. Experimental details: 1 g/L goethite, 25 mM KCl, 50 mM buffer, and 10 μ M mediator.

From Gorski et al. Thermodynamic characterization of Iron Oxide-aqueous Fe^{2+} redox couples. *Environ. Sci. Technol.*, 50, 8538-8547, 2016.

Table 1: Redox potentials and pH measurements at 72h of Fe(III)-oxides suspensions in the absence or presence of aqueous Fe^{2+}

	NaHCO ₃ buffer		MOPS buffer	
	pH	E_h^* (mV)	pH	E_h^* (mV)
Magnetite control	8.1	293	7.7	314
Magnetite + Fe²⁺	7.5	-113	7.5	231
Hematite control	8.1	338	7.7	356
Hematite + Fe²⁺	7.7	-113	7.6	-49
Goethite control	8.3	343	7.7	316
Goethite + Fe²⁺	7.6	-168	7.6	-125
Ferrihydrite control	7.9	319	7.5	317
Ferrihydrite + Fe²⁺	7.3	-79	7.5	-4

*versus Standard Hydrogen electrode (SHE)




Figure 2: Abiotic dechlorination of CLD and iron-reduction coupling

- Dechlorination test at **acid pH**: no degradation showed by measuring chlorides with ionic chromatography.
- Negative redox potentials** are required for dechlorination, and these are mostly conditioned by the pH as derived from the literature (Fig. 3).

With NaHCO₃ buffer, the formation of a FeCO₃ precipitate appears. The E_h measured therefore correspond to the Fe(III)-oxides/FeCO₃ couple which are theoretically less negative than the Fe(III)-oxides/Fe(II) couples (Table 1).

$$E_h(\text{NaHCO}_3) > E_h(\text{MOPS})$$

- Coming soon: dechlorination test in both types of buffer with goethite and then the other Fe(III)-oxides.