



HAL
open science

Metal Actuated Ring Translocation Switches in Water

Hang Yin, Roselyne Rosas, Didier Gigmes, Olivier Ouari, Ruibing Wang,
Anthony Kermagoret, David Bardelang

► **To cite this version:**

Hang Yin, Roselyne Rosas, Didier Gigmes, Olivier Ouari, Ruibing Wang, et al.. Metal Actuated Ring Translocation Switches in Water. *Organic Letters*, 2018, 20 (11), pp.3187–3191. 10.1021/acs.orglett.8b01019 . hal-02091859

HAL Id: hal-02091859

<https://amu.hal.science/hal-02091859>

Submitted on 15 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Metal Actuated Ring Translocation Switches in Water.


Hang Yin,^a Roselyne Rosas,^b Didier Gigmes,^c Olivier Ouari,^c Ruibing Wang,^{a*} Anthony Kermagoret,^{c*} and David Bardelang.^{c*}

^aState Key Laboratory of Quality Research in Chinese Medicine, and Institute of Chinese Medical Sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR, China

^bAix Marseille Univ, CNRS, Spectropole, FR 1739, Marseille, France

^cAix Marseille Univ, CNRS, ICR, Marseille, France

Supporting Information Placeholder



ABSTRACT: Among a series of metal ions in water, silver is the only one to remotely and reversibly switch cucurbit[7]uril (CB[7]) movements (translocation or uptake) on a rigid and linear 3-station viologen-phenylene-imidazole (*V-P-I*) derivative, avoiding undesired pH actuation. ¹H NMR, UV-vis spectroscopy, mass spectrometry, ITC and modeling were combined to show that ring translocation or uptake along a molecular thread is possible in water by Ag⁺ as a metal stimulus.

Molecular switches are increasingly investigated because of their high potential as elementary building-blocks for the next generation of molecular machines.¹ Remarkable examples have been reported by several groups, with designs comprising flexible axle and ring molecules.² Even if water is attractive due to its lack of toxicity, possible enhanced affinity by hydrophobic effects³ and its ability to make molecular switches more amenable for biological applications,⁴ most molecular switches proceed in organic solvents.⁵ Yet, water soluble molecular switches are much less investigated and, even if pH⁶ is an essential stimulus to trigger ring translocation, nature also uses metal ions or ATP hydrolysis, for instance to trigger reversible protein ring contraction in ion channels or to drive ring-shaped motor proteins over DNA strands to achieve essential functions in cells.⁷ To make efficient molecular switches work in water in various contexts, one would ideally be able to choose the best stimulus to actuate ring translocation. Expanding the palette of available stimuli working in water, beyond pH, is thus desirable. For example, when buffers are undesired and H⁺ unusable due to fragile motifs on hosts or guests, metal ions could be valuable stimuli.⁸ However, there are to the best of our knowledge, only a handful of examples of metal actuated ring translocation switches and none work in water.⁹ Reasons for that may be because of difficulties in combining functions recognizable by a metal ion, both in host and guest, or because most metal ions can behave as

Brønsted acids in water and thus affect the pH,¹⁰ rendering unbiased analyses more complicate.

Cucurbit[7]uril (CB[7])¹¹ is a ring shape receptor presenting several advantages to test metal ions as triggers of ring translocation because it is water soluble and possess (i) a versatile hydrophobic cavity and (ii) two carbonyl-laced portals amenable for metal ion binding. In line with our continuous work on CB[*n*],¹² we tested a series of commonly used water soluble metal salts and found that Ag⁺ selectively translocates CB[7] over a simple, rigid molecular axle containing a viologen (*V*), a phenylene (*P*) and an imidazole (*I*) station (*V-P-I*, Figure 1) or pull a 2nd CB[7], without affecting the pH deleteriously.




Figure 1. Molecular structures of *V-P-I* and of CB[7] and representation of the silver actuated molecular switches.




Figure 2. Excerpts of the aromatic region of ¹H NMR spectra for the CB[7] titration of *V-P-I*[Cl₂] (1 mM, a) and *V-P-I*[Cl₂] (1 mM) with AgNO₃ (8 equiv, b) in D₂O with c) proposed structures of the complexes (Ag⁺ ions: red spheres).

Imidazole *V-P-I*[Cl₂] (2 Cl⁻ counter-ions) was prepared in a three-step sequence, and details are given in the Supporting Information (SI). Imidazole *V-P-I*[Cl₂] is water soluble and possesses two pK_a values, as the imidazole station can be either protonated (pK_{a1} = 4.0) or deprotonated (pK_{a2} = 10.4, Figure S6). Therefore, working near neutral pH was essential without buffers to (i) warrant having benzimidazole structures (Figures 1 and 2; i. e. not the protonated or deprotonated forms) and (ii) avoid complications due to cations that are known to interact significantly with CB[7] carbonyl rims.¹³ First, CB[7] complexation on the *V* station was easily monitored by ¹H NMR in D₂O (Figure 2a). Then, NaCl, CsF, AgNO₃, CaCl₂, MgCl₂, CoCl₂, NiCl₂, ZnCl₂, CuSO₄, MnCl₂, FeCl₂, FeSO₄, FeCl₃, GdCl₃, Cr(NO₃)₃, AlCl₃ were assessed for CB[7] translocation. Na⁺, Cs⁺, Ca^{II}, Mg^{II}, Co^{II}, Ni^{II}, Zn^{II}, Mn^{II} and Gd^{III} had essentially no effect on ¹H NMR spectra, even after addition of several equivalents in water (up to 10 equiv Figure S7, pH reduced to 6.10 in the worst case). Conversely, Cr, Al and Fe salts changed the spectrum of *V-P-I*•CB[7] to assignable resonances, while Cu showed broad signals. The obtained spectra for Cr^{III}, Al^{III} and Fe^{III} are very similar to that of *V-P-I*•H⁺•CB[7] (Figure S7) and are probably the result of metal-induced acidification as reflected by the measured pH values of the solutions (Figure S7). The addition of these metallic salts thus resulted in undesired pH triggered CB[7] translocation. Buffering the solutions to limit pH changes triggered by Fe, Cu, Al or Cr would result in precipitates of hydroxides complexes.¹⁴ pH Actuation of the present system is also

interesting, but will be reported elsewhere since it exceeds the scope of this study. However, Ag^I resulted in new NMR spectra without altering too much of the pH. Without Ag⁺, as expected, CB[7] quantitatively binds *V-P-I* on station *V* while increasing further the CB[7] concentration does not change this 1:1 binding mode (Figure 2a). Split chemical resonances for free and complexed *V-P-I* suggest slow exchange on the NMR timescale. The signals of protons 3 and 4 are shifted upfield by ≈ 1.5 ppm in line with CB[7] location on station *V*, with one carbonyl rim downfield-shifting the signals of protons 6 and 7 by 0.37 and 0.13 ppm respectively (<Δδ> ≈ 0.25 ppm, carbonyl rims symmetry broken, Figure S9 and Figure S11, 1 equiv, right). AgNO₃ had a weak, but significant effect on the spectrum of *V-P-I* alone (Figure S12), affecting H₇, H₈ and H₉ that are located nearby the imidazole function and thus reflecting weak binding on the imidazole function (significant effects were also observed by UV-vis spectroscopy, Figure S14). CB[7] Titration of *V-P-I*[Cl₂] with AgNO₃ instead showed CB[7] binding on station *P* (slow exchange on the ¹H NMR timescale, Figure 2b). Silver ions have previously been shown to behave as soft Lewis acids,¹⁵ and imidazoles¹⁶ or the carbonyl rims of cucurbiturils¹⁷ have been reported to be possible ligands for Ag⁺. We postulate a silver-induced host translocation by multiple binding on the guest-imidazole function and by the oxygen carbonyl rims. Two equiv of AgNO₃ were necessary to neutralize the effect of the two chloride ions present in *V-P-I*[Cl₂]. ¹H NMR Titrations of AgNO₃ for a 1:1 *V-P-I*[Cl₂]:CB[7] mixture showed large chemical shift chang-

es up to ~ 10 equiv of Ag^+ for species fast exchanging on the ^1H NMR timescale (Figure S18). This fast-exchange regime was presumably due to the lability of the silver-nitrogen bond upon Ag^+ binding on the imidazole function of $V\text{-}P\text{-}I$ accompanied by the gradual translocation of CB[7] from station V to station P . This allowed the indirect determination of the binding constant for Ag^+ ($K_a = 3470 \pm 550 \text{ M}^{-1}$, fit of experimental points using a 1:1 binding model) toward the $V\text{-}P\text{-}I\text{-}CB[7]_1$ complex and suggested a 1:1:1 stoichiometry for the $V\text{-}P\text{-}I\text{-}CB[7]_1\text{-}Ag^+$ complex (Figures S18 and S27). When $V\text{-}P\text{-}I\text{-}[Cl_2]$ was titrated against CB[7] with enough Ag^+ (8 equiv of AgNO_3 kept constant, Figure 2b), signals of protons 5, 6 and 7 upfield shifted by 1.00, 0.99 and 0.60 ppm respectively up to 1 equiv of CB[7] (ring on station P). While the threaded CB[7] seems to exchange fast on the binding sites with respect to the NMR timescale, the host prefers remaining on station P . However, excess CB[7] completely changed the NMR spectra with large upfield shifts for both sets of protons, 3,4 (1.51 ppm) from station V , and 8,9 ($\langle\Delta\delta\rangle \approx 0.80$ ppm) from station I while the signals of protons 6 and 7 are shifted downfield by approximately double the value as for the 1:1 binding on $V\text{-}P\text{-}I$ ($\langle\Delta\delta\rangle \approx 0.50$ ppm, Figure 2b bottom). This agrees with a CB[7] binding on the V station and a second CB[7] located on the I station. The P station protons experienced significant deshielding effects from two carbonyl rims (one rim from each of the two CB[7]), while no further changes were observed for more than 2.0 equiv of CB[7]. Thus, Ag^+ ions were observed to (i) translocate CB[7] from the V to the P station at 1:1 guest:host ratio and (ii) pull a second CB[7] onto the I station at a 1:2 guest:host stoichiometry. Simultaneous binding of two CB[7] on two vicinal stations (V and P or P and I) was not observed. Mass spectrometry (MS) showed peaks assigned to $V\text{-}P\text{-}I$, and $V\text{-}P\text{-}I\text{-}CB[7]_1$, together with the corresponding H^+ adduct. With AgNO_3 , ion mobility separation prior to analyses allowed us to observe additional peaks at m/z 545.1390 (theo. 545.1390) and m/z 848.7045 (theo. 848.7027) respectively corresponding to the formulas $\text{C}_{66}\text{H}_{62}\text{N}_{32}\text{O}_{14}\text{Ag}_3^{3+}$ and $\text{C}_{66}\text{H}_{62}\text{N}_{33}\text{O}_{17}\text{Ag}_2^{2+}$ in line with complexes $V\text{-}P\text{-}I\text{-}CB[7]_1\text{-}Ag^+$ and $V\text{-}P\text{-}I\text{-}CB[7]_2\text{-}Ag^+\text{-}NO_3^-$ (Figure S19).

DFT minimized structures of $V\text{-}P\text{-}I\text{-}CB[7]_1$ complexes¹⁸ with CB[7] positioned on the V , P or I stations (frequency checked and corrected for BSSE, see SI) showed a marked preference for the complex with CB[7] on station V (Figure 3a), that is in good agreement with ^1H NMR. With a silver ion (1 Ag^+ considered as suggested by NMR and mass spectrometry), the complex with CB[7] on station P is the most stable (Figure 3b, SI), also consistent with ^1H NMR of $V\text{-}P\text{-}I\text{-}CB[7]_1\text{-}Ag^+$, $V\text{-}P\text{-}I\text{-}Ag^+$ and the UV-vis spectra of $V\text{-}P\text{-}I\text{-}Ag^+$ which suggested an interaction between Ag^+ and N -imidazole. In this binding geometry, there are several hydrogen bonds between CB[7] and the guest and the silver ion is 2.20 Å distant from the imidazole nitrogen atom ($\text{Ag}^+\cdots\text{N}$) and there are two additional $\text{Ag}^+\cdots\text{O}$ coordinating bonds (2.32 Å and 2.45 Å) with one CB[7] carbonyl rim (Figure 3b). The combination of these two kinds of interactions seems to be responsible for driving

CB[7] from the V to the P station. The situation is similar for the 1:2 complex, except that one CB[7] is on station V and the 2nd CB[7] is instead on station I (Figure 3c). In this case, the silver ion interacts (i) with the imidazole nitrogen atom (2.20 Å) and (ii) with two carbonyl oxygen atoms of the CB[7] of the I station (2.33 Å and 2.46 Å).




Figure 3. Proposed structures (DFT minimized) in line with ^1H NMR for the complexes (a) $V\text{-}P\text{-}I\text{-}CB[7]_1$, (b) $V\text{-}P\text{-}I\text{-}CB[7]_1\text{-}Ag^+$, and (c) $V\text{-}P\text{-}I\text{-}CB[7]_2\text{-}Ag^+$.

Thus, the imidazole function behaves as a pivot for Ag^+ which in turn attracts CB[7] providing an oxygen-rich environment to further stabilize Ag^+ . Of course, other positions for Ag^+ near a carbonyl rim and around the imidazole station cannot completely be ruled out, but such other positions lead to less stable 1:1:1 complexes by DFT and Ag^+ also significantly affected the ^1H NMR signals of $V\text{-}P\text{-}I$ near the imidazole function and maximum absorbance in UV-vis spectra of $V\text{-}P\text{-}I$ was also significantly shifted (Figures S12 and S14 respectively). The observed silver selectivity for ring translocation could be explained by (i) the charge of the metal ion (a monocationic metal ion is more likely to be bound by a CB[7] carbonyl rim for an overall 2+ charged complex), (ii) the soft lewis acid nature of silver capable of coordinating

both an imidazole function and carbonyl groups of CB[n],¹⁵⁻¹⁷ and (iii) because Ag⁺ cations do not generate insoluble metal oxides at neutral pH.¹⁴

The present system was also investigated by ITC (Figure 4). Without silver ions, the binding affinity between CB[7] and *V-P-I* was determined to be $7.25 \times 10^5 \text{ M}^{-1}$ (Figure 4a). The binding stoichiometry is 1:1, consistent with NMR, DFT and MS. In the presence of Ag⁺, results of ITC titration of *V-P-I* with CB[7] are consistent with a sequential complexation, with apparent binding constant values of $K_{a1} = 6.45 \times 10^6 \text{ M}^{-1}$ and $K_{a2} = 3.75 \times 10^5 \text{ M}^{-1}$.




Figure 4. Microcalorimetric titration of *V-P-I* with CB[7] in aqueous solution, in the absence (a) and in the presence (b) of Ag⁺. Top: thermograms of CB[7] injected into *V-P-I* solutions. Bottom: dependence of ΔH for each injection of CB[7] against the molar ratio between CB[7] and *V-P-I* (the solid line represents the best fit using “one set of binding sites” and “two separate one set of binding sites” models, respectively).

Below 1 equiv of CB[7], corresponding heat changes can tentatively be assigned to 1 Ag⁺-assisted CB[7] binding on *V-P-I* (K_{a1}). The larger binding of CB[7] toward station *P* in the presence of Ag⁺ is well consistent with NMR. The heat changes above 1 equiv of CB[7] are hard to interpret because of the suspected entanglement of two effects: (i) CB[7] translocation back to station *V* and (ii) binding of a second CB[7]. However, ITC confirmed the proposed stoichiometry and the process is both enthalpically and entropically driven (SI). Interestingly, the switching processes were simply reversed by addition of NaCl (Figure 2) precipitating Ag⁺ in the form of AgCl. Reversibility of the molecular switches (1:1 and 1:2 guest:host ratios) was examined over 2 cycles by repetitive alternation of the position of CB[7] on stations *V* and *P* (Figure S25), or repetitive catch-and-release of a 2nd CB[7] (Figure S26). Finally, the role of the anion should not be left unconsidered with a potentially binding imidazole function on the axle and cations ubiquitous in this system but the carbonyl rims of CB[7] are rather expected to behave as anions repellants.

In summary, we report silver-triggered, reversible molecular switches in water based on CB[7] and a simple, rigid Viologen-Phenylene-Imidazole (*V-P-I*) guest axle. Ag⁺ is unique for this system since it selectively and re-

motely controls (i) CB[7] sliding from station *V* to station *P* of the guest or (ii) a second CB[7] uptake and release on guest station *I*. To the best of our knowledge, there is no previous example of a ring molecule reversibly guided (translocated) or pulled (caught) on a rigid axle by a metallic cation in water, but there is still room for improvements (excess Ag⁺ needed) with appropriate design and we anticipate that other guest and host will also exhibit ring movements triggered by metal ions in water. The water solubility of the present system could enlarge the perspectives of biological applications of molecular switches⁴ (i. e. acid-responsive imidazole function). We think that this ternary approach (Ag⁺ / imidazole / CB[7]) is also well suited to open the way to a new family of silver templated interlocked structures¹⁹ and silver actuated molecular switches and machines.

ASSOCIATED CONTENT

Supporting Information

Preparation and characterization of *V-P-I*[Cl₂], 1D-NMR and 2D-NMR spectra, details for mass spectrometry measurements, additional NMR titrations, ITC data, pKa measurements and details of DFT calculations can be found in the Supporting Information. The Supporting Information (pdf file) is available free of charge on the ACS Publications website.

AUTHOR INFORMATION

Corresponding Authors

*E-mail: rwang@umac.mo

*E-mail: anthony.kermagoret@univ-amu.fr

*E-mail: david.bardelang@univ-amu.fr

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

CNRS and Aix-Marseille Université are acknowledged for financial support. Valérie Monnier is gratefully acknowledged for MS analyses (Aix Marseille Univ, CNRS, Centrale Marseille, FSCM, Spectropole, Marseille, France).

REFERENCES

- (1) Leigh, D. A. *Angew. Chem., Int. Ed.* **2016**, *55*, 14506.
- (2) (a) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2000**, *39*, 3348. (b) Green, J. E.; Choi, J. W.; Boukai, A.; Bunimovich, Y.; Johnston-Halperin, E.; Delonno, E.; Luo, Y.; Sheriff, B. A.; Xu, K.; Shin, Y. S.; Tseng, H.-R.; Stoddart, J. F.; Heath, J. R. *Nature* **2007**, *445*, 414. (c) Du, G.; Moulin, E.; Jouault, N.; Buhler, E.; Giuseppone, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 12504.
- (3) (a) Biedermann, F.; Nau, W. M.; Schneider, H.-J. *Angew. Chem. Int. Ed.* **2014**, *53*, 11158. (b) Cao, L.; Sekutor,

- M.; Zavalij, P. Y.; Mlinaric-Majerski, K.; Glaser, R.; Isaacs, L. *Angew. Chem. Int. Ed.* **2014**, *53*, 988.
- (4) (a) Garcia-Lopez, V.; Chen, F.; Nilewski, L. G.; Duret, G.; Aliyan, A.; Kolomeisky, A. B.; Robinson, J. T.; Wang, G.; Pal, R.; Tour, J. M. *Nature* **2017**, *548*, 567. (b) Grunder, S.; McGrier, P. L.; Whalley, A. C.; Boyle, M. M.; Stern, C.; Stoddart, J. F. *J. Am. Chem. Soc.* **2013**, *135*, 17691.
- (5) (a) Erbas-Cakmak, S.; Leigh, D. A.; McTernan, C. T.; Nussbaumer, A. L. *Chem. Rev.* **2015**, *115*, 10081. (b) Kay, E. R.; Leigh, D. A.; Zerbetto, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 72. (c) Feringa, B. L.; van Delden, R. A.; Koumura, N.; Geertsema, E. M. *Chem. Rev.* **2000**, *100*, 1789.
- (6) For representative examples: (a) Mikulu, L.; Michalicova, R.; Iglesias, V.; Yawer, M. A.; Kaifer, A. E.; Lubal, P.; Sindelar, V. *Chem. – Eur. J.* **2017**, *23*, 2350. (b) Farahani, N.; Zhu, K.; Loeb, S. J. *ChemPhysChem* **2016**, *17*, 1875. (c) Zhu, K.; Vukotic, V. N.; Loeb, S. J. *Chem. – Asian J.* **2016**, *11*, 3258. (d) Diniz, A. M.; Basilio, N.; Cruz, H.; Pina, F.; Parola, A. J. *Faraday Discuss.* **2015**, *185*, 361. (e) Zhu, K.; Vukotic, V. N.; Loeb, S. J. *Angew. Chem., Int. Ed.* **2012**, *51*, 2168. (f) Sindelar, V.; Silvi, S.; Parker, S. E.; So-bransingh, D.; Kaifer, A. E. *Adv. Funct. Mater.* **2007**, *17*, 694. (g) Lee, J. W.; Kim, K.; Kim, K. *Chem. Commun.* **2001**, 1042. (h) Bissell, R. A.; Cordova, E.; Kaifer, A. E.; Stoddart, J. F. *Nature* **1994**, *369*, 133. (i) Mock, W. L.; Pierpont, J. *Chem. Commun.* **1990**, 1509.
- (7) (a) Thomsen, N. D.; Lawson, M. R.; Witkowsky, L. B.; Qu, S.; Berger, J. M.; *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E7691. (b) Ma, W.; Schulten, K. *J. Am. Chem. Soc.* **2015**, *137*, 3031. (c) Liu, S.; Chistol, G.; Bustamante, C. *Biophys. J.* **2014**, *106*, 1844. (d) Javaherian, A. D.; Yusifov, T.; Pantazis, A.; Franklin, S.; Gandhi, C. S.; Olcese, R. *J. Biol. Chem.* **2011**, *286*, 20701. (e) Patel, S. S.; Donmez, I. *J. Biol. Chem.* **2006**, *281*, 18265.
- (8) (a) Beves, J. E.; Blight, B. A.; Campbell, C. J.; Leigh, D. A.; McBurney, R. T. *Angew. Chem., Int. Ed.* **2011**, *50*, 9260. (b) Dietrich-Buchecker, C. O.; Sauvage, J.-P. *Chem. Rev.* **1987**, *87*, 795. (c) Xue, M.; Yang, Y.; Chi, X.; Yan, X.; Huang, F. *Chem. Rev.* **2015**, *115*, 7398. (d) Linke, M.; Chambron, J.-C.; Heitz, V.; Sauvage, J.-P.; Semetey, V. *Chem. Commun.* **1998**, 2469. (e) Ling, X.; Masson, E. *Org. Lett.* **2012**, *14*, 4866.
- (9) (a) Altmann, P. J.; Pöthig, A. *Angew. Chem., Int. Ed.* **2017**, *56*, 15733. (b) Baggi, G.; Loeb, S. J. *Chem. – Eur. J.* **2017**, *23*, 14163. (c) Davidson, G. J. E.; Sharma, S.; Loeb, S. J. *Angew. Chem., Int. Ed.* **2010**, *49*, 4938. (d) Sharma, S.; Davidson, G. J. E.; Loeb, S. J. *Chem. Commun.* **2008**, 582. (e) Dietrich-Buchecker, C. O.; Hemmert, C.; Khemiss, A.-K.; Sauvage, J.-P. *J. Am. Chem. Soc.* **1990**, *112*, 8002. (f) Marlin, D. S.; Cabrera, D. G.; Leigh, D. A.; Slawin, A. M. Z. *Angew. Chem., Int. Ed.* **2006**, *45*, 77. (g) Marlin, D. S.; Cabrera, D. G.; Leigh, D. A.; Slawin, A. M. Z. *Angew. Chem., Int. Ed.* **2006**, *45*, 1385. (h) Zhou, W.; Li, J.; He, X.; Li, C.; Lv, J.; Li, Y.; Wang, S.; Liu, H.; Zhu, D. *Chem. – Eur. J.* **2008**, *14*, 754.
- (10) (a) Wang, T.; Glasper, J. A.; Shanks, B. H. *Appl. Catal., A* **2015**, *498*, 214. (b) Muller, P. *Pure and Appl. Chem.* **1994**, *66*, 1077.
- (11) (a) Assaf, K. I.; Nau, W. M. *Chem. Soc. Rev.* **2015**, *44*, 394. (b) Barrow, S. J.; Kasera, S.; Rowland, M. J.; del Barrio, J.; Scherman, O. A. *Chem. Rev.* **2015**, *115*, 12320. (c) Lagona, J.; Mukhopadhyay, P.; Chakrabarti, S.; Isaacs, L. *Angew. Chem., Int. Ed.* **2005**, *44*, 4844. (d) Masson, E.; Ling, X.; Joseph, R.; Kyeremeh-Mensah, L.; Lu, X. *RSC Adv.* **2012**, *2*, 1213.
- (12) (a) Bardelang, D.; Casano, G.; Poulhès, F.; Karoui, H.; Filippini, J.; Rockenbauer, A.; Rosas, R.; Monnier, V.; Siri, D.; Gaudel-Siri, A.; Ouari, O.; Tordo, P. *J. Am. Chem. Soc.* **2014**, *136*, 17570. (b) Casano, G.; Poulhès, F.; Tran, T. K.; Ayhan, M. M.; Karoui, H.; Siri, D.; Gaudel-Siri, A.; Rockenbauer, A.; Jeschke, G.; Bardelang, D.; Tordo, P.; Ouari, O. *Nanoscale* **2015**, *7*, 12143. (c) Li, S.; Chen, H.; Yang, X.; Bardelang, D.; Wyman, I. W.; Wan, J.; Lee, S. M. Y.; Wang, R. *ACS Med. Chem. Lett.* **2015**, *6*, 1174. (d) Bardelang, D.; Banaszak, K.; Karoui, H.; Rockenbauer, A.; Waite, M.; Udachin, K.; Ripmeester, J. A.; Ratcliffe, C. I.; Ouari, O.; Tordo, P. *J. Am. Chem. Soc.* **2009**, *131*, 5402.
- (13) (a) Dutta, S.; Mohanty, C. J.; Pal, H.; Bhasikuttan, A. C. *J. Am. Chem. Soc.* **2010**, *132*, 1395. (b) Tang, H.; Fuent-ealba, D.; Ko, Y. H.; Selvapalam, N.; Kim, K.; Bohne, C. J. *Am. Chem. Soc.* **2011**, *133*, 20623.
- (14) Feitknecht, W.; Schindler, P. *Pure Appl. Chem.* **1963**, *6*, 125.
- (15) (a) Matsuzawa, A.; Mashiko, T.; Kumagai, N.; Shibasaki, M. *Angew. Chem., Int. Ed.* **2011**, *50*, 7616. (b) Ihara, T.; Ishii, T.; Araki, N.; Wilson, A. W.; Jyo, A. *J. Am. Chem. Soc.* **2009**, *131*, 3826. (c) Laube, T.; Weidenhaupt, A.; Hunziker, R. *J. Am. Chem. Soc.* **1991**, *113*, 2561.
- (16) Chen, S.; Guo, Z.; Zhu, S.; Shi, W.-e.; Zhu, W. *ACS Appl. Mater. Interfaces* **2013**, *5*, 5623.
- (17) (a) Chen, K.; Wang, Y.; Hua, Z.-Y.; Xu, J.; Chen, M.-D.; Kong, Q.-G. *Inorg. Chem. Commun.* **2017**, *84*, 72. (b) Wu, Y.; Lan, Y.; Liu, J.; Scherman, O. A. *Nanoscale* **2015**, *7*, 13416. (c) Lu, X.; Masson, E. *Langmuir* **2011**, *27*, 3051. (d) Zeng, J.-P.; Zhang, S.-M.; Zhang, Y.-Q.; Tao, Z.; Zhu, Q.-J.; Xue, S.-F.; Wei, G. *Cryst. Growth Des.* **2010**, *10*, 4509. (e) Lu, X.; Masson, E. *Org. Lett.* **2010**, *12*, 2310. (f) Park, K.-M.; Whang, D.; Lee, E.; Heo, J.; Kim, K. *Chem. – Eur. J.* **2002**, *8*, 498.
- (18) No single crystal could be obtained despite several attempts at 1:1 and 1:2 guest:host ratios, without and with AgNO₃.
- (19) (a) Yang, Y.-D.; Fan, C.-C.; Rambo, B. M.; Gong, H.-Y.; Xu, L.-J.; Xiang, J.-F.; Sessler, J. L. *J. Am. Chem. Soc.* **2015**, *137*, 12966. (b) Goldup, S. M.; Leigh, D. A.; Lusby, P. J.; McBurney, R. T.; Slawin, A. M. Z. *Angew. Chem., Int. Ed.* **2008**, *47*, 6999.