



Hyperspectral recognition of the urban vegetation using spectral library. Example of Kaunas (Lithuania)

Sébastien Gadal

► To cite this version:

Sébastien Gadal. Hyperspectral recognition of the urban vegetation using spectral library. Example of Kaunas (Lithuania). GIS Yakutsk 2019: GIS for Digital Development, NEFU, Apr 2019, Yakutsk, Russia. hal-02098302

HAL Id: hal-02098302

<https://amu.hal.science/hal-02098302>

Submitted on 16 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hyperspectral recognition of the urban vegetation using spectral library. Example of Kaunas (Lithuania)

Sébastien GADAL

Context and goals

Goals

- Characterize, monitor and map urban vegetation over Kaunas city (Lithuania).
- Propose practical solutions for city actors and urban architects in terms of vegetation mapping and inventory gathering.



Geographic context and study zone choice:

- High Vegetation diversity in urban areas.
- Complex distribution of vegetation species within the city.
- Various Morphologies and urban structures: 15e-17e centuries, 19e century, Art-Nouveau, Soviet, post-soviet, etc.
- Canopy state (green, dry, sick).







Airborne hyperspectral data (Rikola)

- Two images acquired over Kaunas city, in July 2015 and September 2016.
- 16 and 64 bands 500nm-900nm (Vis-NIR).
- Spatial resolution of 0,7 m and 0,5 m.
- Covered area: 5km².

Kaunas center July 2015 (**16 bands**)




Kaunas center September 2016 (**64 bands**)



Methodological framework

- Based on the spectral libraries
- Automatic (based on the machine learning)
- Based on the vegetation database (Kaunas tree database and field survey)

Methodology: spectral libraries (1)



Spectral library: laboratory based training samples




Hyperspectral laboratory imager features:

- Brand: Themis Vision VNIR400H (Themis Vision systems LLC).
- Spectral Range: Vis-NIR, 400nm to 1000 nm.
- Spectral resolution: 0,6nm
- Sensor type: CCD.
- Bands number: 955.
- Light: 2 halogen lamps of 100W
- Image Scan: line by line
- Measurements conditions: black room

Oak sample




Spectral measurement



Sébastien Gadal, Walid Ouerghemmi, 2016-2018


Vegetation Mapping and recognition of species using image library training samples and SAM classifier (results)




Species	SAM (16 bands)	SAM (64 bands)
H. Chestnut	6.6	8.7
E. Beech	7.9	33.1
Linden	10.5	11.6
Maple	12.0	11.0
M.Ash	30.0	15.7
Oak	11.5	10.2
N. Spruce	15.0	16.6
Juniperus horizontalis	0.3	0.0
S. Pine	27.7	30.1
Arborvitae	50.0	0.7
Grass	59.4	36.0
O.A. (%)	16.2	21.1

- «slight» to «fair» accuracy ($0\% < O.A. < 40\%$)
- Better identification rate given by **64 bands** image (40% more pixels identified)
- High variability per species

Vegetation Mapping and recognition of group of species using image library training samples and SVM classifier (results)




■ Deciduous
■ Coniferous
■ Grass



Species\Classifier	SVM (16 bands)	SVM (64 bands)
Deciduous	57.9	63.2
Coniferous	45.6	35.0
Grass	56.8	31.3
O.A. (%)	57.0	49.8

- «Moderate» accuracy for both case (40%<O.A.<60%)
- 64 bands image: better estimation of deciduous, under-estimation of coniferous
- 16 bands image: over-estimation of coniferous over grass

Vegetation Mapping and recognition of species using image library training samples and regularization process (results)




Walid Ouerghemmi, Sébastien Gadal, 2016-2018

a) RGB composite test portion over Kaunas city, **b)** classification by laboratory spectral library, **c)** classification by image spectral library, and **d)** regularization of the generated map

- **Poor identification rate and statistical accuracy using laboratory spectral library for classification training (b)**
- **Good identification rate and significant enhancement of statistical accuracy using image spectral library for classification training (c)**
- **Enhancement and smoothing of the classification map using regularization graph-cut algorithm (d)**

Recognition of the species based on the spectral librairy and SAM classifier




Accuracy Results (compilation)


Veg. species	Classif. accuracy (16 bands image)						Classif. accuracy (64 bands image)							
	Classifier Training	Image acquired spectra			laboratory spectra			Image acquired spectra			laboratory spectra			
		10	50	100	10	50	100	10	50	100	10	50	50	
H. Chestnut	0.0	3.7	5.4	20.1	6.4	4.3	2.3	8.3	8.7	20.0	3.2	1.1	3.3	3.2
E. Beech	21.0	7.9	7.9	2.6	10.0	11.1	20.2	38.7	33.1	0.2	4.0	5.2	20.0	19.9
Linden	4.3	17.5	9.2	0.7	0.0	0.0	3.7	2.4	11.6	0.0	0.0	0.0	8.5	8.8
Maple	30.4	14.8	12.4	0.3	0.0	0.1	13.3	12.9	11.0	0.0	0.0	0.0	29.5	28.9
M.Ash	10.2	11.8	22.5	0.0	59.8	59.6	15.2	21.0	15.7	0.0	26.2	20.7	0.0	0.0
Oak	33.8	34.3	17.1	2.1	6.8	8.7	8.8	11.7	10.2	1.6	1.7	3.97	21.9	21.9
N. Spruce	11.3	43.3	21.7	4.9	3.2	4.6	5.8	11.6	16.6	0.4	1.2	2.7	0.0	0.0
Lime glow	0.0	0.3	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
S. Pine	21.5	24.0	22.8	0.0	0.0	0.1	63.8	51.5	30.1	0.0	0.0	0.0	0.4	0.4
Arborvitae	11.8	25.9	46.1	0.0	0.0	4.9	0.7	0.5	0.7	0.0	0.0	0.0	0.0	0.0
Grass	30.4	56.9	60.8	0.0	0.0	0.4	34.1	31.3	36.0	0.0	0.0	0.0	16.6	16.5
O.A. (%)	19.4	18.2	15.6	2.7	3.0	2.9	19.7	19.1	21.1	1.5	1.1	0.9	16.8	16.7
Kappa × 10 ⁻³	70	70	70	-4	-0.9	-0.4	50	100	110	-3	-0.6	-0.1	14	11
Deciduous	--	57.9	--	--	78.0	--	--	63.2	--	--	36.2	--	71.2	71.2
Coniferous	--	45.6	--	--	5.4	--	--	35.0	--	--	1.0	--	3.7	3.7
Grass	--	56.8	--	--	0.0	--	--	31.3	--	--	0.0	--	16.6	16.5
O.A. (%)	--	57.0	--	--	67.2	--	--	49.8	--	--	20.7	--	47.0	46.0
Kappa × 10 ⁻³	--	170	--	--	-20	--	--	190	--	--	-7	--	-3	-7

- «Slight» accuracy of **individual species** recognition (overall accuracy of 21% at best)
- «Moderate» to «Substantial» accuracy of **group of species** recognition (overall accuracy of 67% at best)
- Image spectral library based mapping more accurate than laboratory spectral library based mapping.
- Enhancement of mapping accuracy when increasing band number for **individual species**
- No enhancement of mapping accuracy when increasing band number for **group of species**
- Band selection enhanced accuracy of individual and group of species recognition


Methodology: machine learning (2)





Original spectrum



NMF transformed spectrum
(transformation domain)




Noise free reconstructed spectrum
(original domain)




Spectral library : image based training samples


Kaunas Center **16 bands**, RGD
composite image -**July 2015** -



Kaunas Centre **64 bands**, RGB
composite image - **September 2016** -





Trees inventory over Kaunas city



Statistics

- 371 trees varieties (coniferous and deciduous)
 - 64 dominants varieties (occurrence > 100)
 - 17 very dominant varieties (occurrence > 1000) (Linden, Mountain ash, pine, Maple, etc.)
 - Most dominant variety: small leafed Linden
 - ~90% of deciduous trees, ~10% of coniferous

Vegetation Mapping per species using Reflectance MNF images (SAM and SVM comparison over 64 bands image)




H.Chestnut
Linden
M.Ash
Oak
N.Spruce
S.Pine
Thuja
Grass

Species	SAM (64 bands)	SVM (64 bands)
H. Chestnut	12.1	28.0
Linden	21.1	19.2
M.Ash	49.0	38.8
Oak	61.1	72.5
N. Spruce	26.9	51.6
S. Pine	10.5	18.1
Thuja	10.9	27.6
Grass	21.8	85.5
O.A. (%)	22.6	45.7

- “Fair” ($20\% < O.A. < 40\%$) to “moderate” ($40\% < O.A. < 60\%$) accuracies
- Better **mapping accuracy** given by **64 bands** image
- **Underestimation of grass, and overestimation of other species (N.Spruce and Oak) (SAM)**

Vegetation Mapping per species using Reflectance MNF images and SVM classifier (16 and 64 bands images comparison)





H.Chestnut
Linden
M.Ash
Oak
N.Spruce
S.Pine
Thuja
Grass

Species	SVM (16 bands)	SVM (64 bands)
H. Chestnut	19.1	28.0
Linden	36.0	19.2
M.Ash	38.4	38.8
Oak	36.5	72.5
N. Spruce	15.0	51.6
S. Pine	47.3	18.1
Thuja	50.1	27.6
Grass	91.2	85.5
O.A. (%)	41.2	45.7

- “Moderate” ($40\% < O.A. < 60\%$) accuracy
- Better **mapping accuracy** given by **64 bands** image
- Coniferous better identified using **16 bands** image

Vegetation Mapping per species using Non Reflectance MNF images and SVM classifier (16 and 64 bands images comparison)



- H.Chestnut**
- Linden**
- M.Ash**
- Oak**
- N.Spruce**
- S.Pine**
- Thuja**
- Grass**

Species	SVM (16 bands)	SVM (64 bands)
H. Chestnut	23.3	22.0
Linden	35.8	17.3
M.Ash	46.2	34.0
Oak	45.8	62.2
N. Spruce	19.2	40.3
S. Pine	59.3	17.7
Thuja	48.3	26.8
Grass	92.6	95.8
O.A. (%)	43.6	46.1

- “Moderate” ($40\% < O.A. < 60\%$) accuracy
- Better statistical accuracy given by 64 bands image
- Better expert accuracy given by 16 bands image
- Overestimation of grass specie using 64 bands non reflectance image
- Feature selection over 64 bands MNF image affected badly the mapping accuracy (information loss)

Accuracy Results (compilation)

- **MNF (1):** feature selection in the original domain (reflectance)
- **MNF (2):** feature selection in the transformed domain (non reflectance)

Veg. species	Classif. accuracy (16 bands image)						Classif. accuracy (64 bands image)								
	Classifier Training			SAM			SVM			SAM					
Samples	No NMF	MNF (1)	MNF (2)	No MNF	MNF (1)	MNF (2)	No NMF	MNF (1)	MNF (2)	No MNF	MNF (1)	MNF (2)	No NMF	MNF (1)	MNF (2)
deciduous	H. Chestnut		16.4	16.5	21.5	19.2	19.1	23.3	11.5	12.1	27.0	28.1	28.0	22.0	
	Linden		15.4	16.0	24.5	33.6	36.0	35.8	20.5	21.1	13.1	19.3	19.2	17.3	
	M.Ash		16.7	11.9	25.8	37.5	38.4	46.2	46.3	49.0	50.1	38.5	38.8	34.0	
	Oak		41.0	40.2	49.7	36.2	36.5	45.8	62.3	61.1	72.9	69.9	72.5	62.2	
	N. Spruce		10.0	9.8	21.1	16.4	15.0	19.2	26.8	26.9	16.0	50.7	51.6	40.3	
coniferous	S. Pine		49.8	47.7	55.4	49.1	47.3	59.3	10.7	10.5	18.7	17.9	18.1	17.7	
	Thuja		35.7	31.9	33.7	52.7	50.1	48.3	10.4	10.9	6.7	28.5	27.6	26.8	
	Grass		45.2	45.4	93.9	92.5	91.2	92.6	22.3	21.8	65.0	85.1	85.5	95.8	
	O.A. (%)		22.6	22.4	37.6	40.7	41.2	43.6	21.9	22.6	34.5	45.5	45.7	46.1	
	Kappa		0.10	0.10	0.28	0.29	0.30	0.33	0.11	0.11	0.24	0.34	0.34	0.34	

- “Fair” ($20\% < O.A. < 40\%$) to “moderate” ($40\% < O.A. < 60\%$) accuracies for all studied cases
- Enhancement of mapping accuracy when increasing band number from **16 to 64 bands**.
- **MNF feature selection** enhanced the mapping accuracy (original and transformed domains)
- Mapping over the **transformed domain** (non reflectance images, MNF 2) gives better statistical accuracies than **original domain** (reflectance images, MNF 1)

Conclusions

- The use for **VIS-Nir hyperspectral airborne** imagery, could be considered as a reliable and practical solution for **vegetation mapping** over **urban** areas
- **Limited** spectral samples were used for **classifiers training** (100 samples) due to limited **ground truth** data
- (**SVM**) classifier offer **more accurate** mapping than (**SAM**) classifier (**6% to 23%** better in term of overall accuracy)
- vegetation **species detection** was enhanced when **increasing the bands number** (from **16** to **64 bands**) (**1%** to **5%** better in term of overall accuracy)
- The use of **MNF feature selection** enhanced the mapping accuracy
- **MNF feature selection** over the **transformed domain** (non reflectance images) offer **better** statistical accuracy than over the **original domain** (reflectance images), but **reliability** is not ensured in case of **under-selection** of features
- **Issues and improvements:** improve ground truth data, use other feature selection strategies, use other classifier training strategies

Methodology: vegetation database (3)



Land field GIS database

- Individual trees (371 species)
- Heterogeneous groups (up to 10 species within group)
- Homogeneous groups (1 or 2 major species)

) Recognition of sets en vegetation using SVM (100% and 50% sample objects database) (16 spectral bands)



Romain Barlatier, Sébastien Gadal, Walid Ouerghemmi, 2017-2018

100%



50%

Recognition of sets en vegetation using SVM (100% and 50% of the sample object database) (64 spectral bands)



100% => 50%

Romain Barlatier, Sébastien Gadal, Walid Ouerghemmi, 2017-2018

Vegetation mapping using SVM with 100% and 50% of the sample object database (16 spectral bands)



Romain Barlatier, Sébastien Gadal, Walid Ouerghemmi, 2017-2018

100%



Classification SVM (50% échantillons) par espèces de végétation - Kaunas - Juillet

2015 (Source : HIS)

50%

LEGENDE	
■	Unclassified
■	azuolas paprastasis
■	berzas karpotasis
■	eglé paprastoji
■	gluonis trapusis
■	kastonas paprastasis
■	klevas paprastasis
■	klevas uosialapis
■	liepa mazalape
■	peulouse
■	puisis kalmiņē
■	robinija baltaziede
■	tuja vakaninē
■	uosis paprastasis

Vegetation mapping using SVM with 100% and 50% of the sample object database (64 spectral bands)



LEGENDE

■	Unclassified
■	azuolas paprastasis
■	berzas karpotasis
■	eglé paprastoji
■	gluosnis trapusis
■	kastonas paprastasis
■	klevas paprastasis
■	klevas uosialapis
■	liepa mazalape
■	peulouse
■	pusis kalnинé
■	robinija baltaziede
■	tuja vakariné
■	uosis paprastasis

Classification SVM (50% échantillons) par espèces de végétation - Kaunas -


Septembre 2016 (Source : HIS)

Romain Barlatier, Sébastien Gadal, Walid Ouerghemmi, 2017-2018

Espèces végétales	Image 16 bandes		Image 64 bandes	
	100 échantillons	50,00%	100 échantillons	50,00%
Azuolas paprastasis (%)	0	0,21	0	1,56
Berzas karpotasis (%)	0	0	13,51	44,66
Eglé paprastoji (%)	71,32	43,38	64,41	0
Gluosnis trapusis (%)	0	0	0	0
Kastonas paprastasis (%)	0	2,29	12,8	12,98
Klevas paprastasis (%)	55,18	77,9	12,06	80,39
Klevas uosialapis (%)	28,85	45,45	0,57	16,38
Liepa mazalape (%)	0	23,89	0	75,65
Pelouse (%)	31,98	87,14	22,78	97,9
Pusis kalniné (%)	19,18	0	46,6	0
Robinija baltaziede (%)	42,13	38,07	45,66	40,98
Tuja vakariné (%)	0	14,38	9,09	29,09
Uosis paprastasis (%)	4,29	6,75	16,39	15,57
OA (%)	27,39	63,22	14,72	73,29
Coeff de kappa	0,18	0,47	0,1	0,65
Feuillu (%)	28,42	39,16	15,62	69,48
Conifère (%)	31,27	22,54	36,01	12,44
Pelouse (%)	31,98	87,14	22,78	97,9
OA (%)	30,51	65,75	18,89	79,28
Coeff de kappa	0,12	0,47	0,08	0,65

Classification SVM par espèces et par groupes d'espèces sur les images HIS

Multi-temporal (seasonal) analysis on the Sentinel 2A and Landsat 8 OLI



Performances of recognition of vegetation groups

Espèce végétale	S2A Printemps	S2A Été	S2A Automne	S2A Hiver	Landsat 8
Feuillue (%)	100	98,38	66,41	0	69,42
Conifère (%)	96,67	99,3	97,57	100	98,43
Pelouse (%)	94,74	90,53	94,74	0	100
OA (%)	98,04	97,53	89,1	96,32	93,32
Coeff de kappa	0,97	0,96	0,8	0	0,84

Publications

Gintautas Mozgeris, Vytautė Juodkienė, Donatas Jonikavičius, Lina Straigytė, Sébastien Gadal, et al.. **Ultra-Light Aircraft-Based Hyperspectral and Colour-Infrared Imaging to Identify Deciduous Tree Species in an Urban Environment.** *Remote Sensing*, MDPI, 2018, 10 (10), [⟨10.3390/rs10101668⟩](https://doi.org/10.3390/rs10101668)

Walid Ouergemmi, Sébastien Gadal, Gintautas Mozgeris, Donatas Jonikavičius. **Urban Vegetation Mapping by Airborne Hyperspectral Imagery: Feasability and Limitations.** WHISPER 2018 : 9th Workshop on Hyperspectral Image and Signal Processing: Evolution in Remote Sensing, Sep 2018, Amsterdam, Netherlands. pp.245-249, 2018,

Walid Ouergemmi, Sébastien Gadal, Gintautas Mozgeris. **Urban Vegetation Mapping using Hyperspectral Imagery and Spectral Library.** *IEEE International Geoscience and Remote Sensing Symposium (IGARSS) 2018*, Jul 2018, Valencia, Spain. pp.1632-1635, [⟨https://www.igarss2018.org/⟩](https://www.igarss2018.org/)



ASU



Hyperspectral Imagery for Environmental Urban Planning