



Opposing Effects of Plant-Community Assembly Maintain Constant Litter Decomposition over Grasslands Aged from 1 to 25 Years

Lou Barbe, Andreas Prinzing, Cendrine Mony, Benjamin J. Abbott, M.
Santonja, Kevin Hoeffner, Sarah Guillocheau, Daniel Cluzeau, André-Jean
Francez, Nathalie Le Bris, et al.

► To cite this version:

Lou Barbe, Andreas Prinzing, Cendrine Mony, Benjamin J. Abbott, M. Santonja, et al.. Opposing Effects of Plant-Community Assembly Maintain Constant Litter Decomposition over Grasslands Aged from 1 to 25 Years. *Ecosystems*, 2020, 23 (1), pp.124-136. 10.1007/s10021-019-00392-8 . hal-02147166

HAL Id: hal-02147166

<https://amu.hal.science/hal-02147166>

Submitted on 4 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Opposing effects of plant-community assembly maintain constant litter decomposition over grasslands aged from 1 to 25 years

L. Barbe¹, A. Prinzing¹, C. Mony¹, B. W. Abbott^{1,2}, M. Santonja¹, K. Hoeffner¹, S. Guillocheau¹, D. Cluzeau¹, A.-J. Francez¹, N. Le Bris¹ and V. Jung¹

¹ECOBIO, OSUR, CNRS, Université de Rennes 1, 35000 Rennes, France

²Brigham Young University, Department of Plant and Wildlife Sciences, Provo, USA

Emails: lou.barbe@univ-rennes1.fr; andreas.prinzing@univ-rennes1.fr; cendrine.mony@univ-rennes1.fr; benabbo@gmail.com; mathieu.santonja@univ-rennes1.fr; kevin.hoeffner@univ-rennes1.fr; sarah.guillocheau@univ-rennes1.fr; daniel.cluzeau@univ-rennes1.fr; andre-jean.francez@univ-rennes1.fr; nathalie.lebris@univ-rennes1.fr; vincent.jung@univ-rennes1.fr

Type of article: Research Article

Abstract word count/Total word count 250/4545, number of figures/tables/appendices 4/0/3

Highlights :

- Plant-community assembly impacts plant afterlife traits and decomposer communities
- Plant-community assembly drives litter decomposition at a given successional stage
- Effects of traits and decomposers on decomposition mutually cancel out each other

Short title: Community assembly drives litter decomposition

Corresponding author: Lou Barbe (+33 6 67 44 06 56)

Authors' contributions: All authors contributed to the conception of the experiment and the data collection. LB analyzed the data, with help of AP, CM and VJ. LB wrote the manuscript, with help of all authors.

SUMMARY

Litter decomposition is central to ecosystem functioning and depends, under constant abiotic conditions, on litter quality and decomposer activity. During the assembly of a plant community following disturbance, litter quality is expected to decrease, due to an increasing proportion of resource-conservative species, whereas decomposer activity is expected to increase, due to the establishment of decomposer populations and their response to local vegetation ("Home-Field Advantage", HFA). To date, the combined effect of these processes remains poorly understood. We studied 27 semi-natural grasslands in western France, ranging from 1 to 25 years since last cultivation. We measured the functional composition of plant communities using litter traits (Specific Leaf Area, Leaf Dry Matter Content, C:N ratio, phenolics), characterized the entire community of decomposers (macrofauna, mesofauna, microbes) and performed reciprocal litter transplants to quantify HFA. We found that, overall, decomposition was rapid, and HFA was not evident. While there was substantial among-grassland variation in decomposition and HFA, neither changed with grassland age. Litter quality and decomposer efficiency also remained, overall, unchanged. However, grassland age determined all measured litter traits, and caused soil microbial C:N ratio to decline. While these changes impacted decomposition individually, together they cancelled out each other, resulting in constant decomposition across the chronosequence. Our results suggest that processes driving decomposition differ during grassland succession, and suggest that HFA may be lower in communities with high litter quality. Moreover, simultaneous assembly processes have opposing, and therefore stabilizing effects on decomposition, possibly explaining the outstanding resilience of primary production in temperate grassland ecosystems.

Key-words: plant-community assembly, decomposers and detritivores, Home-Field Advantage, grasslands, litter decomposition, plant functional traits

INTRODUCTION

The decomposition of plant litter is one of the main bottlenecks regulating carbon storage and nutrient cycling in terrestrial ecosystems (Swift and others 1979). Under a given set of abiotic conditions, litter decomposition is controlled by two main biotic parameters and their interaction (Coûteaux and others 1995): litter quality (Cornelissen and others 1999; Makkonen and others 2012), decomposer community (Petersen and Luxton 1982; Bardgett and van der Putten 2014), and response of decomposers to local litter quality (Austin and others 2014; Veen and others 2015). These biotic factors contribute to differences in decomposition among biomes, and among successional stages (e.g., grassland versus forest), however, it remains unclear whether and how they operate during the assembly of a given successional stage. Grasslands are the most widespread successional stage of temperate terrestrial biomes, and the largest terrestrial ecosystems in the globe. Although the successional stage of grasslands is maintained by agricultural practices (grazing or mowing), community assembly processes still operate, from young, species-poor grasslands dominated by the sown species to old grasslands dominated by spontaneous species (i.e. grassland succession, Cramer and others 2008). In grasslands, between 50 and 90% of plant primary production ends up as litter (Cebrian 1999). In contrast to temperate forest ecosystems, the production of litter in temperate grasslands may be sustained through the year, due to continuous growth and senescence of graminoid species. Consequently, litter decomposition is a central ecosystem process in natural and semi-natural grasslands, providing many ecosystem services, including nutrient mineralization for plant regrowth.

Throughout the assembly of a grassland plant community, litter decomposition might decrease, due to decreasing litter quality of plant species. Litter quality is determined by the digestibility and the nutritional quality of litter for decomposers, which depends on functional traits of plants (Cornelissen and Thompson 1997). For instance, litter quality decreases with

high phenol content or high C:N ratio (Hättenschwiler and Vitousek 2000; Quested and others 2007; Bakker and others 2011). Along a succession sequence, these functional traits are likely to differ among successional stages, because traits are involved in plant-plant interactions and community assembly (Hättenschwiler and Vitousek 2000; Garnier and others 2004; Violle and others 2009; Barbe and others 2017). Specifically, resource-conservative traits, which cause litter quality to decline (for instance, high C:N or high leaf dry matter content [LDMC]) may only be expressed in late-succession stages, where plant species are already established and competition is high (Wright and others 2004). During the community assembly of a given successional stage, similar mechanisms might occur and plant species present in older communities might display resource-conservative traits rather than resource-acquisitive traits. These dynamics could result in litter quality, and hence litter decomposition, decreasing with grassland community assembly.

The abundance of grassland detritivores and decomposers and their ability to decompose local litter might increase with time because of processes such as adaptation, plasticity, or ecological sorting, thereby increasing litter decomposition. Both detritivore macrofauna (earthworms) and mesofauna (Collembola, Acari and Enchytraeidae) fragment litter, accelerating the mineralization and respiration of carbohydrates by fungi and bacteria (Petersen and Luxton 1982; Coûteaux and others 1995; Bardgett and van der Putten 2014). Many decomposers are largely immobile and have slow population growth rates. At the onset of grassland community assembly, low decomposer abundance might limit litter decomposition, because grasslands might inherit few decomposers from prior land use: the abundance of detritivores and decomposers decreases considerably in row-crop agriculture, which is considered a major anthropogenic disturbance (Ponge and others 2003, 2013; Chauvat and others 2007). Consequently, time might be required to recover decomposer populations, from the same patch or from adjacent patches (Decaëns and others 2008).

Moreover, decomposer communities are able to adapt to locally-derived litter: the communities sometimes break down litter more efficiently from "Home" compared to other places, with this phenomenon being called Home-Field Advantage (HFA, Freschet and others 2012; Austin and others 2014; Veen and others 2015). This phenomenon occurs due to species sorting or selecting particular genotypes, which increase the decomposition rate of a particular type of litter through time. However, there is mixed evidence about the consistency and prevalence of HFA (Giebelmann and others 2011; Veen and others 2015). Of note, the response of decomposers to litter quality may have a lag time, depending on particular litter traits or decomposer assemblages, and might more likely to occur in plant communities with low litter quality (Milcu and Manning 2011; Veen and others 2018). Overall, the net effect of changes to decomposer abundance and response to litter quality might therefore depend on the duration of the assembly of grassland plant community, and might increase litter decomposition in old grasslands, counteracting the effects of changes to litter quality.

In this study, we investigated how litter traits, decomposer community, and their interactions affect litter decomposition during community assembly using semi-natural grasslands as a model system. We used a 25-years grassland chronosequence to test whether and how litter decomposition changes with grassland age. Specifically, we tested the following hypotheses: (i) grassland age slows down litter decomposition because litter quality declines and, (ii) grassland age accelerates litter decomposition because decomposer abundance increases or HFA emerges or increases. The chronosequence consisted of 27 grasslands in an agricultural landscape in western France (Brittany), ranging from 1 to 25 years since the last crop. We measured four functional traits aggregated at the plant community level; namely, specific leaf area (SLA), LDMC, C:N ratio, and phenolic concentration. We characterized the entire decomposer community (macrofauna, mesofauna and microbes). We performed a reciprocal

litter transplant experiment, to quantify the response of decomposer communities to local litter quality (HFA).

MATERIALS AND METHODS

Study Site and Grassland Selection

We carried out the study at the Long Term Ecological Site (LTER) «*Zone Atelier Armorique*», which is a 150 km² research area in Brittany, France (48°36'N, 1°32'W). Row crops and pastureland cover approximately 90% of the landscape, which is intersected by a well-developed hedgerow network ranging from 50 to 100 m.ha⁻¹ (Thomas and others 2016). We used ground-truthed aerial photos, which were taken every year since 1990, to construct a detailed land-use history for all sites, allowing us to determine the time since cultivation for each grassland precisely. Based on this land-use history and verification with grassland owners, we selected 27 grasslands ranging from one to 25 years since last row cropping. The grasslands were similar with respect to the initial species that were sown (*Lolium perenne* (L.) and *Trifolium repens* or *pratensis*) and management practices (annual mowing and extensive grazing, approximately 60 cows for two days once per month). The grasslands were also similar regarding environmental conditions (mesophilic grasslands) and the type of soil (brown soil that drained freely with well-developed organic and mineral horizons). The total soil C:N ratio of parcels, measured on the top 15 cm of parcels with a 20 cm diameter stainless steel hand corer, was 13.4 ± 1.7 . Parcels were $3 \text{ ha} \pm 1.4 \text{ ha}$.

Characterization of Plant Community and Plant Functional Traits

We characterized the percentage coverage of plant species in each grassland during spring 2015 using 10 quadrats of 1m² that were evenly distributed within the grassland. We identified the most abundant species(*i.e.* species accounting for 80% of the total abundance of the community; Pakeman and Quested 2007), which represented from two to five species for

each grassland (total richness from 8 to 27). For these species, we measured four functional traits related to litter quality; namely, specific leaf area (SLA), leaf dry matter content (LDMC), leaf C:N ratio and leaf phenolic concentration. These traits mainly reflect the physical and chemical properties of the litter (Hättenschwiler and Vitousek 2000; Pérez-Harguindeguy and others 2000; Santiago 2007; Quested and others 2007). To obtain a representative mean value for each trait from each species, we measured the traits of 10 individuals that were collected from randomly selected grasslands in our experiment. Phenolic concentration was measured colorimetrically, according to the Folin-Ciocalteu colorimetric method (Folin and Denis 1915) modified by Santonja and others (2015), using gallic acid as a standard. A 0.25 g litter sample was dissolved in an aqueous methanol solution, that was shaken for 1-h and then filtered. The filtered extract was mixed with Folin-Ciocalteu reagent, a color-reaction stabilizer (Na_2CO_3), and distilled water. After 1-h, phenolics concentration was measured at 765 nm on a spectrophotometer (Thermo Scientific®, USA). We measured SLA and LDMC following the standard protocols of Pérez-Harguindeguy and others (2013), and we measured C:N ratio using an elemental analyzer (FLASH EA 1112 Thermo Finnigan, Waltham, Massachusetts, USA). For each grassland, we aggregated the values of species traits to obtain an abundance-weighted mean for each trait (see Appendix Table S1 for species trait values). Correlations between traits (Appendix Table S2), with only two significant correlations being detected among the six trait correlations that we tested for, indicates low multicollinearity among traits.

Characterization of the Decomposer Community

We sampled detritivore macrofauna (*i.e.* earthworms) in each grassland during spring 2016, according to the method of Ponge and others (2013) and Cluzeau and others (2012). In brief, three 1x1m quadrats were watered three times with 10 l formalin at increasing concentration every 15 min (0.075, 0.075 and 0.12% formaldehyde solution), and a soil block (25x25cm and

20cm depth) was extracted at the end of sampling and hand sorted. Earthworms that were expelled to the surface by the irritant solution, or were recovered during the sorting of the soil block, were collected. Identification was done in the laboratory following the identification key of Cluzeau and others (unpublished, available upon request), based on Bouché (1972). Earthworms were characterized by abundance and biomass (fresh weight) and were grouped into ecological categories: epigeic or anecic – we excluded endogeic earthworms as they do not directly influence decomposition of surface litter (Bouché 1977). Then, we sampled soil detritivore mesofauna using a cylindrical soil corer (5 cm diameter \times 8 cm depth; three soil cores per grassland on two sampling dates). Mesofauna was extracted from the soil cores using the Tullgren funnel method (Berlese 1905) over a 10-day period. Collected organisms were stored in 95% ethanol, counted using a binocular scope and separated into three groups: Acari (Oribatids), Collembola and Enchytraeidae. Finally, we measured soil microbial C:N ratio, as a proxy of the composition of the microbial community, illustrating dominance of fungi or dominance of bacteria (Paul and Clark 1996; Van Elsas and others 2006). We measured the microbial C:N ratio using the fumigation-extraction method from Brookes and others (1985) and Vance and others (1987). We collected and aggregated 15 soil cores from the top 10 cm layer for each grassland to obtain a representative soil sample. From this sample, we used a 30 g soil sample that was passed through a 2-mm sieve, and was then hand-sorted to remove any visible organic material (plant roots, litter fragments). Before and after fumigation-extraction, we measured microbial C concentration using a Wet Oxidation Total Carbon Analyzer 1010 (OI-Analytical, USA) and microbial N concentration using a continuous flux AutoAnalyzer (measuring total dissolved N, AA3 Bran & Luebbe, USA). The microbial C and N concentrations were obtained by subtracting the values obtained before fumigation-extraction from the values obtained after. Microbial C:N ratio was then calculated by dividing microbial C concentration with microbial N concentration.

198 *Abiotic Parameters and Local Landscape and Management*

199 Abiotic parameters such as soil moisture and soil pH can also affect plant traits, decomposer
200 community and litter decomposition (Coûteaux and others 1995). Thus, we quantified soil
201 moisture and pH in each grassland during spring 2016 using soil samples from the top 15 cm
202 layer collected with a 20 cm diameter stainless steel hand corer. After the manual removal of
203 rocks and roots > 2 mm, soil moisture (ω) was determined by mass loss at 105°C for 24 h
204 according to the formula: $\omega = \frac{(W-D)}{D} \times 100$, where W is the wet soil mass on the sampling
205 date and D is the dry soil mass. Soil pH was determined with a 1:10 soil, de-ionized water
206 suspension, following standard methods (Robertson and others 1999). We also accounted for
207 the local landscape surrounding the selected grasslands, which might influence plant traits and
208 decomposer community. Using the database of historical land-use data, we quantified the
209 percentage of each grassland perimeter that was in contact with grasslands older than 5 years,
210 which are adjacent grasslands that represent a significant source of propagules and fauna for
211 the selected grasslands.

212 *Litter Decomposition and Transplant Experiment*

213 In each grassland, we collected litter in the fall of 2015 for the 2–5 species accounting for
214 80% of total plant abundance. We did not collect stems. For all species, we collected green
215 material for litter as in semi-natural grasslands the main portion of plant material contributing
216 to the litter pool is green material, not senesced litter (due to mowing and grazing, Mazzanti
217 and others 1994; Sanaullah and others 2010). The duration of leaf decomposition is very short
218 for the main grassland species in our experiment (especially *Lolium perenne* and legumes
219 from the genera *Trifolium*, see Results for mass losses) such their leaves decompose before
220 achieving complete senescence. Litter was cleaned, air-dried, and placed into 8x8 cm mesh
221 bags. Litterbags had 2 mm mesh on the upper side to avoid contamination by allochthonous

litter, and 5 mm mesh on the lower side to allow numerous detritivores to shuttle between the soil and litter to freely access the litter – during the period of exposure. Litter always remained moist and was never brittle. Litterbags were positioned on bare ground by gently pushing aside grasses (*i.e.* without removing any plants). Each litterbag contained 1g of litter, that was oven-dried equivalent (air-dry/oven-dry ratio calculated from subsamples that were oven-dried but not exposed to decomposition), and were constructed with the relative proportion of species from the grassland. We constructed 15 litterbags for each grassland, five of which were placed in the original grassland (“Home” decomposition) and 10 of which we randomly placed in 10 other grasslands of the experiment (“Away” decomposition) to measure HFA. We performed decomposition experiments during early spring 2016 and litterbags were collected 10 days later, when they reached 30-60% mass loss. Mass loss was measured on all 15 samples per grassland after cleaning the litter, and were oven-dried at 65 °C for three days. Mass loss (%) was calculated as $(1 - \frac{m_1}{m_0}) \times 100$, where m_0 is the initial oven-dried equivalent dry weight and m_1 is the oven-dry weight at collection.

We calculated HFA, litter quality and soil ability (*i.e.* abiotic conditions and decomposer efficiency) for each grassland by running the Decomposer Ability Regression Test (Keiser and others 2014) using SAS University Edition (SAS Institute, Cary, NC). This analysis disentangles, for a given mass loss, the effects of HFA, litter quality, and soil ability, estimated respectively by the parameters η_h , β_l and γ_s in the following equation:

$$Y_i = \alpha + \sum_{l=1}^N \beta_l \text{Litter}_{l_i} + \sum_{s=1}^M \gamma_s \text{Soil}_{s_i} + \sum_{h=1}^K \eta_h \text{Home}_{h_i} + \varepsilon_i$$

where Y_i is the decomposition of observation i , and Litter_l , Soil_s , and Home_h are dummy variables that equal 1 or 0, respectively, depending on the presence or absence of the litter mixture (from the litter mixture 1 to N), soil community (from soil community 1 to M) and

home combination (from home combinations 1 to K). The average decomposition in the data set is α (*i.e.* the intercept of the model) and ε is the error term. β_1 and γ_s were restricted to prevent perfect collinearity. The value of a given parameter that was estimated (η_h , β_1 and γ_s) indicated, as a single percentage per grassland, the difference to the mean decomposition of the dataset due to this parameter. These single parameters per grassland will be further correlated with grassland age, because there is no variation of age within grasslands.

To explore whether and how HFA is driven by litter traits and detritivore and decomposer descriptors, we calculated the HFA for the five “Home” replicates of each grassland, because plant traits vary within grasslands (similar to Veen and others 2018). We used the following equations (Giebelmann and others 2011, Pérez and others 2013):

$$(1) \text{HFA}_{a1} = \text{HDD}_{a1} - \text{ADD}_a - H$$

$$(2) \text{HDD}_{a1} = \sum (\text{D}_{a1a} - \text{D}_{ia1})$$

$$(3) \text{ADD}_a = \sum (\text{D}_{ai} - \text{D}_{ii})$$

$$(4) H = \sum \text{HDD} / (n - 1)$$

HFA_{a1} represents the Home-Field Advantage for litter replicate 1 of grassland a. Positive values indicated, as a percentage, faster home decomposition than expected, while negative values indicated slower home decomposition than expected. HDD_{a1} represents the Home Decomposition Difference for litter replicate 1 from grassland a. This value was the sum of the differences between the decomposition of this litter in its original grassland a (D_{a1a}) and the decomposition in this grassland of litters from other grasslands i (D_{ia1}). ADD_a represents the Away Decomposition Difference for grassland a. This value was the sum of the differences between the decomposition of litter from grassland a in other grasslands i and the home decomposition of grasslands i. H represents the sum of all HDD for all litters divided by amount of litters minus one. Overall, we quantified HFA per litter bag, as established in the

literature (see above), however, we stress that quantification per grassland (*i.e.* using means) led to very similar results and are hence not presented.

To explore the role of litter traits and decomposer community on litter decomposition *per se*, we calculated three mass loss values for each grassland, by averaging mass loss within the three treatment types: home litter at home, home litter away, and away litter at home. This approach allowed us to test for the numerous effects of plant traits and decomposer communities, that may vary within a grassland (similar to Veen *et al.* 2018). Analyses based on all values within grasslands led to the same conclusions (except that explained variance was slightly smaller and *P*-values were even more significant, due to the larger sample) and are therefore not presented.

Statistical Analyses

Before all the statistical tests, we center-scaled all variables (*i.e.* transformed variables by subtracting their mean and dividing by their standard deviation) to permit comparisons of regression coefficients within and among models. We used multiple ordinary least square (OLS) regression models to test the effect of grassland age on plant functional traits (SLA, LDMC, C:N ratio, phenolics) and decomposer community (soil microbial C:N ratio, abundance and biomass of epigeic earthworms and anecic earthworms, and abundance of Acari, Collembola and Enchytraeidae), accounting for abiotic parameters and local landscape (soil moisture, soil pH and percentage of adjacent grasslands). We used simple OLS regression models to test the effect of grassland age on HFA, soil ability, litter quality, and litter mass loss. We used multiple OLS regression models to test the effect of plant functional traits and decomposer community on litter mass loss and HFA (as calculated in Gießelmann and others 2011, Pérez and others 2013) also accounting for abiotic parameters, local landscape and grassland age. Of note, in the model explaining mass loss we also included the

transplantation treatment (home litter at home, home litter away, away litter at home), however, this factor was not significant and we do not present the results. To summarize the results of the individual regression analyses, we conducted a path analysis (Wright 1934), which allowed us to visualize the strength of the indirect effects of grassland age on litter mass loss and HFA, which were effects mediated by plant functional traits and decomposer community. For each indirect effect, we calculated a compound path by multiplying: (i) the standardized regression coefficient from the model that related grassland age to plant functional traits or decomposer community with (ii) the standardized regression coefficient from the model that related plant functional traits or decomposer community to litter mass loss. Finally, we compared, for litter mass loss and HFA, the sum of the indirect effects of grassland age with the single overall effect of grassland age. We graphically examined residuals using probability plots and predicted versus residual plots, indicating that all residual distributions fulfilled the assumption of normality and homogeneity. All statistical analyses were performed with R 3.0.3 (R Development Core Team 2016).

RESULTS

Plant Functional Traits and Decomposer Community Depend on Grassland Age

We observed large differences in the functional traits of the plant community and in the composition of the decomposer community among entire grasslands. Grassland age strongly explained all functional traits of the plant community (simple regression analyses, Figure 1). Specifically, age increased leaf C:N ratio ($F_{18}=11.07$, $P=7.10^{-4}$, $\text{adj-R}^2=0.50$), leaf phenolic concentration ($F_{18}=4.19$, $P=0.03$, $\text{adj-R}^2=0.24$), and SLA ($F_{19}=3.71$, $P=0.04$, $R^2=0.20$), while it decreased LDMC ($F_{25}=31.64$, $P=7.10^{-6}$, $\text{adj-R}^2=0.52$). Old grasslands hence harbored a plant community with mainly thin, moist leaves; however, these leaves contained lower nitrogen concentration and higher phenolic content. Grassland age also increased soil

microbial C:N ratio ($F_{20}=5.32$, $P=0.03$, $R^2=0.17$, Figure 1), as well as the abundance of Acari ($F_{23}=9.20$, $P=6.10^{-3}$, $R^2=0.25$, Figure 1). Earthworm abundance, as well as the abundance of Collembola and Enchytraeidae, was not associated with grassland age; however, the percentage of adjacent grasslands increased the abundance of anecic earthworms (Figure S1). Despite multiple relationships between age and either litter traits or decomposer community, we found one significant correlation between litter traits and detritivore and decomposer community (Acari vs LDMC, $r=-0.54$, $P<0.05$, see Table S2).

Litter Decomposition and Home-Field Advantage Differ Among Grasslands and Depend on Plant Functional Traits and the Decomposer Community

We observed large among-grassland differences in litter mass loss and HFA (Figure S2). Overall, across all of the litters in our experiment, we did not observe a positive mean HFA (no significant difference from a mean of 0, $t=1.41$, $df=133$, $P=0.16$). Grassland age did not directly influence litter mass loss ($F_{25}=0.29$, $P=0.60$, Figure 2), HFA ($F_{25}=0.41$, $P=0.53$, Figure 2), soil ability ($F_{25}=0.71$, $P=0.41$, Figure 2) nor litter quality ($F_{25}=0.22$, $P=0.4$, Figure 2). However, several plant functional traits and the soil microbial C:N ratio, related to grassland age, strongly explained litter mass loss ($F_{56}=12.43$, $P=7.10^{-9}$, $\text{adj-}R^2=0.53$, Figure 3): the leaf C:N ratio decreased litter mass loss, as did LDMC and SLA (Figure 3). Decrease in soil microbial C:N was associated with decrease in litter mass loss (Figure 3). Moreover, soil moisture decreased litter mass loss (Figure S3). None of the plant functional traits influenced HFA, but HFA increased when anecic earthworms were more abundant (Figure S1, $F_{101}=6.37$, $P=0.044$, semi-partial $R^2=0.06$) and when soil moisture was low (Figure S3, $F_{101}=6.37$, $P=0.013$, semi-partial $R^2=0.03$). We also note that, when explaining litter mass loss by functional traits and decomposer community, a positive, direct effect of grassland age on litter mass loss was detected after removal of all other significant effects (Figure S4). Finally, as plant functional traits and soil microbial C:N were influenced by grassland age, we

obtained the following indirect effects of grassland age on litter mass loss (Figure S4): -0.42 via leaf C:N ratio, -0.26 via SLA, 0.53 via LDMC, -0.15 via the soil microbial C:N ratio and 0.22 via an unmeasured parameter. The sum of all indirect effects of grassland age on litter mass loss was -0.08, indicating no significant total effect, which was consistent with the lack of an overall effect of grassland age on litter mass loss that we initially observed (Figure 2).

DISCUSSION

Grassland age influenced all measured functional traits of the plant community, as well as the soil microbial C:N ratio. These changes, in turn, impacted litter mass loss, but they cancelled each other out, resulting in constant litter decomposition across the chronosequence. Litter overall mass loss, litter quality (*i.e.* litter decomposability), decomposer efficiency, and HFA did not change with grassland age.

Plant Functional Traits Change with Age, but not towards Resource-Conservatism

The four functional traits that we measured on plant communities strongly responded to grassland age. Specifically, the plant communities of old grasslands displayed a higher leaf C:N ratio and phenolic concentration, along with higher SLA and lower LDMC. Thus, grassland assembly favored resource-conservative trait values as much as resource-acquisitive trait values. This result contrasted with what we expected from other studies (Wright and others 2004; Quested and others 2007). These changes in functional traits were probably due to plant turnover during grassland assembly, with new species establishing and dominating initial species that had higher C:N and phenolic concentration (for instance, *Ranunculus repens* or *Dactylis glomerata*, Figure S5). These characteristics might enhance the resistance of these new species against herbivores (Bernays and others 1989). Moreover, high SLA and low LDMC might facilitate high resource acquisition and hence regrowth to compensate for herbivory (Briske 1996). This observed plant succession during grassland assembly might be

attributed to the immigration of new plant species from the surrounding landscape or by the germination of seeds already present in the soil seed bank. Overall, these results show that shifts to plant afterlife traits arise during the assembly of a successional stage (*i.e.* grasslands), not only across successional stages, for example after grassland abandonment (Kahmen and Poschlod 2004; Quested and others 2007).

Offsetting Effects of Plant Traits on Litter Decomposition

Individually, each of these changes to plant traits (except changes to phenolics) influenced litter decomposition, but they did not reduce litter decomposition, which remained unchanged with grassland age overall: an increase in SLA and C:N ratio decreased litter decomposition, but this was compensated by a decrease in LDMC which increased litter decomposition. While the effects of the C:N ratio and LDMC are consistent with the literature (Bakker and others 2011; Pakeman and others 2011), the negative effect of SLA on mass loss was inconsistent with the literature (Santiago 2007; Makkonen and others 2012). However, studies on SLA usually focus on SLA values distinctly smaller than ours. This negative relationship might be explained by the higher compaction of litter with very thin leaves (*i.e.* high SLA), slowing down the colonization of litters by detritivores and decomposers. Alternatively, compensatory feeding (*i.e.* enhanced consumption rate on poor-quality litter to ensure sufficient resource assimilation to meet metabolic needs; Gessner and others 2010) on litter with low SLA might also explain the negative effect of SLA on mass loss. Finally, the lack of influence of leaf phenolic concentration on decomposition was possibly due to the lower leaf phenolic concentration of grassland plants compared to other systems (Hättenschwiler and Vitousek 2000) which might have prevented the effects from being detectable. Taken together, our results suggest, contrary to what we expected, that community assembly processes result in similar litter quality during grassland succession, likely due to compensatory effects on afterlife traits and litter quality.

391 *Little Impact of Grassland Age on Detritivores and Decomposers*

392 Only some components of decomposer communities responded to grassland age, partially
393 supporting our predictions. Anecic and epigeic earthworms (macrofauna) did not respond to
394 grassland age; however, the abundance of anecic earthworms increased in grasslands that
395 shared more edges with other grasslands, suggesting the immigration of anecics from the
396 surrounding landscape, consistently with their high dispersal ability (Caro and others 2013).
397 Considering mesofauna and microbes, we observed two shifts: the abundance of Oribatids
398 (Acari, detritivore mesofauna) increased with grassland age, as did the soil microbial C:N
399 ratio. The increase in the abundance of Oribatids might be attributed to the recovery of
400 populations of these soil organisms after land-use turnover and the disruptive agricultural
401 practices before the sowing of grasslands (Hülsmann and Wolters 1998). Recovery might be
402 permitted by the immigration of new species from an adjacent landscape, or by the
403 demographic development of the remaining initial species. Finally, the increase in soil
404 microbial C:N ratio with grassland age might be attributed to a reduction in mineral N input
405 after cropping, ultimately causing the depletion of microbial N (Wardle 1992).

406 *Strong Functional Redundancy of Detritivores and Decomposers in Grassland Ecosystems?*

407 Unexpectedly, the abundance of macrofauna and mesofauna did not influence decomposition
408 (Milcu and others 2008), suggesting that microbial and fungal decomposition were not limited
409 by the fragmentation of leaves. The only decomposer-mediated effect we could identify was
410 the increase in soil microbial C:N ratio that was associated with decreasing litter mass loss.
411 Possibly, high soil microbial C:N also reflected a change in the composition of the microbial
412 community, from a dominance of bacteria to a dominance of fungi (Paul and Clark 1996; Van
413 Elsas and others 2006), which might be less efficient at decomposing labile litter (Santonja
414 and others 2018). This slight, microbial C:N-mediated, decrease in litter mass loss was

probably compensated for by a slight increase of litter mass loss due to a statistically direct effect of grassland age in the full model; this effect was necessarily that of age on one or several parameters that we did not measure (and it may be a biologically indirect effect, like the effects of plant traits and decomposers that we observed), and it explains why soil ability, which represents decomposer efficiency and abiotic conditions, remained globally similar among grasslands. Overall, the lack of relationships between detritivores and decomposers with decomposition might be attributed to the high functional redundancy of these groups for decomposing grassland litter (Gessner and others 2010). Despite variation in the abundance of detritivores and decomposers in our grasslands, there was no effect of any particular group on decomposition. Given the high decomposability of litter in grasslands, the effects of detritivores and decomposers on decomposition might be particularly redundant, with this feature being very stable across time.

No Impact of Grassland Age on Home-Field Advantage, Due to High Litter Decomposability?

We found, overall, a low HFA, with a substantial variation across grasslands. We found only one significant relationship between the numerous plant traits and detritivores and decomposers that we sampled, not indicating ecological sorting or abundance shifts of detritivores and decomposers in response to litter traits. Overall, we found no relationship between grassland age and HFA: grassland age did not influence HFA, nor did plant functional traits and soil microbial C:N ratio. Besides, litter decomposition rates were very high in our experiment, resulting from the very high decomposability of the green litter of graminoids and *Trifolium sp.* (Sanaullah and others 2010 for similar decomposition rates). High decomposition also resulted from the humid and relatively warm, frost-free winter in the study region which increases decomposer activity (this rapidity is also illustrated by the absence of a distinct litter layer in the grasslands). Therefore, our results support that strong HFA may only arise when home litter is very recalcitrant or very dissimilar to other litter

(Milcu and Manning 2011; Li and others 2017; Lu and others 2017; Palozzi and Lindo 2018; Hoyos-Santillan and others 2018). Only increased abundance of anecic earthworms triggered some HFA. This interesting phenomenon might reflect, for instance, the response of digestive enzymes of anecic earthworms to local litter (Eisenhauer and others 2012): the digestive metabolism of earthworms could acclimate to home litter, and then secrete more enzymes that are adapted to home litter, which would increase digestion and hence litter consumption. Alternatively, in their permanent galleries, the mucus of anecic earthworms might have been a durable, high quality substrate for decomposer microorganisms, giving them time to adapt to local litter, ultimately triggering HFA. Finally, high soil moisture triggered a slight Home-Field Disadvantage. This might be due to waterlogging triggering anoxia in the top soil, decreasing bacterial and fungal activity (thus decreasing mass loss per se) especially in soft, easily hydratable and compactable litters produced within the home environment. Overall, these results show that grassland maturation did not drive HFA, and that HFA might only occur when litter is very recalcitrant or dissimilar, which was not the case in our system. Hence, HFA did not require experience of local decomposers with local plants, but rather environments favorable for macrofauna and aerobic decomposition. Earthworms might have a role in this process, despite being quite generalist organisms, with microbes not being the only biotic drivers of HFA (Freschet and others 2012; Austin and others 2014). Further studies, including the sampling of detritivore and decomposer traits, and including the exclusion of fauna, are required to elucidate the precise role of fauna and microbes on HFA.

CONCLUSIONS

We found that community assembly maintained similar litter decomposition along a 25-years chronosequence, and that HFA was not associated with plant community assembly or age. HFA was quite low and was only associated with the macrofauna community and soil moisture, which were not related to grassland age. However, our results showed that several

plant functional traits and the soil microbial C:N ratio changed during community assembly. In turn, these changes acted on decomposition, but cancelled out each other, resulting in the same decomposition rate across succession. These results also suggest that in temperate grassland ecosystems, where decomposition is very fast, numerous and important changes in plant and decomposer communities lead to the same levels of ecosystem functions, possibly explaining the outstanding resilience of many grassland ecosystems despite being often a transitional stage.

ACKNOWLEDGEMENTS

We thank Aurélien Pierre for measuring the soil microbial C:N ratio, and we acknowledge Albin Fertil, Daniel Cylly, André Bastin, Valentin Blanchard, Romain Georges, Olivier Jambon, Jérémy Guy, Nathan Vannier and Stéphanie Llopis for help with sampling earthworms. We are very grateful to Mathilde Le Moing for earthworm identification. We thank Jean-Sébastien Pierre for helping with the SAS statistical analyses. The work benefited from the support of the LTER site “ZA Armorique”.

REFERENCES

- Austin AT, Vivanco L, González-Arzac A, Pérez LI. 2014. There's no place like home? An exploration of the mechanisms behind plant litter-decomposer affinity in terrestrial ecosystems. *New Phytologist* 204: 307–314.
- Bakker MA, Carreño-Rocabado G, Poorter L. 2011. Leaf economics traits predict litter decomposition of tropical plants and differ among land use types: Leaf economics traits and decomposition. *Functional Ecology* 25: 473–483.
- Barbe L, Jung V, Prinzing A, Bittebiere A-K, Butenschoen O, Mony C. 2017. Functionally dissimilar neighbors accelerate litter decomposition in two grass species. *New Phytologist* 214: 1092–1102.
- Bardgett RD, van der Putten WH. 2014. Belowground biodiversity and ecosystem functioning. *Nature* 515: 505–511.
- Berlese A. 1905. Apparichio per raccogliere presto ed in gran numero di piccolo artropodi. *Redia* 2: 85–89.
- Bernays EA, Driver GC, Bilgener M. 1989. Herbivores and plant tannins. *Advances in ecological research* 19: 263–302.
- Bouché, MB. 1972. *Lombriciens de France: écologie et systématique*, Paris.
- Bouché MB. 1977. Stratégies lombriciennes. *Ecological Bulletins* 25: 122–132.
- Briske DD. 1996. Strategies of plant survival in grazed systems: a functional interpretation. *The ecology and management of grazing systems* 37–67.

507 Brookes PC, Landman A, Pruden G, Jenkinson DS. 1985. Chloroform fumigation and the
 508 release of soil nitrogen: a rapid direct extraction method to measure microbial biomass
 509 nitrogen in soil. *Soil Biology and Biochemistry* 17: 837–842.

510 Caro G, Decaëns T, Lecarpentier C, Mathieu J. 2013. Are dispersal behaviours of earthworms
 511 related to their functional group? *Soil Biology and Biochemistry* 58: 181–187.

512 Cebrian J. 1999. Patterns in the fate of production in plant communities. *The American*
 513 *Naturalist* 154: 449–468.

514 Chauvat M, Wolters V, Dauber J. 2007. Response of collembolan communities to land-use
 515 change and grassland succession. *Ecography* 30: 183–192.

516 Cluzeau D, Guernion M, Chaussod R, Martin-Laurent F, Villenave C, Cortet J, Ruiz-
 517 Camacho N, Pernin C, Mateille T, Philippot L, Bellido A, Rougé L, Arrouays D, Bispo A,
 518 Pérès G. 2012. Integration of biodiversity in soil quality monitoring: Baselines for microbial
 519 and soil fauna parameters for different land-use types. *European Journal of Soil Biology* 49:
 520 63–72.

521 Cornelissen JHC, Pérez-Harguindeguy N, Diaz S, Grime JP, Marzano B, Cabido M,
 522 Vendramini F, Cerabolini B. 1999. Leaf structure and defence control litter decomposition
 523 rate across species and life forms in regional floras on two continents. *New Phytologist* 143:
 524 191–200.

525 Cornelissen JHC, Thompson K. 1997. Functional leaf attributes predict litter decomposition
 526 rate in herbaceous plants. *New Phytologist* 135: 109–114.

527 Coûteaux M-M, Bottner P, Berg B. 1995. Litter decomposition, climate and litter quality.
 528 *Trends in Ecology & Evolution* 10: 63–66.

529 Cramer VA, Hobbs RJ, Standish RJ. 2008. What's new about old fields? Land abandonment
530 and ecosystem assembly. *Trends in Ecology & Evolution* 23: 104–112.

531 Decaëns T, Margerie P, Aubert M, Hedde M, Bureau F. 2008. Assembly rules within
532 earthworm communities in North-Western France—A regional analysis. *Applied Soil Ecology*
533 39: 321–335.

534 Eisenhauer N, Reich PB, Isbell F. 2012. Decomposer diversity and identity influence plant
535 diversity effects on ecosystem functioning. *Ecology* 93: 2227–2240.

536 Folin O, Denis W. 1915. A colorimetric method for the determination of phenols (and phenol
537 derivatives) in urine. *Journal of Biological Chemistry* 22: 305–308.

538 Freschet GT, Aerts R, Cornelissen JHC. 2012. Multiple mechanisms for trait effects on litter
539 decomposition: moving beyond home-field advantage with a new hypothesis: Substrate-
540 matrix quality interactions in decay. *Journal of Ecology* 100: 619–630.

541 Garnier E, Cortez J, Billès G, Navas M-L, Roumet C, Debussche M, Laurent G, Blanchard A,
542 Aubry D, Bellmann A, Neill C, Toussaint J-P. 2004. Plant functional markers capture
543 ecosystem properties during secondary succession. *Ecology* 85: 2630–2637.

544 Gessner MO, Swan CM, Dang CK, McKie BG, Bardgett RD, Wall DH, Hättenschwiler S.
545 2010. Diversity meets decomposition. *Trends in ecology & evolution* 25: 372–380.

546 Giebelmann UC, Martins KG, Brändle M, Schädler M, Marques R, Brandl R. 2011. Lack of
547 home-field advantage in the decomposition of leaf litter in the Atlantic Rainforest of Brazil.
548 *Applied Soil Ecology* 49: 5–10.

549 Güsewell S, Gessner MO. 2009. N : P ratios influence litter decomposition and colonization
550 by fungi and bacteria in microcosms. *Functional Ecology* 23: 211–219.

551 Hättenschwiler S, Vitousek PM. 2000. The role of polyphenols in terrestrial ecosystem
 552 nutrient cycling. *Trends in Ecology & Evolution* 15: 238–243.

553 Hülsmann A, Wolters V. 1998. The effects of different tillage practices on soil mites, with
 554 particular reference to Oribatida. *Applied Soil Ecology* 9: 327–332.

555 Hoyos-Santillan J, Lomax BH, Turner BL, Sjögersten S. 2018. Nutrient limitation or home
 556 field advantage: Does microbial community adaptation overcome nutrient limitation of litter
 557 decomposition in a tropical peatland? *Journal of Ecology* 106: 1558–1569.

558 Kahmen S, Poschlod P. 2004. Plant functional trait responses to grassland succession over 25
 559 years. *Journal of Vegetation Science* 15: 21–32.

560 Keiser AD, Keiser DA, Strickland MS, Bradford MA. 2014. Disentangling the mechanisms
 561 underlying functional differences among decomposer communities. *Journal of Ecology* 102:
 562 603–609.

563 Li YB, Li Q, Yang JJ, Lü XT, Liang WJ, Han XG, Bezemer TM. 2017. Home-field
 564 advantages of litter decomposition increase with increasing N deposition rates: a litter and soil
 565 perspective. *Functional Ecology* 31: 1792–1801.

566 Lu W, Liu N, Zhang Y, Zhou J, Guo Y, Yang X. 2017. Impact of vegetation community on
 567 litter decomposition: Evidence from a reciprocal transplant study with ¹³C labeled plant litter.
 568 *Soil Biology and Biochemistry* 112: 248–257.

569 Makkonen M, Berg MP, Handa IT, Hättenschwiler S, van Ruijven J, van Bodegom PM, Aerts
 570 R. 2012. Highly consistent effects of plant litter identity and functional traits on
 571 decomposition across a latitudinal gradient. *Ecology Letters* 15: 1033–1041.

572 Mazzanti A, Lemaire G, Gastal F. 1994. The effect of nitrogen fertilization upon the herbage
 573 production of tall fescue swards continuously grazed with sheep. II-herbage consumption.
 574 Grass and Forage Science 49: 352–359.

575 Milcu A, Manning P. 2011. All size classes of soil fauna and litter quality control the
 576 acceleration of litter decay in its home environment. Oikos 120: 1366–1370.

577 Milcu A, Partsch S, Scherber C, Weisser WW, Scheu S. 2008. Earthworms and legumes
 578 control litter decomposition in a plant diversity gradient. Ecology 89: 1872–1882.

579 Pakeman RJ, Eastwood A, Scobie A. 2011. Leaf dry matter content as a predictor of grassland
 580 litter decomposition: a test of the “mass ratio hypothesis”. Plant and Soil 342: 49–57.

581 Pakeman RJ, Quested HM. 2007. Sampling plant functional traits: What proportion of the
 582 species need to be measured? Applied Vegetation Science 10: 91–96.

583 Palozzi JE, Lindo Z. 2018. Are leaf litter and microbes team players? Interpreting home-field
 584 advantage decomposition dynamics. Soil Biology and Biochemistry 124: 189–198.

585 Paul, EA, Clark FE. 1996. Soil Microbiology and Biochemistry. Academic Press, California.
 586 340p.

587 Perez G, Aubert M, Decaëns T, Trap J, Chauvat M. 2013. Home-field advantage: a matter of
 588 interaction between litter biochemistry and decomposer biota. Soil Biology and Biochemistry
 589 67: 245–254.

590 Pérez-Harguindeguy N, Diaz S, Cornelissen JHC, Vendramini F, Cabido M, Castellanos A.
 591 2000. Chemistry and toughness predict leaf litter decomposition rates over wide spectrum of
 592 functional types and taxa in central Argentina. Plant and Soil 218: 21–30.

593 Pérez-Harguindeguy N, Díaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte
 594 MS, Cornwell WK, Craine JM, Gurvich DE, Urcelay C, Veneklaas EJ, Reich PB, Poorter L,
 595 Wright IJ, Ray P, Enrico L, Pausas JG, de Vos AC, Buchmann N, Funes G, Quétier F,
 596 Hodgson JG, Thompson K, Morgan HD, ter Steege H, Sack L, Blonder B, Poschlod P,
 597 Vaieretti MV, Conti G, Staver AC, Aquino S, Cornelissen JHC. 2013. New handbook for
 598 standardised measurement of plant functional traits worldwide. *Australian Journal of Botan*,
 599 61: 167–234.

600 Petersen H, Luxton M. 1982. A Comparative Analysis of Soil Fauna Populations and Their
 601 Role in Decomposition Processes. *Oikos* 39: 288–388.

602 Ponge J-F, Gillet S, Dubs F, Fedoroff E, Haese L, Sousa JP, Lavelle P. 2003. Collembolan
 603 communities as bioindicators of land use intensification. *Soil Biology and Biochemistry* 35:
 604 813–826.

605 Ponge J-F, Pérès G, Guernion M, Ruiz-Camacho N, Cortet J, Pernin C, Villenave C,
 606 Chaussod R, Martin-Laurent F, Bispo A, Cluzeau D. 2013. The impact of agricultural
 607 practices on soil biota: A regional study. *Soil Biology and Biochemistry* 67: 271–284.

608 Quested H, Eriksson O, Fortunel C, Garnier E. 2007. Plant traits relate to whole-community
 609 litter quality and decomposition following land use change. *Functional Ecology* 21: 1016–
 610 1026.

611 R Core Team (2016) R: A language and environment for statistical computing. R Foundation
 612 for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

613 Robertson GP. 1999. Standard soil methods for long-term ecological research (Vol. 2),
 614 Oxford University Press.

615 Sanaullah M, Chabbi A, Lemaire G, Charrier X, Rumpel C. 2010. How does plant leaf
 616 senescence of grassland species influence decomposition kinetics and litter compounds
 617 dynamics? *Nutrient Cycling in Agroecosystems* 88: 159–171.

618 Santiago LS. 2007. Extending the leaf economics spectrum to decomposition: evidence from
 619 a tropical forest. *Ecology* 88: 1126–1131.

620 Santonja M, Fernandez C, Gauquelin T, Baldy V. 2015. Climate change effects on litter
 621 decomposition: intensive drought leads to a strong decrease of litter mixture interactions.
 622 *Plant and Soil* 393: 69–82.

623 Santonja M, Foucault Q, Rancon A, Gauquelin T, Fernandez C, Baldy V, Mirleau P. 2018.
 624 Contrasting responses of bacterial and fungal communities to plant litter diversity in a
 625 Mediterranean oak forest. *Soil Biology and Biochemistry* 125: 27–36.

626 Swift MJ, Heal OW, Anderson JM. 1979. *Decomposition in terrestrial ecosystems* (Vol. 5),
 627 Univ of California Press.

628 Thomas Z, Abbott BW, Troccaz O, Baudry J, Pinay G. 2016. Proximate and ultimate controls
 629 on carbon and nutrient dynamics of small agricultural catchments. *Biogeosciences* 13: 1863–
 630 1875.

631 Vance ED, Brookes PC, Jenkinson DS. 1987. An extraction method for measuring soil
 632 microbial biomass C. *Soil Biology and Biochemistry* 19: 703–707.

633 Van Elsas JD, Trevors JT, Jansson JK, Nannipieri P. 2006. *Modern soil microbiology*. CRC
 634 Press, Florida. 683p.

635 Veen GFC, Freschet GT, Ordonez A, Wardle DA. 2015. Litter quality and environmental
 636 controls of home-field advantage effects on litter decomposition. *Oikos* 124: 187–195.

- Veen GFC, Keiser AD, van der Putten WH, Wardle DA. 2018. Variation in home-field advantage and ability in leaf litter decomposition across successional gradients. *Functional Ecology* 32 :1563–1574.
- Violle C, Garnier E, Lecoœur J, Roumet C, Pothier C, Blanchard A, Navas M-L. 2009. Competition, traits and resource depletion in plant communities. *Oecologia* 160: 747–755.
- Wardle DA. 1992. A comparative assessment of factors which influence microbial biomass carbon and nitrogen levels in soil. *Biological reviews* 67: 321–358.
- Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F, Cavender-Bares J, Chapin T, Cornelissen JHC, Diemer M, Flexas J, Garnier E, Groom PK, Gulias J, Hikosaka K, Lamont BB, Lee T, Lee W, Lusk C, Midgley JJ, Navas M-L, Niinemets Ü, Oleksyn J, Osada N, Poorter H, Poot P, Prior L, Pyankov VI, Roumet C, Thomas SC, Tjoelker MG, Veneklass EJ, Villar R. 2004. The worldwide leaf economics spectrum. *Nature* 428: 821–827.
- Wright S. 1934. The method of path coefficients. *The Annals of Mathematical Statistics* 5: 161–215.

Figure captions


Figure 1. Grassland age significantly influenced all aggregated functional traits of the plant community (phenolics, leaf C:N ratio, Leaf Dry Matter Content, and Specific Leaf Area) as well as the abundance of soil Acari (mesofauna) and soil microbial C:N ratio. Green points represent plant trait values, orange points represent values of decomposer and detritivore indicators. See Results for all model parameters.

Figure 2. Grassland age did not influence Home-Field Advantage (HFA), soil ability (illustrating soil conditions and decomposer efficiency), litter quality (*i.e.* litter decomposability) nor litter overall mass loss (calculated as the mean per grassland across all litters). See Material and Methods for the transplantation treatment and calculations.

Figure 3. Three aggregated functional traits of plant community and soil microbial C:N ratio decreased litter mass loss. Each graph presents partial residuals, so accounts for the simultaneous effect of the other variables in the model. SP- R^2 represents semi-partial residuals. Green points represent plant trait values, and orange points represent values of decomposer and detritivore indicators. The three data points per grassland are the mean of the three treatments (home litter exposed at home, home litter exposed away, and away litter exposed at home; the factor containing these treatments did not significantly interact with age in the statistical model explaining mass loss). See Material and Methods for the transplantation treatment and calculations, and Results for all model parameters.

681 **FIGURES**

682



683

684 **Figure 1.**




Figure 2.




Figure 3.

SUPPORTING INFORMATION

Table S1. Mean values of functional traits measured on plant species representing 80% of the grassland communities in our experimental design (see Material and Methods) and that were accounted for in the decomposition experiment. Phenolics and LDMC (Leaf Dry Matter Content) are in %, SLA (Specific Leaf Area) is in $\text{mm}^2.\text{g}^{-1}$, and C:N ratio has no unit.

Species	Functional traits							
	Phenolics		C:N ratio		LDMC		SLA	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>Agrostis stolonifera</i>	3.32	0.19	10.78	1.35	21.67	2.09	36.22	3.38
<i>Agrostis tenuis</i>	3.02	0.07	10.36	0.07	32.90	0.66	22.60	1.23
<i>Dactylis glomerata</i>	3.58	0.27	10.47	0.30	16.81	1.19	37.48	8.03
<i>Holcus lanatus</i>	2.92	0.09	9.79	0.16	19.92	2.40	43.53	9.50
<i>Lolium italicum</i>	2.56	0.12	9.30	0.09	22.11	1.78	36.25	5.65
<i>Lolium perenne</i>	2.80	0.10	9.17	0.05	25.00	2.78	22.81	2.71
<i>Ranunculus repens</i>	3.58	0.30	12.22	0.51	14.79	2.62	31.70	7.61
<i>Trifolium pratense</i>	3.82	0.15	9.36	0.20	21.31	1.45	18.90	4.59
<i>Trifolium repens</i>	3.76	0.16	8.93	0.09	20.22	2.01	29.45	5.64

Table S2. Correlations between plant functional traits, between decomposers and between plant functional traits and decomposers (Leaf Dry Matter Content, LDMC; Specific Leaf Area, SLA). For each correlation, the given parameter is the Pearson's or Spearman's coefficient (depending on the normality of variable distribution) with its significance (*: $P<0.05$; ***: $P<0.001$).

		<i>Plant functional traits</i>				<i>Decomposer community</i>				
		LDMC	SLA	Leaf C:N ratio	Phenolics	Microbial C:N ratio	Collembola	Acari	Enchytraeidae	Anecis
<i>Plant functional traits</i>	SLA	-0.13								
	Leaf C:N ratio	-0.07	0.45*							
	Phenolics	0.01	0.20	0.79***						
	Microbial C:N ratio	-0.34	0.26	0.26	0.18					
<i>Decomposer community</i>	Collembola	-0.22	-0.24	0.07	0.23	0.15				
	Acari	-0.54*	0.14	0.39	0.35	0.43*	0.54*			
	Enchytraeidae	-0.13	-0.16	-0.03	0.31	-0.09	-0.09	0.07		
	Anecis	0.01	-0.34	-0.10	0.22	-0.04	0.30	0.17	0.25	
	Epigeics	0.08	0.04	0.02	-0.01	-0.17	-0.23	-0.10	0.40	0.14




Figure S1. Percentage of adjacent grasslands increased the abundance of anecic earthworms, which, in turn, increased Home-Field Advantage (HFA). SP-R² represents semi-partial residuals. See Figure S4 for parameters of models, and see Material and Methods for the transplantation treatment and calculations of HFA.




Figure S2. Boxplot of mass loss and Home-Field Advantage (HFA) for all litters from the grasslands in our experiment. Grasslands are ordered by age, from the youngest (A, one year old) to the oldest (AA, 25 years old). We observe that, while there is substantial intra-grassland variation, there is still important inter-grassland variation for mass loss (ANOVA with grassland as explicative factor: $P < 10^{-16}$, $r^2 = 0.27$) and HFA (ANOVA with grassland as explicative factor: $P = 4.9 \times 10^{-4}$, $r^2 = 0.23$).






Figure S3. Soil moisture decreased litter mass loss and Home-Field Advantage (HFA). SP-R² represents semi-partial residuals. See Results and Figure S4 the model parameters, and see Material and Methods for the transplantation treatment and calculations of HFA.



750

751 **Figure S4.** Path analysis model for the indirect effects of grassland age on litter mass loss and Home-Field Advantage. These effects are
 752 mediated by the functional traits of plant community and by the decomposer community. The figure also presents the effects of abiotic and
 753 landscape variables accounted for in the models. Effects are indicated as the standardized regression coefficients of the model, that is the sign and
 754 magnitude of the effect. Explained variances are indicated in with each dependent variable, with adj-R². The dotted line represents a statistically
 755 direct effect of grassland age on litter decomposition in the model (*i.e.* a biological indirect effect of grassland age that was not mediated by any
 756 of the parameters that we measured). The relationship between soil pH and phenolics might be the least causal within the analysis.



757

758 **Figure S5.** Trend lines of mean cover percentage of the nine species used in the
 759 decomposition experiment across grassland ageing. These species represent 80% of the total
 760 abundance of the plant community. See Table S1 for species traits.