



HAL
open science

Leaf Nutrients and Macroinvertebrates Control Litter Mixing Effects on Decomposition in Temperate Streams

Mathieu Santonja, Hector Rodriguez-Perez, Nathalie Le Bris, Christophe Piscart

► **To cite this version:**

Mathieu Santonja, Hector Rodriguez-Perez, Nathalie Le Bris, Christophe Piscart. Leaf Nutrients and Macroinvertebrates Control Litter Mixing Effects on Decomposition in Temperate Streams. *Ecosystems*, 2020, 23 (2), pp.400-416. 10.1007/s10021-019-00410-9 . hal-02165628

HAL Id: hal-02165628

<https://amu.hal.science/hal-02165628>

Submitted on 9 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **TITLE:** Leaf nutrients and macroinvertebrates control litter mixing effects on decomposition in
2 temperate streams

3

4 **RUNNING HEAD:** Litter mixture decomposition in temperate streams

5

6 **AUTHORS:** Mathieu Santonja^{1,2*}, Héctor Rodríguez-Pérez¹, Nathalie Le Bris¹, Christophe
7 Piscart¹

8

9 **ADDRESSES**

10 **1.** Univ Rennes, CNRS, ECOBIO UMR 6553, F-35000 Rennes, France.

11 **2.** Aix Marseille Univ, Avignon Université, CNRS, IRD, IMBE, Marseille, France.

12

13 ***CORRESPONDING AUTHOR**

14 Email: mathieu.santonja@gmail.com

15 ORCID: 0000-0002-6322-6352

16

17 **ABSTRACT**

18 Plant litter decomposition is an essential ecosystem function in temperate streams. Both
19 riparian vegetation and decomposer communities are major determinants of the decomposition
20 efficiency and the interactions occurring within litter mixtures. However, the extent to which
21 such litter mixture interactions are affected by combined shifts in litter traits and decomposer
22 community is not well understood. We used leaf litter from 10 European tree species in order
23 to study litter decomposition and litter mixture effects occurring in two-species litter mixtures
24 in a temperate forested stream of northwestern France. The study distinguished between (i)
25 decomposition involving microorganisms alone or together with invertebrates, and (ii)
26 decomposition involving litter mixtures of similar or dissimilar nutrient content. Increasing
27 mean litter nutrient concentration favored both microbial activity and litter decomposition rate.
28 Surprisingly, the highest litter mixture effects occurred in mixtures containing two nutrient-rich
29 litters and occurred mainly in macroinvertebrate presence. Both the “mass-ratio hypothesis”,
30 expressed as the community weighted mean traits ($Trait_{CWM}$), and the “niche complementarity
31 hypothesis”, expressed as the functional dissimilarity of litter traits ($Trait_{FD}$), contributed to
32 explain litter mixture effects. However, $Trait_{CWM}$ was found to be a better predictor than $Trait_{FD}$.
33 Finally, when evaluating the individual contributions of litter nutrients, calcium and magnesium
34 appeared as important drivers of litter mixture effects. Our findings suggest that the mass-ratio
35 hypothesis overrules the niche complementarity hypothesis as a driver of litter diversity effects.
36 Our study highlights the key importance of macroinvertebrates and of leaf nutrients, such as Ca
37 and Mg, which are often neglected in decomposition studies in streams.

38

39 **KEYWORDS:** Biodiversity-ecosystem functioning; community weighted means; functional trait
40 dissimilarity; litter traits; litter decomposition; litter nutrients; temperate stream

41 **HIGHLIGHTS**

42 - Macroinvertebrates and leaf nutrients drive microbial activity and litter decomposition
43 rate

44 - Macroinvertebrate presence and nutrient concentration control litter mixing effect
45 intensity

46 - Mean nutrient concentration is more important than concentration dissimilarity within
47 litter mixture

48

49 INTRODUCTION

50 Litter decomposition is an essential ecosystem function controlling the carbon (C) and
51 nutrient cycles in both terrestrial (Swift and others 1979; Cadish and Giller 1997) and aquatic
52 (Cummins 1974; Wallace and others 1997) ecosystems. Litter decomposition rate is jointly
53 affected by the litter traits (e.g. Cornwell and others 2008; Garcia-Palacios and others 2016),
54 the environmental conditions, such as the temperature (e.g. Fierer and others 2005; Follstad
55 Shah and others 2017), and the decomposer communities (e.g. Hättenschwiler and others 2015;
56 Gessner and others 2010; Jabiol and others 2013; Santonja and others 2018a). Rates of litter
57 decomposition are essentially controlled by litter chemistry (Cornwell and others 2008; Garcia-
58 Palacios and others 2016). In addition to secondary compounds (e.g. tannins) and fiber
59 components (e.g. lignin), concentrations of nutrients such as nitrogen (N) and phosphorus (P)
60 are useful litter traits for the prediction of the decomposition rates of single-species litter
61 (Gessner and Chauvet 1994; Cornwell and others 2008; Schindler and Gessner 2009).
62 Nevertheless, as mainly reported in terrestrial ecosystems, other nutrients such as calcium (Ca),
63 magnesium (Mg), potassium (K), sodium (Na), and sulfur (S), can also affect the decomposition
64 efficiency (Kaspari and others 2008; Makkonen and others 2012; Garcia-Palacios and others
65 2016). However, the potential importance of these nutrients is often neglected in decomposition
66 studies in streams compared to N and P.

67 In the majority of natural ecosystems, litter material from different plant species
68 decomposes together. Given that forested stream food webs are dependent upon allochthonous
69 leaf litter (Vannote and others 1980; Wallace and others 1997; Gessner and others 1999),
70 understanding the relationship between riparian tree species composition (and its associated
71 litter traits) and the decomposition rates in streams draining forested watersheds is of
72 considerable ecological importance (Swan and Palmer 2004; Kominoski and others 2007).
73 Changing plant species composition can modify litter trait control over decomposition in two

74 ways. First, according to the “mass-ratio hypothesis” (Grime 1998; Garnier and others 2004),
75 litter trait control over decomposition changes along community-weighted mean (CWM) trait
76 values (Fig. 1a, Quested and others 2007; Mokany and others 2008; Laughlin 2011). Second,
77 according to the “niche complementarity hypothesis” (Petchey and Gaston 2006; Diaz and
78 others 2007), a change in the functional dissimilarity (FD) of litter trait-values affects the degree
79 to which resource-use complementary occurs within the decomposer community, and its
80 capacity to induce non-additive litter mixing effects on decomposition (i.e. litter mixtures that
81 decompose at different rates than predicted by the mass-ratio hypothesis) (Fig. 1b, Wardle and
82 others 1997; Vos and others 2013; Handa and others 2014). These two mechanisms can operate
83 simultaneously by affecting both decomposer community and decomposition efficiency as a
84 result of plant (litter) composition change (e.g. Garcia-Palacios and others 2017; Santonja and
85 others 2018b). However, our knowledge about the relative importance of these two mechanisms
86 in controlling the decomposer communities and the efficiency of the decomposition process in
87 streams is very limited.

88 Mechanisms involved in positive litter mixing effects likely include nutrient transfer,
89 such as N or P, from nutrient-rich to nutrient-poor litter species. For example, nutrients may be
90 transported through fungal hyphae connecting two different leaf litter species (Schimel and
91 Hättenschwiler 2007; Gessner and others 2010; Lummer and others 2012), in which case an
92 acceleration of decomposition in species mixtures is expected since the nutrients are used more
93 efficiently overall. According to the literature (Wardle and others 1997; Sanpera-Calbet et al.
94 and others 2009; Santonja and others 2015a, 2015b; Santschi and others 2018), positive mixing
95 effects are most likely expected in the mixtures including one poor-quality litter species and a
96 high-quality one (Fig 1b; but see Frainer and others (2015) for contrasting results). However,
97 whether, and to what extent, mixing litter affects decomposition rates in streams remains much
98 debated, since litter mixture effects appear to be idiosyncratic (Schindler and Gessner 2009;

99 [Gessner and others 2010](#); [Cardinale and others 2011](#); [Lecerf and others 2011](#)). Previous studies
100 highlighting contrasting results have focused on the effects of N and P in explaining such litter
101 mixture effects (e.g. [Frainer and others 2015](#); [Santschi and others 2018](#)). However, other key
102 nutrients, such as Mg or Ca, could potentially affect decomposition rates in litter mixtures.

103 Additionally, it is also not clear to what extent macroinvertebrates contribute to
104 decomposition in litter mixtures ([Hättenschwiler and Gasser 2005](#); [Swan and Palmer 2006b](#);
105 [Swan 2011](#)). Shredder macroinvertebrates can play a prominent role in the decomposition
106 process in stream ecosystems ([Anderson and Sedell 1979](#); [Cummins and Klug 1979](#); [Handa and
107 others 2014](#); [Garcia-Palacios and others 2016](#)), by consuming and fragmenting the litter
108 material ([Allan 1996](#); [Graça 2001](#)), by stimulating microbial decomposition ([Wetzel 1995](#);
109 [Villanueva and others 2012](#)), and by mediating plant diversity effects on litter decomposition
110 ([Lecerf and others 2005](#); [Kominoski and others 2007](#); [Sanpera-Calbet and others 2009](#)).
111 According to [Cummins and others \(1989\)](#), such a role played by macroinvertebrates would be
112 higher in litter mixtures of distinct degradability than in more homogeneous litter mixtures.
113 Indeed, this type of litter mixture could simultaneously create a suitable microhabitat (more
114 refractory litter) and a food resource (more labile litter). Additionally, litter mixtures could
115 promote an increase in both growth rate ([Swan and Palmer 2006a](#)) and macroinvertebrate
116 population due to their aggregative behavior ([Presa-Abós and others 2006](#)), enhancing therefore
117 the decomposition of litter mixtures ([Sanpera-Calbet and others 2009](#)).

118 In order to address these gaps, we used leaf litter from 10 common European tree species
119 to evaluate how litter diversity effects are affected by litter nutrients (Ca, K, Mg, N, Na, P, and
120 S) in a temperate stream. Moreover, in order to elucidate which organisms may be responsible
121 for mediating such litter diversity effects, we distinguished between decomposition involving
122 microorganisms alone (with decomposition in fine-mesh litterbags) and both microorganisms
123 and macroinvertebrates (with decomposition in coarse-mesh litterbags) ([Boulton and Boon](#)

124 1991). Five of the plant species showed nutrient-rich litter, while the other five showed nutrient-
125 poor litter (Fig. 2). Mixtures of two litter species were created in all possible pairwise
126 combinations, in order to distinguish between litter mixtures of similar nutrient content (i.e. in
127 mixing two nutrient-poor or two nutrient-rich litters) and litter mixtures of dissimilar nutrient
128 content (i.e. in mixing one nutrient-poor litter and one nutrient-rich litter). Despite the fact that
129 both litter traits, litter mass loss and microbial activity are continuous variables, such artificial
130 distinction into two discrete nutrient categories was necessary to better understand where and
131 why litter diversity effects occur. We evaluated (i) the litter decomposition rate and the
132 microbial activity (i.e. CO₂ production) in the two-species litter mixtures, and (ii) the litter
133 diversity effects occurring on the decomposition rate and on the microbial activity in these litter
134 mixtures after 30 days of immersion in a temperate forested stream in northwestern France
135 (Piscart and others 2009).

136 First, according to the “mass-ratio hypothesis” (Grime 1998; Garnier and others 2004),
137 the increase in both litter decomposition rate and microbial activity would be positively
138 correlated to the increase in CWM scores in nutrient concentration within the litter mixture
139 (Cornwell and others 2008; Foucreau and others 2013). Therefore, in our first hypothesis, we
140 predicted higher litter decomposition rate and microbial activity in the mixture of two nutrient-
141 rich litters (Fig. 1a). Second, according to the “niche complementarity hypothesis” (Petchey
142 and Gaston 2006; Diaz and others 2007), mixtures of very dissimilar litter species yield the
143 highest FD scores and are related to higher litter diversity effects (Wardle and others 1997;
144 Lecerf and others 2011; Santschi and others 2018). Thus, in our second hypothesis, we
145 predicted the highest litter diversity effects in the mixtures of nutrient-poor and nutrient-rich
146 litters (Fig. 1b), which would be shown by an increase in litter decomposition rate and microbial
147 activity due to synergistic effects between the two litters in mixtures in the present study.
148 Finally, macroinvertebrates may contribute up to 6 times more influence than microorganisms

149 on litter decomposition in temperate unaltered forested streams (Piscart and other 2009), favor
150 microbial community development (Wetzel 1995; Villanueva and others 2012), and mediate
151 litter diversity effects (Lecerf and others 2005), we hypothesized that litter decomposition rate,
152 microbial activity, and litter diversity effects would be enhanced by macroinvertebrate presence
153 (Figs. 1a and 1b).

154

155 MATERIALS AND METHODS

156

157 *Study site and material collection*

158 The experiment was conducted in the Hermitage stream, located in the Villecartier
159 Forest in northwestern France (48°28' N, 1°33' W). The stream bed was dominated by sand
160 and leaf litter (site H1 in Piscart and others 2009). The stream water was circumneutral, well-
161 oxygenated, and possessed moderate nutrient concentrations (Piscart and others 2009): 11 ± 1
162 mg l^{-1} dissolved oxygen, $49 \pm 18 \mu\text{g N l}^{-1}$ ammonium, $510 \pm 19 \mu\text{g N l}^{-1}$ nitrate, $10 \pm 8 \mu\text{g N l}^{-1}$
163 nitrite, $19 \pm 8 \mu\text{g P l}^{-1}$ phosphate. The macroinvertebrate community of the Hermitage stream
164 was composed of species belonging to the Amphipoda, Isopoda and Trichoptera orders, with
165 crustaceans representing 74% of this macroinvertebrate community (Supplementary Table S1).
166 *Fagus sylvatica* was the dominant tree species in the forested watershed, but the riparian
167 vegetation was composed of diverse deciduous tree species, including those employed in the
168 experiment.

169 The leaf litter of 10 common European tree species was collected: *Acer platanoides*,
170 *Alnus glutinosa*, *Betula pendula*, *Castanea sativa*, *Corylus avellana*, *Carpinus betulus*, *Fagus*
171 *sylvatica*, *Quercus robur*, *Salix atrocinerea* and *Tilia cordata*, hereafter referred to by their
172 genus name. These 10 species were selected using the existing literature (e.g. Lecerf and others
173 2007; Schindler and Gessner 2009; Santonja and others 2018a), based on the nutrient

174 concentrations of their leaves, to represent five species with nutrient-rich litter (NRL), and five
175 with nutrient-poor litter (NPL) (Fig. 2; a cluster analysis well discriminates these two groups).
176 Freshly abscised leaves were collected over the entire period of maximum litter fall, from
177 October to November 2015. They were dried at room temperature, and stored until the
178 beginning of the experiment.

179

180 *Litter decomposition experiment*

181 Leaf litter decomposition was studied for a period of 30 days, using the litterbag method
182 (Boulton and Boon 1991). Coarse- and fine-mesh litterbags (5 mm and 0.5 mm mesh size,
183 respectively) were used and filled with 2 g of dry leaves. The leaf litter enclosed in fine-mesh
184 litterbags was only accessible to microorganisms, whereas the coarse-mesh bags also allowed
185 access to macroinvertebrates. The litterbags contained either a single species (10 treatments) or
186 a mixture of two species in all possible pairwise combinations (45 treatments). The mixed-
187 species litterbags received equal amounts of both species (i.e. 1 g). A total of 440 litterbags ([10
188 single-species + 45 two-species mixtures] × 2 mesh sizes × 4 replicates) were used for the
189 experiment.

190 In December 2016, the litterbags were immersed for 30 days in the Hermitage stream.
191 After removal, the litterbags were immediately sealed in plastic bags to prevent the loss of litter
192 material and were transported to the laboratory. The leaves were separated by species, which
193 was possible even with small fragments of litter (owing to marked morphological differences
194 among the species) and were carefully cleaned under water to remove macroinvertebrates and
195 attached sediment particles.

196 Avoiding the central veins, six leaf discs (10 mm-diameter) for the single-species litters,
197 or three disks for each species in the two-species litter mixtures, were cut for the purpose of
198 microbial activity measurement. The remaining leaf material was frozen at -20 °C.

199

200 *Microbial activity measurement*

201 Microbial CO₂ production measurements, as a measure of overall heterotrophic
202 microbial activity, were performed in order to evaluate the activity of the microbial
203 communities that colonized the leaves after 30 days of decomposition. We adapted the protocol
204 proposed by [Anderson and Domsch \(1973\)](#). Briefly, the six leaf disks taken in each litterbag
205 were placed in 125 ml glass bottles with 25 ml of filtered water (GF/F glass microfiber filter,
206 WhatmanTM) from the Hermitage stream and then preincubated for 12 h at 20 °C in the dark
207 allowing the microbial respiration to saturate the water with CO₂. In the matter of fact, at
208 constant temperature, it is only possible to increase the CO₂ concentration in the air phase of
209 the respiration chamber if the aqueous phase is oversaturated in CO₂. After a night of remaining
210 open to the atmosphere, the respiration chambers were hermetically sealed and the first sample
211 of air was taken, the second air sample was sampled after a 4 h incubation at 20 °C in the dark.
212 CO₂ production in a given time (i.e. 4 h) was then calculated as the difference between the final
213 CO₂ concentration and the initial CO₂ concentration. Preliminary studies with our experimental
214 conditions demonstrated that the CO₂ production was linear and, since all the process of
215 incubation occurred in the dark, there was no photosynthesis and thus no alteration of the
216 dissolved CO₂ partial pressure. In total, 440 respiration chambers, corresponding to the 440
217 litterbags, were prepared. At the beginning and at the end of the incubation period, 1 ml of air,
218 taken with a syringe, was injected into a gas chromatograph (µGC SRA A 3000) in order to
219 estimate the microbial activity (i.e. CO₂ air content). The leaf discs were then dried at 65 °C for
220 72 h and the CO₂ production was calculated as µg C-CO₂ per h and per g of litter dry mass (±
221 0.1 mg).

222 In parallel to the measurement of the microbial activity, the leaves remaining in the 440
223 litterbags were dried at 65 °C for 72 h, and then weighed to the nearest 0.1 mg. After weighing

224 the leaf disks and remaining leaf litter, all the litter material from a given litterbag was combined
225 and ground to a fine powder using a ball mill, before measuring the litter ash content. We
226 obtained ash-free dry mass by burning the combined sample at 550 °C for 4 h. Additional
227 samples were also used to estimate ash-free dry mass of the initial litter material. The percentage
228 of ash-free dry mass data was used to correct both the initial and the remaining leaf material
229 before the computation of (i) the percentage of litter mass loss after 30 days of field
230 decomposition and (ii) the microbial CO₂ production per h and per g of litter.

231

232 *Litter trait measurement*

233 The initial litter traits were determined from four samples of each of the 10 litter species
234 ([Supplementary Table S2](#)). Prior to the chemical analysis, each litter sample was ground into
235 powder using a ball mill. The carbon (C), nitrogen (N), and sulfur (S) concentrations were
236 determined by thermal combustion, using a Vario Pyro cube CNS analyzer (Elementar France
237 SARL, Lyon, France). The phosphorus (P) concentration was measured colorimetrically using
238 the molybdenum blue method ([Grimshaw and others 1989](#)). To 80 mg of ground litter sample
239 8 ml of HNO₃ and 2 ml of H₂O₂ were added and the mixture was heated at 175 °C for 40 min
240 using microwaves (Ethos One, Milestone SRL, Italy). After this mineralization step, the sample
241 was diluted to a total of 50 ml. A hundred µl of sample, 100 µl of NaOH, 50 µl of mixed reagent
242 (emetic tartar and ammonium molybdate solution), and 50 µl of ascorbic acid were mixed
243 directly in a 96 well microplate. After 30 min at 40 °C, the reaction was completed, and the P
244 concentration was measured at 720 nm using a microplate reader (Victor, PerkinElmer,
245 Singapore). Following the mineralization step (i.e. the same as for P analysis), calcium (Ca),
246 magnesium (Mg), potassium (K), and sodium (Na) concentrations were measured using an
247 atomic absorption spectrometer (AAS, iCE 3000 series, ThermoScientific, China).

248 To assess the “mass-ratio hypothesis”, we calculated the community-weighted mean
249 (CWM) trait values of litter mixtures as the average trait values of litter mixtures following
250 [Garnier and others \(2004\)](#) as: $Trait_{CWM} = \sum_{i=1}^n p_i \times trait_i$, where p_i is the relative abundance
251 for species i in the litter mixture and $trait_i$ is the trait value for species i . These calculations were
252 performed for each of the 8 litter traits. The highest scores of CWM were reached for litter
253 mixtures containing two litter species with the highest nutrient concentrations ([Fig. 1a](#)). Since
254 we predicted that both microbial activity and litter decomposition rate respond to the “mass-
255 ratio hypothesis”, we expected the increase in both decomposition rate and microbial activity
256 to be positively correlated to the increase in CWM values.

257 To assess the “niche complementarity hypothesis”, we calculated the functional
258 dissimilarity (FD) of litter mixtures according to Rao’s quadratic entropy ([Botta Dukat 2005](#);
259 [Epps and others 2007](#)) as: $Trait_{FD} = \sum_{i=1}^n \sum_{j=1}^n p_i p_j * d_{ij}$, where p_i and p_j are the relative
260 abundance for species i and j in the litter mixture, and d_{ij} the Euclidian distance between species
261 i and j for the trait considered. These calculations were performed for each of the 8 litter traits.
262 The highest scores of FD were reached for litter mixtures containing two species with very
263 dissimilar nutrient concentrations ([Fig. 1b](#)). Since we predicted that the litter diversity effects
264 respond to the “niche complementarity hypothesis”, we expected the increase in relative litter
265 mixture effects to be positively correlated to the increase in FD values.

266

267 ***Statistical analyses***

268 All of the statistical analyses were conducted using R software ([R Core Team 2013](#)),
269 with significance levels indicated as * for $P < 0.05$, ** for $P < 0.01$, and *** for $P < 0.001$.

270 A principal component analysis (PCA) was conducted using the values of the eight
271 measured litter traits of the 10 tree species ([Supplementary Table S2](#)) in order to discriminate
272 the five tree species with nutrient-rich litter (NRL) from the other five tree species with nutrient-

273 poor litter (NPL). The differences in the initial litter traits were assessed using one-way
274 ANOVAs, followed by Tukey's tests to carry out post-hoc pairwise comparisons.

275 For the single-species litter, three-way ANOVAs, followed by Tukey's post hoc tests,
276 were used to test for the effects of litter type (separated in NPL vs. NRL), litter species identity
277 (10 litters), macroinvertebrate presence (fine-mesh bag [FMB] vs. coarse-mesh bag [CMB]),
278 and their interactions on litter decomposition rate and microbial activity.

279 In order to further test whether litter decomposition and microbial activity differed
280 between litter mixtures and single litter species, the relative litter mixture effects (RME) on
281 litter decomposition rate and microbial activity were calculated. The RME was calculated as
282 the relative difference between the observed litter decomposition rate/microbial activity (O)
283 from the litter mixtures compared to those expected based on the respective single litter species
284 treatments (E), following the formula $(O - E) / E \times 100\%$ (Wardle and others 1997). One-
285 sample Student's *t*-tests were used to test whether the RME were significantly different from
286 zero.

287 For the two-species litter mixtures, two-way ANOVAs, followed by Tukey's post hoc
288 tests, were used to test for the effects of litter mixing (NPL-NPL, NPL-NRL and NRL-NRL),
289 macroinvertebrate presence (FMB vs. CMB), and their interactions i) on litter decomposition
290 rate and microbial activity and ii) on the RME on litter decomposition rate and on microbial
291 activity.

292 For a more detailed understanding of how the mixture of leaf litter affected the litter
293 decomposition rate and microbial activity, we evaluated the effects of the mean litter traits
294 ($Trait_{CWM}$) and functional litter trait dissimilarities ($Trait_{FD}$) of the eight measured litter traits
295 (Supplementary Table S2). First, a principal component analysis (PCA) was conducted using
296 the CWM or the FD values of the eight measured litter traits across the 45 two-species litter
297 mixtures. CWM1 and CWM2, and FD1 and FD2 represented the two first components of the

298 PCAs conducted using the CWM or the FD values across the litter mixtures, respectively.
299 Second, multiple linear regression models were performed in order to decipher the relative
300 contributions of Trait_{CWM} and Trait_{FD}. In these models the effects of the Traits_{CWM} (i.e. CWM1
301 and CWM2), Traits_{FD} (i.e. FD1 and FD2), macroinvertebrate presence (FMB vs. CMB), and
302 their interactions were tested i) on litter decomposition rate and microbial activity and ii) on the
303 RME on litter decomposition rate and on microbial activity.

304

305 **RESULTS**

306

307 *Litter traits*

308 The PCA based on the element concentrations showed that the first PCA axis explained
309 50.2% of the variation and discriminated between the nutrient-rich litters (NRL) on the left and
310 the nutrient-poor litters (NPL) on the right of the axis (Fig. 2).

311

312 *Litter species incubated individually*

313 The NRL exhibited two times more litter mass loss and microbial activity than the NPL
314 (Table 1; Fig. 3). The presence of macroinvertebrates increased the litter mass loss and the
315 microbial activity, but this effect was dependent on the litter type (litter type × mesh size
316 interaction, Table 1). This significant interaction was explained by a stronger increase in litter
317 mass loss and microbial activity in the NRL category (+59% litter mass loss and +33%
318 microbial activity) compared to the NPL category (+23% litter mass loss and +28% microbial
319 activity) (Fig. 3).

320 In addition, litter mass loss and microbial activity were significantly affected by litter
321 species identity (Table 1). Within the NRL category, *Alnus* showed the highest litter mass loss
322 compared to the four other species (Supplementary Fig. S1a), as well as a higher microbial

323 activity than *Carpinus*, *Acer*, and *Tilia* (Supplementary Fig. S1b). Within the NPL category,
324 *Salix* and *Betula* showed both a higher litter mass loss and a higher microbial activity than the
325 three other species (Supplementary Fig. S1a and S1b). The positive effect of the
326 macroinvertebrate presence on the litter mass loss also varied according to the litter species
327 identity (litter species \times mesh size interaction, Table 1), as the effects ranged from +68% for
328 *Alnus* to an absence of effect for *Fagus* (Supplementary Fig. S1a).

329

330 ***Effects of mixing low- and high-quality litter***

331 Litter mass loss and microbial activity increased according to the gradient NPL-NPL <
332 NPL-NRL < NRL-NRL mixtures (Table 2; Fig. 4a and 4c), with two times more litter mass
333 loss and microbial activity in mixtures of two nutrient-rich litters compared to the mixtures of
334 two nutrient-poor litters. The relative mixture effects (RME) on litter mass loss and on
335 microbial activity showed a similar trend, with an increase in RME according to the same
336 gradient (Table 2; Fig. 5). The NPL-NPL mixtures lost -3.5% litter mass compared with the
337 expected values from the respective single litter species (Fig. 5a), while the observed microbial
338 activity of the NPL-NPL mixtures did not differ significantly from the expected values (Fig.
339 5c). The NPL-NRL mixtures lost +5.3% litter mass than expected from the respective single
340 litter species (Fig. 5a), while the observed microbial activity of the NPL-NRL mixtures did not
341 differ from the expected values (Fig. 5c). The NRL-NRL mixtures showed +8.9% litter mass
342 and +16.3% microbial activity than expected from the respective single litter species (Figs. 5a
343 and 5c).

344 Litter mass loss and microbial activity were respectively 41% and 30% higher with
345 macroinvertebrates than without (Table 2; Fig. 4b and 4d). The RME on litter mass loss and on
346 microbial activity were also higher with macroinvertebrates than without (Table 2).
347 Specifically, the litter mixtures exhibited synergistic effects in the presence of

348 macroinvertebrates, with +7.3% litter mass loss and +9.9% microbial activity than expected
349 from the respective single litter species (Figs. 5b and 5d).

350

351 *CWM- versus FD-trait control over litter decomposition rate and microbial activity*

352 The PCA of the CWM traits showed that the first PCA axis (CWM1), explaining 50.2%
353 of the variation, was determined by high scores of P and Ca concentrations, and, to a lower
354 extent, by the scores of the K, Mg, Na, and S concentrations (Supplementary Fig. S2a). The
355 low scores of the second PCA axis (CWM2), explaining 21.9% of the variation, were related
356 to high values of N and C concentrations (Supplementary Fig. S2a). Regarding functional trait
357 dissimilarity, high scores of the first PCA axis (FD1), explaining 28.5% of the variation, were
358 related to the increase in dissimilarity in the Mg, S, and C concentrations, while the low scores
359 were related to the increase in dissimilarity in the N and Ca concentrations (Supplementary Fig.
360 S2b). The low scores along the second axis (FD2), explaining 22.0% of the variation, were
361 largely dependent on the increase in dissimilarity in the K, Na, and P concentrations
362 (Supplementary Fig. S2b).

363 When simultaneously evaluating the effects of the CWM and FD traits (Table 3), we
364 found that both litter mass loss and microbial activity were mainly controlled by the CWM
365 values (i.e. increasing mean nutrient concentrations within the litter mixture) compared to the
366 FD values (i.e. increasing dissimilarity in nutrient concentrations within the litter mixture). In
367 fact, CWM values (CWM1 + CWM2) explained respectively 9 and 15 times more of the overall
368 variance in litter mass loss and in microbial activity than FD values (FD1 + FD2) (Table 3).

369 When simultaneously evaluating the effects of the CWM and FD traits on RME, we
370 found that the RME on litter mass loss was controlled by the CWM (CWM1), the presence of
371 macroinvertebrates, and the interaction between the FD and the presence of macroinvertebrates
372 (FD2 × mesh size) (Table 3). Increasing CWM1 scores was related to higher RME (Fig. 6a).

373 The significant interaction between FD2 and mesh size (Table 3) showed that increasing FD in
374 the initial K, Na, and P concentrations stimulated microbial-driven RME (Fig. 6d). Concerning
375 the RME on microbial activity, it was found to be significantly affected only by CWM1 and the
376 presence of macroinvertebrates (Table 3). Similar to what we observed for litter mass loss,
377 increasing CWM1 scores was related to higher RME on microbial activity (Fig. 6e).

378 The RME on litter mass loss appeared to be more strongly correlated with the increase
379 in P, Ca, Mg and Na concentrations than with the K, N, or S concentrations, and these
380 relationships were more marked in the coarse-mesh litterbags in which macroinvertebrates were
381 present, than in the fine-mesh litterbags in which macroinvertebrates were absent (Table 4).
382 The RME on microbial activity appeared to be more strongly correlated with the increase in
383 Ca, K, and Mg concentrations than in N and P concentrations, and these relationships were also
384 more marked in the presence than in the absence of macroinvertebrates (Table 4).

385

386 **DISCUSSION**

387

388 *Niche complementarity hypothesis vs. mass-ratio hypothesis for explaining litter diversity* 389 *effects*

390 We found evidence of leaf litter diversity effects on decomposition rates, including
391 additive, synergistic, and antagonistic effects, as previously shown in other studies (Kominoski
392 and others 2007; Srivastava and others 2009; Lecerf and others 2011; Handa and others 2014).
393 Surprisingly, the pattern observed in our study highlighted that the largest mixture effects
394 occurred in mixtures containing the combination of two nutrient-rich litters (NRLs), rather than
395 in mixtures of one nutrient-rich litter (NRL) and one nutrient-poor litter (NPL), as predicted in
396 our second hypothesis (Fig. 1b). Such a finding is in agreement with the mass-ratio hypothesis,
397 and emphasizes the fact that increasing the mean value of the nutrient pool in litter mixtures

398 favors the occurrence of litter diversity effects. In any case our results pointed out that the litter
399 diversity effects on decomposition were strongly litter-quality dependent. This finding is
400 congruent with the recent study of [Jabiol and Chauvet \(2015\)](#) in which higher synergistic effects
401 on litter decomposition occurred when *Alnus glutinosa* was combined with *Fraxinus*
402 *angustifolia* (i.e. two NRLs) than when *Alnus glutinosa* was combined with *Quercus ilex* (i.e.
403 one NRL and one NPL).

404 Surprisingly, $Trait_{CWM}$ proved to be a better predictor of litter diversity effects than
405 $Trait_{FD}$, for both the coarse- and fine-mesh litterbags, contrasting with our second hypothesis.
406 Indeed, based on the well-developed literature on the drivers of the litter decomposition process
407 (e.g. [Hättenschwiler and others 2005](#); [Cornwell and others 2008](#); [Gessner and others 2010](#);
408 [Makkonen and others 2012](#); [Vos and others 2013](#); [Handa and others 2014](#); [Garcia-Palacios and](#)
409 [others 2017](#)), we expected that litter diversity effects occur when there are increases in the
410 nutrient concentration dissimilarity of the litter mixtures (i.e. according to the niche
411 complementary hypothesis, [Fig. 1b](#)), while the mean nutrient concentration within the litter
412 mixture must only explain the litter decomposition rate (i.e. according to the mass-ratio
413 hypothesis, [Fig. 1a](#)). The results from the coarse-mesh litterbags showed no evidence of nutrient
414 dissimilarity effect ($Trait_{FD}$), conforming with the conclusion of [Frainer and others \(2015\)](#),
415 which also highlighted, over a lower range of litter traits (N, P, and lignin), that no litter
416 dissimilarity effect was involved in the litter diversity effects occurring in two-species mixtures.
417 In contrast, $Trait_{FD}$ was linked to the litter diversity effects for the fine-mesh litterbags. In this
418 case, litter mixtures with contrasting nutrient concentrations may have improved the availability
419 of different nutrient sources for microorganisms ([Schimel and Hättenschwiler 2007](#); [Handa and](#)
420 [others 2014](#)), as suggested by the niche complementarity hypothesis. Filamentous fungi,
421 including the aquatic hyphomycetes that dominate fungal communities on decomposing leaves
422 in streams ([Krauss and others 2011](#)), can indeed extend their hyphae over considerable distances

423 in order to acquire remote resources that they transport to the locations of active hyphal growth
424 (Ritz 2006). It has previously been suggested that fungi-driven N transfer among litter species
425 varying in their initial N concentration may contribute to litter mixture effects (Schimel and
426 Hättenschwiler 2007; Vos and others 2013; Handa and others 2014). Thus, nutrients from a
427 nutrient-rich litter may be translocated to another litter depleted in nutrient (i.e. a nutrient-poor
428 litter; Schimel and Hättenschwiler 2007; Handa and others 2014). Therefore, our findings
429 highlighted that both Trait_{CWM} and Trait_{FD} contributed to explain litter diversity effects driven
430 by microorganisms alone. Interestingly, Trait_{FD} did not explain litter diversity effects driven by
431 microorganisms and macroinvertebrates together. In this case, it could be hypothesized that the
432 additional nutrients provided by macroinvertebrate feces alleviated the nutrient limitation for
433 microorganisms (Wetzel 1995; Joyce and Wotton 2008), and consequently the Trait_{FD}
434 contribution to litter diversity effects.

435

436 *Macroinvertebrates control over decomposition and litter diversity effects*

437 Across all of the 45 litter mixtures, litter mixture decomposition was not found to be
438 enhanced in the fine-mesh litterbags (i.e. additive effects), while the litter mixtures in the
439 coarse-mesh litterbags lost on average 7.3% more mass than the litter in the single-species
440 treatments (i.e. synergistic effects). The absence of litter mixture effects when
441 macroinvertebrates were excluded suggests that the activity of microbial decomposers alone
442 did not induce any effect of litter mixture on decomposition in the studied temperate stream.
443 Previous studies performed in lotic systems, where decomposition primarily involves
444 microorganisms, have also reported a lack of a synergism (Ferreira and others 2012; Bruder
445 and others 2014). For instance, similar to our findings, Jabiol and Chauvet (2015) reported that
446 no effect of litter mixture was observed when detritivores were excluded during the litter
447 decomposition in a Mediterranean stream in southern France. In a laboratory experiment, Swan

448 [and Palmer \(2006b\)](#) also reported that litter mixture effects were contingent on the feeding
449 activity of the isopod *Caecidotea communis*.

450 In homogenous species mixtures comprising litters of a single litter category,
451 macroinvertebrates have little choice to select among different qualities of leaves, as opposed
452 to the choice offered between leaf litter species in heterogeneous litter mixtures ([Swan and
453 Palmer 2006b](#)). However, as macroinvertebrates can simultaneously exploit multiple litter
454 species to meet their elemental demands ([Leroy and Marks 2006](#)), combining two NRLs that
455 nutritionally complement one another might stimulate the feeding activity of detritivores ([Vos
456 and others 2013](#)). For example, the mixing of one P-rich with one N-rich species in the present
457 study (such as *Acer platanoides* and *Alnus glutinosa*, respectively) increased litter mass loss of
458 the mixture by 14%, probably due to the increased nutritional value of such litter mixing. In
459 contrast, mixing two P-rich species (e.g. *Acer platanoides* and *Carpinus betulus*) increased litter
460 mass loss of the mixture by only 4%, while mixing two P-poor species (e.g. *Castanea sativa*
461 and *Fagus sylvatica*) decreased litter mass loss of the mixture by 10%. Alternatively in
462 NRL+NPL mixtures, high detritivore density may be promoted by the co-occurrence of a
463 suitable resource, such as that provided by the nutrient-rich litter, and a complex and structured
464 habitat provided by the nutrient-poor litter ([Sanpera-Calbet and others 2009](#); [Jabiol and others
465 2014](#)).

466

467 ***Litter nutrients control over decomposition and litter diversity effects***

468 In agreement with our predictions and the literature, nutrient-rich litters decomposed
469 faster than nutrient-poor litters, confirming that nutrient concentration is an important
470 determinant of leaf litter decomposition ([Kaspari and others 2008](#); [Makkonen and others 2012](#);
471 [Garcia-Palacios and others 2016](#)). The rates of litter decomposition observed in the present
472 study were similar to those reported in other studies concerning temperate streams (e.g. [Lecerf](#)

473 and others 2007; Schindler and Gessner 2009; Frainer and others 2015). For instance, Lecerf
474 and others (2007) reported 80% and 42% mass loss for *Alnus glutinosa* (i.e. a nutrient-rich
475 litter) and *Quercus robur* (i.e. a nutrient-poor litter), respectively, after 34 days of
476 decomposition in a temperate stream in central France. The results of the present study were
477 also similar to the 16.5% mass loss for *Fagus sylvatica* (i.e. a nutrient-poor litter) after 30 days
478 of decomposition obtained in the same Petit Hermitage stream in February 2005 (Piscart and
479 others 2009).

480 Moreover, our results also highlighted the important role of nutrients not usually
481 considered in litter diversity experiments. The first principal component of the CWM-trait PCA
482 was the main driver of litter diversity effects in both fine- and coarse-mesh litterbags, and was
483 driven by P, Ca, K, Mg, Na, and S. Their relative content in the leaves is closely related to one
484 other (Garcia-Palacios and others 2016), hampering a straightforward interpretation of
485 combination of elements. Nevertheless, when evaluating the individual contributions of litter
486 nutrients, Ca and Mg appeared to be important drivers of litter diversity effects compared to N
487 (and to P only for microbial activity), suggesting for the first time that these two nutrients may
488 play an important role in litter diversity effects in addition to, or independently of, the P or N
489 content. The Ca content is known to positively affect the growth and activity of aquatic
490 hyphomycetes, by enhancing the fungal capacity to transfer N between distinct litter types
491 (Jenkins and Suberkropp 1995). Meanwhile, Ca and Mg are known to be essential elements in
492 the diet of macroinvertebrates, since they are required for enzymatic reactions, nerve
493 connections, muscle function, and skeleton formation (National Research Council 2005).
494 Moreover, in some temperate streams, crustaceans could represent up to 95% of shredder
495 biomass (Piscart and others 2011), and the Ca content of leaves could be a significant source of
496 Ca for their cuticles (Cairns and Yan 2009), especially in streams with a low Ca content in the
497 water (Glazier 1998), such as in Brittany, where the present study was conducted.

498

499 **CONCLUSION**

500 As expected, increasing mean nutrient concentration and macroinvertebrate presence
501 favored both microbial activity and litter decomposition rate. In addition, we experimentally
502 demonstrated for the first time that the mass-ratio hypothesis (i.e. Trait_{CWM}) overrules the niche
503 complementarity hypothesis (Trait_{FD}) as a driver of litter diversity effects in a temperate stream.
504 In fact, in strong contrast to our expectations, the combination of two nutrient-rich litters yielded
505 the highest litter mixture effects. In addition, the synergistic effects of litter mixing were mainly
506 evident in the presence of macroinvertebrates. Both community weighted mean traits
507 (Trait_{CWM}) and the functional dissimilarity of litter traits (Trait_{FD}) contributed to explain the
508 litter mixture effects. There was no support for Trait_{FD} explaining litter diversity effects in the
509 presence of macroinvertebrates, while both Trait_{CWM} and Trait_{FD} contributed to explain litter
510 mixture effects driven by microorganisms alone. Finally, Ca and Mg, which are often neglected
511 in decomposition studies in streams, were found to be important drivers of litter mixture effects.

512

513 **ACKNOWLEDGMENTS**

514 The chemical analyses were performed at the Plateforme d'Analyses Chimiques en
515 Ecologie (PACE, LabEx Centre Méditerranéen de l'Environnement et de la Biodiversité,
516 Montpellier, France), as well as at the PLateforme AnalYtique (PLAY, UMR CNRS 6553
517 ECOBIO, Rennes, France). We thank Raphaëlle Leclerc, Bruno Buatois and Nicolas Barthes
518 for technical assistance during the chemical analyses as well as Pierre Mariotte for his reviewing
519 of the English. This study was funded by the Selune project (project number 1 053 864) and by
520 the Rennes Metropole's project AIS 2015. The study was also supported by the LTSER France
521 Zone Atelier Armorique. Finally, the authors declare no conflict of interest.

522

523 **AUTHORS' CONTRIBUTIONS**

524 M.S., H.R.P., N.L.B. and C.P. conceived and performed the experiments. M.S. analyzed
525 the data and led the writing of the manuscript. All authors contributed critically to the drafts
526 and gave final approval for publication.

527

528 **REFERENCES**

529 Allan DJ. 1996. Stream Ecology. Chapman and Hall, New York.

530 Anderson, JPE, Domsch KH. 1973. Quantification of bacterial and fungal contributions to soil
531 respiration. Archiv für Mikrobiologie 93:113-127. Doi: 10.1007/BF00424942

532 Anderson NH, Sedell JR. 1979. Detritus processing by macroinvertebrates in stream
533 ecosystems. Annual Review of Entomology 24:351-377. Doi:
534 10.1146/annurev.en.24.010179.002031

535 Botta Dukát Z. 2005. Rao's quadratic entropy as a measure of functional diversity based on
536 multiple traits. Journal of Vegetation Science 16:533-540. Doi: 10.1658/1100-
537 9233(2005)16[533:RQEAAM]2.0.CO;2

538 Boulton AJ, Boon PI. 1991. A review of methodology used to measure leaf litter decomposition
539 in lotic environments: time to turn over an old leaf? Australian Journal of Marine Freshwater
540 Resources 42:1-43. Doi: 10.1071/MF9910001

541 Bruder A, Schindler MH, Moretti MS, Gessner MO. 2014. Litter decomposition in a temperate
542 and a tropical stream: the effects of species mixing, litter quality and shredders. Freshwater
543 Biology 59:438-449. <https://doi.org/10.1111/fwb.12276>

544 Cadish G, Giller KE. 1997. Driven by Nature: Plant Litter Quality and Decomposition. CAB
545 international, Wallingford.

546 Cairns A, Yan N. 2009. A review of the influence of low ambient calcium concentrations on
547 freshwater daphniids, gammarids, and crayfish. *Environmental Reviews* 17:67-79. Doi:
548 10.1139/A09-005

549 Cardinale BJ, Matulich KL, Hooper DU, Byrnes JE, Duffy E, Gamfeldt L, ... Gonzalez A.
550 2011. The functional role of producer diversity in ecosystems. *American Journal of Botany*
551 98:572-592. Doi: 10.3732/ajb.1000364

552 Cornwell WK, Cornelissen JH, Amatangelo K, Dorrepaal E, Eviner VT, Godoy O, ... Westoby
553 M. 2008. Plant species traits are the predominant control on litter decomposition rates within
554 biomes worldwide. *Ecology letters* 11:1065-1071. Doi: 10.1111/j.1461-0248.2008.01219.x

555 Cummins KW, Klug MJ. 1979. Feeding ecology of stream invertebrates. *Annual review of*
556 *ecology and systematics* 10:147-172. Doi: 10.1146/annurev.es.10.110179.001051

557 Cummins KW, Wilzbach MA, Gates DM, Perry JB, Taliferro WB. 1989. Shredders and riparian
558 vegetation: Leaf litter that falls into streams influences communities of stream invertebrates.
559 *BioSciences* 39:24-30 Doi: 10.2307/1310804

560 Cummins KW. 2012. Structure and function of stream ecosystems. *Bioscience* 24:631-641.
561 <https://doi.org/10.2307/1296676>

562 Diaz S, Lavorel S, De Bello F, Quetier F, Grigulis K, Robson TM. 2007. Incorporating plant
563 functional diversity effects in ecosystem service assessments. *Proceedings of the National*
564 *Academy of Sciences of the USA* 104:20684-20689. Doi: 10.1073/pnas.0704716104

565 Epps KY, Comerford NB, Reeves III JB, Cropper Jr WP, Araujo QR. 2007. Chemical diversity-
566 highlighting a species richness and ecosystem function disconnect. *Oikos* 116:1831-1840.
567 Doi: 10.1111/j.0030-1299.2007.15853.x

568 Ferreira V, Encalada AC, Graça MA. 2012. Effects of litter diversity on decomposition and
569 biological colonization of submerged litter in temperate and tropical streams. *Freshwater*
570 *Science* 31:945-962. Doi: 10.1899/11-062.1

571 Fierer N, Craine JM, McLauchlan K, Schimel JP. 2005. Litter quality and the temperature
572 sensitivity of decomposition. *Ecology* 86:320-326. Doi: 10.1890/04-1254

573 Follstad Shah JJ, Kominoski JS, Ardón M, Dodds WK, Gessner MO, Griffiths NA, ... Manning
574 DW. 2017. Global synthesis of the temperature sensitivity of leaf litter breakdown in streams
575 and rivers. *Global Change Biology* 23:3064-3075. Doi: 10.1111/gcb.13609

576 Foucreau N, Puijalón S, Hervant F, Piscart C. 2013. Effect of leaf litter characteristics on leaf
577 conditioning and on consumption by *Gammarus pulex*. *Freshwater Biology* 58:1672-1681.
578 Doi: 10.1111/fwb.12158

579 Frainer A, Moretti MS, Xu W, Gessner MO. 2015. No evidence for leaf-trait dissimilarity
580 effects on litter decomposition, fungal decomposers, and nutrient dynamics. *Ecology*
581 96:550-561. Doi: 10.1890/14-1151.1

582 Garcia-Palacios P, McKie BG, Handa IT, Frainer A, Hättenschwiler S. 2016. The importance
583 of litter traits and decomposers for litter decomposition: a comparison of aquatic and
584 terrestrial ecosystems within and across biomes. *Functional Ecology* 30:819-829. Doi:
585 10.1111/1365-2435.12589

586 Garcia-Palacios P, Shaw EA, Wall DH, Hättenschwiler S. 2017. Contrasting mass-ratio vs.
587 niche complementarity effects on litter C and N loss during decomposition along a regional
588 climatic gradient. *Journal of Ecology* 105:968-978. Doi: 10.1111/j.1466-8238.2011.00651.x

589 Garnier E, Cortez J, Billès G, Navas ML, Roumet C, Debussche M, ... Toussaint JP. 2004.
590 Plant functional markers capture ecosystem properties during secondary
591 succession. *Ecology* 85:2630-2637. Doi: 10.1890/03-0799

592 Gessner MO, Chauvet E. 1994. Importance of stream microfungi in controlling breakdown rates
593 of leaf litter. *Ecology* 75:1807-1817. Doi: 10.2307/1939639

594 Gessner MO, Chauvet E, Dobson M. 1999. A perspective on leaf litter breakdown in streams.
595 *Oikos* 85:377-384. Doi: 10.2307/3546505

596 Gessner MO, Swan CM, Dang CK, Mckie BG, Bardgett RD, Wall DH, Hättenschwiler S. 2010.
597 Diversity meets decomposition. *Trends in Ecology and Evolution* 25:372-380. Doi:
598 10.1016/j.tree.2010.01.010

599 Glazier DS. 1998. Springs as model systems for ecology and evolutionary biology: a case study
600 of *Gammarus minus* Say (Amphipoda) in mid-Appalachian springs differing in pH and ionic
601 content. In: Botosaneanu L (eds) *Studies in crenobiology: the biology of springs and*
602 *springbrooks*, Backhuys Publishers, Leiden, pp 41-54.

603 Graça MAS. 2001. The role of invertebrates on leaf litter decomposition in streams-a review.
604 *International Review of Hydrobiology* 86:383-393. Doi: 10.1002/1522-
605 2632(200107)86:4/5<383::AID-IROH383>3.0.CO;2-D

606 Grime JP. 1998. Benefits of plant diversity to ecosystems: immediate, filter and founder effects.
607 *Journal of Ecology* 86:902-910. Doi: 10.1046/j.1365-2745.1998.00306.x

608 Grimshaw HM, Allen SE, Parkinson JA. 1989. Nutrient elements. In: Allen SE (eds) *Chemical*
609 *analysis of ecological materials*, 2nd Edition, Blackwell Scientific, Oxford, pp 81-159.

610 Handa IT, Aerts R, Berendse F, Berg MP, Bruder A, Butenschoen O, ... Hättenschwiler S.
611 2014. Consequences of biodiversity loss for litter decomposition across
612 biomes. *Nature* 509:218-221. Doi: 10.1038/nature13247

613 Hättenschwiler S, Gasser P. 2005. Soil animals alter plant litter diversity effects on
614 decomposition. *Proceedings of the National Academy of Sciences*, 102:1519-1524.
615 <https://doi.org/10.1073/pnas.0404977102>

616 Hättenschwiler S, Tiunov AV, Scheu S. 2005. Biodiversity and litter decomposition in
617 terrestrial ecosystems. *Annual Review of Ecology, Evolution and Systematics* 36:191-218.
618 Doi: 10.1146/annurev.ecolsys.36.112904.151932

619 Jabiol J, McKie BG, Bruder A, Bernadet C, Gessner MO, Chauvet E. 2013. Trophic complexity
620 enhances ecosystem functioning in an aquatic detritus-based model system. *Journal of*
621 *Animal Ecology* 82:1042-1051. Doi: 10.1111/1365-2656.12079

622 Jabiol J, Cornut J, Danger M, Jouffroy M, Elger A, Chauvet E. 2014. Litter identity mediates
623 predator impacts on the functioning of an aquatic detritus-based food web. *Oecologia*,
624 176:225-235. Doi: 10.1007/s00442-014-2990-y

625 Jabiol J, Chauvet E. 2015. Biodiversity and litter decomposition: a case study in a
626 Mediterranean stream. *Freshwater Science* 34:423-430. Doi: 10.1086/680094

627 Jenkins CC, Suberkropp K. 1995. The influence of water chemistry on the enzymatic
628 degradation of leaves in streams. *Freshwater Biology* 33:245-253. Doi: 10.1111/j.1365-
629 2427.1995.tb01165.x

630 Joyce P, Wotton RS. 2008. Shredder fecal pellets as stores of allochthonous organic matter in
631 streams. *Journal of the North American Benthological Society* 27:521-528. Doi: 10.1899/07-
632 102.1

633 Kaspari M, Wright J, Yavitt J, Harms K, Garcia M, Santana M. 2008. Multiple nutrients limit
634 litterfall and decomposition in a tropical forest. *Ecology Letters* 11:35-43. Doi:
635 10.1111/j.1461-0248.2007.01124.x

636 Kominoski JS, Pringle CM, Ball BA, Bradford MA, Coleman DC, Hall DB, Hunter MD. 2007.
637 Nonadditive effects of leaf litter species diversity on breakdown dynamics in a detritus-based
638 stream. *Ecology* 88:1167-1176. Doi: 10.1890/06-0674

639 Krauss GJ, Solé M, Krauss G, Schlosser D, Wesenberg D, Bärlocher F. 2011. Fungi in
640 freshwaters: ecology, physiology and biochemical potential. *FEMS microbiology reviews*
641 35:620-651. Doi: 10.1111/j.1574-6976.2011.00266.x

642 Laughlin DC. 2011. Nitrification is linked to dominant leaf traits rather than functional
643 diversity. *Journal of Ecology* 99:1091-1099. Doi: 10.1111/j.1365-2745.2011.01856.x

644 Lecerf A, Dobson M, Dang CK, Chauvet E. 2005. Riparian plant species loss alters trophic
645 dynamics in detritus-based stream ecosystems. *Oecologia* 146:432-442. Doi:
646 10.1007/s00442-005-0212-3

647 Lecerf A, Risnoveanu G, Popescu C, Gessner MO, Chauvet E. 2007. Decomposition of diverse
648 litter mixtures in streams. *Ecology* 88:219-227. [https://doi.org/10.1890/0012-](https://doi.org/10.1890/0012-9658(2007)88[219:DODLMI]2.0.CO;2)
649 [9658\(2007\)88\[219:DODLMI\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88[219:DODLMI]2.0.CO;2)

650 Lecerf A, Marie G, Kominoski JS, LeRoy CJ, Bernadet C, Swan CM. 2011. Incubation time,
651 functional litter diversity, and habitat characteristics predict litter-mixing effects on
652 decomposition. *Ecology* 92:160-169. Doi: 10.1890/10-0315.1

653 Leroy CJ, Marks JC. 2006. Litter quality, stream characteristics and litter diversity influence
654 decomposition rates and macroinvertebrates. *Freshwater biology* 51:605-617. Doi:
655 10.1111/j.1365-2427.2006.01512.x

656 Lummer D, Scheu S, Butenschoen O. 2012. Connecting litter quality, microbial community
657 and nitrogen transfer mechanisms in decomposing litter mixtures. *Oikos* 121:1649-1655.
658 Doi: 10.1111/j.1600-0706.2011.20073.x

659 Makkonen M, Berg MP, Handa IT, Hättenschwiler S, van Ruijven J, van Bodegom PM, ...
660 Aerts R. 2012. Highly consistent effects of plant litter identity and functional traits on
661 decomposition across a latitudinal gradient. *Ecology Letters* 15:1033-1041. Doi:
662 10.1111/j.1461-0248.2012.01826.x

663 Mokany K, Ash J, Roxburgh S. 2008. Functional identity is more important than diversity in
664 influencing ecosystem processes in a temperate native grassland. *Journal of Ecology* 96:884-
665 893. Doi: 10.1111/j.1365-2745.2008.01395.x

666 National Research Council. 2005. *Mineral Tolerance of Animals*, 2nd ed. National Academic
667 Press, Washington.

668 Petchey OL, Gaston KJ. 2006. Functional diversity: back to basics and looking forward.

669 Ecology letters 9:741-758.doi: 10.1111/j.1461-0248.2006.00924.x

670 Piscart C, Genoel R, Dolédec S, Chauvet E, Marmonier P. 2009. Effects of intense agricultural
671 practices on heterotrophic processes in streams. *Environmental Pollution* 157:1011-1018.
672 Doi: 10.1016/j.envpol.2008.10.010

673 Piscart C, Navel S, Maazouzi C, Montuelle B, Cornut J, Mermillod-Blondin F, Creuse des
674 Chatelliers M, Simon L, Marmonier P. 2011. Leaf litter recycling in benthic and hyporheic
675 layers in agricultural streams with different types of land use. *Science of the Total
676 Environment* 409:4373-4380. Doi: 10.1016/j.scitotenv.2011.06.060

677 Presa Abos C, Lepori F, McKie BG, Malmqvist B. 2006. Aggregation among resource patches
678 can promote coexistence in stream-living shredders. *Freshwater Biology* 51:545-553.
679 <https://doi.org/10.1111/j.1365-2427.2006.01509.x>

680 Quested H, Eriksson O, Fortunel C, Garnier E. 2007 Plant traits relate to whole-community
681 litter quality and decomposition following land use change. *Functional Ecology* 21:1016-
682 1026. Doi: 10.1111/j.1365-2435.2007.01324.x

683 R Core Team. 2013. R: A Language and Environment for Statistical Computing. R Foundation
684 for Statistical Computing, Vienna, Austria.

685 Ritz K. 2006. Fungal roles in transport processes in soils. In MG Gadd eds, *Fungi in
686 biogeochemical cycles*. Cambridge University Press, Cambridge, UK, pp. 51-73.

687 Sanpera-Calbet I, Lecerf A, Chauvet E. 2009. Leaf diversity influences in-stream litter
688 decomposition through effects on shredders. *Freshwater Biology* 54:1671-1682. Doi:
689 10.1111/j.1365-2427.2009.02216.x

690 Santonja M, Baldy V, Fernandez C, Balesdent J, Gauquelin T. 2015a. Potential shift in plant
691 communities with climate change in a Mediterranean Oak forest: consequence on nutrients
692 and secondary metabolites release during litter decomposition. *Ecosystems* 18:1253-1268.
693 Doi: 10.1007/s10021-015-9896-3

694 Santonja M, Fernandez C, Gauquelin T, Baldy V. 2015b. Climate change effects on litter
695 decomposition: intensive drought leads to a strong decrease of litter mixture interactions.
696 *Plant and Soil* 393:69-82. Doi: 10.1007/s11104-015-2471-z

697 Santonja M, Pellan L, Piscart C. 2018a. Macroinvertebrate identity mediates the effects of litter
698 quality and microbial conditioning on the leaf litter recycling in temperate streams. *Ecology*
699 *and Evolution* 8:2542-2553. Doi: 10.1002/ece3.3790

700 Santonja M, Foucault Q, Rancon A, Gauquelin T, Fernandez C, Baldy V, Mirleau P. 2018b.
701 Contrasting responses of bacterial and fungal communities to plant litter diversity in a
702 Mediterranean oak forest. *Soil Biology and Biochemistry* 125:27-36. Doi:
703 10.1016/j.soilbio.2018.06.020

704 Santschi F, Gounand I, Harvey E, Altermatt F. 2018. Leaf litter diversity and structure of
705 microbial decomposer communities modulate litter decomposition in aquatic systems.
706 *Functional Ecology* 32:522-532. Doi: 10.1111/1365-2435.12980

707 Schimel JP, Hättenschwiler S. 2007. Nitrogen transfer between decomposing leaves of different
708 N status. *Soil Biology and Biochemistry* 39:1428-1436. Doi: 10.1016/j.soilbio.2006.12.037

709 Schindler MH, Gessner MO. 2009. Functional leaf traits and biodiversity effects on litter
710 decomposition in a stream. *Ecology* 90:1641-1649. Doi: 10.1890/08-1597.1

711 Srivastava DS, Cardinale BJ, Downing AL, Duffy JE, Jouseau C, Sankaran M, Wright JP. 2009.
712 Diversity has stronger top-down than bottom-up effects on decomposition. *Ecology*
713 90:1073-1083. Doi: 10.1890/08-0439.1

714 Swan CM, Palmer MA. 2004. Leaf diversity alters litter breakdown in a piedmont stream.
715 *Journal of the North America Benthological Society* 23:15-28.

716 Swan CM, Palmer MA. 2006a. Composition of speciose leaf litter alters stream detritivore
717 growth, feeding activity and leaf breakdown. *Oecologia* 147:469-478. Doi: 10.1007/s00442-
718 005-0297-8

719 Swan CM, Palmer MA. 2006b. Preferential feeding by an aquatic consumer mediates non-
720 additive decomposition of speciose leaf litter. *Oecologia* 149:107-114. Doi:
721 10.1007/s00442-006-0436-x

722 Swan C M. 2011. Consumer presence and resource diversity independently induce stability of
723 ecosystem function in a Piedmont stream. *Ecosphere* 2:136. Doi: 10.1890/ES11-00274.1

724 Swift MJ, Heal OW, Anderson JM. 1979. *Decomposition in terrestrial ecosystems*. Berkeley,
725 University of California press.

726 Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE. 1980. The river continuum
727 concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137. Doi: 10.1139/f80-
728 017

729 Villanueva VD, Albarino R, Canhoto C. 2012. Positive effect of shredders on microbial
730 biomass and decomposition in stream microcosms. *Freshwater Biology* 57:2504-2513. Doi:
731 10.1111/fwb.12023

732 Vos VC, van Ruijven J, Berg MP, Peeters ET, Berendse F. 2013. Leaf litter quality drives litter
733 mixing effects through complementary resource use among detritivores. *Oecologia* 173:269-
734 280. Doi: 10.1007/s00442-012-2588-1

735 Wallace JB, Eggert SL, Meyer JL, Webster JR. 1997. Multiple trophic levels of a forest stream
736 linked to terrestrial litter inputs. *Science* 277:102-104. Doi: 10.1126/science.277.5322.102

737 Wardle DA, Bonner KI, Nicholson KS. 1997. Biodiversity and plant litter: Experimental
738 evidence which does not support the view that enhanced species richness improves
739 ecosystem function. *Oikos* 79:247-258. Doi: 10.2307/3546010

740 Wetzel RG. 1995. Death, detritus, and energy flow in aquatic ecosystems. *Freshwater Biology*
741 33:83-89. Doi: 10.1111/j.1365-2427.1995.tb00388.x

742

743 **TABLES**

744

745 **Table 1.** Effects of litter quality (NPL vs. NRL), litter species identity (10 species), litterbag
 746 mesh size (fine-mesh vs. coarse-mesh litterbags), and their interactions on litter decomposition
 747 rate and microbial activity in the single-species litterbags. d.f. = degrees of freedom, %SS =
 748 percentage of type III sums of squares. *F*-values and associated *P*-values (with the respective
 749 symbols * for $P < 0.05$, ** for $P < 0.01$, and *** for $P < 0.001$) are indicated.

750

	d.f.	Litter decomposition		Microbial activity	
		%SS	<i>F</i> -value	%SS	<i>F</i> -value
Litter quality (LQ)	1	59.1	854.8 ***	60.0	239.7 ***
Litter species (LS)	8	12.6	22.8 ***	11.3	5.6 ***
Mesh size (MS)	1	15.8	228.9 ***	11.1	44.3 ***
LQ × MS	1	6.6	95.9 ***	1.4	5.7 *
LS × MS	8	1.6	3.0 **	1.1	0.6
Residuals	60	4.2		15.0	

751

752

753 **Table 2.** Effects of litter quality mixing (NPL-NPL, NPL-NRL and NRL-NRL), litterbag mesh
754 size (fine-mesh *vs.* coarse-mesh litterbags), and their interactions i) on litter decomposition rate
755 and microbial activity and ii) on relative mixture effects (RME) on litter decomposition rate
756 (LD) and on microbial activity (MA). d.f. = degrees of freedom, %SS = percentage of type III
757 sums of squares. *F*-values and associated *P*-values (with the respective symbols * for $P < 0.05$,
758 ** for $P < 0.01$, and *** for $P < 0.001$) are indicated.

759

	d.f.	Litter decomposition		Microbial activity		RME on LD		RME on MA	
		%SS	<i>F</i> -value	%SS	<i>F</i> -value	%SS	<i>F</i> -value	%SS	<i>F</i> -value
Litter quality (LQ)	2	49.1	617.5 ***	55.5	506.2 ***	21.6	58.0 ***	19.7	51.9 ***
Mesh size (MS)	1	30.8	776.2 ***	23.1	421.8 ***	11.5	61.5 ***	13.4	70.7 ***
LQ × MS	2	6.0	75.3 ***	2.1	18.6 ***	0.8	2.3	0.0	0.1
Residuals	354	14.1		19.3		66.0		66.8	

760

761

762 **Table 3.** Effects of community-weighted mean traits (CWM), functional trait dissimilarities
763 (FD), litterbag mesh size (fine-mesh vs. coarse-mesh litterbags), and their interactions i) on
764 litter decomposition rate and microbial activity and ii) on relative mixture effects (RME) on
765 litter decomposition (LD) and on microbial activity (MA). CWM1 and CWM2, and FD1 and
766 FD2 represented the two first components of the PCAs conducted using the CWM or the FD
767 values in [Supplementary Fig. S2](#). d.f. = degrees of freedom, %SS = percentage of type III sums
768 of squares. *F*-values and associated *P*-values (with the respective symbols * for $P < 0.05$, **
769 for $P < 0.01$, and *** for $P < 0.001$) are indicated.

770

	d.f.	Litter decomposition		Microbial activity		RME on LD		RME on MA	
		%SS	F-value	%SS	F-value	%SS	F-value	%SS	F-value
CWM1	1	48.8	2693.2 ***	55.1	1344.1 ***	15.4	81.8 ***	23.1	129.3 ***
CWM2	1	1.6	86.4 ***	1.6	38.1 ***	0.3	1.4	0.3	1.7
FD1	1	3.7	202.3 ***	1.8	43.2 ***	0.2	0.8	0.2	1.1
FD2	1	1.9	107.1 ***	1.4	33.9 ***	3.7	19.9 ***	0.5	2.9
Mesh size (MS)	1	30.8	1704.0 ***	22.9	559.5 ***	11.5	60.8 ***	13.3	74.6 ***
CWM1 × MS	1	6.0	332.0 ***	1.9	46.5 ***	0.3	1.3	0.0	0.3
CWM2 × MS	1	0.1	3.4	0.3	7.1 **	0.7	3.6	0.1	0.3
FD1 × MS	1	0.7	40.1 ***	0.7	17.5 ***	0.4	2.0	0.2	1.2
FD2 × MS	1	0.1	5.3 *	0.1	2.7	1.6	8.3 **	0.0	0.2
Residuals	350	6.3		14.3		66.0		62.2	

771

772

773 **Table 4.** Relationships between CWM values of chemical element concentrations and relative
 774 mixture effects (RME) on litter decomposition rate and on microbial activity in fine-mesh and
 775 coarse-mesh litterbags. Adjusted R^2 and associated P -values (* $P < 0.05$, ** $P < 0.01$, *** $P <$
 776 0.001) are indicated.
 777

	C	N	P	Ca	K	Mg	Na	S
RME on litter decomposition								
Fine-mesh litterbags	0.19 **	0.19 **	0.28 ***	0.26 ***	0.14 **	0.21 **	0.18 **	0.09 *
Coarse-mesh litterbags	0.17 **	0.12 *	0.27 ***	0.35 ***	0.24 ***	0.30 ***	0.33 ***	0.15 **
RME on microbial activity								
Fine-mesh litterbags	0.12 *	0.02	0.13 *	0.27 **	0.26 ***	0.31 ***	0.27 ***	0.21 ***
Coarse-mesh litterbags	0.07	0.16 **	0.27 ***	0.41 ***	0.30 ***	0.43 ***	0.16 **	0.16 **

778


779

780 **FIGURES**

781

782 **Fig. 1.** (a) Hypothetical relationship between the different categories of litter mixtures and the
783 litter decomposition rate of these litter mixtures. According to the mass-ratio hypothesis, the
784 decomposition rate increases with the increase in CWM values in nutrient concentration within
785 a litter mixture. (b) Hypothetical relationship between the different categories of litter mixtures
786 and the intensity of litter diversity effects occurring in these litter mixtures. According to the
787 niche complementary hypothesis, the intensity of litter diversity effects increases with the
788 increase in FD values in nutrient concentration, which would be shown by an increase in litter
789 decomposition rate in the present study. More precisely, this increase in decomposition rate,
790 due to synergistic effects between the two litters in mixtures, will be higher in litter mixtures
791 exhibiting intermediate nutrient concentration compared to litter mixtures exhibiting low or
792 high nutrient concentrations (Fig. 1a). NPL = nutrient-poor litter, NRL = nutrient-rich litter.
793 For both relationships, red dotted lines illustrate hypothetical increases in both litter
794 decomposition rate and litter diversity effects mediated by macroinvertebrate presence.

795




796

797

798 **Fig. 2.** Principal component analysis (PCA) based on the chemical element concentrations of
799 the 10 litter species. Variance explained by each principal component and associated
800 eigenvalues are shown in brackets.


801



802

803


804 **Fig. 3.** Litter decomposition rate (a) and microbial activity (b) in the single-species litterbags
805 according to litter quality (nutrient-poor litter [NPL] vs. nutrient-rich litter [NRL]) and to
806 litterbag mesh size (fine-mesh vs. coarse mesh litterbags). Each bar represents the mean value
807 \pm SE; n= 20. The litter decomposition rate is indicated in percent mass loss relative to the initial
808 mass. The microbial activity is expressed as $\mu\text{g C-CO}_2$ production per h and per g of litter.
809



810

811

812 **Fig. 4.** Litter decomposition rate (a, b) and microbial activity (c, d) according to litter quality
813 mixing (a, c) and to litterbag mesh size (b, d). Each bar represents the mean value \pm SE. NPL
814 = nutrient-poor litter, NRL = nutrient-rich litter, FMB = fine-mesh litterbag, CMB = coarse-
815 mesh litterbag. Different letters denote significant differences between treatments with $a < b <$
816 c.
817




818

819

820 **Fig. 5.** Relative mixture effects (RME) on litter decomposition rate (a, b) and on microbial
 821 activity (c, d) according to litter quality mixing (a, c) and to litterbag mesh size (b, d). Each bar
 822 represents the mean value \pm SE. NPL = nutrient-poor litter, NRL = nutrient-rich litter, FMB =
 823 fine-mesh litterbag, CMB = coarse-mesh litterbag. The RME on litter decomposition rate and
 824 on microbial activity are indicated as the relative difference between the observed and the
 825 expected values from the respective single litter species treatments. Different letters denote
 826 significant differences between treatments with $a < b < c$.


827



828

829

830 **Fig. 6.** Relative mixture effects (RME) on litter decomposition rate (panels a, b, c and d) and
831 on microbial activity (panels e, f, g and h) as a function of community weighted mean traits
832 (CWM1 and CWM2 from the PCA using the CWM values, [Supplementary Fig. S2a](#)) and
833 functional trait dissimilarities (FD1 and FD2 from the PCA using the FD values, [Supplementary](#)
834 [Fig. S2b](#)) in fine-mesh (white symbol) and coarse-mesh (grey symbol) litterbags. The RME on
835 litter decomposition rate and on microbial activity are indicated as the relative difference
836 between the observed and the expected values from the respective single litter species
837 treatments. Significant relationships are indicated with dotted (fine-mesh litterbags) or grey
838 (coarse-mesh litterbags) lines.
839



840

841