



**HAL**  
open science

## **Incorporating phylogeography for modelling the distribution of the carob tree (*Ceratonia siliqua*, Leguminosae) in future climate change**

Alex Baumel, Eleonora Potenza, Valentine Frelon, Juan Viruel, Gonzalo Nieto-Feliner, Magda Bou Dagher Kharrat, Lahcen Ouahmane, Hervé Sanguin, Stefano La Malfa, Mario Diguardo, et al.

### ► To cite this version:

Alex Baumel, Eleonora Potenza, Valentine Frelon, Juan Viruel, Gonzalo Nieto-Feliner, et al.. Incorporating phylogeography for modelling the distribution of the carob tree (*Ceratonia siliqua*, Leguminosae) in future climate change. Genetics to the rescue: Managing forests sustainably in a changing world. GENTREE final conference, Jan 2020, Avignon, France. . hal-02460403

**HAL Id: hal-02460403**

**<https://amu.hal.science/hal-02460403v1>**

Submitted on 30 Jan 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Incorporating phylogeography for modelling the distribution of the carob tree (*Ceratonia siliqua*, Leguminosae) in future climate change.

Alex Baumel, Eleonora Potenza, Valentine Frelon, Juan Viruel, Gonzalo Nieto-Feliner, Magda Bou Dagher Kharrat, Lahcen Ouahmane, Hervé Sanguin, Stefano La Malfa, Mario Diguardo, Samuel Pironon, Jean-Pierre Suc, Nicolas Le Galliot, Frédéric Médail, Agathe Leriche.

The carob is an evergreen thermophilous fruit tree widely harvested throughout the Mediterranean for food and forage since antiquity. Currently, the carob is found in cultivation in orchards or in association with other crops, and it has expanded towards industrial, agricultural and soil restoration purposes. Wild populations are found in shrublands, forests as well as rocky outcrops and temporary river banks.




### Carob distribution modelling with Maxent

759 points, 6 climatic variables, linear, quadratic and interaction features


Selected climatic variables (Worldclim DB):  
 Annp: Mean annual temperature  
 Maxtvm: Maximum temperature warmest month  
 Mintcm: Minimum temperature coldest month  
 Psea: Precipitation seasonality  
 Twetq: Mean temperature of wettest quarter  
 Tsea: Temperature seasonality

Red : probability of presence 0.8-1  
 Grey: probability of presence 0.25-1  
 Green points: carob occurrences




### Pessimistic forecasts?

What happens if the climate niche is divided according to phylogeographic structure?




Carob phylogeography from Viruel et al. in press \*

Two distinctive lineages diverged after a strong historical bottleneck, which subsequently split in four genetic clusters across the Mediterranean (SM, South Morocco; SS, South Spain; CM, Central Mediterranean; EM, East Mediterranean). Admixture is frequent due to human dissemination, such as in the North Moroccan populations (NM) which show asymmetric introgression between Eastern and Western genetic clusters.



### New forecasting done with Maxent in respect to phylogeography and averaging results of 3 GCMs for RCP 4.5



A range contraction in the Mediterranean is predicted for carob tree due to climate change. Overlap analysis of climatic niches as well as Maxent modelling indicate that incorporating phylogeography may improve carob distribution modelling and forecasting. In contrast with forecast done at the species level, the analyses done at the genetic groups level support SM, and EM as the most likely persistent areas whereas the most affected would be SS.



\* Viruel J, et al. A strong east-west Mediterranean divergence supports a new phylogeographic history of the carob tree (*Ceratonia siliqua*, Leguminosae) and multiple domestications from native populations. *J Biogeogr.* 2019;00:1-12. <https://doi.org/10.1111/jbi.13726>