



HAL
open science

Assessing the effectiveness of bird rehabilitation: temporarily captive-reared Little Owls (*Athene noctua*) experience a similar recruitment rate as wild birds

Olivier Hameau, Alexandre Millon

► To cite this version:

Olivier Hameau, Alexandre Millon. Assessing the effectiveness of bird rehabilitation: temporarily captive-reared Little Owls (*Athene noctua*) experience a similar recruitment rate as wild birds. *Journal für Ornithologie = Journal of Ornithology*, 2019, 160 (2), pp.581-585. 10.1007/s10336-019-01633-2 . hal-02472247

HAL Id: hal-02472247

<https://amu.hal.science/hal-02472247>

Submitted on 10 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Assessing the effectiveness of bird rehabilitation: temporarily**
2 **captive-reared Little Owls (*Athene noctua*) experience similar**
3 **recruitment rate to wild birds.**

4
5 Olivier HAMEAU¹ & Alexandre MILLON²

6
7 ¹ Ligue pour la Protection des Oiseaux Provence Alpes Côte d'Azur (LPO PACA), villa Saint
8 Jules, 6 avenue Jean Jaurès, 83400 Hyères, France

9 ² Aix Marseille Université, CNRS, IRD, Avignon Université, Institut Méditerranéen
10 Biodiversité & Ecologie (IMBE), Technopôle Arbois-Méditerranée, 13545 Aix-en-Provence,
11 France

12
13 Corresponding author: alexandre.millon@imbe.fr +33 442908474

14 AM orcid ID: 0000-0002-9475-4123

15 Keywords: owls, bird care centre, population reinforcement, breeding success

16 Running title: Measuring the effectiveness of owl rehabilitation

17
18 For consideration in *Journal of Ornithology* as a short note

19
20 **Acknowledgments**

21 We thank the Ligue pour la Protection des Oiseaux (LPO) team from the bird care centre of
22 Buoux and the network of volunteers dedicated to bird rescue in Provence. The care center
23 received fundings from Vaucluse and Bouches-du-Rhône councils, Provence-Alpes-Côte
24 d'Azur region and Direction Régionale de l'Environnement, de l'Aménagement et du
25 Logement-PACA. Owl monitoring received initial support from a LEADER scheme in
26 partnership with Luberon natural park and Fondation Nature & Découvertes. Ringing was
27 achieved under a licence delivered by the Centre de Recherche sur la Biologie des
28 Populations d'Oiseaux. Finally, we thank M. Gruebler and an anonymous reviewer for their
29 helpful and relevant comments.

30

31 **Abstract:** A large amount of young birds are caught each year, shortly after having left the
32 nest, and brought to care centers by people. Those birds are temporarily hand-raised before
33 release. The effectiveness of this action remains however largely unassessed. Here we
34 monitored the fate of 119 rehabilitated Little Owls (*Athene noctua*) and found a recruitment
35 rate similar to wild birds (11.8% of 119 rehabilitated birds vs. 10.7% of 382 wild fledglings).
36 Whether rehabilitated birds were released in autumn, or in following spring, seems not to
37 affect recruitment probabilities, although the latter showed a tendency for reduced breeding
38 success and dispersal compared to wild birds, suggesting autumn releases may be favoured.

39 **Introduction**

40 The period after fledging is a critical phase during which young birds leave their nest with
41 only limited flight skills (Cox et al. 2014). During this period, they are highly vulnerable to
42 predation, but can also be collected by unaware people and brought to bird care centres.
43 There, they will be hand-reared and usually kept in conditions where they can improve flight
44 skills, before being released into the wild.

45 This ex-situ conservation action is widespread, especially for nocturnal raptor species in
46 which chicks leave their nest well before being able to fly. In France for instance, a total of
47 2333 young owls of seven species have been brought to eight bird care centres between 2009
48 and 2015, a collect which was unnecessary in 78% of the cases (A.-L. Dugué & Ligue pour la
49 Protection des Oiseaux, *pers. comm.*). The effectiveness of this action has however been
50 rarely assessed and dedicated studies often suffer from data paucity and/or the lack of a proper
51 control group (Joys et al. 2003). Whether temporary captivity at young age affect bird
52 probability to recruit into the wild population and successfully reproduce has not been
53 properly investigated to date (Ellis et al. 2000, Goldsworthy et al. 2000).

54 Here we assessed the rehabilitation effectiveness of young Little Owls (*Athene noctua*) by
55 monitoring the fate of birds released in an intensively-monitored study area and comparing
56 recruitment probability, dispersal and annual breeding success to wild birds. We further tested
57 whether releasing rehabilitated birds in the next spring, instead of in the autumn, can reduce
58 overwinter mortality and therefore could enhance the efficacy of reinforcement/reintroduction
59 schemes (Van Nieuwenhuysse et al. 2008, Mitchell et al. 2011).

60

61 **Methods**

62 *Study species, study area and population monitoring*

63 The Little Owl is a small-sized nocturnal raptor occurring in temperate and Mediterranean
64 regions of the Western Palearctic. There the species inhabits open farmland, including
65 vineyards and orchards. Chicks leave their nest at 28-32 days, with limited flight skills (Van
66 Nieuwenhuysen et al. 2008). Another 10-14 days are required to the owlets for flying properly
67 (Schönn et al. 1991).

68 We monitored a wild population of Little Owls over 100 km² in the Apt valley
69 (43°54'11"N 5°17'37"E), Luberon natural park, south-eastern France (Fig. S1). The area
70 consists in a mix of farming areas including vineyards, orchards, cereals (74%) with wooded
71 areas (21%). Between 97 and 115 nest-boxes have been monitored annually between 2006
72 and 2017, according to a standardised protocol (see online resource). The number of nest-box
73 occupied by a breeding pair increased from 5 to 24 between 2006 and 2009, then oscillated
74 between 25 and 34 from 2010 onwards. An unknown number of pairs bred outside nest-
75 boxes, in natural cavities or buildings, and were therefore not monitored. A peak of vocal
76 activities occurred in March-April and the median date of first-egg laying was April 29th ($N =$
77 204). Nest-box monitoring included the capture and ringing of breeding adults and of all
78 chicks when 15-20 days old (see online resource for a detailed protocol). We considered an
79 owl as a recruit when it has been caught in a nest-box containing eggs or chicks. Dispersal
80 distances were calculated between the birth nest-box (wild birds), or the release nest-box
81 (rehabilitated birds), and the nest-box used for prime reproduction.

82 *Captive-rearing techniques and release*

83 Between 5 and 25 fledglings (mean = 14 ± 6 individuals), typically 4-6 weeks old, were
84 brought annually to the bird care centre held by LPO-PACA in Buoux (43°49'55"N
85 5°22'42"E), 5 km from the centre of the study area. Birds were brought from the Provence-
86 Alpes-Côte-d'Azur region (PACA), within a radius of ca. 100 km around Buoux. Owls were
87 kept indoor for a week in a box to ascertain they were able to feed by themselves. They were

88 then transferred to small outdoor aviaries (dimensions L×W×H: 8×6×3m) for four weeks and
89 finally into a larger pre-release aviary (30×6×2.5m). Contacts with humans were limited to a
90 unique daily feeding event. Food items were a mix of dead 1-day-old chicken and mice (2
91 prey.day⁻¹). No live prey were given.

92 A total of 119 Little Owls have been released between 2008 and 2015 (Table 1). Birds
93 were fitted with a metal ring (Museum Paris) and sexed using molecular techniques (see
94 online resource). Thirty-two birds have been additionally fitted with a radio-transmitter
95 weighing 2.5g glued on central tail feathers (see online resource for details). Releases
96 systematically consisted in one female and one male (one exception in 2013, see Table 1) put
97 together in a nest-box that was unoccupied during the previous breeding season. No food was
98 provided in the nest-box. Four cohorts (birds born in 2007-2010; $N = 74$) were released in
99 March of the subsequent year $t+1$, i.e. at the start of the breeding season. The rationale behind
100 this was to allow the owls to spend the winter under benign conditions, fed *ad libitum*, for
101 reducing overwinter mortality. Then, three cohorts (birds born in 2013-2015; $N = 45$) were
102 released in September of their birth year. This period precedes the autumn peak of vocal
103 activity in Little Owls, when dispersal and territory acquisition takes place (Exo 1988). No
104 owls born in 2011-2012 were released in the study area.

105 *Statistical analyses*

106 Statistical analyses were run using R 3.4.3 (R Development Core Team 2017). Recruitment
107 probabilities were modelled using generalised linear mixed models with binomial distribution
108 of error and year as random factor (function *glmmPQL*). Dispersal data were modelled using
109 linear models with log₁₀-transformed distances (adding the minimal non-zero recorded
110 distance, δ). Breeding success was measured as the number of fledglings raised by a female
111 Little Owl (male data were too sparse for conducting similar analyses) minus the annual mean
112 number of fledgling per pair, to account for among-year variability. Relative breeding success

113 was then modelled using mixed linear models with female identity as random factor.
114 Individual age was included as an explanatory covariate (log-transformed). Residuals from
115 Gaussian models (dispersal and breeding success) were checked for normality and
116 homoscedasticity. Regression coefficients (β) were shown ± 1 SE.

117

118 **Results & Discussion**

119 *Recruitment probability*

120 Overall, owls passed through the care centre had a recruitment probability similar to wild
121 birds from the same cohorts (14 recruits out of 119 rehabilitated birds, 11.8% vs. 41/382 wild
122 birds, 10.7%; $\beta = 0.10 \pm 0.33$, $P = 0.76$; Table 1). Annual recruitment rates for the two groups
123 were slightly correlated ($r = 0.74$, $N = 7$, $P = 0.057$), suggesting similar processes were
124 governing temporal variation in recruitment. Transmitters did not seem to affect the
125 probability of an owl to recruit (6 recruits out of 32 birds with transmitters, 3 out of 42
126 without transmitters; $\beta = 0.68 \pm 0.48$, $P = 0.15$). This result is not surprising given most birds
127 had lost their transmitters before the start of the breeding season, therefore limiting the burden
128 of carrying extra-weight (for details see Fig. S2).

129 *What is the best season for releasing rehabilitated owls?*

130 To answer this question, we compared the fate of birds released in autumn t vs. spring $t+1$.
131 Although we acknowledge the optimal setting would have been to release birds from a same
132 cohort at the two seasons, the within-cohort comparison of recruitment probabilities between
133 rehabilitated and wild birds provides nevertheless some relevant information. Among the 79
134 birds kept in captivity during their first winter, 5 died overwinter (6.3%), indicating captivity
135 strongly reduced winter mortality, apparent survival of first-year Little Owls ranges between 8
136 and 30% (Exo and Hennes 1980, Schaub et al. 2006, Le Gouar et al. 2011). However,

137 recruitment probabilities were not higher than for birds released during the autumn of their
138 first year of life (12.2% vs. 11.1%; Table 1). Overall, recruitment probability of wild birds did
139 not differ from rehabilitated birds released in either period ($\beta = 0.19 \pm 0.42$, $P = 0.66$; $\beta =$
140 -0.40 ± 0.54 , $P = 0.46$, for spring and autumn release respectively).

141 Dispersal distances between natal or release and breeding nest-boxes ranged from 0 to
142 14010 m (median = 1960, $N = 86$). Females dispersed slightly further than males (log10-
143 transformed values + $\delta = 190$; $\beta = 0.20 \pm 0.08$, $P = 0.02$). Rehabilitated birds released in
144 spring showed shorter dispersal distance compared to wild birds ($\beta = -0.56 \pm 0.14$, $P <$
145 0.001), while there was no such difference when release took place in autumn ($\beta = -0.01 \pm$
146 0.17 , $P = 0.95$; Fig. 1).

147 Breeding success of female Little Owls increased with age (log-transformed age; $\beta = 0.61$
148 ± 0.26 , $P = 0.02$, $N = 114$ breeding events from 60 known-age females, including rehabilitated
149 birds). Rehabilitated females released in spring $t+1$ tended to have lower breeding success
150 than wild females ($\beta = -1.21 \pm 0.70$, $P = 0.09$, $N = 5$ females for 6 breeding events vs. 50/99),
151 while those released in autumn t did not suffer from such a reduction ($\beta = -0.30 \pm 0.62$, $P =$
152 0.63 , $N = 4/9$).

153 *Towards efficient owl release techniques*

154 Here we took advantage of the intensive monitoring of a wild population to accurately record
155 the recruitment of rehabilitated young Little Owls and released into the wild. The fate of
156 rehabilitated birds is rarely assessed and usually focus on survival, ignoring recruitment (i.e.
157 survival till effective reproduction), although this measurement is crucial to evaluate the
158 effectiveness of bird care centre (Van Nieuwenhuysse et al. 2008). Despite a hard-release
159 protocol (Haase 1993, Mitchell et al. 2011), our results showed that temporarily captive-raised
160 Little Owls had recruitment probabilities similar to wild birds and reproduced successfully.

161 However, we cannot rule out the possibility that wild birds have actually higher recruitment
162 rate than rehabilitated ones, associated to a higher propensity to successfully disperse outside
163 the study area (Amar et al. 2000). Our dispersal results, however, did not provide evidence for
164 this hypothesis. Contrary to our expectations, birds kept in captivity and thus provided with
165 food throughout winter, did not show higher recruitment rates than birds released in autumn.
166 This result suggest that mortality of juveniles may occur to a large extent shortly after
167 fledging, rather than throughout the winter season (Exo and Hennes 1980, Coles and Petty
168 1997, Cox et al. 2014, Perrig et al. 2017). Rehabilitated birds, whatever their timing of
169 release, escaped this critical period. Furthermore, evidences for reduced breeding success and
170 dispersal for birds released in spring suggest it may be preferable to release rehabilitated Little
171 Owls in autumn, during the dispersal phase, rather than in next year's spring. Under such
172 conditions, breeding success of rehabilitated birds did not significantly differ from wild birds.
173 Spring release might however be of interest in a reintroduction program for setting birds
174 locally.

175 In conclusion, while people education should be implemented to reduce the unnecessary
176 collect of young birds, our results demonstrate that simple hand-rearing and release
177 techniques are appropriate for rehabilitating young Little Owls.

178

179 **References**

180 Amar A, Arroyo, BE, Bretagnolle V (2000) Post-fledging dependence and dispersal in hatched
181 and wild Montagu's Harriers *Circus pygargus*. *Ibis* 142: 21–28.

182 Coles CF, Petty SJ (1997) Dispersal behaviour and survival of juvenile tawny owls (*Strix*
183 *aluco*) during the low point in a vole cycle. In Duncan, J. R. et al. (eds), *Biology and*
184 *conservation of owls of the northern hemisphere*. USDA Forest Service, North Central

185 Research Station, General Technical Report NC-190, pp. 111–118.

186 Cox AW, Thompson FRI, Cox AS, Faaborg J (2014) Post-fledging survival in passerine birds
187 and the value of post- fledging studies to conservation. *J Wildl Manage* 78: 183–193.

188 Ellis DH, Sullivan KA, Thomas NJ (2000) Post-realease survival of hand-reared and parent-
189 reared Mississippi Sandhill cranes. *Condor* 102: 104–112.

190 Exo KM (1988) Jahreszeitliche ökologische Anpassungen des Steinkauzes (*Athene noctua*). *J*
191 für Ornithol 129: 393–415.

192 Exo KM, Hennes R (1980) Beitrag zur Populationsökologie des Steinkauzes (*Athene noctua*)
193 Eine analyse deutscher und niederländischer ringfunde. *Vogelwarte* 30: 162–179.

194 Goldsworthy SD, Giese M, Gales RP, Brothers N, Hamill J (2000) Effects of the Iron Baron
195 oil spill on little penguins (*Eudyptula minor*). II. Post-release survival of rehabilitated
196 oiled birds. *Wildl Res* 27: 573–582.

197 Haase P (1993) Zur situation und brutbiologie des steinkauzes *Athene n. noctua* SCOP., 1769
198 im Westhaveland. *Naturshcutz und Landschaftspfl. Brand.* 2: 29–37.

199 Joys AC, Clark JA, Clark NA, Robinson RA (2003) An investigation of the effectiveness of
200 rehabilitation of birds as shown by ringing recoveries. *British Trust for Ornithology*
201 report, 49p.

202 Le Gouar PJ, Schekkerman H, van der Jeugd HP, Boele A, van Harxen R, Fuchs P, Stroeken
203 P, van Noordwijk AJ (2011) Long-term trends in survival of a declining population: the
204 case of the little owl (*Athene noctua*) in the Netherlands. *Oecologia*: 369–379.

205 Mitchell AM, Wellicome TI, Brodie D, Cheng KM (2011) Captive-reared burrowing owls
206 show higher site-affinity, survival, and reproductive performance when reintroduced
207 using a soft-release. *Biol Conserv* 144: 1382–1391.

208 Perrig M, Grübler MU, Keil H, NaefDaenzer B (2017) Post-fledging survival of Little Owls
209 *Athene noctua* in relation to nestling food supply. *Ibis* 159: 532–540.

210 R Development Core Team (2017) R: A language and environment for statistical computing.
211 R Foundation for Statistical Computing, Vienna, Austria.

212 Schaub M., Ullrich B, Knötzsch G, Albrecht P, Meisser C. (2006) Local population dynamics
213 and the impact of scale and isolation: a study on different little owl populations. *Oikos*
214 115: 389–400.

215 Schönn S, Scherzinger W, Exo KM, Ille R (1991) Der Steinkauz. Die Neue Brehm-Bücherei.
216 Ziemsen Verlag.

217 Van Nieuwenhuyse D, Genot J-C, Johnson DH (2008) The little owl, Conservation, Ecology
218 and Behavior of *Athene noctua*. Cambridge University Press.

219

220

221 **Table 1:** Numbers of young Little Owls released annually after being brought to bird care
 222 centre and later recaptured as breeder, to be compared with wild birds monitored in the same
 223 study area (Luberon natural park). Rehabilitated birds were released either in autumn of their
 224 first year of life (autumn t) or in next spring (spring $t+1$), after a winter kept in captivity.

225


Cohort	Release period	Number of birds released	Number of rehabilitated birds recruited	%	Number of ringed wild birds	Number of wild birds recruited	%
2007	Spring $t+1$	20	1	5%	16	1	6.3%
2008	Spring $t+1$	14	3	21.4%	37	7	18.9%
2009	Spring $t+1$	18	3	16.7%	63	8	12.7%
2010	Spring $t+1$	22	2	9.1%	59	9	15.3%
Total in spring $t+1$		74	9	12.2%	175	25	14.3%
2013	Autumn t	13	2	15.4%	65	7	10.8%
2014	Autumn t	12	1	8.3%	71	5	7%
2015	Autumn t	20	2	10%	71	4	5.6%
Total in autumn t		45	5	11.1%	207	16	7.7%
Grand total		119	14	11.8%	382	41	10.7%

226

227

228 Figure 1: Dispersal distances (± 1 SD) between natal or release nest-boxes and nest-boxes
229 where first breeding was recorded for Little Owls according to sex and bird type (rehabilitated
230 birds released in autumn [$N = 5$ females, 1 male] or in spring [$N = 5, 4$], and wild birds [$N =$
231 32, 39]).

232



233