



**HAL**  
open science

## Resistance to antibiotics of bacteria in tropical countries

Didier Raoult

► **To cite this version:**

Didier Raoult. Resistance to antibiotics of bacteria in tropical countries. *Lancet Planetary Health*, 2019, 3 (6), pp.e238-e239. <10.1016/S2542-5196(19)30092-0>. <hal-02512546>

**HAL Id: hal-02512546**

**<https://amu.hal.science/hal-02512546v1>**

Submitted on 25 Oct 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

1 **Resistance to antibiotics of bacteria in tropical countries**

2

3 Didier Raoult<sup>1\*</sup>

4

5 1. Aix Marseille Université, MEPHI, IRD, IHU Méditerranée Infection, Marseille, France

6

7 \* **Corresponding author:** Prof. Didier Raoult

8 Email: [didier.raoult@gmail.com](mailto:didier.raoult@gmail.com)

9 Phone: (+33) 4 13 73 24 01

10 Fax: (+33) 4 13 73 24 02

11

12 **Key words:** Antibiotic resistance, tropical countries, natural antibiotic compounds

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27 The identification of the source of resistance of bacteria to antibiotics is, in my opinion, too naïve and  
28 too much based on the guilt of human-related activity. This perception of the world yields analyses  
29 that claim to establish an exclusive link between the use of man-made antibiotics and the resistance  
30 of microorganisms to antibiotics. This illusion neglects the importance of the source of natural  
31 antibiotics. Most antibiotics are secreted naturally by microorganisms, bacteria, of the actinomycetes  
32 phylum, or fungi(1). These antibiotics are naturally present in the environment and the antibiotic  
33 resistance of microorganisms existed long before their medical use. Beta-lactamases are found in  
34 genetic sequences of microorganisms from the Middle Ages, and the resistance of *Staphylococcus*  
35 *aureus* in the 19th century has recently been demonstrated (2). Antibiotics are assembled in  
36 microorganisms by non-ribosomal protein synthases and polyketides synthases, which are the only  
37 known translation devices outside the ribosome. Their existence may have preceded that of  
38 ribosomes and may be archaic translation machineries. Since biodiversity in general is much higher in  
39 the humid intertropical zone, it is also likely to be the place where the most diverse microorganisms  
40 are found, as evidenced by the richness of the human microbiota in rural intertropical areas  
41 compared to urban or temperate areas(3). This biodiversity can be accompanied by a biodiversity of  
42 molecules with antibiotic activity. Finally, the molecules secreted by non-ribosomal protein synthase  
43 and polyketide synthase have multiple activities, not only antimicrobial but also antifungal. Against  
44 eukaryotes, some molecules are used as anti-cancer treatments. Some human beta lactamases have  
45 anticancer drug inhibitory activity as with mitomycin(4). The spectrum of activity of beta lactamases  
46 far exceeds the anti-beta lactamine activity, playing a role in the digestion of nucleic acid (4).

47

48 Antibiotic resistance is a global concern and, interestingly, the source of resistance is often found in  
49 the world's intertropical or hottest regions (5). This specificity has not been satisfactorily explained  
50 and has just been confirmed by a very extensive study carried out in this journal(6) which shows that  
51 a country's overall antibiotic consumption is not correlated with the level of bacterial resistance. On  
52 the contrary, there is an inverse relationship, the lower the prescription rate, the higher the

53 resistance level. We have confirmation of this in France, where antibiotic consumption is one of the  
54 highest in the world, and where the level of resistance is one of the lowest in the world, particularly  
55 in Marseille(7).The same is true in the United States, where difficult-to-treat Gram-negative bacteria  
56 represent a very small proportion of the bacteria isolated from the blood cultures of hospitalized  
57 patients(8). The reason for these differences remains poorly explained. One of the explanations  
58 proposed was the poor management of antibiotics prescribing in humans, which could explain this  
59 discrepancy(5). This seems to overlook the idea that man-made antibiotics are the only source of  
60 antibiotics, neglecting those produced in the environment. Antibiotics can eventually be used as  
61 food by other microorganisms, which in turn develop enzymes to digest them for consumption(9). It  
62 has been evaluated that most families of antibiotics can be used as growth factors for soil  
63 microorganisms, and the digestion of penicillin and its use as a sole source of carbon has been  
64 demonstrated in *Escherichia coli* transformed in vitro. This shows that antibiotics do not only kill  
65 microorganisms, but can also serve as a nutrient, especially in areas particularly rich in environmental  
66 microorganisms. In addition, it has recently been shown that some arthropods are capable of  
67 secreting penicillins and cephalosporins, leading to a less simplistic view of antibiotics, of selection  
68 pressure exclusively from human-made drugs, and resistance(10). Thus, beta-lactam resistance genes  
69 are detected in all vertebrates, including humans, but also in archaea, which are not sensitive to  
70 beta-lactams (11).The work conducted in this journal on Kenya shows that in a city where antibiotic  
71 prescribing is extremely low, the level of resistance of *Escherichia coli* is particularly high, not only in  
72 humans but also in wildlife. This study must put into perspective a much less anthropocentric  
73 approach but integrate the fact that antibiotics are probably more than 3 billion years old, as are  
74 antibiotic resistance enzymes, and that we can only understand, and therefore combat antibiotic  
75 resistance, if we have a more scientific vision of this world.

76 In conclusion, public health considerations lead to a significant simplification, in particular for  
77 sending simple messages that can be understandable to all, but partially false. This needs to be  
78 acknowledged. Control measures should not prevent the integration of recent scientific data for fear

79 of misuse of antibiotics. Such a line of thought might end up being dangerous for humans and more  
80 emphasis should be put on the complexity of the relationships between antibiotics and the enzymes  
81 that, among others, digest them.

82

### 83 **Funding**

84 No funding was received for this work.

85

### 86 **Competing interests**

87 There are no conflicts of interest.

88

### 89 **Reference List**

90

- 91 (1) Angelakis E, Merhej V, Raoult D. Related actions of probiotics and antibiotics on gut  
92 microbiota and weight modification. *Lancet Infect Dis* 2013 October;13(10):889-99.
- 93 (2) Raoult D, Rolain JM. The Living Croquet Theory: The *Staphylococcus aureus* Paradigm. *Int J*  
94 *Antimicrob Agents* 2019 April 12.
- 95 (3) Angelakis E, Bachar D, Yasir M, Musso D, Djossou F, Melenotte C et al. Comparison of the gut  
96 microbiota of obese individuals from different geographic origins. *New Microbes New Infect*  
97 2019 January;27:40-7.
- 98 (4) Lee SY, Brem J, Pettinati I, Claridge TD, Gileadi O, Schofield CJ et al. Cephalosporins inhibit  
99 human metallo beta-lactamase fold DNA repair nucleases SNM1A and SNM1B/apollo. *Chem*  
100 *Commun (Camb)* 2016 May 10;52(40):6727-30.
- 101 (5) Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. Anthropological and  
102 socioeconomic factors contributing to global antimicrobial resistance: a univariate and  
103 multivariable analysis. *Lancet Planet Health* 2018 September;2(9):e398-e405.
- 104 (6) Hassell JM, Ward MJ, Muloi D, Bettridge JM, Robinson TP, Kariuki S et al. Epidemiology of  
105 Clinically Relevant Antimicrobial Resistance at the Wildlife-Livestock-Human Interface in  
106 Nairobi. *The Lancet Global Health*. In press 2019.
- 107 (7) Le Page S., Dubourg G, Baron SA, Rolain JM, Raoult D. No global increase in resistance to  
108 antibiotics: a snapshot of resistance from 2001 to 2016 in Marseille, France. *Eur J Clin*  
109 *Microbiol Infect Dis* 2018 December 4.
- 110 (8) Raoult D, Rolain JM. The paradigm of the Shadoks and antibiotic resistance. *Clin Infect Dis*  
111 2019 March 6.

- 112 (9) Hibbing ME, Fuqua C, Parsek MR, Peterson SB. Bacterial competition: surviving and thriving  
113 in the microbial jungle. *Nat Rev Microbiol* 2010 January;8(1):15-25.
- 114 (10) Pontarotti P, Raoult D. Why do arthropods secrete beta-lactams? *Int J Antimicrob Agents*  
115 2019 April;53(4):370.
- 116 (11) Keshri V, Panda A, Levasseur A, Rolain JM, Pontarotti P, Raoult D. Phylogenomic Analysis of  
117 beta-Lactamase in Archaea and Bacteria Enables the Identification of Putative New  
118 Members. *Genome Biol Evol* 2018 April 1;10(4):1106-14.  
119  
120