



**HAL**  
open science

## SARS-CoV-2: fear versus data

Yanis Roussel, Audrey Giraud-Gatineau, Marie-Therese Jimeno, Jean-Marc Rolain, Christine Zandotti, Philippe Colson, Didier Raoult

### ► To cite this version:

Yanis Roussel, Audrey Giraud-Gatineau, Marie-Therese Jimeno, Jean-Marc Rolain, Christine Zandotti, et al.. SARS-CoV-2: fear versus data. *International Journal of Antimicrobial Agents*, 2020, pp.105947. 10.1016/j.ijantimicag.2020.105947 . hal-02517904

**HAL Id: hal-02517904**

**<https://amu.hal.science/hal-02517904>**

Submitted on 22 Aug 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## Coronavirus SARS-CoV-2: Fear versus Data

Yanis ROUSSEL<sup>\*1,2</sup>, Audrey GIRAUD-GATINEAU<sup>\*1,3,4,5</sup> Marie-Thérèse JIMENO<sup>5</sup>, Jean-Marc  
ROLAIN<sup>1,2</sup>, Christine ZANDOTTI<sup>1,2</sup> Philippe COLSON<sup>1,2</sup>, Didier RAOULT<sup>1,2</sup>

1 Institut Hospitalo-universitaire Méditerranée Infection, Marseille, France.

2 Aix Marseille Université, Institut de Recherche pour le Développement (IRD), Assistance Publique-  
Hôpitaux de Marseille, Evolution Phylogénie et Infections (MEPHI).

3 Aix Marseille Université, Institut de Recherche pour le Développement (IRD), Assistance Publique-  
Hôpitaux de Marseille, Service de Santé des Armées, Microbes Vecteurs Infections Tropicales et  
Méditerranéennes (VITROME).

4 Centre d'Epidémiologie et de Santé Publique des Armées (CESPA), Marseille, France.

5 Assistance Publique – Hôpitaux de Marseille (AP-HM).

### Corresponding author

Didier Raoult,

IHU Méditerranée Infection, Faculté de Médecine,

Aix-Marseille Université, 27 Boulevard Jean Moulin,

13385 Marseille CEDEX 5, France.

Tel: +33 (0)4 91 32 43 75; fax: + 33 (0)4 91 38 77 72. Email: [didier.raoult@gmail.com](mailto:didier.raoult@gmail.com)

\* These two authors contributed equally to the work

28 **Abstract**

29 The novel coronavirus from China (SARS-CoV2) is spreading in the World, creating a huge reaction  
30 despite its current low incidence outside China and the far East. Moreover, it is not compared to the 4  
31 current circulating coronaviruses that cause millions of case in the World. Here, we compare incidence  
32 and death rate of these 4 common coronaviruses to that of SARS-COV2 in OECD countries. We  
33 conclude that the problem of SARS-CoV2 is probably overestimated so far, as 2,6 millions people die  
34 annually of respiratory infections versus less than 4000 for SARS-CoV2 so far.

35

36 **Keywords:** COVID-19; coronavirus; SARS-CoV-2

37

## 38 **Introduction**

39 *Coronaviridae* represent a very important family of animal and human viruses (1; 2) that circulated  
40 permanently. These four common human coronaviruses (HKU1, NL63, OC43, E229) represent  
41 between 10 to 20% of respiratory infections worldwide and are present in all continents (3; 4; 5; 6; 7;  
42 8; 9; 10; 11; 12; 13; 14) (Table 3). Mortality is poorly assessed, but it is clear that there are chronic  
43 carriers as well as asymptomatic carriers. There are even two studies, one carried out by our team  
44 showing that there are as many asymptomatic carriers as symptomatic patients, and a second carried  
45 out in Ghana that shows that the number of asymptomatic carriers was not significantly different from  
46 the number of symptomatic carriers (3; 9). Since then, three epidemic episodes of emerging  
47 coronaviruses have been reported. The first, the SARS virus, has had very little impact on global  
48 morbidity and mortality since there have been more than 8000 recognized cases and 774 deaths (15;  
49 16). The second, MERS-Coronavirus, remained localized in Saudi Arabia, with a small epidemic of  
50 mainly nosocomial infections in South Korea (17). The MERS-Coronavirus, like SARS, have  
51 highlighted the major danger of nosocomial transmission to healthcare personnel who must absolutely  
52 be preserved in these epidemics (18). Finally, SARS-CoV-2 that appeared in December has expanded  
53 and has now affected more than 90,000 people worldwide (2; 19; 20). However, there is a significant  
54 number of cases only in the Far East so far. It is incontestably contagious since the example of a quasi-  
55 experimental study on the Diamond Princess cruise-ship that showed that if one locked up infected  
56 patients with uninfected patients, uninfected patients very quickly become infected in a confined  
57 world, and thus more 700 cases were diagnosed on this cruise-ship (21). However, commonly-  
58 circulating coronaviruses remain globally very predominant in the global situation because of their  
59 global distribution and their non-negligible mortality (22; 14). In this study, we wanted to share the  
60 experience of a reference laboratory representing around 1% of serious and diagnosed forms of  
61 respiratory infections, particularly seasonal, in France. This will allow evaluating the relative mortality  
62 of different human coronaviruses presented in hospitals in Marseille, compared to that of the Chinese  
63 coronavirus as it has appeared for the moment in countries with the same economic development,  
64 including France.

65

66 **Material and methods**

67 Assistance Publique – Hôpitaux de Marseille (AP-HM) is administrating all public hospitals in  
68 Marseille, France, including four university hospitals: La Timone hospital, Conception Hospital, North  
69 Hospital and South hospital, corresponding to 3,400 beds and 125,000 admissions each year (23). IHU  
70 Méditerranée Infection diagnosis laboratory is testing all samples from AP-HM in which respiratory  
71 viruses are suspected. The diagnosis is performed through molecular biology analysis. The results are  
72 monitored by a weekly automated surveillance system in IHU Méditerranée Infection, coupled to a  
73 laboratory information system (Nexlabs) (23). SARS-CoV-2 epidemiological data were obtained  
74 through an online platform gathering data from public agencies (24). Statistical analyses were  
75 performed by using BiostatGV online-software.

76

77

78 **Results**

79 In 2016, a total of 594 000 people deceased in France, among which 59,2% deceased in a care  
80 establishment (25). In 2016, a total of 2854 patients deceased in AP-HM hospitals. We can therefore  
81 estimate that each year, around 0,8% of people that deceased in care establishments in France died in  
82 AP-HM. This estimate allows us to have an idea of the number of peoples affected by a pathogen in  
83 France according to the number of people who died each year at the AP-HM.

84 From January, 1<sup>st</sup> 2013 to December, 31<sup>th</sup> 2019, 21 662 samples were tested by the AP-HM diagnosis  
85 laboratory. Among these, 770 samples were positive for coronavirus, of which 8 died (Mortality rate:  
86 1%). Among identified coronaviruses, 63 were identified as Coronavirus HKU1 (1 died, Mortality rate  
87 1.6%), 74 were identified as Coronavirus NL63 (2 died, Mortality rate 2.7%), 92 as Coronavirus E229  
88 (1 died, Mortality rate 1.1%) and 160 as Coronavirus OC43 (4 died, mortality rate 2.5%). 381  
89 Coronavirus, diagnosed before 2017, were not assigned to any of these four strains (Table 1).

90 We performed systematic testing (molecular biology) in the context of SARS-CoV-2 from January, 1<sup>st</sup>  
91 2020 to March, 2<sup>nd</sup> 2020, 7059 samples from patients presenting viral infections symptoms were  
92 tested by AP-HM diagnosis laboratory. Among them, 543 samples were positive to Coronavirus with  
93 2 deaths (Mortality rate: 0,36%): 277 were positive to Coronavirus HKU1, 146 Coronavirus NL63, 77  
94 to Coronavirus OC43, and 43 to Coronavirus 229E. No SARS-CoV-2 have been identified among  
95 these samples. One patient died with Coronavirus OC43 (Mortality rate 1,3%) and 1 patient died with  
96 Coronavirus HKU1 (Mortality rate 0,36%). No patients died with Coronavirus NL63 and E229 during  
97 this period.

98 During the same period, IHU Méditerranée Infection has been the regional center for the detection of  
99 the novel coronavirus SARS-CoV-2. A total of 596 analyses were performed on suspect patients since  
100 the emergence of the novel pathogen, among which 4 were positive to the SARS-CoV-2, and 709 were  
101 performed on French returnees from the Hubei province, all negative.

102 Until the 2<sup>nd</sup> of March 2020, a total of 90 307 patients were tested positive to SARS-CoV-2 in the  
103 world, among which 3086 died (mortality rate 3,4%). Including OECD countries only, 7476 patients  
104 were tested positive to SARS-CoV-2, among which 96 died (Mortality rate 1.3%) (Table 2). In France,  
105 191 people were tested positive, 3 died (Mortality rate 1,6%).

106 We compared the mortality rate of Coronavirus SARS-CoV-2 in OECD countries (1,3%) with the  
107 mortality rate of common coronaviruses identified in AP-HM patients (0,8%) from January 1st, 2013  
108 to March 2nd, 2020. We performed a  $\chi^2$  test on these values. The p-value was 0.11 (not significant).

109

110

111 **Discussion**

112 In this work, we highlight that the mortality of common Coronavirus infections is of 0.8% in France.  
113 This is to be compared with the mortality of the SARS-CoV-2 in the European or American developed  
114 countries of comparable economic level, which is of 1,3% (Table 2). If our extrapolation of deaths in  
115 AP-HM is true, in metropolitan France this would represent  $543 / 0.8 * 100 = 67,875$  cases of patients  
116 hospitalized with a respiratory infection with common coronaviruses in two months, almost as much  
117 as SARS-CoV-2 in the world. In fact, mortality from respiratory infections is extremely dependent on  
118 the quality of care and access to care, and severe forms have a better prognosis in countries with  
119 superior medical infrastructures. Under these conditions, there does not seem to be a significant  
120 difference in the mortality of Coronavirus SARS-CoV-2 in OECD countries with an economic level  
121 comparable to that of common coronaviruses ( $\chi^2$  test  $p=0.11$ ). Of course, the major flaw in this study  
122 is that the percentage of death attributable to the virus is not determined, but this is the case for all of  
123 the studies reporting respiratory virus infections, including those of SARS-CoV-2. Indeed, viral  
124 infections are ecosystem infections where the outcome depends on the inoculums and the surrounding  
125 microbiota (26). Thus, certain bacteria seem to be associated with symptomatic manifestations, such as  
126 *Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Staphylococcus aureus*, which are known to  
127 be the bacteria causing an excess of mortality due to secondary infection. Finally, seasonality,  
128 geographic location, heat, humidity are co-factors, as are age, gender and underlying pathologies.  
129 Under these conditions and all other things being equal, the disease with SARS-CoV-2 cannot be  
130 described as being statistically more severe than that linked to currently circulating other  
131 coronaviruses .

132 Finally, in particular in OECD countries, the SARS-CoV-2 does not seem to be particularly deadlier  
133 than other circulating viruses. In addition to coronaviruses, there are 16 endemic viruses commonly  
134 circulating in developed countries (Adenovirus, Bocavirus, Cytomegalovirus, Enterovirus, Influenza A  
135 H1N1 virus, Influenza A H3N2 virus, Influenza B virus, Metapneumovirus, Parainfluenzae virus 1,  
136 Parainfluenzae virus 2, Parainfluenzae virus 3, Parainfluenzae virus 4, Parechovirus, Picornavirus,  
137 Rhinovirus, Syncytial respiratory virus) and 2.6 million deaths from respiratory infections (apart from  
138 tuberculosis) per year have been noted in recent years in the world (27). There is little chance that the



139 emergence of SARS-CoV-2 could change this statistic significantly. Fear could have a larger impact  
140 than the virus itself: a first case of suicide motivated by the fear of SARS-COV-2 has been reported in  
141 India (28).

142 In addition, coronaviruses that have rarely been tested systematically around the world may persist in  
143 the pharynx of asymptomatic people being a potential source of population immunity (29).  
144 Furthermore, it should be noted that in systematic studies that have been carried out for other  
145 coronaviruses (but not yet for SARS-CoV-2), the percentage of asymptomatic carriers is equal or even  
146 higher as compared to symptomatic patients. We may soon have the same data for SARS-CoV-2,  
147 which will further reduce the relative risk associated with this specific pathology.

148

#### 149 **Declarations**

150 **Funding:** This work was supported by the French Government under the “Investments for the Future”  
151 programme managed by the National Agency for Research (ANR), Méditerranée-Infection 10-IAHU-  
152 03.

153 **Competing Interests :** No

154 **Ethical Approval :** Not applicable. Testing of repatriates was approved by the ethical board of the  
155 Committee for the Protection of Persons (CPP Ile de France VI dated 06–02–2020).

156 **References**

- 157 (1) To KK, Hung IF, Chan JF, Yuen KY. From SARS coronavirus to novel animal and human  
158 coronaviruses. *J Thorac Dis.* 2013;5 Suppl 2(Suppl 2):S103–S108
- 159 (2) Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease  
160 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center  
161 for Disease Control and Prevention [published online ahead of print, 2020 Feb 24]. *JAMA.*  
162 2020;10.1001/jama.2020.2648.
- 163 (3) Le-Viet N, Le VN, Chung H, et al. Prospective case-control analysis of the aetiologies of acute  
164 undifferentiated fever in Vietnam. *Emerg Microbes Infect.* 2019;8(1):339–352
- 165 (4) Hand J, Rose EB, Salinas A, et al. Severe Respiratory Illness Outbreak Associated with Human  
166 Coronavirus NL63 in a Long-Term Care Facility. *Emerg Infect Dis.* 2018;24(10):1964–1966
- 167 (5) Vandroux D, Allou N, Jabot J, et al. Intensive care admission for Coronavirus OC43 respiratory  
168 tract infections. *Med Mal Infect.* 2018;48(2):141–144
- 169 (6) Kanwar A, Selvaraju S, Esper F. Human Coronavirus-HKU1 Infection Among Adults in  
170 Cleveland, Ohio. *Open Forum Infect Dis.* 2017;4(2):ofx052. Published 2017 Mar 25
- 171 (7) Trombetta H, Faggion HZ, Leotte J, Nogueira MB, Vidal LR, Raboni SM. Human coronavirus and  
172 severe acute respiratory infection in Southern Brazil. *Pathog Glob Health.* 2016;110(3):113–118
- 173 (8) Lau SK, Woo PC, Yip CC, et al. Coronavirus HKU1 and other coronavirus infections in Hong  
174 Kong. *J Clin Microbiol.* 2006;44(6):2063–2071
- 175 (9) Owusu M, Annan A, Corman VM, et al. Human coronaviruses associated with upper respiratory  
176 tract infections in three rural areas of Ghana. *PLoS One.* 2014;9(7):e99782. Published 2014 Jul 31
- 177 (10) Zhang SF, Tuo JL, Huang XB, et al. Epidemiology characteristics of human coronaviruses in  
178 patients with respiratory infection symptoms and phylogenetic analysis of HCoV-OC43 during 2010-  
179 2015 in Guangzhou. *PLoS One.* 2018;13(1):e0191789. Published 2018 Jan 29
- 180 (11) Zeng ZQ, Chen DH, Tan WP, et al. Epidemiology and clinical characteristics of human  
181 coronaviruses OC43, 229E, NL63, and HKU1: a study of hospitalized children with acute respiratory  
182 tract infection in Guangzhou, China. *Eur J Clin Microbiol Infect Dis.* 2018;37(2):363–369

- 183 (12) Sipulwa LA, Ongus JR, Coldren RL, Bulimo WD. Molecular characterization of human  
184 coronaviruses and their circulation dynamics in Kenya, 2009-2012. *Virology*. 2016;13:18. Published  
185 2016 Feb 1
- 186 (13) Kiyuka PK, Agoti CN, Munywoki PK, et al. Human Coronavirus NL63 Molecular Epidemiology  
187 and Evolutionary Patterns in Rural Coastal Kenya. *J Infect Dis*. 2018;217(11):1728–1739.
- 188 (14) Killerby ME, Biggs HM, Haynes A, et al. Human coronavirus circulation in the United States  
189 2014-2017. *J Clin Virol*. 2018;101:52–56
- 190 (15) Peiris JS, Yuen KY, Osterhaus AD, Stöhr K. The severe acute respiratory syndrome. *N Engl J*  
191 *Med*. 2003;349(25):2431–2441
- 192 (16) de Wit E, van Doremalen N, Falzarano D, Munster VJ. SARS and MERS: recent insights into  
193 emerging coronaviruses. *Nat Rev Microbiol*. 2016;14(8):523–534.
- 194 (17) Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel  
195 coronavirus from a man with pneumonia in Saudi Arabia [published correction appears in *N Engl J*  
196 *Med*. 2013 Jul 25;369(4):394]. *N Engl J Med*. 2012;367(19):1814–1820
- 197 (18) Hui DS, Azhar EI, Kim YJ, Memish ZA, Oh MD, Zumla A. Middle East respiratory syndrome  
198 coronavirus: risk factors and determinants of primary, household, and nosocomial  
199 transmission. *Lancet Infect Dis*. 2018;18(8):e217–e227
- 200 (19) WHO. Coronavirus disease (COVID-19) outbreak. Available at:  
201 <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> [Accessed: 04.03.2020]
- 202 (20) Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. *Zhonghua Liu Xing*  
203 *Bing Xue Za Zhi*. 2020;41(2):145–151
- 204 (21) Rocklöv J, Sjödin H, Wilder-Smith A. COVID-19 outbreak on the Diamond Princess cruise ship:  
205 estimating the epidemic potential and effectiveness of public health countermeasures [published online  
206 ahead of print, 2020 Feb 28]. *J Travel Med*. 2020;taaa030.
- 207 (22) Gagneur A, Vallet S, Talbot PJ, et al. Outbreaks of human coronavirus in a pediatric and neonatal  
208 intensive care unit. *Eur J Pediatr*. 2008;167(12):1427–1434

209 (23) Abat C, Chaudet H, Colson P, Rolain JM, Raoult D. Real-Time Microbiology Laboratory  
210 Surveillance System to Detect Abnormal Events and Emerging Infections, Marseille, France. *Emerg*  
211 *Infect Dis.* 2015;21(8):1302–1310.

212 (24)  
213 [https://docs.google.com/spreadsheets/d/1f3LGuqwzegr7ZdGlzPOCDAYFk8RTaLTmMLF\\_K\\_5EVCC](https://docs.google.com/spreadsheets/d/1f3LGuqwzegr7ZdGlzPOCDAYFk8RTaLTmMLF_K_5EVCC/edit#gid=783518927)  
214 [/edit#gid=783518927](https://docs.google.com/spreadsheets/d/1f3LGuqwzegr7ZdGlzPOCDAYFk8RTaLTmMLF_K_5EVCC/edit#gid=783518927) Consulted on 03 march 2020, 9.00 GMT.

215 (25) INSEE, 594 000 personnes décédées en France en 2016, pour un quart d’entre elles à leur  
216 domicile, INSEE FOCUS, N°95, published 12/10/2017, <https://www.insee.fr/fr/statistiques/3134763>,  
217 published 12/10/2017, <https://www.insee.fr/fr/statistiques/3134763>

218 (26) Edouard S, Million M, Bachar D, et al. The nasopharyngeal microbiota in patients with viral  
219 respiratory tract infections is enriched in bacterial pathogens. *Eur J Clin Microbiol Infect Dis.*  
220 2018;37(9):1725–1733. doi:10.1007/s10096-018-3305-8

221 (27) GBD 2017 Mortality Collaborators. Global, regional, and national age-sex-specific mortality and  
222 life expectancy, 1950-2017: a systematic analysis for the Global Burden of Disease Study 2017  
223 [published correction appears in *Lancet.* 2019 Jun 22;393(10190):e44]. *Lancet.*  
224 2018;392(10159):1684–1735.

225 (28) Goyal K, Chauhan P, Chhikara K, Gupta P, Singh MP. Fear of COVID 2019: First suicidal case  
226 in India ! [published online ahead of print, 2020 Feb 27]. *Asian J Psychiatr.* 2020;49:101989.  
227 doi:10.1016/j.ajp.2020.101989

228 (29) Raoult D, Zumla A, Locatelli F, et al. Coronavirus infections: Epidemiological, clinical and  
229 immunological features and hypotheses. *Cell Stress*, in Press, [published online ahead of print, 2020  
230 Feb 24].  
231  
232

233

234 Table 1: Results for AP-HM diagnosis laboratory

Agent	Location	Confirmed cases	Deaths	Rate
<b>Coronavirus*</b>				
Coronavirus OC43	AP-HM, Marseille, France	160	4	0,0250
Coronavirus NL63		74	2	0,0270
Coronavirus HKU1		63	1	0,0159
Coronavirus E229		92	1	0,0109
<b>Coronavirus**</b>				
Coronavirus OC43	AP-HM, Marseille, France	77	1	0,0130
Coronavirus NL63		146	0	0
Coronavirus HKU1		277	1	0,0036
Coronavirus E229		43	0	0

\* From 2013-01-01 to 2019-12-31

\*\* From 2020-01-01 to 2020-03-02

235 Table 2: SARS-CoV-2 international epidemiological situation (OECD countries)

236 Consulted on 03 march 2020, 9.00 GMT

Countries	Confirmed cases	Deaths	Rate
Germany	157	0	0,0%
Australia	30	1	3,3%
Austria	18	0	0,0%
Belgium	8	0	0,0%
Canada	24	0	0,0%
South Korea	4335	28	0,6%
Denmark	4	0	0,0%
Spain	123	0	0,0%
Estonia	1	0	0,0%
United States	100	6	6,0%
Finland	7	0	0,0%
France	191	3	1,6%
Greece	7	0	0,0%

Ireland	1	0	0,0%
Iceland	3	0	0,0%
Israel	10	0	0,0%
Italy	2036	52	2,6%
Japan	274	6	2,2%
Latvia	1	0	0,0%
Lithuania	1	0	0,0%
Luxembourg	1	0	0,0%
Mexico	5	0	0,0%
Norway	25	0	0,0%
New Zealand	1	0	0,0%
Netherlands	18	0	0,0%
Portugal	2	0	0,0%
Sweden	15	0	0,0%
Switzerland	38	0	0,0%
United Kingdom	40	0	0,0%
TOTAL	7476	96	1,3%

237

238

239 Table 3: List of studies that tested Coronavirus in fever cases in various countries

REFERENCE	COUNTRY	TESTED CASES	Diagnosis of Coronavirus
Trombetta	Brazil	775	7.6%
Zang	Guangzou China	13 048	244 (2.25%)
Zeng	Guangzou China	11 399	489 (4.3%)
Killerby	USA	18 806	2.2%
Kiyuka	Kenya	5 573	10.1%
Owusu	Ghana	593 cases	13.7%
Dube	South Africa	620 controls 214 (TB)	10.5% 8%
Sipulwa	Kenya	417	8.4%
Subramoney	South Africa	860	4.8%
Nunes	South Africa	1 026	15%
Le Viet	Viet-Nam	378	4 (1,05%)