



Platelets, Thrombo-Inflammation, and Cancer: Collaborating With the Enemy

Ana-Luisa Palacios Acedo, Diane Mege, Lydie Crescence, Francoise
Dignat-George, Dubois Christophe, Panicot-Dubois Laurence

► To cite this version:

Ana-Luisa Palacios Acedo, Diane Mege, Lydie Crescence, Francoise Dignat-George, Dubois Christophe, et al.. Platelets, Thrombo-Inflammation, and Cancer: Collaborating With the Enemy. *Frontiers in Immunology*, 2019, 1, 10.3389/fimmu.2019.01805 . hal-02544297

HAL Id: hal-02544297

<https://amu.hal.science/hal-02544297>

Submitted on 16 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Platelets, Thrombo-Inflammation, and Cancer: Collaborating With the Enemy

Ana Luisa Palacios-Acedo¹, Diane Mège^{1,2}, Lydie Crescence¹, Françoise Dignat-George¹, Christophe Dubois^{1*} and Laurence Panicot-Dubois¹

¹ Aix Marseille Univ, INSERM 1263, INRA 1260, Center for CardioVascular and Nutrition Research (C2VN), Marseille, France,

² Department of Digestive Surgery, Timone University Hospital, Marseille, France

OPEN ACCESS

Edited by:

Mirta Schattner,
National Council for Scientific and
Technical Research
(CONICET), Argentina

Reviewed by:

Paola Patrignani,
Università degli Studi G. d'Annunzio
Chieti e Pescara, Italy
Anna-Karin Olsson,
Uppsala University, Sweden

*Correspondence:

Christophe Dubois
christophe.dubois@univ-amu.fr

Specialty section:

This article was submitted to
Molecular Innate Immunity,
a section of the journal
Frontiers in Immunology

Received: 15 April 2019

Accepted: 17 July 2019

Published: 31 July 2019

Citation:

Palacios-Acedo AL, Mège D,
Crescence L, Dignat-George F,
Dubois C and Panicot-Dubois L (2019)
Platelets, Thrombo-Inflammation, and
Cancer: Collaborating With the
Enemy. *Front. Immunol.* 10:1805.
doi: 10.3389/fimmu.2019.01805

Platelets are small anucleate cells present in the blood stream, their typical role in primary hemostasis has been well-described. However, new evidence suggests that they have critically important roles in cancer progression and inflammation. Cancer cells can activate platelets, thus using them as physical shields from blood shear forces and natural killer (NK) cells. The activated platelets may also regulate hematopoietic and immune cell migration toward the tumor site; therefore, contributing to the cancer-associated inflammation. The activation of platelets by cancer cells may also contribute to metastasis and cancer progression by stimulating deep venous thrombosis and neutrophil extracellular trap formations (NETs) that “hide” cancer cells. We strived to review the current literature to dissect the role of platelets in cancer-associated thrombosis and tumor microenvironment inflammation.

Keywords: platelets, thrombosis, inflammation, cancer, NETs

PLATELET MORPHOLOGY AND PHYSIOLOGY

Platelets were first described by Bizzozzo in 1882 who described them microscopically and established that platelets were the first component of the blood to adhere to damaged blood vessel walls *in vivo* and *in vitro* (1). Since his discovery, platelets have been traditionally linked to hemostasis and thrombus formation (1–4). However, recent studies have shown that they are key players in tumor progression and metastasis, inflammation, atherogenesis, and antimicrobial host defense (1–4).

Platelets are arguably the most beautiful cells in the human body as they have an extraordinary capacity for morphological change and powerful secretion properties (5). As small, anucleate, discoid cells they are the smallest in blood circulation; measuring 2–5 μm in diameter with a thickness of 0.5 μm and a mean cell volume of 6–10 fl (5–7). Platelets are originated from big, nucleated cells called megakaryocytes that reside in the bone marrow and are part of the hemopoietic cell line (6). Platelets have an average lifespan of 5–7 days in the blood stream; where they endure such harsh conditions that as they age, they are reduced in size (5, 8). The average healthy human has 150,000–400,000 platelets per microliter (platelets/ μl) in circulation at any given time, and changes in total platelet count and mean platelet volume are often related to pathological conditions and are used as an acute inflammatory marker (5, 9, 10).

The platelet membrane is covered in glycoproteins like GPIIb-IX-V, GPVI, and GP α IIb β III; which are essential for complete platelet aggregation and adhesion (11). The membrane also has protease activated receptors: PAR-1, PAR-4, and the P2Y family receptors that mediate activation

and aggregation (11). Platelets also contain three different kind of secretory granules: α -granules, dense granules, and lysosomal granules (12, 13).

The α -granules are the most represented and contain membrane-associated and soluble proteins that are expressed in the platelet membrane when it is activated (13). These membrane markers are involved in various processes; including cell adhesion, coagulation, inflammation, cell growth, and host defense (5, 13). They include P-selectin, fibrinogen, vonWillenbrand factor, epidermal growth factor, vascular endothelium growth factor, platelet-derived growth factor, and complement C3 and C4 precursors; to name a few (3, 7, 10).

Dense granules, on the other hand, are slightly rarer with just three to eight per human platelet (7, 13). They contain high concentrations of adenine nucleotides, specifically ADP and ATP; along with serotonin and histamine, which are released upon platelet activation (7, 13, 14). The third granule group, or lysosomes, is the least common with only 1–2 per cell (7, 13). They contain protein degrading enzymes like cathepsins, elastases, collagenases, and glucosidases as well as LAMP-1, LAMP-2, and CD63 (7, 13, 14).

Platelet activation may occur through contact with different agonists, the most predominant being: thrombin, ADP, von Willenbrand factor, and collagen (15). Thrombin is the most powerful platelet agonist; it acts on the GPIb-IX-V and the PAR receptors (7, 15). The PAR receptors have 4 subgroups: PAR2 is not present in the platelets and PAR3 functions only as a cofactor to PAR4 activation (15, 16). PAR1 is the most potent receptor for thrombin induced platelet activation, as it more sensitive to lower thrombin levels than PAR4 (15, 16).

Another strong platelet activator is adenosine diphosphate (ADP); it can be exogenous or released from the dense granules of activated platelets themselves; constituting an activation loop between converging platelets (10, 17). The ADP receptors in the platelets are the P2Y protein family (17). P2Y1 initiates ADP-induced platelet aggregation and is responsible for platelet shape change and P2Y12 amplifies and stabilizes the aggregation response (17). As the alpha granules contain ADP, this can constitute an activation loop between platelets that amplifies their aggregation (2, 17).

Von Willenbrand factor (vWF) is a large glycoprotein produced in the Weibel-Palade body of endothelial cells and by megakaryocytes; it is present in the platelet alpha granules and subendothelial connective tissue (7, 18, 19). It plays an essential role in primary and secondary hemostasis; as a mediator of platelet adhesion, and as a carrier for coagulation factor FVIII (18). The vWF is exposed in activated endothelial cells where it interacts with platelet GPIb α and supports platelet translocation to the subendothelium (19). The platelet α IIb β III integrin also

interacts with vWF, causing a cross linking of platelets that enables platelet aggregation and plug formation (18, 19).

When platelets have been activated, they expose negatively charged phosphatidylserine (PS) on their membrane through activation of scramblase (e.g., TMEM16F) (20). This acts as an anchor for the assembly of the prothrombinase complex which converts fibrinogen to fibrin (19). Activated platelets also contribute to the intrinsic pathway of coagulation by secreting Poly-P in their dense granules that activates FXII (19). Meanwhile, coagulation in itself will also activate platelets, as thrombin will cleave and activate PARs on the platelets; thus creating a positive feedback loop that greatly amplifies the hemostasis/coagulation process (20, 21).

PLATELETS, THROMBOSIS, AND CANCER

Tumor Cells Can Activate Platelets

The association between cancer and thrombosis has been known since 1865 when Armand Trousseau first described that localized cancers can induce venous thrombus formation at distant sites (21, 22). This malignant-associated thrombosis is one of the most common clinical manifestations in cancer patients and is associated to worse prognosis and survival (23). The major reason for the high thrombotic risk in cancer patients is that cancer cells can activate platelets and stimulate aggregation through direct and indirect mechanisms (19, 21). The tumor-cell induced platelet aggregation (TCIPA) has been demonstrated in various cell lines like pancreatic, colorectal, and kidney (24–26). Additionally, the TCIPA has been correlated to higher metastatic potential (19). There are several mechanisms involved in in platelet activation and TCIPA (20).

An important mechanism of TCIPA is the cancer cell secretion of thrombin (15, 27). Thrombin is a serine protease that converts fibrinogen to fibrin, but also activates coagulation factors V, VIII, XI, and XIII and the PAR receptors on platelets themselves (15, 19). Pancreatic and lung cancer in specific have been proven to activate platelets via thrombin secretion as well as thromboxane A2 secretion (28, 29). Tumor cells also express ADP, which activates platelets via the P2Y1 and P2Y12 receptors, making platelets release more ADP from their dense granules and thus activating other nearby platelets (30, 31). Interestingly; colon, prostate, and breast cancer cells can bind platelet Fc γ RIIIa and induce dense granule secretion in the platelets (11). Different cancer types like squamous and germinal have also been proven to express podoplanin which binds to platelet-expressed CLEC-2 and induces platelet activation (32).

Tissue factor is arguably the main activator of the coagulation cascade once it comes into contact with activated factor VIIa in the blood stream (2, 5, 30). Tissue factor is often expressed in cancer cells and cancer derived microparticles (2, 6, 27). Elevated levels of TF in the serum has been evidenced in several types of cancer and in chemotherapy-induced thrombosis (28). Platelet as well as cancer derived microparticles have also been described to express Tissue Factors in their membrane, and thus contribute to platelet activations and cancer-related thrombosis (33–35).

There are other indirect mechanisms of platelet activation by the cancer cells. For example, cancer-cell expressed mucins can

Abbreviations: ADP, adenosine diphosphate; CLEC 2, C-type Lectin Like 2 receptors; CXCL, chemokine; F, coagulation factor; IL, interleukin; MHC, major histocompatibility complex I; MP, microparticle; NET, neutrophil extracellular traps; PAR, proteinase activated receptor; PDGF, platelet-derived growth factor; PECAM, platelet-endothelial cell adhesion molecule; PSGL-1, P-Selectin protein ligand 1; TF, tissue factor; TGF, transforming growth factor; TLR, toll like receptor; VEGF, vascular endothelial growth factor.

force platelets and granulocytes to interact (36). Subsequently, there is bidirectional signaling and Cathepsin G release by the granulocytes, which cleaves the platelet protease activated receptor-4 (PAR4) and activating G proteins (G_q and $G_{12/13}$) to induce shape change and platelet activation (36, 37). There are also malignancy-linked deficiencies of the vWF cleaving protease: ADAM13. Its deficiency causes large vWF multimers to circulate which can in turn activate platelets (11, 36, 38). A correlation between the presence of metastatic tumors and the concentration of vWF multimers in circulation, as well as aberrant ADAM13 in circulation has been previously demonstrated (39, 40).

The activation of platelets by cancer cells has a myriad of pro-cancerous effects like stimulating tumor growth, preparing the metastatic niche, and helping the metastatic cells survive in circulation. The induction of a cyclooxygenase 2 (COX-2) mediated paracrine signaling between the stromal and epithelial cells in the adenoma mediated by activated platelets can give the ensuing cancer cells a more aggressive phenotype (41–43). However, it has been shown that low-dose aspirin can have an antimetastatic effect by inhibiting COX-1 (43–45). This inhibition would decrease the cancer-mediated platelet activation and aggregation; thus, having an anti-metastatic effect on the tumor cells (41, 44).

Platelets Influence Tumor Growth

Platelets have a myriad of growth factors stored in their alpha (α) granules (5–7, 13). They are present in the tumor microenvironment outside of the vasculature where they can come into direct contact with the malignant cells (31, 46). When activated, they secrete transforming growth factor beta (TGF- β), vascular endothelial growth factor (VEGF), and platelet derived growth factor (PDGF) (47, 48). These factors not only induce tumor growth, but also promote angiogenesis and tumoral neovascularization (14).

It is important to also take into account that platelets also have anti-cancerous effects. Recently, platelet-derived microRNA has been identified as an important regulator of tumor development (49). Platelet-derived microparticles transfers miR-24 into cancer cells which targets mt-Nd2 and Snora75; triggering mitochondrial regulation and inhibiting tumor growth (49). This shows that platelet function and effects on cancer progression may be stage and context dependent (21, 49).

Platelet Receptors Mediate Distant Pre-metastatic Niche Preparation

Platelets are covered in membrane receptors that promote heterotypic cell interactions (27, 30, 31). These interactions play a crucial role in tumor growth and metastatic spread (11, 30, 31). Cancer cells that enter blood circulation during the metastatic process are exposed to high shear stress and to the immune system; to survive they use activated platelets to shield themselves (10, 27).

P-selectin is expressed on the surface of activated platelets and endothelial cells and is an important adhesion molecule (27). Cancer cells can bind to platelet P-selectin through TCIPA and form aggregates to protect themselves from the blood circulation

and “hide” from NK cells (50). It has been proposed that platelet α Ib β 3 integrin can link fibrin with tumor α V β 3 integrin (19, 46, 47). The role of thrombin and integrin signaling is also very important in the platelet-cancer cell bonding mechanism (4, 27, 51). Thrombin increases the mRNA and protein levels of α V β 3 integrin and serves as a ligand to this receptor; it also increases the secretion of vascular-endothelial growth factor (VEGF) in human prostatic cancer cells (51).

Platelets also have an important effect on the preparation of the pre-metastatic niche (31, 52, 53). Primary tumors secrete metastasis-related proteins to the target organ that stimulate the migration of bone marrow-derived cells to create this pre-metastatic niche and stimulate neo-vasculogenesis (52, 53). Platelets have a role in managing the pre-metastatic communications; they secrete CXCL5 and CXCL7 upon contact with tumor cells to recruit granulocytes for the formation of the early metastatic niche (52–54). Activated platelets also release growth factors from their α granules, as well as metalloproteases that contribute to the degradation of the extracellular matrix and the preparation of the aforementioned metastatic niche (13, 54, 55).

Circulating Tumor Cell Survival and Arrest Is Mediated by Platelets

It is now widely accepted that increased platelet counts enhance cancer's metastatic power; while thrombocytopenia (low platelet count) may hinder the process (56). As previously stated, platelets have many adhesion molecules including integrins (α Ib β 3), selectins (P-selectin), leucine rich glycoproteins (P-selectin glycoprotein ligand -PSGL-1- and GPIb/V/IX), and immunoglobulin superfamily proteins (platelet-endothelial adhesion molecule -PECAM-1) (30, 31). These molecules allow them to form aggregates with cancer cells to protect them from the shear forces that would otherwise destroy their membranes (31, 50). These aggregates also serve to stabilize cancer cell arrest on the endothelial wall (27, 53, 55).

The TCIPA results in platelets coating the cancer cell and thus protecting it from the natural killer (NK) cells in the blood stream (57). They can also impair the NK cell mediated cytolytic/tumorlytic activity by secreting platelet TGF- β (57). The TGF- β impairs NK granule mobilization and interferon- γ secretion by downregulation the NKG2D immunoreceptor (50, 57). Another way that platelets aid the cancer cells escape the immune system is by membrane protein transfer (47, 48). In the midst of the platelet aggregate; the cancer cells can co-express platelet markers as well major histocompatibility (MHC) molecules to further camouflage themselves (58).

Platelets support cancer cell arrest in the same manner as it contributes to leucocyte arrest: by selectin (P-selectin) dependent rolling/tethering and integrin dependent adhesion (α Ib β 3, GP-Ib α , and vWF all contribute to firm adhesion) (21, 22). It is also important to note that many cancer cells express “platelet receptors” like α Ib β 3, α V β 3, or GP-Ib α (31, 58). These receptors not only help cancer cells escape the immune response but also mediate direct cancer-endothelial and cancer-leucocyte

interactions that promote cancer cell extravasation and prepare them for the colonization of the target tissue (31, 59).

Another important TCIPA effect on cancer endothelial transmigration is the release of ADP from the activated platelets' dense granules (31, 60, 61). ADP interacts with the endothelial PY2 receptor (P2Y1R), causing endothelial cell junctions to become laxer and enabling cancer cells to pass through more easily (60–62). Serotonin is also contained in the dense granules, and experimental studies have demonstrated that by blocking its receptor metastatic spread was inhibited (31, 63). Cancer patients that tend to have higher than average serotonin levels in the blood have a worse prognosis and survival (63).

PLATELETS, INFLAMMATION, AND CANCER

Malignant tumors have often been described as wounds that do not heal (64). Two of the most important tumor characteristics are their constitutive angiogenesis and perennial inflammation as well as the fibroblast infiltration and constant stroma regeneration (64). The vasculature in tumors is often fenestrated, facilitating the trans-endothelial transport, and exposing subendothelial factors like collagen and TF (14, 65). As we have previously stated, cancer cells can activate platelets

through the various TCIPA mechanisms; with the added effect of the exposed subendothelial procoagulant factors there is a continuous platelet-activation loop (5, 19, 66). The activated platelets release their granule content that modulates the tumor microenvironment, including pro-inflammatory cytokines (67).

The proinflammatory cytokines released by the platelets are powerful recruiters and activators of leucocytes (67). These molecules include CXCL1, CXCL4, CXCL5, CXCL7, CXCL12 (SDF-1), and interleukin-8 (IL8) (67, 68). The CXCL12 chemokine attracts hematopoietic cells to the tumor site, stimulating tumor growth, and angiogenesis (69). Macrophages are also CXCR4 positive cells that are recruited to the tumor site by the platelet-expressed CXCL12 (67). On the other hand, CXCL5, and CXCL7 platelet secretion in distant sites to the primary tumor recruit granulocytes to prepare the pre-metastatic niche (5, 50, 52–54).

The activated platelets also express IL-1 β (synthesized in the platelet from pre-mRNA) (70). The IL-1 β induces TF expression in endothelial cells and stimulates expression of endothelial-leucocyte adhesion molecules (70). IL-1 β also promotes platelet activation in an autocrine manner via the IL-1 receptor (67).

Transforming growth factor β (TGF- β) expressed and secreted by activated platelets in the tumor microenvironment has immunosuppressive properties and aids in the cancer cell escape




FIGURE 1 | Schematic review of the different mechanisms of platelet activation that can lead to thrombo-inflammation in cancer. Figure created using Servier Medical Art available at <http://smart.servier.com/>. Copyright Ana Luisa Palacios-Acedo.

from immune system recognition (71). TGF- β is also partially responsible for the transformation of the neutrophils toward a pro-tumorigenic phenotype (67).

PLATELETS AND NET FORMATION IN CANCER

Neutrophils are the body's first line of defense and have been traditionally characterized by two modes of action: pathogen engulfment and anti-microbial substance secretion (72). In recent years a new function has been identified: neutrophil extracellular traps (NETs) (72). The NETs are the result of the neutrophils' chromatin and granular content being expelled from the nucleus to form a web-like structure (67, 72). This structure can physically entrap and kill pathogens (67, 72). There are recent studies that suggest that NETs may also be involved in tumor progression, metastasis, and cancer-associated thrombosis (73).

Platelet TLR4 can trigger NETosis in activated neutrophils; and histones 3 and 4 released during the process can in turn, activate the platelets in a continuous loop (73, 74). The extracellular DNA in the NETs is capable of binding and activating coagulation factor XII as well as activating platelets directly (75). Additionally, activated platelet P-selectin can prime neutrophils through P-selectin glycoprotein ligand-1 (PSGL-1) activation and trigger NET formation (76, 77). These activation routes suggest that NETs are indeed a procoagulant factor as they provide a strong stimulus as well as a scaffold for thrombus formation (78). NETs promote fibrin deposition, recruit red blood cells and enhance platelet activation, and in turn, platelet activation promotes NET formation (76, 78).

Indeed, the link between NET formation and venous thromboembolism has long been established. In a baboon model of occlusion induced iliac thrombosis, researchers demonstrated an increase in circulation of NETs after 48 h that was maintained for 6 consecutive days; along with the presence of DNA markers in the thrombus (74). Another group demonstrated that plasma DNA is elevated in patients with deep vein thrombosis vs. healthy patients (79).

It is interesting to note that neutrophils originated from cancer patients are more prone to NETosis when exposed to PMA than those from healthy patients (80). This may be partially explained by the NET-activating properties of granulocyte colony-stimulating factor (G-CSF), and IL-8; which are locally secreted by tumor cells (67).

It has been proposed that tumor educated platelets may exert a pro-NETosis effect on the tumor-microenvironment neutrophils (80, 81). Cancer cells can allegedly use the NETs to protect themselves from shear stress in the circulation and from the immune system during the metastatic process (67, 81). The NET-induced platelet activation might play an important role in cancer progression, enhancing TCIPA, and the pro-thrombotic state (76). However, further research and information is needed to shed light on the contribution of platelets to the generation of NETs and their involvement in cancer progression.

CONCLUSION

Platelets are small but very powerful cells that interact with all components of the circulatory system. They are the main player in primary hemostasis but contribute to the secondary wave as well. As of recently, their involvement in the immune response was described, showing their power in regulating their environment. Their interactions with cancer cells and the tumor microenvironment are very complex and seem to have dual behaviors: pro and anti-cancerous, with the pro-cancerogenic effect out-numbering the anti-cancerous effects (Figure 1). However, it may seem that tumor education of platelets recruits them to the cancer cause, making them an ideal ally of tumor progression. This in turn, causes platelets to be continuously activated enhancing their thrombotic power and augmenting cancer-associated thrombosis. More research is needed in order to be able to establish the true power of these cells.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

REFERENCES

1. Brewer DB. Max Schultze (1865), G. Bizzozero (1882) and the discovery of the platelet. *Br J Haematol.* (2006) 133:251–8. doi: 10.1111/j.1365-2141.2006.06036.x
2. Lindemann S, Krämer B, Seizer P, Gawaz M. Platelets, inflammation, and atherosclerosis. *J Thromb Haemost.* (2007) 5 (Suppl. 1):203–11. doi: 10.1111/j.1538-7836.2007.02517.x
3. Engemann B, Massberg S. Thrombosis as an intravascular effector of innate immunity. *Nat Rev Immunol.* (2013) 13:34–45. doi: 10.1038/nri3345
4. Gay LJ, Felding-Habermann B. Contribution of platelets to tumour metastasis. *Nat Rev Cancer.* (2011) 11:123–34. doi: 10.1038/nrc3004
5. Holinstat, M. Normal platelet function. *Cancer Metastasis Rev.* (2017) 36:195–8. doi: 10.1007/s10555-017-9677-x
6. Machlus KR, Italiano JE. The incredible journey: from megakaryocyte development to platelet formation. *J Cell Biol.* (2013) 201:785–96. doi: 10.1083/jcb.201304054
7. Gremmel T, Frelinger A, Michelson A. Platelet physiology. *Semin Thromb Hemost.* (2016) 42:191–204. doi: 10.1055/s-0035-1564835
8. Thompson CB, Jakubowski JA, Quinn PG, Deykin D, Valeri CR. Platelet size and age determine platelet function independently. *Blood.* (1984) 63:1372–5.
9. Sayit AT, Gunbey PH, Terzi Y. Is the mean platelet volume in patients with acute cholecystitis an inflammatory marker? *J Clin Diagn Res.* (2015) 9:TC05–7. doi: 10.7860/JCDR/2015/12028.6061
10. Menter DG, Kopetz S, Hawk E, Sood AK, Loree JM, Gesele P, et al. Platelet “first responders” in wound response, cancer, and metastasis. *Cancer Metastasis Rev.* (2017) 36:199–213. doi: 10.1007/s10555-017-9682-0
11. Bambace NM, Holmes CE. The platelet contribution to cancer progression. *J Thromb Haemost.* (2011) 9:237–49. doi: 10.1111/j.1538-7836.2010.04131.x

12. Thon JN, Italiano JE. Platelets: production, morphology and ultrastructure. *Handb Exp Pharmacol.* (2012) 210:3–22. doi: 10.1007/978-3-642-29423-5_1
13. Pagel O, Walter E, Jurk K, Zahedi RP. Taking the stock of granule cargo: platelet releasate proteomics. *Platelets.* (2017) 28:119–28. doi: 10.1080/09537104.2016.1254762
14. Wojtukiewicz MZ, Sierko E, Hempel D, Tucker SC, Honn KV. Platelets and cancer angiogenesis nexus. *Cancer Metastasis Rev.* (2017) 36:249–62. doi: 10.1007/s10555-017-9673-1
15. De Candia E. Mechanisms of platelet activation by thrombin: a short history. *Thromb Res.* (2012) 129:250–6. doi: 10.1016/j.thromres.2011.11.001
16. Morrow DA, Braunwald E, Bonaca MP, Ameriso SF, Dalby AJ, Fish MP, et al. Vorapaxar in the secondary prevention of atherothrombotic events. *N Engl J Med.* (2012) 366:1404–13. doi: 10.1056/NEJMoa1200933
17. Hardy AR, Jones ML, Mundell SJ, Poole AW. Reciprocal cross-talk between P2Y1 and P2Y12 receptors at the level of calcium signaling in human platelets. *Blood.* (2004) 104:1745–52. doi: 10.1182/blood-2004-02-0534
18. Randi AM, Laffan MA. Von Willebrand factor and angiogenesis: basic and applied issues. *J Thromb Haemost.* (2017) 15:13–20. doi: 10.1111/jth.13551
19. Xu XR, Zhang D, Oswald BE, Carrim N, Wang X, Hou Y, et al. Platelets are versatile cells: new discoveries in hemostasis, thrombosis, immune responses, tumor metastasis and beyond. *Crit Rev Clin Lab Sci.* (2016) 53:409–30. doi: 10.1080/10408363.2016.1200008
20. Gyobu S, Ishihara K, Suzuki J, Segawa K, Nagata S. Characterization of the scrambling domain of the TMEM16 family. *Proc Natl Acad Sci USA.* (2017) 114:6274–9. doi: 10.1073/pnas.1703391114
21. Schlesinger M. Role of platelets and platelet receptors in cancer metastasis. *J Hematol Oncol.* (2018) 11:125. doi: 10.1186/s13045-018-0669-2
22. Varki A. Trousseau's syndrome: multiple definitions and multiple mechanisms. *Blood.* (2007) 110:1723–9. doi: 10.1182/blood-2006-10-053736
23. Ferroni P, Martini F, Portarena I, Grenga I, Riondino S, La Farina F, et al. An activated protein C-dependent thrombin generation assay predicts chemotherapy-associated venous thromboembolism in cancer patients. *Thromb Haemost.* (2011) 105:931–2. doi: 10.1160/TH10-11-0757
24. Heinmöller E, Weinl RJ, Heidtmann HH, Salge U, Seitz R, Schmitz I, et al. Studies on tumor-cell-induced platelet aggregation in human lung cancer cell lines. *J Cancer Res Clin Oncol.* (1996) 122:735–44. doi: 10.1007/BF01209121
25. Heinmöller E, Schropp T, Kisker O, Simon B, Seitz R, Weinl RJ. Tumor cell-induced platelet aggregation *in vitro* by human pancreatic cancer cell lines. *Scand J Gastroenterol.* (1995) 30:1008–16. doi: 10.3109/00365529509096346
26. Pearlstein E, Salk PL, Yogeewaran G, Karparkin S. Correlation between spontaneous metastatic potential, platelet-aggregating activity of cell surface extracts, and cell surface sialylation in 10 metastatic-variant derivatives of a rat renal sarcoma cell line. *Proc Natl Acad Sci USA.* (1980) 77:4336–9. doi: 10.1073/pnas.77.7.4336
27. Mezouar S, Frère C, Darbousset R, Mege D, Crescence L, Dignat-George F, et al. Role of platelets in cancer and cancer-associated thrombosis: experimental and clinical evidences. *Thromb Res.* (2016) 139:65–76. doi: 10.1016/j.thromres.2016.01.006
28. Adesanya MA, Maraveyas A, Madden LA. PO-27 - Thrombin generation in pancreatic cancer and multiple myeloma with use of calibrated automated thrombography. *Thromb Res.* (2016) 140:S186. doi: 10.1016/S0049-3848(16)30160-8
29. Coughlin SR. Protease-activated receptors in hemostasis, thrombosis and vascular biology. *J Thromb Haemost.* (2005) 3:1800–14. doi: 10.1111/j.1538-7836.2005.01377.x
30. Menter DG, Tucker SC, Kopetz S, Sood AK, Crissman JD, Honn KV. Platelets and cancer: a casual or causal relationship: revisited. *Cancer Metastasis Rev.* (2014) 33:231–69. doi: 10.1007/s10555-014-9498-0
31. Li N. Platelets in cancer metastasis: To help the “villain” to do evil. *Int J Cancer.* (2016) 138:2078–87. doi: 10.1002/ijc.29847
32. Suzuki-Inoue K, Fuller GL, García A, Eble JA, Pöhlmann S, Inoue O, et al. A novel Syk-dependent mechanism of platelet activation by the C-type lectin receptor CLEC-2. *Blood.* (2006) 107:542–9. doi: 10.1182/blood-2005-05-1994
33. Lechner D, Weltermann A. Chemotherapy-induced thrombosis: a role for microparticles and tissue factor? *Semin Thromb Hemost.* (2008) 34:199–203. doi: 10.1055/s-2008-1079261
34. Mezouar S, Mege D, Darbousset R, Farge D, Debourdeau P, Dignat-George F, et al. Involvement of platelet-derived microparticles in tumor progression and thrombosis. *Semin Oncol.* (2014) 41:346–58. doi: 10.1053/j.seminoncol.2014.04.010
35. Thomas GM, Brill A, Mezouar S, Crescence L, Gallant M, Dubois C, et al. Tissue factor expressed by circulating cancer cell-derived microparticles drastically increases the incidence of deep vein thrombosis in mice. *J Thromb Haemost.* (2015) 13:1310–9. doi: 10.1111/jth.13002
36. Shao B, Wahrenbrock MG, Yao L, David T, Coughlin SR, Xia L, et al. Carcinoma mucins trigger reciprocal activation of platelets and neutrophils in a murine model of Trousseau syndrome. *Blood.* (2011) 118:4015–23. doi: 10.1182/blood-2011-07-368514
37. Faraday N, Schunke K, Saleem S, Fu J, Wang B, Zhang J, et al. Cathepsin G-dependent modulation of platelet thrombus formation *in vivo* by blood neutrophils. *PLoS ONE.* (2013) 8:e71447. doi: 10.1371/journal.pone.0071447
38. Oleksowicz L, Bhagwati N, DeLeon-Fernandez M. Deficient activity of von Willebrand's factor-cleaving protease in patients with disseminated malignancies. *Cancer Res.* (1999) 59:2244–50.
39. Koo BH, Oh D, Chung SY, Kim NK, Park S, Jang Y, et al. Deficiency of von Willebrand factor-cleaving protease activity in the plasma of malignant patients. *Thromb Res.* (2002) 105:471–6. doi: 10.1016/S0049-3848(02)00053-1
40. Kumar S, Rao N, Ge R. Emerging roles of ADAMTSs in angiogenesis and cancer. *Cancers.* (2012) 4:1252–99. doi: 10.3390/cancers4041252
41. Patrignani P, Patrono C. Aspirin, platelet inhibition, and cancer prevention. *Platelets.* (2018) 29:779–85. doi: 10.1080/09537104.2018.1492105
42. Dovizio M, Bruno A, Tacconelli S, Patrignani P. Mode of action of aspirin as a chemopreventive agent. *Recent Results Cancer Res.* (2013) 191:39–65. doi: 10.1007/978-3-642-30331-9_3
43. Bruno A, Dovizio M, Tacconelli S, Patrignani P. Mechanisms of the antitumoural effects of aspirin in the gastrointestinal tract. *Best Pract Res Clin Gastroenterol.* (2012) 26:e1–13. doi: 10.1016/j.bpg.2012.10.001
44. Guillem-Llobat P, Dovizio M, Bruno A, Ricciotti E, Cufino V, Sacco A, et al. Aspirin prevents colorectal cancer metastasis in mice by splitting the crosstalk between platelets and tumor cells. *Oncotarget.* (2016) 7:32462–77. doi: 10.18632/oncotarget.8655
45. Patrignani P, Patrono C. Aspirin and cancer. *J Am Coll Cardiol.* (2016) 68:967–76. doi: 10.1016/j.jacc.2016.05.083
46. Karparkin S, Ambrogio C, Pearlstein E. The role of tumor-induced platelet aggregation, platelet adhesion and adhesive proteins in tumor metastasis. *Prog Clin Biol Res.* (1988) 283:585–606.
47. Wang S, Li Z, Xu R. Human cancer and platelet interaction, a potential therapeutic target. *Int J Mol Sci.* (2018) 19:1246. doi: 10.3390/ijms19041246
48. Waldmann TA. Cytokines in cancer immunotherapy. *Cold Spring Harb Perspect Biol.* (2018) 10:a028472. doi: 10.1101/cshperspect.a028472
49. Michael JV, Wurtzel JGT, Mao GF, Rao AK, Kolpakov MA, Sabri A, et al. Platelet microparticles infiltrating solid tumors transfer miRNAs that suppress tumor growth. *Blood.* (2017) 130:567–80. doi: 10.1182/blood-2016-11-751099
50. Egan K, Cooke N, Kenny D. Living in shear: platelets protect cancer cells from shear induced damage. *Clin Exp Metastasis.* (2014) 31:697–704. doi: 10.1007/s10585-014-9660-7
51. Tsopanoglou NE, Maragoudakis ME. Role of thrombin in angiogenesis and tumor progression. *Semin Thromb Hemost.* (2004) 30:63–9. doi: 10.1055/s-2004-822971
52. Kerr BA, McCabe NP, Feng W, Byzova TV. Platelets govern pre-metastatic tumor communication to bone. *Oncogene.* (2013) 32:4319–24. doi: 10.1038/ncr.2012.447
53. Labelle M, Begum S, Hynes RO. Platelets guide the formation of early metastatic niches. *Proc Natl Acad Sci USA.* (2014) 111:E3053–61. doi: 10.1073/pnas.1411082111
54. Rolli M, Fransvea E, Pilch J, Saven A, Felding-Habermann B. Activated integrin $\alpha_3\beta_1$ cooperates with metalloproteinase MMP-9 in regulating migration of metastatic breast cancer cells. *Proc Natl Acad Sci USA.* (2003) 100:9482–7. doi: 10.1073/pnas.1633689100
55. Buerger D, Wenz F, Groden C, Brockmann MA. Tumor-platelet interaction in solid tumors. *Int J Cancer.* (2012) 130:2747–60. doi: 10.1002/ijc.27441
56. Gasic GJ, Gasic TB, Stewart CC. Antimetastatic effects associated with platelet reduction. *Proc Natl Acad Sci USA.* (1968) 61:46–52. doi: 10.1073/pnas.61.1.46

57. Nieswandt B, Hafner M, Echtenacher B, Männel DN. Lysis of tumor cells by natural killer cells in mice is impeded by platelets. *Cancer Res.* (1999) 59:1295–300.
58. Placke T, Örgel M, Schaller M, Jung G, Rammensee HG, Kopp HG, et al. Platelet-derived MHC class I confers a pseudonormal phenotype to cancer cells that subverts the antitumor reactivity of natural killer immune cells. *Cancer Res.* (2012) 72:440–8. doi: 10.1158/0008-5472.CAN-11-1872
59. Steeg PS. New insights into the tumor metastatic process revealed by gene expression profiling. *Am J Pathol.* (2005) 166:1291–4. doi: 10.1016/S0002-9440(10)62348-6
60. Li WH, Qiu Y, Zhang HQ, Liu Y, You JF, Tian XX, et al. P2Y2 receptor promotes cell invasion and metastasis in prostate cancer cells. *Br J Cancer.* (2013) 109:1666–75. doi: 10.1038/bjc.2013.484
61. Ballerini P, Dovizio M, Bruno A, Tacconelli S, Patrignani P. P2Y12 receptors in tumorigenesis and metastasis. *Front Pharmacol.* (2018) 9:66. doi: 10.3389/fphar.2018.00066
62. Schumacher D, Strlic B, Sivaraj KK, Wetschurck N, Offermanns S. Platelet-derived nucleotides promote tumor-cell transendothelial migration and metastasis via P2Y2 receptor. *Cancer Cell.* (2013) 24:130–7. doi: 10.1016/j.ccr.2013.05.008
63. Leoncik V, Wu H, Ward LT, Kierzek AM, Plant NJ. Generation of 2,000 breast cancer metabolic landscapes reveals a poor prognosis group with active serotonin production. *Sci Rep.* (2016) 6:19771. doi: 10.1038/srep19771
64. Byun JS, Gardner K. Wounds that will not heal: pervasive cellular reprogramming in cancer. *Am J Pathol.* (2013) 182:1055–64. doi: 10.1016/j.ajpath.2013.01.009
65. Carmeliet P, Jain RK. Angiogenesis in cancer and other diseases. *Nature.* (2000) 407:249–57. doi: 10.1038/35025220
66. Goubran HA, Burnouf T, Radosevic M, El-Ekiaby M. The platelet–cancer loop. *Eur J Intern Med.* (2013) 24:393–400. doi: 10.1016/j.ejim.2013.01.017
67. Olsson AK, Cedervall J. The pro-inflammatory role of platelets in cancer. *Platelets.* (2018) 29:569–73. doi: 10.1080/09537104.2018.1453059
68. Gleissner CA, von Hundelshausen P, Ley K. Platelet chemokines in vascular disease. *Arterioscler Thromb Vasc Biol.* (2008) 28:1920–7. doi: 10.1161/ATVBAHA.108.169417
69. Sun X, Cheng G, Hao M, Zheng J, Zhou X, Zhang J, et al. CXCL12/CXCR4/CXCR7 chemokine axis and cancer progression. *Cancer Metastasis Rev.* (2010) 29:709–22. doi: 10.1007/s10555-010-9256-x
70. Brown GT, Narayanan P, Li W, Silverstein RL, McIntyre TM. Lipopolysaccharide stimulates platelets through an IL-1 autocrine loop. *J Immunol.* (2013) 191:5196–203. doi: 10.4049/jimmunol.1300354
71. Neuzillet C, Tijeras-Raballand A, Cohen R, Cros J, Faivre S, Raymond E, et al. Targeting the TGF β pathway for cancer therapy. *Pharmacol Ther.* (2015) 147:22–31. doi: 10.1016/j.pharmthera.2014.11.001
72. Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, et al. Neutrophil extracellular traps kill bacteria. *Science.* (2004) 303:1532–5. doi: 10.1126/science.1092385
73. Cools-Lartigue J, Spicer J, Najmeh S, Ferri L. Neutrophil extracellular traps in cancer progression. *Cell Mol Life Sci.* (2014) 71:4179–94. doi: 10.1007/s00018-014-1683-3
74. Kimball AS, Obi AT, Diaz JA, Henke PK. The emerging role of NETs in venous thrombosis and immunothrombosis. *Front Immunol.* (2016) 7:236. doi: 10.3389/fimmu.2016.00236
75. Jorch SK, Kubes P. An emerging role for neutrophil extracellular traps in noninfectious disease. *Nat Med.* (2017) 23:279–87. doi: 10.1038/nm.4294
76. Cedervall J, Hamidi A, Olsson A-K. Platelets, NETs, and cancer. *Thromb Res.* (2018) 164:S148–52. doi: 10.1016/j.thromres.2018.01.049
77. Etulain J, Martinod K, Wong SL, Cifuni SM, Schattner M, Wagner DD. P-selectin promotes neutrophil extracellular trap formation in mice. *Blood.* (2015) 126:242–6. doi: 10.1182/blood-2015-01-624023
78. Fuchs TA, Brill A, Duerschmied D, Schatzberg D, Monestier M, Myers DD, et al. Extracellular DNA traps promote thrombosis. *Proc Natl Acad Sci USA.* (2010) 107:15880–5. doi: 10.1073/pnas.1005743107
79. Diaz JA, Fuchs TA, Jackson TO, Kremer Hovinga JA, Lämmle B, Henke PK, et al. Plasma DNA is elevated in patients with deep vein thrombosis. *J Vasc Surg Venous Lymphat Disord.* (2013) 1:341. doi: 10.1016/j.jvsv.2012.12.002
80. Demers M, Krause DS, Schatzberg D, Martinod K, Voorhees JR, Fuchs TA, et al. Cancers predispose neutrophils to release extracellular DNA traps that contribute to cancer-associated thrombosis. *Proc Natl Acad Sci USA.* (2012) 109:13076–81. doi: 10.1073/pnas.1200419109
81. Erpenbeck L, Schön MP. Neutrophil extracellular traps: protagonists of cancer progression? *Oncogene.* (2017) 36:2483–90. doi: 10.1038/onc.2016.406

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Palacios-Acedo, Mège, Crescence, Dignat-George, Dubois and Panicot-Dubois. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.