



**HAL**  
open science

## Unilateral vestibular deafferentation impairs embodied spatial cognition

Diane Deroualle, Liliane Borel, Brandon Tanguy, Laurence Bernard-Demanze, Arnaud Devèze, Marion Montava, Jean-Pierre Lavieille, Christophe Lopez

► **To cite this version:**

Diane Deroualle, Liliane Borel, Brandon Tanguy, Laurence Bernard-Demanze, Arnaud Devèze, et al.. Unilateral vestibular deafferentation impairs embodied spatial cognition. *European Journal of Neurology*, 2019, 1, 10.1007/s00415-019-09433-7 . hal-02553545

**HAL Id: hal-02553545**

**<https://amu.hal.science/hal-02553545v1>**

Submitted on 24 Apr 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Click here to view linked References](#)

# 1 **Unilateral vestibular deafferentation impairs** 2 **embodied spatial cognition**

3  
4  
5  
6  
7  
8  
9 Diane Deroualle<sup>1</sup>, Liliane Borel<sup>1</sup>, Brandon Tanguy<sup>1</sup>, Laurence Bernard-Demanze<sup>2</sup>,  
10 Arnaud Devèze<sup>3</sup>, Marion Montava<sup>2</sup>, Jean-Pierre Lavieille<sup>2</sup>, and Christophe Lopez<sup>1(\*)</sup>  
11  
12  
13  
14  
15

16  
17 <sup>1</sup>Aix Marseille Univ, CNRS, LNSC, FR3C, Marseille, France

18  
19 <sup>2</sup>Department of Otorhinolaryngology, Head and Neck Surgery, Hôpital La Conception,  
20 APHM, Marseille, France

21  
22 <sup>3</sup>Ramsay Générale de Santé, Hôpital Privé Clairval, Department of Otorhinolaryngology,  
23 Head and Neck Surgery, Marseille, France  
24  
25  
26  
27  
28  
29  
30

## 31 AUTHORS' ORCID:

32 Diane Deroualle = [orcid.org/0000-0001-5710-9419](https://orcid.org/0000-0001-5710-9419)

33  
34 Marion Montava = [orcid.org/0000-0003-1347-9412](https://orcid.org/0000-0003-1347-9412)

35  
36 Christophe Lopez = [orcid.org/0000-0001-9298-2969](https://orcid.org/0000-0001-9298-2969)  
37  
38  
39  
40  
41  
42  
43  
44

45 (\*) Address for correspondence:

46 Dr Christophe Lopez

47  
48 Laboratoire de Neurosciences Intégratives et Adaptatives - UMR 7260

49  
50 Aix Marseille Univ & Centre National de la Recherche Scientifique (CNRS)

51  
52 Centre Saint-Charles, Fédération de Recherche 3C – Case B

53  
54 3, Place Victor Hugo

55  
56 13331 Marseille Cedex 03, France

57  
58 Tel: +33 4 13 55 08 41 Fax: + 33 4 13 55 08 44

59  
60 email: [christophe.lopez@univ-amu.fr](mailto:christophe.lopez@univ-amu.fr)  
61  
62  
63  
64  
65

35 **Abstract**

1  
2 36 A growing number of studies indicate that cognitive complaints are common in patients with peripheral  
3  
4 37 vestibular disorders. A better understanding of how vestibular disorders influence cognition in these  
5  
6 38 patients requires a clear delineation of the cognitive domains affected by vestibular disorders. Here, we  
7  
8 39 compared the consequences of left and right vestibular neurectomy on third-person perspective taking  
9  
10 40 – a visuo-spatial task requiring **mainly** own-body mental imagery, and on 3D objects mental rotation  
11  
12 41 imagery – requiring object-based mental imagery but no perspective taking. Patients tested one week  
13  
14 42 after a unilateral vestibular neurectomy and a group of age- and gender-matched healthy participants  
15  
16 43 played a virtual ball-tossing game from their own first-person perspective (1PP) and from the  
17  
18 44 perspective of a distant avatar (third-person perspective, 3PP). Results showed larger response times in  
19  
20 45 the patients with respect to their controls for the 3PP taking task, but not for the 1PP task and the 3D  
21  
22 46 objects mental imagery. In addition, we found that only patients with left vestibular neurectomy  
23  
24 47 presented altered 3PP taking abilities when compared to their controls. This study suggests that  
25  
26 48 unilateral vestibular loss affects mainly **own-body mental transformation** and that only left vestibular  
27  
28 49 loss seems to impair **this cognitive process**. Our study also brings further evidence that vestibular signals  
29  
30 50 contribute to the sensorimotor bases of social cognition and strengthens the connections between the so  
31  
32 51 far distinct fields of social neuroscience and human vestibular physiology.

33 52

34 53

35 54

36 55 **Keywords**

37 56 Perspective taking, Vestibular system, Unilateral vestibular loss, Mental imagery, 3D object mental  
38 57 rotation

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

## 58 Introduction

1  
2 59 A growing number of studies indicate that cognitive complaints are common in patients with vestibular  
3  
4 60 disorders (reviewed in [1, 2]). For example, a recent epidemiological study in US adults revealed that  
5  
6 61 patients with vestibular vertigo have an eight-fold increase in the odds of reporting impaired memory  
7  
8 62 and attention, limiting their activities [3]. A better understanding of how vestibular disorders influence  
9  
10 63 cognition in these patients requires a clear delineation of the cognitive domains affected by vestibular  
11  
12 64 disorders. Yet, the paucity of publications reporting negative findings makes it difficult to identify the  
13  
14 65 cognitive domains impaired and spared in patients with vestibular disorders.

15  
16 66 Several studies have compared populations of patients with vestibular disorders and controls,  
17  
18 67 with mixed findings depending on the aetiology, on unilateral vs bilateral vestibular loss, history of the  
19  
20 68 disease and the cognitive task. Popp *et al.* [4], for example, showed that bilateral vestibular loss impaired  
21  
22 69 visuospatial abilities, processing speed, short-term memory and executive functions, whereas unilateral  
23  
24 70 vestibular loss impaired only visuospatial abilities and processing speed. Spatial memory was impaired  
25  
26 71 in navigation studies using locomotion [5] and virtual reality tasks [6]. Vestibular disorders also  
27  
28 72 impaired space representation when participants were not moving or displaced at all, such as during  
29  
30 73 mental rotation of bodies and non-corporeal objects [7–10]. We note that it remains unclear from these  
31  
32 74 studies if vestibular disorders impair more egocentric (*i.e. own-body*) mental imagery or object-based  
33  
34 75 mental imagery.

35  
36 76 Another debated question concerns the consequences of left vs right peripheral vestibular  
37  
38 77 disorders on cognition. A rationale for this question is the anatomical and functional asymmetry within  
39  
40 78 the vestibulo-thalamo-cortical system. Neuroimaging studies have reported a predominance of  
41  
42 79 vestibular processing in the right cerebral hemisphere in right-handed participants, and an overall  
43  
44 80 dominance of the projections from one labyrinth to the ipsilateral cerebral hemisphere [11, 12]. As  
45  
46 81 various cognitive tasks involve differently lateralized brain areas, different effects of left and right  
47  
48 82 vestibular disorders on cognition can be predicted. Only few studies have tackled directly this question,  
49  
50 83 and overall their findings were controversial, showing stronger impairments in left [13] or right [14]  
51  
52 84 vestibular disorders, or equal impairments in left and right vestibular disorders [4].

53  
54 85 The present study aims at clarifying the vestibular contributions to visuo-spatial perspective  
55  
56 86 taking. While the environment is usually experienced from an egocentric *first-person perspective* (1PP),  
57  
58 87 humans can simulate someone else’s viewpoint, referred to as *third-person perspective* (3PP) taking.  
59  
60 88 3PP taking is an important component of social cognition which contributes to understand others and  
61  
62 89 anticipate their actions. There is evidence suggesting that 3PP taking is a strongly embodied process  
63  
64 90 involving egocentric mental imagery [15]. Here, we compared patients one week after right or left  
65  
66 91 vestibular neurectomy and healthy participants involved in a virtual ball-tossing game [16, 17] played  
67  
68 92 either from their own viewpoint (1PP), or from the viewpoint of a distant avatar – thus requiring 3PP  
69  
70 93 taking. Patients were tested in the acute stage of a unilateral vestibular deafferentation, when brain

94 metabolism disorganization is maximal [18], to document the impact of the vestibular tone imbalance  
95 on 3PP taking abilities. Performances were compared to those from 3D objects mental rotation, requiring  
96 object-based mental imagery, but no perspective-taking. Given evidence of a vestibular contribution to  
97 the perspectival experience [16, 19, 20], and overlapping neural mechanisms for vestibular information  
98 processing and perspective-taking [21], we hypothesized that unilateral vestibular deficits would impair  
99 3PP taking more than object-based mental imagery.

## 102 **Methods**

### 103 **Patients**

104 We tested 23 patients (13 women; mean age  $\pm$  SD:  $51 \pm 12$  years) before and 7 days after a unilateral  
105 vestibular neurectomy. Cognitive tests were performed 7 days after the surgery. All patients were right-  
106 handed (Edinburgh Handedness inventory:  $79 \pm 22\%$ ). Twenty-one patients were operated on for an  
107 acoustic neuroma (grade 2, 3 or 4 from Koos classification [22]) and two patients for Menière's disease  
108 (**Table 1**). Menière's disease patients had a pure unilateral vestibular pathology, with normal video head  
109 impulse test (vHIT) and hearing on their contralesional ear. Twelve patients received a left neurectomy  
110 and 11 a right neurectomy using the retrosigmoid approach ( $n = 20$ ) or the translabyrinthine approach  
111 ( $n = 3$ ). Depending on the size of the neuroma and the operating approach, the neurectomy aimed at  
112 removing the vestibular nerve totally ( $n = 19$ ) or partially ( $n = 4$ ). Before and 7 days after neurectomy,  
113 patients received standard otoneurological examinations including a pure tone audiometry (SCR  
114 Electroniques, France), a bithermal caloric test and a vHIT (Synapsys, France), as well as measurement  
115 of vestibulo-ocular responses on a rotating chair (Synapsys, France) and of the subjective visual vertical  
116 (Framiral, France). Patients were compared to 23 age- and gender-matched healthy volunteers (13  
117 women,  $52 \pm 12$  years). They were right-handed ( $86 \pm 25\%$ ) and had no history of otoneurological and  
118 psychiatric disease.

### 120 **Ethics statement**

121 All participants were informed about the study and gave their written informed consent. Experimental  
122 procedures were approved by the local Ethics Committee (CPP Sud-Méditerranée II, 2011-A01221-40)  
123 and followed the Declaration of Helsinki.

### 125 **Visual stimuli and tasks**

126 **3PP taking task.** Participants were involved in a virtual ball-tossing game [16]. Visual stimuli showed  
127 6 avatars around a circle (see **Figure 1A** and the legend, for details). Participants were instructed to  
128 adopt the perspective of the avatar holding the ball (identified by a blue T-shirt) and from this position  
129 to toss the ball to the avatar wearing a red T-shirt. They indicated as quickly and as accurately as possible

130 whether they had to toss the ball to the right or left, by pressing on a respond pad [16]. There was no  
131 time limit to answer. The experiment was a 5 *Angle of visuo-spatial perspective taking* (5 positions of  
132 the avatar holding the ball:  $\pm 60^\circ$ ,  $\pm 120^\circ$  and  $180^\circ$ )  $\times$  4 *positions of the avatar* receiving the ball ( $\pm 60^\circ$   
133 and  $\pm 120^\circ$  with respect to the avatar holding the ball) design, with four repetitions of each of the 20  
134 stimuli. Participants completed 80 trials divided into 4 blocks of 20 trials presented in a random order.

135  
136 **1PP task.** Five avatars were shown in the visual scene (**Figure 1B**) and the hands holding the ball in the  
137 front of the scene indicated that participants had to toss the virtual ball from their 1PP [16]. Participants  
138 indicated as quickly and accurately as possible whether they had to toss the ball to their right or left (to  
139 the avatar located at  $\pm 60^\circ$ ,  $\pm 120^\circ$ ) without time limit. Each of the four visual stimuli was repeated four  
140 times in one block of 16 trials presented in a random order.

141  
142 **Visual detection task.** A visual detection task [16] was used to control for potential attentional and visual  
143 differences between groups of participants (**Figure 1C**). In 10 trials, one avatar held a ball, while in 10  
144 other trials, no ball was shown. Participants were instructed to detect as quickly and as accurately as  
145 possible that there was a ball in the visual scene, without time limit. Participants completed one block  
146 of 20 trials presented in a random order.

147  
148 **3D objects mental rotation task.** Twenty out of 23 patients were involved in a 3D objects mental rotation  
149 task [9]. Pairs of identical or different 3D objects were presented side by side on a computer screen  
150 (**Figure 1D**). When identical, the second object was rotated with respect to the first object by  $20^\circ$ ,  $40^\circ$   
151 or  $80^\circ$ . We instructed participants to imagine a rotation of one of the objects to align it with the other  
152 and to indicate as quickly and as accurately as possible whether the objects were identical or different.  
153 There was no time limit to answer. There were two blocks of 36 pairs of objects, of which 24 were  
154 identical and 12 were different. Pairs of objects were presented randomly within each block, and the  
155 presentation of the blocks was counterbalanced.

## 156 157 **Experimental procedures**

158 Participants were tested while comfortably sitting on a chair with their head maintained by a forehead  
159 and chin rest. SuperLab 4.5 (Cedrus Corporation, USA) was used to present visual stimuli on a computer  
160 screen ( $64.4 \times 49.4$  cm) placed 50 cm in front of the participants. Thirteen patients and controls first  
161 completed the 1PP task and then the 3PP taking task, while the others started with the 3PP taking task.  
162 All participants finished with the visual detection task and the 3D objects mental rotation task.  
163 Participants took short breaks after each block and task. Before the experiment, they trained to each task  
164 with a random selection of 10 trials. **After the experiment, participants filled out a questionnaire about  
165 the strategies they used. They indicated whether it happened during the experiment they imagined**

166 rotating the entire visual scene with the avatars to solve the 3PP taking task. If it was the case, they were  
167 asked to rate how often they did it, using a 7-point Likert scale ranging from ‘rarely’ to ‘all the time’.

## 170 Data analysis

171 For each participant and task, we calculated the mean response times (RTs) and percentage of errors.  
172 Trials yielding incorrect answers were discarded from the analysis of the RTs and we excluded trials for  
173 which RTs exceeded 2 standard deviations of the participant’s grand average. As the Shapiro-Wilk test  
174 revealed not normally distributed RTs for most of the dependent variables, we used a log transformation  
175 and applied repeated-measures ANCOVA when this procedure led to normal distribution. Variables that  
176 remained non-normally distributed and the percentage of errors were analysed using non-parametric  
177 procedures (SPSS 25, IBM Corporation, USA). Differences were considered significant when  $P < 0.05$ .  
178 One outlier participant was excluded for the 3PP taking task (RTs > 16 s), as well as one participant for  
179 the 1PP and visual detection tasks, resulting in a total of 22 patients and 22 controls in the final analysis.

180 We used partial correlations to calculate the relation between 13 clinical parameters (Table 1)  
181 and performance in the cognitive tasks. To minimize the risk of false positives, we used Bonferroni-  
182 corrected  $\alpha$  level of  $P < 0.05/13$  for statistical significance ( $P < 0.0039$ ).

## 185 Results

### 186 3PP taking task

187 **Response times.** RTs were first analysed using a repeated-measures ANCOVA with Angle of 3PP taking  
188 as within-subject factor, Group and Gender as between-subject factors, and Age as covariate. Age was  
189 significantly related to RTs ( $F_{1,39} = 12.19, P < 0.005$ ), whereas Gender was not ( $F_{1,39} = 0.08, P = 0.77$ ).  
190 Importantly, we found a significant effect of the Group after controlling for the effect of Age ( $F_{1,39} =$   
191 4.40,  $P < 0.05$ ). **Figure 2A** shows significantly larger RTs for 3PP taking in patients than in controls.  
192 As expected, the analysis shows a significant effect of the Angle of 3PP taking ( $F_{3,107} = 2.78, P < 0.05$ ),  
193 with longer RTs for larger angles of perspective taking (Greenhouse-Geisser correction,  $\epsilon = 0.69$ ). There  
194 was no significant interaction between Angle and Group, or with the covariate. RTs were not  
195 significantly related to the grade of the neuroma (Kruskal-Wallis test,  $P > 0.05$ ) (**Figure 2B**).

196 In a second analysis, we explored how the side of neurectomy influenced 3PP taking. We found  
197 that patients with left neurectomy showed significantly larger RTs than their age and gender-matched  
198 controls ( $F_{1,17} = 6.81, P < 0.05$ ) (**Figure 2A**). There was no other significant main effect or interaction,  
199 but Age was significantly related to RTs ( $F_{1,17} = 8.04, P < 0.05$ ). By contrast, patients with right  
200 neurectomy did not differ from their controls ( $F_{1,17} = 0.04, P = 0.84$ ) (**Figure 2A**). The effect of Angle  
201 of 3PP taking was not significant after Greenhouse-Geisser correction ( $F_{2,38} = 2.53, P = 0.09$ ), but the



202 effect of Age was significant ( $F_{1,17} = 7.74, P < 0.05$ ). Direct comparison of patients' performance with  
203 left and right neurectomy did not reveal significant differences in RTs ( $F_{1,17} = 2.42, P = 0.14$ ). Thus, the  
204 effect of Group observed in the whole-population analysis seems to arise from altered 3PP taking mainly  
205 in patients with left neurectomy. We note that this result cannot be accounted for by different **vestibular**  
206 **functions** in patients with left and right neurectomy (see **Table 1**).

207  
208 **Errors.** Friedman's tests revealed an effect of the Angle of 3PP taking on the percentage of errors in  
209 patients ( $\chi^2_{(4)} = 25.7, P < 0.001$ ) and controls ( $\chi^2_{(4)} = 37.8, P < 0.001$ ) (**Figure 2C**). However, Mann-  
210 Whitney tests did not reveal a significant effect of the Group ( $-60^\circ: P = 0.13; -120^\circ: P = 0.16; 180^\circ: P$   
211  $= 0.33; +120^\circ: P = 0.44; +60^\circ: P = 0.08$ ).

212  
213 **Questionnaire about 3PP taking strategies.** 18/22 patients (82%) and 19/22 controls (86%) reported  
214 they never imagined rotating the entire visual scene to solve the task, whereas 4/22 patients (18%) and  
215 3/22 controls (14%) reported they "sometimes" imagined rotating the visual scene. No participant  
216 reported having used this strategy "all the time". Strategies were reported in similar proportion in the  
217 whole sample of patients and controls ( $\chi^2 = 0.17, P = 0.68$ ), and in similar proportion in patients with  
218 left and right vestibular neurectomy ( $\chi^2 = 1.22, P = 0.27$ ).

### 219 **First-person perspective task**

220  
221 RTs between the whole sample of patients and the controls did not differ (Mann-Whitney test,  $U = 244,$   
222  $P = 0.55$ ) (**Figure 2D**). Similar analyses for patients with left and right neurectomy revealed no  
223 differences between the groups (left:  $U = 22, P = 0.15$ ; right:  $U = 40, P = 0.48$ ). As the percentages of  
224 errors were very low, they were not analysed.

### 225 **Visual detection task**

226  
227 RTs were significantly longer in the whole sample of patients when compared to the controls (Mann-  
228 Whitney test,  $U = 287, P < 0.05$ ) (**Figure 2D**). RTs differed between the patients with left neurectomy  
229 and their controls ( $U = 113, P < 0.05$ ), but they were not different between the patients with right  
230 neurectomy and their controls ( $U = 40, P = 0.63$ ). As the percentages of errors were very low, they were  
231 not analysed.

### 232 **3D objects mental rotation**

233  
234 **Response times.** A repeated-measures ANCOVA revealed a significant main effect of Angular disparity  
235 (Greenhouse-Geisser correction,  $\epsilon = 0.67: F_{1,3,47} = 18.95, P < 0.0001$ ), with longer RTs for larger angles  
236 of mental rotation. In contrast with the 3PP taking task, there was no significant effect of the Group  
237 ( $F_{1,35} = 0.15, P = 0.71$ ) (**Figure 3A**). We found no significant effect of Age ( $F_{1,35} = 3.07, P = 0.09$ ) and



238 Gender ( $F_{1,35} = 0.47, P = 0.50$ ), but there was a significant interaction of Angular disparity  $\times$  Age ( $F_{1,3,47}$   
239  $= 3.01, P < 0.05$ ). Similar analyses conducted separately for patients with left and right neurectomy  
240 revealed no effect of the Group (left:  $F_{1,15} = 0.09, P = 0.78$ ; right:  $F_{1,15} = 0.03, P = 0.87$ ) (**Figure 3A**).

241  
242 **Errors.** Friedman's tests revealed a significant effect of Angular disparity in patients ( $\chi^2_{(2)} = 27.3, P <$   
243  $0.0001$ ) and in controls ( $\chi^2_{(2)} = 22.7, P < 0.0001$ ), with more errors for larger angles of mental rotation  
244 (**Figure 3B**). Mann-Whitney tests showed a statistical trend for higher percentage of errors in patients  
245 than in controls for  $20^\circ$  ( $U = 269, P = 0.06$ ), but not for  $40^\circ$  ( $U = 249, P = 0.19$ ) and  $80^\circ$  ( $U = 260, P =$   
246  $0.11$ ).

### 247 **Clinical data and their correlation with visuo-spatial cognition**

248 **Table 1** shows that overall the two groups of patients did not differ with respect to their clinical  
249 status, with the exception of worse postural performance in patients with right than left vestibular  
250 loss, and a larger proportion of cophosis at D+7 in the patients with left vestibular neurectomy. There  
251 was an important variability in the presurgical vestibular functions, owing to the different locations  
252 and grades of the neuroma [22]. Thus, we calculated partial correlations between clinical parameters  
253 and cognitive performance, while controlling for the effects of age (as RTs for 3PP taking were  
254 related to age). There was no significant correlation between clinical parameters and cognitive  
255 performance at a Bonferroni-corrected  $\alpha$  level. The only statistical trend was a positive correlation  
256 between the deficit at the caloric test measured at D-1 and the average RTs (log transform) for the  
257 3PP taking task ( $r = 0.596, P = 0.007$ ). No such trend was found for the correlation between the other  
258 cognitive tasks and clinical parameters recorded at D-1 and D+7.

## 260 **Discussion**

### 261 **Different effect of vestibular disorders on egocentric vs object-based mental imagery**

262 The present data show that patients tested one week after unilateral vestibular neurectomy had impair  
263 3PP taking abilities, as revealed by longer RTs in patients than in controls. Interestingly, no deficit was  
264 found when patients were required to toss the virtual ball from their 1PP, that is, when no own-body  
265 mental transformation was necessary. This indicates that deficits in 3PP taking were not merely related  
266 to attentional or oculomotor deficits to explore the visual scene and to discriminate left/ right orientations  
267 (**see below**). Rather, our results speak in favour of a more specific vestibular contribution to perspective  
268 taking and highlight the specific role of vestibular information in embodied spatial cognition [23].  
269 Another recent study, in which patients with objective balance deficits were required to imitate lateral  
270 body tilts of an avatar, showed longer movement latencies and more errors in the patients [7]. Our results

274 are also in line with a previous study in healthy participants using the same 3PP taking task, showing  
275 that natural vestibular stimulation (on a rotating chair) helped to adopt the perspective of a distant avatar  
276 when the directions of the physical rotation and simulated viewpoint were congruent [16].

277 Although our participants were trained to use explicitly own-body mental imagery during the  
278 3PP taking task, we cannot exclude that some of them used other strategies, such as simulating a ‘bird  
279 perspective’ of the visual scene or mixed egocentric/allocentric strategies depending on the avatar’s  
280 position. Debriefing questionnaires confirmed that most participants complied to the instructions and  
281 used mainly an egocentric strategy. Importantly, as mixed egocentric/allocentric strategies were reported  
282 in similar proportions in the patients and the controls, strategies cannot account for the different 3PP  
283 taking abilities between the groups.

284 In contrast with the 3PP taking task, the unilateral vestibular neurectomy had no effect on 3D  
285 object mental imagery. There are only few studies about the consequences of vestibular disorders on  
286 object-based mental imagery and they showed divergent results. Impaired object-based mental imagery  
287 seems to depend on the extent of the vestibular loss, as patients with bilateral loss were the most impaired  
288 [8, 9], and on the compensation stage, as deficits were found only in the early stage after vestibular  
289 neurectomy [9, 10].

290 Altogether, our data suggest that acute unilateral vestibular deficits affect differently visuo-  
291 spatial tasks involving mostly egocentric mental imagery (3PP taking) and object-based mental imagery.  
292 We propose that shared neural networks are involved in encoding own-body rotations (vestibular cortical  
293 network) and in simulating own-body and viewpoint rotations (3PP taking). This proposition is first  
294 supported by our previous results in healthy participants showing that whole-body rotations on a rotating  
295 chair influenced 3PP taking in a direction-specific way, but not object-based mental imagery [16].  
296 Second, a recent fMRI study [21] showed that clockwise/counterclockwise mental rotation of the  
297 observer’s viewpoint (as used in our 3PP taking task) and galvanic vestibular stimulation both activated  
298 the bilateral posterior insular cortex and the right parietal operculum (OP2), two core areas of the  
299 vestibular cortical network.

### 300 **Different effect of right vs left vestibular disorders on perspective taking**

301 A main result of the present study is that 3PP taking abilities depended on the side of vestibular  
302 deafferentation. Only patients with left neurectomy showed larger RTs in the 3PP taking task, when  
303 compared to their controls. This was not the case when visuo-spatial judgements were performed from  
304 their 1PP. However, patients with left neurectomy were also slower for the visual detection task. This  
305 indicates that left vestibular loss is more likely to disturb the mechanisms of egocentric own-body mental  
306 imagery and alertness than the mechanisms of visuo-spatial tasks that do not require egocentric mental  
307 imagery (i.e., 1PP task and 3D object mental rotation). It is difficult to compare our data with previous  
308 studies as only few of them have investigated spatial cognition with respect to the side of the vestibular  
309 loss. In line with our findings, Saj *et al.* [13] found that only left vestibular neurectomy significantly  
310

311 deviated the perceived straight ahead. Left cathodal galvanic vestibular stimulation also impaired mental  
1 312 imagery in healthy participants [24], while only left-warm caloric vestibular stimulation distorted their  
2 313 body schema [25]. However, opposite data have been reported, with slower recovery of the subjective  
3 314 visual vertical in right vestibular neuritis [26], and a **trend** for poorer spatial navigation in right vestibular  
4 315 deficits [14]. **Another study reported no effect of the side of the vestibular loss on cognitive deficits [4].**  
5 316 The discrepancy between these data may be related to the diversity of the spatial functions tested,  
6 317 involving different brain areas.

11 318 Why would left vestibular neurectomy impair 3PP taking? Neuroimaging studies have  
12 319 emphasized a right hemispheric dominance of vestibular processing in right-handed participants [11] –  
13 320 as were our participants. Accordingly, a stronger impact of right vestibular neurectomy on 3PP taking  
14 321 can be expected. Our results do not support this hypothesis. However, they are in line with neuroimaging  
15 322 data showing different compensation patterns in acute right and left vestibular neuritis, with a shift of  
16 323 the dominant ascending input from the ipsilateral to the contralateral pathways [18]. Alternatively, based  
17 324 on results from a diffusion tensor imaging study of the vestibulo-thalamo-cortical pathways [12], we  
18 325 predict that right vestibular neurectomy would predominantly disrupt multisensory processing in the  
19 326 right parieto-insular cortex, whereas left neurectomy is more likely to disrupt multisensory processing  
20 327 *bilaterally* in this area. We propose that left vestibular neurectomy evoked deficits in 3PP taking by  
21 328 disrupting bilateral areas underpinning 3PP taking. Of note, an fMRI study [17] described the neural  
22 329 correlates of 3PP taking in a virtual ball-tossing game similar to that used in the present study. 3PP  
23 330 taking, when compared to 1PP, activated bilaterally the anterior insula and medial frontal cortices, as  
24 331 well as the left inferior parietal lobe. As 3PP taking involves a bilateral network of insular and parieto-  
25 332 frontal areas, we anticipate that it is very likely more affected by predominantly bilateral vestibulo-  
26 333 thalamo-cortical projections from the left vestibular receptors [12].

### 38 334 **Role of visuo-spatial attention and alertness**

40 335 **Patients had longer RTs than controls in the visual detection task, and RTs for the 3PP taking task and**  
41 336 **the visual detection task were larger after left than right vestibular neurectomy. As previous studies have**  
42 337 **related vestibular loss to poorer visuo-spatial ability and attention [27], we cannot exclude that larger**  
43 338 **attentional deficits in patients with left neurectomy contributed to larger RTs for 3PP taking. However,**  
44 339 **as RTs were not larger for the 1PP task and the 3D-object mental imagery task in these patients, we can**  
45 340 **rule out a general effect of attentional deficits in all tasks after left neurectomy. We interpret increased**  
46 341 **RTs in the visual detection task as resulting from deficits of alertness (because RTs were short, around**  
47 342 **750 ms, whereas RTs were around 3000 ms for 3PP taking). Visual detection assessed more alertness**  
48 343 **(reaction times) than high-level visuo-spatial cognition. Our results are consistent with data showing**  
49 344 **that patients with unilateral vestibular loss performed worse than healthy controls in a processing speed**  
50 345 **task [4].**

## 348 **Relation between clinical and behavioural data**

349 We only found a statistical trend for slower 3PP taking in patients with larger deficits at the presurgical  
350 caloric test. This relation suggests that information from the lateral semicircular canals, or the normal  
351 functioning of the structures processing this information, is involved in own-body mental rotation in the  
352 yaw plane. The caloric test delivers low-frequency stimulation of the canals and assesses vestibular  
353 pathways that are likely relevant for the cognitive functions tested here. Interestingly, stimulation of the  
354 lateral canals by whole-body rotation influenced RTs for 3PP taking in a direction-dependent manner  
355 [16] and caloric vestibular stimulation improved own-body mental imagery in healthy participants [28].  
356 Our data are congruent with observations in patients with unilateral vestibular disorders showing that  
357 lower caloric response correlated with deficits in processing speed, memory and executive function [4].  
358 Finally, we found no correlation between hearing loss and performance in the cognitive tasks, congruent  
359 with recent findings showing similar performance in deaf signers and hearing nonsigners engaged in  
360 visuo-spatial perspective-taking, 3D object mental rotation and spatial orientation tasks [29].

## 362 **Limitations of the study and perspectives**

363 A limitation of the present study is the absence of detailed neuropsychological evaluation of the  
364 participants, which could have accounted for some variability in our data. However, the mean age in our  
365 sample was not high, controls were selected to carefully match the patients, and our statistical analyses  
366 controlled for the effect of age.

367 The present study documents deficits of spatial cognition only in the acute stage of a unilateral  
368 vestibular neurectomy. This was motivated by the need to better understand the domains of cognition  
369 affected and spared when a vestibular tone imbalance is present, i.e. when compensatory mechanisms  
370 are not fully expressed. There is evidence of brain disorganization in the acute stage of a unilateral  
371 vestibular loss, both in humans [18] and animal [30] models. It is unknown if the deficits in 3PP taking  
372 reported here vanish when patients have compensated from their unilateral vestibular disorder. Future  
373 studies should endeavor to explore the postlesional recovery of spatial cognition deficits after unilateral  
374 vestibular deafferentation and compare its time course to that of functional recovery.

375 As in most studies of this kind, a limitation of the present study is the sample size, which may  
376 explain some discrepancies in the literature regarding the effects of left and right vestibular loss on  
377 cognition [4, 13, 14, 26], in addition to the variety of tasks and procedures used. Studies remain to be  
378 conducted in large samples of patients to determine the effects of left vs right vestibular loss on a large  
379 range of spatial cognitive functions.

## 381 **Conclusions**

382 The present study shows that acute unilateral vestibular deafferentation impairs embodied spatial  
383 cognition by disrupting visuo-spatial tasks involving mostly egocentric mental imagery (3PP taking),  
384 but not object-based mental imagery. These effects were observed only in patients with left vestibular

385 loss. We believe that the present investigation helps understand the cognitive complaints of patients with  
1 386 vestibular disorders. Finally, our data brings further evidence that vestibular signals contribute to several  
2 387 sensorimotor bases of social cognition, including 3PP taking, and strengthens the connections between  
3 388 the so far distinct fields of social neuroscience and human vestibular physiology (for a perspective, see  
4 389 [23]).  
5 390  
6 391

## 11 392 **Acknowledgements**

13  
14 393 The research leading to these results has received funding from the People Programme (Marie Curie  
15 394 Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under REA grant  
16 395 agreement number 333607 ('BODILYSELF, vestibular and multisensory investigations of bodilyself-  
17 396 consciousness').  
18 397  
19 398

## 23 399 **Compliance with ethical standards**

24 400 **Conflicts of interest.** C. Lopez serves as an associate editor of *Frontiers in Integrative Neuroscience*  
25 401 and guest editor of *Frontiers in Neurology*.  
26 402  
27 403

28 404 **Ethical standard.** Ethical approvals have been obtained from the local ethics committees (Comité de  
29 405 Protection des Personnes Sud-Méditerranée II, 2011-A01221-40) and participants provided informed  
30 406 consent in accordance with the Declaration of Helsinki.  
31 407  
32 408  
33 409  
34 410  
35 411  
36 412  
37 413  
38 414  
39 415  
40 416  
41 417  
42 418  
43 419  
44 420  
45 421  
46 422  
47 423  
48 424  
49 425  
50 426  
51 427  
52 428  
53 429  
54 430  
55 431  
56 432  
57 433  
58 434  
59 435  
60 436  
61 437  
62 438  
63 439  
64 440  
65 441

## Figure legends

**Figure 1.** Visual stimuli and tasks. (A) Example of a visual scene for the 3PP taking task showing six avatars located around a circle (adapted from Ref. [16]). One avatar was facing the participants ( $180^\circ$ ), one was seen from the back ( $0^\circ$ ), two were on their right ( $60^\circ$  and  $120^\circ$ ), and two on their left ( $-60^\circ$  and  $-120^\circ$ ). From one trial to another, the colour of the avatars' T-shirt could change (white, red or blue). Participants were instructed to simulate the visuo-spatial perspective of the avatar holding the ball (identified by a blue T-shirt) and from this position to toss the ball to the avatar wearing a red T-shirt. The avatar with the blue T-shirt was located either at  $\pm 60^\circ$ ,  $\pm 120^\circ$  or  $180^\circ$  (but never at  $0^\circ$ , to avoid confusion with the 1PP task), resulting in a total of 5 angles of 3PP taking. The avatar wearing the red T-shirt was located  $60^\circ$  or  $120^\circ$  to the right or left of this avatar. Participants pressed a key on a response pad (RB-830, Cedrus Corporation, San Pedro, USA) with their right middle finger to toss the ball to the right, or their right index finger to toss the ball to the left. A fixation cross was presented during 1 s before each visual scene. In the example presented, the correct response is to toss the ball to the left. (B) Example of a visual scene for the 1PP task showing five avatars located around a circle. One avatar was facing the participants ( $180^\circ$ ), two were on their right ( $60^\circ$  and  $120^\circ$ ), and two on their left ( $-60^\circ$  and  $-120^\circ$ ). Participants were instructed to toss the ball to the avatar wearing a red T-shirt, which was located at  $\pm 60^\circ$  or  $\pm 120^\circ$  from their viewpoint (4 visual stimuli). Participants pressed a key with their right middle finger to toss the ball to the right, or their right index finger to toss the ball to the left. A fixation cross was presented during 1 s before each visual scene. The correct response here is to toss the ball to the right. (C) Example of a visual scene for the visual detection task. Visual stimuli were the same as for the 3PP taking task, except that all avatars wore a white T-shirt. Thirteen participants had to press a button with their right middle finger as soon as they detected a ball, or with their right index finger when they detected there was no ball, and vice versa for the other participants. A fixation cross was presented during 1 s before each visual scene. The correct response here is "yes", as there is a ball in the scene. (D) Example of a visual stimulus for the 3D objects mental rotation. Half of the participants pressed a button with their right middle finger if objects were identical, or with their right index finger if they were different, and vice versa for the others. The correct response here is "no", as the objects were different.

**Figure 2.** Influence of unilateral vestibular deafferentation on visuo-spatial perspective taking. (A) Mean response times for the 3PP taking task are plotted separately for the whole population of patients (left part), for the patients with left vestibular neurectomy (central part) and the patients with right vestibular neurectomy (right part) together with their age- and gender-matched control participants. Error bars represent the standard error of mean. \* indicates statistical significance between the patients and their controls ( $P < 0.05$ ). (B) Mean response times for the 3PP taking task are illustrated for the patients as a function of the vestibular pathology. (C) Box-and-whisker plots illustrate the percentage of

456 errors in the whole sample of participants. The top and bottom ends of the whisker represent the 95<sup>th</sup> and  
1 457 5<sup>th</sup> percentiles of the distribution and the horizontal line inside the box represents the median. (D)  
2  
3 458 Response times for the 1PP task and the visual detection task are plotted for the whole sample of patients.  
4  
5 459 Same conventions as for Parts C.

6 460  
7  
8 461 **Figure 3.** Influence of unilateral vestibular deafferentation on 3D objects mental rotation. (A) Mean  
9  
10 462 response times for the mental rotation of 3D objects are plotted separately for the whole population of  
11 463 patients (left part), for the patients with left vestibular neurectomy (central part) and the patients with  
12  
13 464 right vestibular neurectomy (right part) together with their age- and gender-matched control participants.  
14  
15 465 Error bars represent the standard error of mean. (B) Box-and-whisker plots illustrate the percentage of  
16 466 errors in the whole sample of participants. The top and bottom ends of the whisker represent the 95<sup>th</sup> and  
17  
18 467 5<sup>th</sup> percentiles of the distribution and the horizontal line inside the box represents the median.

19 468  
20  
21 469  
22  
23 470  
24  
25 471  
26  
27 472  
28  
29 473  
30  
31 474  
32  
33 475  
34  
35 476  
36  
37 477  
38  
39 478  
40  
41 479  
42  
43 480  
44  
45 481  
46  
47 482  
48  
49 483  
50  
51 484  
52  
53 485  
54  
55 486  
56  
57 487  
58  
59 488  
60  
61  
62  
63  
64  
65




## References

1. Smith PF (2017) The vestibular system and cognition. *Curr Opin Neurol* 30:84–89. <https://doi.org/10.1097/WCO.0000000000000403>
2. Bigelow RT, Agrawal Y (2015) Vestibular involvement in cognition: Visuospatial ability, attention, executive function, and memory. *J Vestib Res Equilib Orientat* 25:73–89. <https://doi.org/10.3233/VES-150544>
3. Bigelow RT, Semenov YR, du Lac S, et al (2016) Vestibular vertigo and comorbid cognitive and psychiatric impairment: the 2008 National Health Interview Survey. *J Neurol Neurosurg Psychiatry* 87:367–372. <https://doi.org/10.1136/jnnp-2015-310319>
4. Popp P, Wulff M, Finke K, et al (2017) Cognitive deficits in patients with a chronic vestibular failure. *J Neurol* 264:554–563. <https://doi.org/10.1007/s00415-016-8386-7>
5. Péruch P, Borel L, Gaunet F, et al (1999) Spatial performance of unilateral vestibular defective patients in nonvisual versus visual navigation. *J Vestib Res* 9:37–47
6. Brandt T, Schautzer F, Hamilton DA, et al (2005) Vestibular loss causes hippocampal atrophy and impaired spatial memory in humans. *Brain* 128:2732–41
7. Allum JHJ, Langewitz W, Sleptsova M, et al (2017) Mental body transformation deficits in patients with chronic balance disorders. *J Vestib Res Equilib Orientat* 27:113–125. <https://doi.org/10.3233/VES-170613>
8. Grabherr L, Cuffel C, Guyot JP, Mast FW (2011) Mental transformation abilities in patients with unilateral and bilateral vestibular loss. *Exp Brain Res* 209:205–14. <https://doi.org/10.1007/s00221-011-2535-0>
9. Péruch P, Lopez C, Redon C, et al (2011) Vestibular information is necessary for maintaining metric properties of representational space: Evidence from mental imagery. *Neuropsychologia* 49:3136–3144
10. Candidi M, Micarelli A, Viziano A, et al (2013) Impaired mental rotation in benign paroxysmal positional vertigo and acute vestibular neuritis. *Front Hum Neurosci* 7:783. <https://doi.org/10.3389/fnhum.2013.00783>
11. Dieterich M, Bense S, Lutz S, et al (2003) Dominance for vestibular cortical function in the non-dominant hemisphere. *Cereb Cortex* 13:994–1007
12. Dieterich M, Kirsch V, Brandt T (2017) Right-sided dominance of the bilateral vestibular system in the upper brainstem and thalamus. *J Neurol* 264:55–62. <https://doi.org/10.1007/s00415-017-8453-8>
13. Saj A, Honoré J, Bernard-Demanze L, et al (2013) Where is straight ahead to a patient with


- 523 unilateral vestibular loss? Cortex 49:1219–1228.  
1  
2 524 <https://doi.org/10.1016/j.cortex.2012.05.019>
- 3 525 14. Hüfner K, Hamilton DA, Kalla R, et al (2007) Spatial memory and hippocampal volume  
4  
5 526 in humans with unilateral vestibular deafferentation. *Hippocampus* 17:471–85.  
6  
7 527 <https://doi.org/10.1002/hipo.20283>
- 8  
9 528 15. Kessler K, Thomson LA (2010) The embodied nature of spatial perspective taking:  
10  
11 529 embodied transformation versus sensorimotor interference. *Cognition* 114:72–88.  
12  
13 530 <https://doi.org/10.1016/j.cognition.2009.08.015>
- 14 531 16. Deroualle D, Borel L, Devèze A, Lopez C (2015) Changing perspective: The role of  
15  
16 532 vestibular signals. *Neuropsychologia* 79, Part B:175–185.  
17  
18 533 <https://doi.org/10.1016/j.neuropsychologia.2015.08.022>
- 19  
20 534 17. David N, Bewernick BH, Cohen MX, et al (2006) Neural representations of self versus  
21  
22 535 other: visual-spatial perspective taking and agency in a virtual ball-tossing game. *J Cogn*  
23  
24 536 *Neurosci* 18:898–910
- 25 537 18. Becker-Bense S, Dieterich M, Buchholz H-G, et al (2014) The differential effects of acute  
26  
27 538 right- vs. left-sided vestibular failure on brain metabolism. *Brain Struct Funct* 219:1355–  
28  
29 539 1367. <https://doi.org/10.1007/s00429-013-0573-z>
- 30  
31 540 19. Pavlidou A, Ferrè ER, Lopez C (2018) Vestibular stimulation makes people more  
32  
33 541 egocentric. *Cortex J Devoted Study Nerv Syst Behav* 101:302–305.  
34  
35 542 <https://doi.org/10.1016/j.cortex.2017.12.005>
- 36 543 20. Lopez C, Elzière M (2018) Out-of-body experience in vestibular disorders - A prospective  
37  
38 544 study of 210 patients with dizziness. *Cortex J Devoted Study Nerv Syst Behav* 104:193–  
39  
40 545 206. <https://doi.org/10.1016/j.cortex.2017.05.026>
- 41  
42 546 21. Macaуда G, Moisa M, Mast FW, et al (2019) Shared neural mechanisms between imagined  
43  
44 547 and perceived egocentric motion - A combined GVS and fMRI study. *Cortex J Devoted*  
45  
46 548 *Study Nerv Syst Behav* 119:20–32. <https://doi.org/10.1016/j.cortex.2019.04.004>
- 47 549 22. Koos WT, Day JD, Matula C, Levy DI (1998) Neurotopographic considerations in the  
48  
49 550 microsurgical treatment of small acoustic neurinomas. *J Neurosurg* 88:506–512.  
50  
51 551 <https://doi.org/10.3171/jns.1998.88.3.0506>
- 52  
53 552 23. Deroualle D, Lopez C (2014) Toward a vestibular contribution to social cognition. *Front*  
54  
55 553 *Integr Neurosci* 8:16. <https://doi.org/10.3389/fnint.2014.00016>
- 56 554 24. Lenggenhager B, Lopez C, Blanke O (2008) Influence of galvanic vestibular stimulation  
57  
58 555 on egocentric and object-based mental transformations. *Exp Brain Res* 184:211–21  
59
- 60 556 25. Lopez C, Nakul E, Preuss N, et al (2018) Distorted own-body representations in patients  
61  
62  
63  
64  
65

- 557 with dizziness and during caloric vestibular stimulation. *J Neurol* 265:86–94.  
1 558 <https://doi.org/10.1007/s00415-018-8906-8>
- 3 559 26. Toupet M, Van Neechel C, Bozorg Grayeli A (2014) Influence of body laterality on recovery  
4 560 from subjective visual vertical tilt after vestibular neuritis. *Audiol Neurootol* 19:248–255.  
5 561 <https://doi.org/10.1159/000360266>
- 7 562 27. Bigelow RT, Semenov YR, Trevino C, et al (2015) Association between visuospatial  
8 563 ability and vestibular function in the Baltimore longitudinal study of aging. *J Am Geriatr*  
9 564 *Soc* 63:1837–1844. <https://doi.org/10.1111/jgs.13609>
- 11 565 28. Falconer CJ, Mast FW (2012) Balancing the mind: vestibular induced facilitation of  
12 566 egocentric mental transformations. *Exp Psychol* 59:332–339.  
13 567 <https://doi.org/10.1027/1618-3169/a000161>
- 15 568 29. Secora K, Emmorey K (2019) Social abilities and visual-spatial perspective-taking skill:  
16 569 deaf signers and hearing Nnonsigners. *J Deaf Stud Deaf Educ.*  
17 570 <https://doi.org/10.1093/deafed/enz006>
- 19 571 30. Stackman RW, Clark AS, Taube JS (2002) Hippocampal spatial representations require  
20 572 vestibular input. *Hippocampus* 12:291–303. <https://doi.org/10.1002/hipo.1112>
- 22 573


**A** Virtual ball-tossing game: 3PP




**B** Virtual ball-tossing game: 1PP




**C** Visual detection task: 1PP



**D** 3D objects mental rotation task








Table1

	Age (years)	Gender	Diagnosis	Examination at D-1				Examination at D+7			
				CVS (% deficit)	VOR (gain)	VHIT (gain) <sup>a</sup>	SVV (°) <sup>b</sup>	CVS (% deficit)	VOR (gain)	VHIT (gain) <sup>a</sup>	SVV (°) <sup>b</sup>
Left neurectomy	49.9 ± 11.6	7 women 5 men	4 grade 2 6 grade 3 2 grade 4	66.2 ± 39.8	0.36 ± 0.14	0.45 ± 0.36	-2.3 ± 2.4	95.6 ± 11.7	0.43 ± 0.21	0.38 ± 0.41	-7.1 ± 4.0
Right neurectomy	53.0 ± 14.0	6 women 5 men	3 grade 2 3 grade 3 3 grade 4 2 MD	41.9 ± 34.3	0.44 ± 0.22	0.69 ± 0.36	0.1 ± 1.0	80.0 ± 25.2	0.41 ± 0.22	0.25 ± 0.28	-6.2 ± 4.2
	<i>P</i> = 0.57 <sup>c</sup>	<i>P</i> = 0.86 <sup>d</sup>	<i>P</i> = 0.63 <sup>e</sup>	<i>P</i> = 0.19 <sup>c</sup>	<i>P</i> = 0.35 <sup>c</sup>	<i>P</i> = 0.18 <sup>c</sup>	<i>P</i> = 0.01 <sup>c</sup>	<i>P</i> = 0.16 <sup>c</sup>	<i>P</i> = 0.90 <sup>c</sup>	<i>P</i> = 0.56 <sup>c</sup>	<i>P</i> = 0.63 <sup>c</sup>

**Table 1.** Comparison of clinical data in patients with left and right vestibular neurectomy. Means ± standard deviations are reported before (D-1) and after (D+7) surgery for each group of patients. MD: Menière's disease, CVS: caloric vestibular stimulation, VOR: vestibulo-ocular reflex, VHIT: video head impulse test, SVV: subjective visual vertical.

<sup>a</sup> Gain for the lateral semicircular canal on the side of the disease

<sup>b</sup> Negative signs indicate that the subjective visual vertical was tilted to the side of the disease (ipsilesional tilt)

<sup>c</sup> Independent-samples t-tests

<sup>d</sup> Chi-square

<sup>e</sup> Chi-square for the grade of neuroma



		Left neurectomy	Right neurectomy	Statistics
<b>Age (years)</b>		49.9 ± 11.6	53.0 ± 14.0	<i>P</i> = 0.57 <sup>a</sup>
<b>Gender</b>		7 women, 5 men	6 women, 5 men	<i>P</i> = 0.86 <sup>b</sup>
<b>Diagnosis and acoustic neuroma grade (Koos classification)</b>		4 grade 2 6 grade 3 2 grade 4	3 grade 2 3 grade 3 3 grade 4 2 Menière's disease	<i>P</i> = 0.63 <sup>c</sup>
<b>Unilateral vestibular neurectomy</b>		2 partial, 10 total	2 partial, 9 total	<i>P</i> = 0.92 <sup>b</sup>
<b>Examination at D-1</b>	CVS (% deficit)	66.2 ± 37.8	39.9 ± 31.1	<i>P</i> = 0.10 <sup>a</sup>
	VOR (gain)	0.38 ± 0.14	0.44 ± 0.22	<i>P</i> = 0.52 <sup>a</sup>
	vHIT ipsi. (gain)			
	Anterior canal	0.81 ± 0.15	0.88 ± 0.24	<i>P</i> = 0.49 <sup>a</sup>
	Lateral canal	0.45 ± 0.36	0.69 ± 0.36	<i>P</i> = 0.18 <sup>a</sup>
	Posterior canal	0.59 ± 0.37	0.63 ± 0.36	<i>P</i> = 0.84 <sup>a</sup>
	vHIT contra. (gain)			
	Anterior canal	0.95 ± 0.09	0.93 ± 0.07	<i>P</i> = 0.50 <sup>a</sup>
	Lateral canal	0.88 ± 0.10	0.90 ± 0.11	<i>P</i> = 0.56 <sup>a</sup>
	Posterior canal	0.73 ± 0.32	0.90 ± 0.11	<i>P</i> = 0.16 <sup>a</sup>
	Posture (mm/s) <sup>d</sup>			
	Eyes open	7.2 ± 3.3	12.3 ± 5.5	<b><i>P</i> = 0.015 <sup>a</sup></b>
	Eyes closed	9.2 ± 4.0	15.4 ± 11.0	<i>P</i> = 0.09 <sup>a</sup>
	SVV (°) <sup>e</sup>	-1.9 ± 2.6	-0.5 ± 1.9	<i>P</i> = 0.15 <sup>a</sup>
Pure tone average (dB)				
Ipsi.	1 Cop. (58.7 ± 26.0) <sup>f</sup>	2 Cop. (56.2 ± 35.6) <sup>f</sup>	<i>P</i> = 0.87 <sup>a</sup>	
Contra.	19.6 ± 9.0	18.9 ± 14.4	<i>P</i> = 0.88 <sup>a</sup>	
<b>Examination at D+7</b>	CVS (% deficit)	96.4 ± 9.8	80.0 ± 25.2	<i>P</i> = 0.14 <sup>a</sup>
	VOR (gain)	0.42 ± 0.20	0.41 ± 0.20	<i>P</i> = 0.89 <sup>a</sup>
	vHIT ipsi. (gain)			
	Anterior canal	0.57 ± 0.31	0.60 ± 0.26	<i>P</i> = 0.87 <sup>a</sup>
	Lateral canal	0.41 ± 0.37	0.26 ± 0.27	<i>P</i> = 0.46 <sup>a</sup>
	Posterior canal	0.60 ± 0.42	0.52 ± 0.26	<i>P</i> = 0.74 <sup>a</sup>
	vHIT contra. (gain)			
	Anterior canal	0.99 ± 0.03	0.88 ± 0.13	<i>P</i> = 0.10 <sup>a</sup>
	Lateral canal	0.89 ± 0.06	0.88 ± 0.11	<i>P</i> = 0.86 <sup>a</sup>
	Posterior canal	0.93 ± 0.11	0.84 ± 0.15	<i>P</i> = 0.31 <sup>a</sup>
	Posture (mm/s) <sup>d</sup>			
	Eyes open	7.4 ± 3.7	13.8 ± 12.5	<i>P</i> = 0.11 <sup>a</sup>
	Eyes closed	11.2 ± 6.6	19.9 ± 11.6	<b><i>P</i> = 0.038 <sup>a</sup></b>
	SVV (°) <sup>e</sup>	-7.1 ± 4.0	-6.2 ± 4.2	<i>P</i> = 0.63 <sup>a</sup>
Pure tone average (dB)				
Ipsi.	10 Cop. (12.5) <sup>f</sup>	5 Cop. (49.0 ± 21.3) <sup>f</sup>	<b><i>P</i> = 0.038 <sup>g</sup></b>	
Contra.	17.8 ± 9.6	19.9 ± 13.8	<i>P</i> = 0.69 <sup>a</sup>	
Rotational vertigo	4 positive, 6 negative	1 positive, 7 negative	<i>P</i> = 0.20 <sup>b</sup>	
Spont. nystagmus (°/s)				
Horizontal	4.5 ± 5.2	5.5 ± 7.5	<i>P</i> = 0.75 <sup>a</sup>	
Vertical	0.7 ± 0.7	1.7 ± 2.8	<i>P</i> = 0.30 <sup>a</sup>	
Body tilt	5 positive, 5 negative	3 positive, 5 negative	<i>P</i> = 0.60 <sup>b</sup>	
Past pointing	4 positive, 6 negative	3 positive, 5 negative	<i>P</i> = 0.91 <sup>b</sup>	

**Table 1.** Comparison of clinical data in patients with left and right vestibular neurectomy. Means ± standard deviations are reported before (D-1) and after (D+7) surgery for each group of patients. CVS: caloric vestibular stimulation, VOR:

<sup>a</sup> Independent-samples t-tests

<sup>b</sup> Chi-square

<sup>c</sup> Chi-square for the grade of neuroma

<sup>d</sup> Average velocity of the center of foot pressure measured on a posturography platform with a stable support surface

<sup>e</sup> Negatives signs indicate an ipsilesional tilt of the subjective visual vertical

<sup>f</sup> The number of patients with copiosis (Cop.) is indicated; the mean hearing loss in the other patients without copiosis is indicated in brackets

<sup>g</sup> Chi-square for the number of patients with copiosis

---

vestibulo-ocular reflex, VHIT: video head impulse test, SVV: subjective visual vertical, ipsi.: ipsilesional ear, contra.: contralesional ear, Cop.: cophosis.