



HAL
open science

Litter of mediterranean species as a source of volatile organic compounds

Justine Viros, Catherine Fernandez, Henri Wortham, Jordane Gavinet, Caroline Lecareux, Elena Ormeño

► **To cite this version:**

Justine Viros, Catherine Fernandez, Henri Wortham, Jordane Gavinet, Caroline Lecareux, et al.. Litter of mediterranean species as a source of volatile organic compounds. *Atmospheric Environment*, 2020, 242, pp.117815. <10.1016/j.atmosenv.2020.117815>. <hal-02937601>

HAL Id: hal-02937601

<https://amu.hal.science/hal-02937601v1>

Submitted on 14 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Litter of Mediterranean species as a source of volatile organic compounds**

2 Viros J^{1*}., Fernandez C¹., Wortham H²., Gavinet J¹., Lecareux C¹., Ormeño E¹.

3 ¹CNRS, Aix Marseille Univ, Avignon Univ, IRD, IMBE, Marseille, France.

4 ²Aix Marseille Univ., CNRS, LCE, Laboratoire de Chimie de l'Environnement, Marseille, France

5

6 *Viros Justine corresponding author: justine.viros@imbe.fr ; +33 4 13 55 11 62


7 Aix-Marseille Université – Bâtiment Sciences Naturelles – 3 place Victor Hugo 13003 Marseille

8 Abstract

9 Leaf litter, at the interface between the soil and the atmosphere, releases Biogenic Volatile
10 Organic Compounds (BVOC) which play an important role both in global atmospheric chemistry
11 and plant ecology. Litter is especially important in the Mediterranean region, where it is produced
12 in high amounts and decomposes slowly. Only leaf litter emissions of two dominant species (*Pinus*
13 *halepensis* and *Eucalyptus* spp.) present in the Mediterranean region have been studied so far. The
14 aim of this study is to characterize (quantitatively and qualitatively) the BVOC emissions (C10-
15 C15) from litter (freshly fallen and non-decomposed leaves) under laboratory conditions. Sixteen
16 species frequently found in the Mediterranean region were considered, nine of which possess
17 terpene storage structures. Results show a large diversity of BVOC (87 compounds detected,
18 terpenes, alkanes, alkenes, aldehydes, ketones, benzenoids), with terpene emission being higher
19 than non-terpenic emission in most of the cases. Species were classified within 3 categories:
20 negligible emitters (e.g. *Acer monspessulanum* and *Quercus ilex*, $< 0.10 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$), low emitters
21 (*Pinus pinea*, *Quercus pubescens*, $0.1 - 1.0 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) and moderate emitters (*Thymus vulgaris*,
22 *Cotinus coggygria* and *Pinus halepensis* between 1.5 to $4.7 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$). Litter that possessed
23 terpene storage structures and known to store high terpene concentration did not always release the
24 highest terpene emission rates (e.g. *Eucalyptus globulus*, *Rosmarinus officinalis*). Species that do
25 not possess such structures release only non terpenic emissions (e.g. alkanes, aldehydes).
26 Estimation of the potential contribution of *P. halepensis* and *Q. pubescens* litter to O_3 and
27 secondary organic aerosol (SOA) formation in the atmosphere showed that *P. halepensis* has a
28 larger potential impact on the global air quality than *Q. pubescens*. This study will contribute to
29 build future BVOC emissions inventories on leaf litter for their further integration in atmospheric
30 chemistry models.

31 **Keywords:** BVOC emissions, terpenes, storage structure, air quality, plant functional types

32 **Graphical abstract**



33 **Highlights**

- 34
- Litter emission rate ranges from < 0.1 to $4.7 \mu\text{g}\cdot\text{gDM}^{-1}\cdot\text{h}^{-1}$
- 35
- The highest emitters are *Thymus vulgaris*, *Cotinus coggygria* and *Pinus halepensis*
- 36
- Terpene emission rates are most often higher than non-terpenic emissions
- 37
- Litter BVOC emissions of a species are not necessarily lower than green leaf BVOC
- 38
- Litter BVOC emissions contribute to air quality in a non-negligible way

39

Introduction

40 Living plant species are well known to release a vast bouquet of Biogenic Volatile Organic
41 Compounds (BVOC). Terrestrial living vegetation releases between 700 and 1000 TgC of BVOC
42 every year to the atmosphere, of which 40 to 65% are terpenes, mainly isoprene (309–706 TgC
43 yr⁻¹), followed by monoterpenes (26–156 TgC yr⁻¹) and sesquiterpenes (26 TgC yr⁻¹) (Navarro et
44 al., 2014). Higher BVOC emissions are probably released at the global scale since it has been
45 estimated that soil (bare soil + litter) contributes to 1-20 % of total BVOC emission in different
46 ecosystems (Asensio et al., 2007; Kramshøj et al., 2016; Tang et al., 2019). Quantification of leaf
47 litter emissions at global or regional scales has not been performed using BVOC emission models
48 due to data scarcity (Peñuelas et al., 2014). BVOC emissions from litter are important in ecological-
49 related processes as they modify plant growth and germination, acting as volatile allelochemicals
50 (Santonja et al., 2019) and in atmospheric chemistry by influencing the OH· budget, O₃ and
51 secondary organic aerosol formation (Fehsenfeld et al., 1992; Spittler et al., 2006). As recently
52 reviewed (Tang et al., 2019), a missing OH sink would be due to incomplete characterization of
53 BVOC sources, particularly emissions from litter and soil.

54 Quantification of BVOC emissions at the ecosystem scale is not properly resolved with
55 available models (Peñuelas et al., 2013) where only BVOC emission from the canopy are
56 considered. In order to integrate litter in BVOC emission models, the first step it is to list BVOC
57 emission rates of different plant species litter as previously carried out for living plants (Owen et
58 al., 1997, 2001). The few studies that have reported litter BVOC emissions seem to support that
59 litter is the second major source of BVOC after the canopy (Mäki et al., 2017; Tang et al., 2019)
60 and releases higher emissions than bare soil (Ramirez et al., 2010). BVOC litter-related studies
61 refer either to the layer of dead plant material present on the soil surface (Schade and Goldstein,
62 2001; Gray et al., 2014) or dead plant material that is detached from a living plant (Isidorov et al.,
63 2010; Isidorov et al., 2016), since ecologically the word litter has these two meanings (Krishna and
64 Mohan, 2017).

65 It is difficult to evaluate if leaf litter of a given species releases higher or lower emission
66 rates compared to its green leaves. For example, litter emission rate of *Pinus sylvestris* L., measured
67 with a static system and SPME (Solid Phase Micro-Extraction), is 7.5 µg.g_{DM}⁻¹.h⁻¹ (Isidorov et al.,
68 2010) in good agreement with emissions from green litter (maximal emission rate 6 µg.g_{DM}⁻¹.h⁻¹;
69 Vanhatalo et al., 2018). Emission rates of *Picea abies* (L.) H. Karst litter, also measured with a

70 static system and SPME, is $1.3 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ (Isidorov et al., 2010) contrast with green leaf emission
71 ($\sim 0.1\text{-}0.4 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Bourtsoukidis et al., 2014). Likewise, *Pinus taeda* L. litter showed an
72 average emission rate of $117 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ (Ramirez et al., 2010) after litter conditioning (drying at
73 75°C and incubation for 72 days), which is roughly 100 times higher compared to its green leaf
74 emissions (Helmig et al., 2006). Litter conditioning and handling previous to measurements
75 probably explain such differences. However, all these studies suggest that terpene emissions from
76 litter are not negligible compared to green leaves.

77 Leaf litter emission varies according to decomposition stages (Gray et al., 2010; Isidorov
78 et al., 2010) as terpenes volatilize from terpene pools and microbiota associated to litter releases
79 and consumes BVOC. As a result, comparing leaf litter emission between species requires to
80 consider litter from the same decomposition stage. Using fresh leaf litter that has not undergone
81 microbial decomposition in soil allows to isolate BVOC emission from litter and is also the most
82 accurate way to compare BVOC emission rates from green leaves and leaf litter.

83 Fresh litter could importantly contribute to BVOC emissions and the atmospheric chemistry
84 in the Mediterranean region where climate is characterized by a long and hot summer period (2-5
85 months; Polade et al., 2014) which could favor BVOC emissions from litter (Greenberg et al.,
86 2012). Moreover, litter decomposes slowly (Hart et al., 1992) and many Mediterranean plant
87 species including *Pinus* spp. and *Quercus* spp. produce important litter biomasses (e.g. *Pinus pinea*
88 L., *Quercus coccifera* L., Arianoutsou and Radea, 2000). The Mediterranean area is also highly
89 populated, leading to increases in anthropogenic pollutants, such as NO_x , necessary for
90 tropospheric O_3 formation, in the presence of BVOC and light (Atkinson and Arey, 2003). All
91 these interacting phenomena highlight the importance of focusing on other biogenic sources of
92 BVOC emission such as litter and not only living vegetation.

93 BVOC emissions from litter rely on both, soil microbial activity and specific BVOC
94 reservoirs (such as glandular trichomes, secretory cavities) and cuticle where waxes are
95 accumulated (Bondada et al., 1996). BVOC reservoirs (present only in some species) allow to store
96 a high quantity of compounds resulting in toxic concentration for the other organisms (herbivore
97 or phytopathogenic microorganisms) (Lange, 2015). BVOC reservoirs mostly contain terpenic
98 compounds (Wagner, 1991) (excepting isoprene which is produced *de novo* and never stored in
99 specific storing structures, Sharkey and Yeh, 2001), and are thus referred to as terpene storage
100 structures. These structures can either be internal to the leaf such as the secretory cavities within

101 needles of resinous species or external to the leaf such as the glandular trichome (e.g. *Lamiaceae*
102 spp). Numerous Mediterranean plant species (*Rosmarinus officinalis* L., *Pinus halepensis* Mill.,
103 *Pinus sylvestris*, *Pinus nigra* R. Legay) possess remarkable terpene amounts in these leaf reservoirs
104 (5-30 mg.g_{DM}⁻¹) presumably allowing these terpenes to be released in high amounts from leaf litter.
105 Other species (*Cistus* spp, *P. pinea*) only store small amounts of terpenes (5–64.5 μg.g_{DM}⁻¹) in these
106 structures (Ormeño et al., 2009). While terpene reservoirs are present in some species only, all
107 vascular species possess an upper cuticle layer which mainly stores low weight saturated and
108 unsaturated aliphatic hydrocarbons (< C16) (Bondada et al., 1996; Barik et al., 2004) but also a
109 small fraction of terpene compounds in coniferous species (Joensuu et al., 2016). All these volatiles
110 could thereby plausibly be released by leaf litter cuticle.

111 The main goal of this study is to characterize BVOC emissions from litter of 16
112 Mediterranean species to identify those that could mostly contribute to biosphere-atmosphere
113 interactions. Species belonging to different Plant Functional Types (PFT) are considered, including
114 evergreen trees, deciduous trees and evergreen shrubs. This study considers fresh leaf litter
115 (senescent leaves) that has not undergone decomposition by soil microorganisms and focuses on
116 leaf terpene content (from literature data) to evaluate if this parameter could be an indicator of the
117 potential of litter to release BVOC. Additionally, we aimed to estimate the potential impact of litter
118 BVOC emissions on air quality. For this purpose, we focused on two tree species (*P. halepensis*
119 and *Q. pubescens*) and calculated, for both plant species, three indicators of their influence on air
120 quality: regional litter BVOC emissions, photochemical ozone creation potential (POCP) and
121 secondary organic aerosol production (SOAP). These two plant species were chosen for this
122 modelling exercise since they are major species in the Sud-PACA region where they occupy 197
123 000 ha and 356 000 ha respectively (IGN – Inventory of 2014 to 2018).

124 **2. Material and Methods**

125 *2.1. Plant material*

126 Fresh leaf litter (hereafter referred to as litter) of 16 Mediterranean species was collected in
127 2017 and 2018 in Southern France (Sud-PACA region). Litter consisted of both senescent leaves
128 previous to abscission and senescent leaves freshly fallen from the plant (within the first days
129 following leaf abscission) collected by means of a fine-mesh net previously placed in the field to
130 avoid litter contact with soil (Santonja et al., 2018). The studied species were chosen in order to
131 select a significant number of the most abundant Mediterranean species. We paid a special attention

132 to integrate both, species with leaf terpenic reservoirs and species that do not possess such

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>Q. ilex</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>Q. ilex</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>Q. pygmaea</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>Q. lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>Q. pedunculata</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>Q. pubescens</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno & Llusia et al.
<i>Q. pubescens</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>Q. officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>Q. agrifolia</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>Q. sessiliflora</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejean et al. (2014)
<i>Q. ilex</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>Q. agrifolia</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt et al.
<i>Q. agrifolia</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier et al. (1999)
<i>Q. pubescens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier et al.
<i>Q. agrifolia</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio et al.
<i>Q. agrifolia</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

133 structures, also species belonging to different plant functional types and finally taking into account
134 their litter production. For example, *P. halepensis* is an *Evergreen needle leaf tree*, possesses
135 terpene storage structures, occupies 231 000 ha in France and features a non-negligible litter
136 production ($150\text{-}530 \text{ g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$,

137 Table 1). *Q. pubescens* is a *Deciduous broad leaf tree* free of terpene storage structure,
138 occupies 320 000 ha in the Sud-PACA region and features a moderate litter production ($138\text{-}194$
139 $\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$,

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>...s</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>...folius</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et
<i>...pygria</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>...lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (200)
<i>...cedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen &
<i>...nsis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno & Llusia ar
	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewit
<i>...fficialis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>...aris</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et
<i>...ssulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard- & (2014)
<i>...a</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et
<i>...cifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt &
	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselm & (1999)
<i>...escens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselm & Saunier c
<i>...er</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio e
<i>...rus</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et

141 Litter was collected from 3 to 10 individuals in different locations and months according to
142 the time of the year when litter fall peak of each species occurs (

143 Table 1). Litter collection of each species took place in the same location and was never

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					g.m ⁻² .y ⁻¹	Reference	µg.g ⁻¹ .h ⁻¹	Type	Reference
<i>Cistus</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>foliolus</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>cogygia</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>globulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>oxycedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>halensis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno & Llusia et al.
	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>vulgaris</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>monspessulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejean (2014)
<i>arborea</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>coccifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Kesselmeier
	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier (1999)
<i>pubescens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier
<i>suber</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio
<i>canadensis</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

144 carried out after rain events to avoid excess of humidity. Average litter humidity was 9.6 ± 1.2 %
 145 (\pm standard error).

146 Seven of the studied species do not possess specific terpene storage structures (*Acer*
 147 *monspessulanum* L., *Erica arborea*, L., *Q. coccifera*, *Quercus ilex* L., *Quercus pubescens* Willd.,
 148 *Quercus suber* L., *Ulex parviflorus* Pourr.). Nine species possess terpene storage structures internal
 149 to the leaf (*Eucalyptus globulus* Labill., *Juniperus oxycedrus* L., *P. halepensis*, *P. pinea*) or external
 150 to the leaf (*Cistus albidus* L., *Cistus salviifolius* L., *Cotinus coggygia* Scop., *R. officinalis*, *Thymus*
 151 *vulgaris* L.,

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>...s</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>...folius</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>...pygria</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>...lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>...cycedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>...ensis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Owen & Ormeno & Llusia et al.
	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>...officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>...aris</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>...ssulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejeu & Pio (2014)
<i>...a</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>...cifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Ormeno
	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier et al. (1999)
<i>...escens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier et al.
<i>...er</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio et al.
<i>...rus</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

153 **Table 1** – Species studied, family they belong to, data when litter was collected and location, presence of storage structures in leaves, leaf litter
 154 production, green leaf emission rate and type according to literature. Species are classified according to the presence or absence of specific terpene-
 155 storage structures and alphabetically ordered.

	Species name	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
						g.m ⁻² .y ⁻¹	Reference	µg.g ⁻¹ .h ⁻¹	Type	Reference
TERPENES STORING SPECIES	<i>Cistus albidus</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Hewit (2000); Ormeno et al. (2007a)
	<i>Cistus salviifolius</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al. (2001)
	<i>Cotinus coggygria</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
	<i>Eucalyptus globulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. (2000); Nunes & Pio (2001)
	<i>Juniperus oxycedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Hewit (2000)
	<i>Pinus halepensis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno et al. (2007a); Llusia and Peñuelas (2000)
	<i>Pinus pinea</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. (1997); Owen & Hewit (2000)
	<i>Rosmarinus officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Hewit (2000); Ormeno et al. (2007a)
	<i>Thymus vulgaris</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al. (2004)
NON TERPENES STORING SPECIES	<i>Acer monspessulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Zielinski et al. (2014)
	<i>Erica arborea</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al. 1997
	<i>Quercus coccifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Hewit (2000); Staudt & Lhoutellier (2011)
	<i>Quercus ilex</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier and Staudt (1999)
	<i>Quercus pubescens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier et al. (1999); Saunier et al. (2017b)
	<i>Quercus suber</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. (2005); Sánchez - Osorio et al. (2019)
	<i>Ulex parviflorus</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al. (2001)

156 Iso: isoprene; MT : monoterpenes, ST : sesquiterpenes

157 ^a calculated mean (150 -530 g.m⁻².y⁻¹) from Arianoutsou and Radea (2000)

158 ^b calculated mean (138 -194 g.m⁻².y⁻¹) from Saunier et al. (2018)

159 2.2. Collection of BVOC emissions from litter

160 A dynamic pull-push system was used to collect BVOC released by litter. This system is
161 composed by 2 Pyrex® jars of 1.2 L each and a system that continuously flushes air inside these
162 jars. The air flushed into each jar was previously filtered with both drierite (W.A.
163 Hammond Drierite™ Indicating Absorbents Mesh size 8, USA) to avoid the excess of humidity
164 inside the system, and activated charcoal (Untreated, Mesh 4-8, Sigma Aldrich, USA) to remove
165 VOC from ambient air. Finally, the air goes through a particle filter treated with sodium thiosulfate
166 to remove the O₃ that could interact with BVOC emitted by litter.

167 Ten grams of freshly collected litter were placed inside a jar, leaving the other one empty to
168 check for the absence of VOC in the air flushed through the jars (blank measurement). We used
169 between 3 to 10 replicates per species, and emissions were collected once for each replicate. The
170 emission factor of litter (standardized emission rate at 30 °C; Guenther et al., 2006) was measured
171 by placing the jars inside a stove at 30 °C. Temperature and air humidity were measured with an
172 iButton ® (Hygrochron, Maxim Integrated iButtons, USA) placed inside each jar. Before
173 measurements took place, air inside the jars was renewed (1 L.min⁻¹ during 30 min) to avoid
174 sampling BVOC that would initially be inside the jar. During VOC sampling, flow inside and
175 outside the jar was set up to 0.2 L.min⁻¹ for 20 min. We used Mass Flow Controllers (MFC, EL-
176 Flow select, Bronkhorst France, 0-1 L. min⁻¹) to control the flows. The air coming out of each jar
177 was collected onto glass cartridges (20 cm height; 10.5 mm external diameter, Gerstel®, Germany)
178 filled with two kind of adsorbents : Tenax TA (50 mg; Agilent technologies; 20 - 35 Mesh) and
179 Carbotrap B (120 mg; Carbotrap Graphitized Carbon Black n°20287; Sigma Aldrich). Before use,
180 cartridges were conditioned at 300 °C for 3 h with a constant nitrogen flow (Nitrogen 4.5, purity
181 99.995 %) using a cartridge conditioner (TERA environment; RT-12-CN-3601-01). After BVOC
182 collection, cartridges were sealed and put inside a freezer at -4 °C until being analyzed by GC-MS
183 (see below).

184 Between samplings, the Pyrex jars were rinsed with water, then with ethanol (70 %) and
185 finally put inside a muffle furnace for 4 h at 400 °C to ensure full VOC and plant debris removal
186 that could influence the following samples. Blanks were performed for confirmation.

187 2.3. Chemical analysis

188 BVOC were analyzed using a Gas Chromatography (GC, 6890N, Agilent Technologies®)
189 coupled to a Mass Spectrometer Detector (MSD 5973, Agilent Technologies ®). BVOC trapped
190 in the cartridges were desorbed by a thermodesorber using a ramp of temperature from 50 °C to
191 250 °C at the rate of 3 °C.s⁻¹ for 15 min (TDS3/CIS4, Gerstel®), using helium as carrier gas (50
192 mL.min⁻¹). BVOC were then trapped in the Cooled Injection System (CIS) at 6.8 Psi pressure at
193 - 50 °C using liquid nitrogen. Then, the CIS was rapidly heated (25 s) from -50 °C to 250 °C
194 allowing molecule passage to the chromatographic column (HP5-MS, 30 m length, 0.25 mm
195 diameter, 0.25 µm film thickness, Agilent Technologies®). Molecules were separated using a ramp
196 of temperature from 40 °C to 280 °C with a rate of 3 °C.min⁻¹ with a constant helium flow of 1.0
197 mL.min⁻¹. The method accuracy (101.27 %) and precision (4.7 %) were determined through
198 replicate analyses (dodecane). The calibration linearity or coefficient of determination (R² =
199 0.9935, CV = 0.0019) and the limit of detection (LOD = 0.10 ng) were determined using standard
200 calibrations. Molecule identification in chromatograms (Suppl. Figure 2) was achieved using
201 spectral libraries (NIST library), spectra from injected standards of high purity (> 99 %, Sigma
202 Aldrich®, France, Suppl Table 1 & 2), and by comparison of our experimental and theoretical
203 Kovats index from Adams (2007).

204 2.4. Scanning electron microscopy

205 Since we did not find any data on the type of leaf terpene storage structures in *C. coggygria*,
206 we explored its litter surface (abaxial side) using a Scanning Electron Microscopy (SEM, model
207 XL30 ESEM from FEI). Small surfaces of leaf litter were fastened with adhesive tape on brass
208 stubs and then gold coated in a metallizer using a Cressington 108 Sputter Coater.

209 2.5. Calculation of regional litter emission, global POCP and SOAP of *P. halepensis* and *Q.* 210 *pubescens*

211 In order to assess the impact of litter emission on air quality, we calculated three indicators.
212 First, we estimated the regional litter BVOC emissions (t.y⁻¹) for the Sud-PACA region according
213 to the following equation:

$$214 \text{ Regional litter BVOC emissions} = \frac{ER \times 8760 \times \text{Litter production} \times \text{Surface}}{10^9} \quad (1)$$

215 where, *ER* refers to emission rates (µg.g_{DM}⁻¹.h⁻¹) reported in this study, 8760 refers to the hours in
216 one year, *Litter production* refers to annual litter production (g.m⁻²) noted in Table 1, the *Surface*

217 refers to the forest area (m²) (IGN 2014-2018) occupied by each species in the Sud-PACA region
218 and 10⁹ is used to express results in ton. Second, for the two litter, we have calculated the
219 photochemical ozone creation potential (POCP) and the potential secondary organic aerosol
220 production (SOAP) using the following equations:

$$221 \quad POCP = \sum_{n=i} [contribution (\%)_i \times POCP_i] \quad (2)$$

$$222 \quad SOAP = \sum_{n=i} [contribution (\%)_i \times SOAP_i] \quad (3)$$

223 where, *contribution (%)_i* refers to the relative contribution of the BVOC *i* to the litter emission
224 under study, *POCP_i* refers to the individual POCP of the BVOC *i* as reported in the Suppl. Table
225 6 and *SOAP_i* refers to individual SOAP of the BVOC *i* as reported in the Suppl. Table 6,

226 The POCP for individual VOC were reported from Dominutti et al. (2019). Unfortunately,
227 POCP values were missing for monoterpenes and sesquiterpenes. Thus, as suggested by several
228 authors, the isoprene POCP value (109.2) was also used for monoterpenes, because of their
229 chemical similarity (Huang et al., 2017; Dominutti et al., 2019). The same reasoning was applied
230 in this work for sesquiterpenes for which the isoprene value was used. Likewise, the SOAP for
231 individual VOC were reported from Derwent et al., (2010) but SOAP values were available in
232 literature for only two monoterpenes having similar values (α -pinene: 17.4 and β -pinene: 18.1).
233 They were missing for the other monoterpenes and sesquiterpenes. As suggested by Dominutti et
234 al. (2019), the SOAP value of β -pinene was used in the present work for all the monoterpenes
235 because of their similar chemical properties. The SOAP values of individual sesquiterpenes were
236 estimated from their SOA Mass Yield, using SOAP and SOA Mass Yield of α -pinene and isoprene
237 as reference according to the following equation:

$$238 \quad SOAP_i = \frac{SOA_M_Y_i \times SOAP_{ref}}{SOA_M_Y_{ref}} \quad (4)$$

239 where, *SOAP_i* and *SOA_M_Y_i* are the SOAP and the SOA Mass Yield of the sesquiterpene *i*
240 respectively, *SOAP_{ref}* and *SOA_M_Y_{ref}* are the SOAP and the SOA Mass Yield of the reference
241 VOC. Because several SOA Mass Yields were available in literature for each sesquiterpene and
242 reference VOC, the mean values were used (Suppl. Table 7). The calculated individual SOAP
243 values obtained using the two reference VOC (isoprene and α -pinene) were close and the mean
244 values were used in the present work (Suppl. Table 7)

245 2.6. Statistical analysis

246 Differences and similarities in the BVOC emission profile between litter from different plant
247 species were assessed using a Principal Component Analysis (PCA). The statistical analysis of the
248 data was performed with R software (version 3.4.X, R Core Team 2017) using ade4, factoextra
249 packages.

250 3. Results

251 3.1. BVOC emissions from leaf litter

252 Regarding total emissions, we identified a remarkable diversity with a total of 87 compounds,
253 mainly terpenes (69 different compounds) followed by non-terpenic compounds (18 compounds)
254 including alkanes, alkenes, aldehydes, ketones and benzenoids. Within each species, the bouquet
255 of released compounds was also important for most species ranging from 4 emitted compounds in
256 *Q. ilex* to 46 in *J. oxycedrus* (Suppl. Table 1 and 2). Terpene-storing species emitted an important
257 diversity of compounds (mean: 29 ± 4 compounds) with the most commonly emitted terpenes being
258 limonene, β -caryophyllene, α -pinene, α -humulene and α -terpinolene (Suppl. Table 1). By contrast,
259 the non-terpene storing species emitted on average 8 ± 1 compounds. Quantitatively, the studied
260 species could be classified in 3 different groups based on their total emissions: *i*) negligible emitters
261 ($< 0.1 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) including *A. monspessulanum*, *Q. ilex*, *U. parviflorus*, *Q. suber* and *E. arborea*,
262 *ii*) low emitters ($0.1 < \text{emissions} < 1.0 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) including (in decreasing order) *R. officinalis*,
263 *Q. pubescens*, *P. pinea*, *Q. coccifera*, *C. albidus*, *J. oxycedrus*, *C. salviifolius* and *E. globulus* and
264 , *iii*) moderate emitters ($1.0 < \text{emissions} < 5.0 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) including (in decreasing order) *T.*
265 *vulgaris*, *C. coggygria* and *P. halepensis* (4.74; 3.37 and $1.52 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ respectively; Suppl.
266 Table 3).


267 Terpene emissions only occurred from terpene-storing species. The highest terpene emitters
268 were also *T. vulgaris*, *C. coggygria* and *P. halepensis* (4.74; 3.37; $1.52 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ respectively)
269 whereas the remaining species emitted between 0.22 (*E. globulus*) and $0.71 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ (*R.*
270 *officinalis*). Monoterpenes dominated emission from *T. vulgaris* (α -terpinyl acetate, α -terpineol,
271 limonene, linalool and trans-4-thujanol) and *C. coggygria* (limonene, cis- β -ocimene and myrcene)
272 representing 98 % and 91 % respectively (Suppl. Table 1). By contrast, *P. halepensis* featured
273 sesquiterpene emissions (61 % of total emissions), mainly β -caryophyllene and α -humulene
274 (Suppl. Table 1) and showed the highest sesquiterpene emission rates ($0.92 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) of all
275 species. Within each terpene emitter species, terpenes accounted for the major fraction in all

276 species, excepting *C. albidus* where terpene emissions ($0.18 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) were close to non-
277 terpenic emissions ($0.11 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Suppl. Table 3).

278 All species released non-terpenic emissions (Suppl. Table 3; Suppl. Table 4), saturated
279 aliphatic alkanes ($\text{C}_{14}\text{H}_{30}$, $\text{C}_{13}\text{H}_{28}$), followed by aldehydes (nonanal, decanal), representing the
280 major fraction in most species (Suppl. Table 1 and 2). The most remarkable non-terpenic emissions
281 occurred in *Q. pubescens* > *Q. coccifera* > *C. coggygia* (0.61 , 0.32 and $0.23 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$
282 respectively), followed by *P. halepensis* and *C. albidus* (0.16 and $0.11 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$, respectively).


283 Species that do not possess terpene storage structures showed lower total BVOC emission
284 rates than terpene storing species, with two exceptions, *Q. pubescens* and *Q. coccifera*, with total
285 non-terpenic emission (0.61 and $0.32 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$, respectively) higher than total emissions of the
286 terpene-storing species *C. albidus*, *C. salviifolius* and *J. oxycedrus*. (0.30 , 0.21 and $0.21 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$
287 ¹, respectively).

288 Clear differences occurred between species, even when they belong to the same genus
289 (Figure 1). In *Pinus* spp, *P. halepensis* mainly emitted sesquiterpenes (60 % of total emission),
290 with a large diversity of volatiles (27 compounds) and a total emission rate 3-times higher (1.52
291 $\mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) than *P. pinea* ($0.54 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) which mainly released monoterpenes (60 % of total
292 emissions) with a higher diversity (43 compounds) (Suppl. Table 3). *Quercus* spp.- the main source
293 of non-terpenic emissions - showed a poorer diversity of BVOC (4-10 compounds) compared to
294 *Pinus* spp. *Q. suber* and *Q. ilex* showed lower emission rates ($< 0.1 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) compared to *Q.*
295 *coccifera* and *Q. pubescens* (0.32 and $0.61 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) (Suppl. Table 4). *C. albidus* and *C.*
296 *salviifolius* emitted BVOC in similar rates (Suppl. Table 3), but *C. albidus* mostly released
297 monoterpenes (50 % of total emission) (mainly limonene, Suppl. Table 1), while *C. salviifolius*
298 mainly released sesquiterpenes (71 % of total emission, β -caryophyllene and α -humulene). Litter
299 emission rates also exhibited some quantitative differences between samples of the same species,
300 up to 5-times different in samples of *E. globulus*.



301
 302 **Figure 1** – Principal Component Analysis showing litter BVOC profiles of the different species studied.
 303 Litter samples are represented with symbols and variables are represented with arrows. Variables with the
 304 highest weight in Dim 1 are alkanes and aldehydes on the right part of PC1, and monoterpene derivatives
 305 and monoterpene ketones on the left part of Dim 1. Variables with the highest weight in Dim 2 are
 306 sesquiterpene hydrocarbons and monoterpenes hydrocarbons on the top part of Dim 2. Weights of all
 307 variables is shown in Suppl Table 3.


308 The sum of the most emitted terpenes (representing individually at least 5 % of the total
 309 emission) contribute, at least, to half of the total emissions of a given species. For example, only
 310 two and three terpenes account for ~ 50 % of total emissions from *J. oxycedrus* and the remaining
 311 emissions were also characterized by terpene compounds, indicating that this species emits a large
 312 diversity of terpenes in small quantities. By contrast, only four terpenes represent 90 % of total
 313 emissions from *R. officinalis*, indicating few terpenes dominated emissions from this species
 314 (Figure 2).



3__

316 **Figure 2** – Contribution (%) of the major terpenes to the total emissions of nine plant species that possess
 317 terpene storage structures. BVOC shown in the caption are ordered according to their molecular weight and
 318 represent at least 5% of total emission of the species under study. Values are mean of n=3-10.

319 The contribution of each group of compounds to the total emission varied between species
 320 (Figure 3 and Figure 4). Terpene emissions (monoterpene hydrocarbons, oxygenated
 321 monoterpenes, sesquiterpene hydrocarbons and oxygenated sesquiterpenes) represented most of
 322 the emissions for the species that possess terpene storage structures (Figure 3). For example, they
 323 represent 62 % to 100 % of the total emission (*C. albidus* and *R. officinalis* respectively). By
 324 contrast, for species that do not possess terpene storage structures, non terpenic emissions are the
 325 most emitted compounds with alkanes and aldehydes being the major BVOC. *Quercus* spp. showed
 326 major alkane emissions (54 % to 77 % of total emission for *Q. coccifera* and *Q. pubescens*) while
 327 *A. monspessulanum*, *E. arborea* and *U. parviflorus* mostly emitted aldehydes (52 %, 46 % and
 328 56 % respectively; Figure 4).




329

330

331 **Figure 3** – Contribution (%) of the compound groups (terpene and non-terpenic groups) to the total
 332 emissions of nine species that possess terpene storage structures. Values are mean of n=3-10.

333



334

335 **Figure 4** – Contribution (%) of the compound groups to the total emissions of six species that do not possess
 336 terpene storage structures. Values are mean of n=3-10.

337 *3.2. Potential participation of P. halepensis and Q. pubescens litter BVOC to air quality at regional*
 338 *scale*

339 The potential impact of the two main tree species in the Sud-PACA region on the regional
 340 air quality are assessed according to the three indicators previously described: their regional BVOC
 341 emissions (t.y^{-1}) and their potential to form SOA and O_3 . *P. halepensis* regional emissions represent
 342 almost 3 times those of *Q. pubescens* (8919 and 3158 t.y^{-1} respectively, Table 2) and its POCP is
 343 2 times higher (96.2) than the POCP of *Q. pubescens* (47.7).

344 Despite the lower regional BVOC emissions from *Q. pubescens* litter, the SOAP calculated
 345 for both species is similar (36.9 and 37.5 for *P. halepensis* and *Q. pubescens* respectively). Indeed,
 346 on the one hand, *Q. pubescens* emissions are dominated by alkanes (pentadecane and tetradecane)
 347 which have the highest individual SOAP (54.5, Suppl. Table 6) but this high values are
 348 compensated by the SOAP value (zero) of aldehydes (second major group of BVOC from *Q.*
 349 *pubescens*). On the other hand, *P. halepensis* emissions are mainly constituted by monoterpenes
 350 and sesquiterpenes which have a lower individual SOAP (18 and 51.1 respectively, Suppl. Table
 351 6).

352 **Table 2** – Emission rates (ER) of the litter of *P. halepensis* and *Q. pubescens* and the estimation of their
 353 litter emission at a regional scale (t.y^{-1}), their litter production ($\text{g.m}^{-2}.\text{y}^{-1}$) and the global litter POCP and
 354 SOAP attributed

Plant species	Emission type ^a	ER ($\mu\text{g.gDM}^{-1}.\text{h}^{-1}$) ^b	Litter production ($\text{gDM.m}^{-2}.\text{y}^{-1}$)	ER ($\mu\text{g.gDM}^{-1}.\text{y}^{-1}$)	ER ($\text{g.m}^{-2}.\text{y}^{-1}$) ^e	Assessed regional BVOC emission (t.y^{-1}) ^f	POCP ^g	SOAP ^g
<i>P. halepensis</i>	ST > MT > Ald > Alk	1.5	340 ^c	13300	4,5	8919	96.2	36.9
<i>Q. pubescens</i>	Alk > Ald	0.6	166 ^d	5340	0,9	3158	47.7	37.5

MT: monoterpenes; ST: sesquiterpenes; Ald: Aldehydes; Alk: Alkanes

^a Emissions that represent at least 1% of total emissions

^b Results from this study

^c Arianoutsou & Radea (2000)

^d Saunier et al. (2018)

^e Obtained by multiplying the emission rate ($\mu\text{g.gDM}^{-1}.\text{y}^{-1}$) by litter production

^f Obtained by multiplying emission rates ($\text{g.m}^{-2}.\text{y}^{-1}$) by the surface occupied by the plant species in the Sud-PACA region: 197000 ha for *P. halepensis* and 356000 ha for *Q. pubescens* according to to IGN 2014-2018

^g Refer to material and methods for calculations and Suppl. Table 6 for SOAP and POCP of each compound

355 **4. Discussion**

356 This study shows that fresh leaf litter is a source of BVOC emissions with important
357 differences between species but also within litter of a given species (up to 5-times). BVOC
358 emissions from green leaves are indeed also characterized by an important variability within a
359 species under the same environmental conditions. For example, the study from Genard-Zielinski
360 et al. (2014) reported important inter-canopy differences (up to 4-times) for *Q. pubescens* under
361 very similar conditions of light, temperature and air relative humidity in the field.

362 Results are discussed in the following subsections for all the species grouped according to
363 their PFT as used in BVOC emission models (Guenther et al., 2006; Messina et al., 2016). Before
364 that, it is worth to mention that results in the present work were obtained at an averaged litter
365 humidity of 9.6 % which is close to previous published data for senescent leaves (8 % in Isidorov
366 2003, 5-7 % Romero et al., 2019). Different litter humidity values would have probably led to
367 higher emission rates considering the laboratory results from Greenberg et al (2012) on *Pinus*
368 *ponderosa* needle litter (specific values not reported in their paper). Excluding Greenberg et al
369 (2012), all other litter BVOC-related studies in the laboratory have been performed using a single
370 humidity (8 % litter humidity of *Pinus sylvestris* in Isidorov 2003, 0 % humidity in 12 species in
371 Gray et al., 2010).

372 *4.1. BVOC emissions from litter of evergreen needle leaf tree species*

373 *P. halepensis* showed one of the highest litter emission rates in the present study ($1.52 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$)
374 $^1\cdot\text{h}^{-1}$), in agreement with previous results on *P. sylvestris* non decomposed litter $1.96 \mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$
375 (Isidorov et al., 2010), and contrasting to *P. pinea* (~ 3-times less emission rates). This difference
376 could be partly explained by the terpene content within their leaf litter which is negligible in *P.*
377 *pinea* ($0.064 \text{ mg}\cdot\text{g}_{\text{DM}}^{-1}$) and high in *P. halepensis* ($5 \text{ mg}\cdot\text{g}_{\text{DM}}^{-1}$, Ormeño et al., 2009). The moderate
378 emission rates of *P. halepensis* litter could significantly contribute to BVOC emissions at the scale
379 of the Mediterranean region where it shows a rather important leaf litter production ($150\text{-}530$
380 $\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$),

381 Table 1). *P. halepensis* could also remarkably contribute to total BVOC in local areas such as

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					g.m ⁻² .y ⁻¹	Reference	µg.g ⁻¹ .h ⁻¹	Type	Reference
<i>P. halepensis</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>P. halepensis</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>P. halepensis</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>P. halepensis</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>P. halepensis</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>P. halepensis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno & Llusia ar
<i>P. halepensis</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>P. halepensis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>P. halepensis</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>P. halepensis</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard- & Pio (2014)
<i>P. halepensis</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>P. halepensis</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Ormeno
<i>P. halepensis</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier (1999)
<i>P. halepensis</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier
<i>P. halepensis</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio
<i>P. halepensis</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

382 Southern France (PACA- Sud Region) were *P. halepensis* is especially present (231 000 ha,
383 (Fernandez et al., 2013). Considering all the Western Mediterranean area, *P. halepensis* is more
384 widespread (~3.5 million ha) than *P. pinea* (320 000 ha) (Quézel and Médail, 2003). Thereby, it is
385 likely that litter from *P. halepensis* contributes more importantly to BVOC emissions than *P. pinea*
386 in the Mediterranean region. Contrasting to leaf litter, literature shows higher emission rates from
387 green leaves of *P. pinea* and *P. halepensis*, mainly represented by monoterpene compounds (

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>J. asperifolia</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>J. folius</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>J. pygria</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>J. lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>J. oxycedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>J. pensis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno & Llusia et al.
<i>J. pensis</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>J. officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>J. parisi</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>J. sissulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejeu (2014)
<i>J. sissulanum</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>J. scifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Ormeno
<i>J. scifera</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier (1999)
<i>J. scensens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier et al.
<i>J. serotina</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio et al.
<i>J. sissulanum</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

389 *J. oxycedrus* is a small tree found both in Mediterranean forest and garrigues. This species
390 seems to contribute little to BVOC emissions since leaf litter emission rates ($0.21 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) is
391 low (

392 Table 1). This tree species possesses internal terpene storing structures and their content is

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>J. asperifolia</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>J. folius</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et
<i>J. pygria</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>J. lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. Pio (200
<i>J. oxycedrus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen &
<i>J. vasis</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Ormeno Llusia ar
	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewit
<i>J. officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>J. parisi</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et
<i>J. sissulanum</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard- (2014)
<i>J. a</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et
<i>J. cifer</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt &
	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselm (1999)
<i>J. escens</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselm Saunier c
<i>J. er</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. Osorio e
<i>J. orus</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et

393 often used to produce essential oil. However, terpene emissions from litter contrast with literature
 394 on its needle essential oil content which is characterized by monoterpenes (41 % of total emission;
 395 Dob et al., 2006) while its litter emits mostly sesquiterpenes (57 % of total emission; Figure 3).
 396 The green leaves of *J. oxycedrus* also emits different types of BVOC including isoprene, which
 397 can not be produced by the litter, monoterpenes and sesquiterpenes, its emission can reach 2.3
 398 $\mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$. The difference of emission between green leaves and litter might also be explained by
 399 the thickness of the cuticle of this species (Vasic and Dubak, 2012) that could act as a barrier for
 400 the emission when still intact.

401 4.2. BVOC emissions from litter of evergreen broad leaf tree species

402 Litter of *Q. ilex* and *Q. suber* released non-terpenic compounds only such as fatty acids
403 derivatives (alkanes, alkenes) probably emitted by the cuticle present on top of the leaf litter
404 (Maffei, 1994). Green leaves of these *Quercus* spp. are known to release high monoterpene
405 emission which require photosynthesis to occur, own to the absence of monoterpene reservoirs
406 within their foliage (Kesselmeier and Staudt, 1999). Such feature explains the absence of terpene
407 emissions from their freshly collected leaf litter. It can be concluded that evergreen forests of
408 *Quercus* ssp. are a remarkable source of terpene compounds from the canopy as shown in literature
409 and a minor source of non-terpenic compounds from litter, although this could be moderated by
410 their abundance and litter production.

411 *E. globulus* litter showed a rather low emission rate ($0.20 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) dominated by
412 monoterpenes. Contrastingly, Gray et al. (2010) showed that *Eucalyptus* litter released very high
413 amounts of monoterpenes ($0.4 \mu\text{mol}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$), which represents $54.4 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$. In their study,
414 litter had been previously oven-dried at 60°C and litter leaves were cut into small pieces triggering
415 monoterpene emissions from litter and accumulation in the head space system they used (125 mL
416 jars). It is important to note that the authors explicitly indicate that they aimed to compare litter
417 emission rates between species not to accurately quantify them. *E. globulus* canopy is a major
418 source of isoprene ($38 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Keenan et al., 2009) with some monoterpene emissions too (2.7
419 $\mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$). Moreover, terpene content of green leaves of *E. globulus* is very high ($25 \text{mg}\cdot\text{g}_{\text{DM}}^{-1}$)
420 contrasting with the leaf litter emission we measured. Since several authors (Wilt et al., 1993;
421 Romero et al., 2019) showed a good agreement between terpene content in leaf litter and green
422 leaves of numerous species, one can expect that terpene content in litter of *E. globulus* is also very
423 high. We argue that such discrepancy (between low emission rates found in this study and high
424 terpene content in green leaves reported in literature) is maybe due to the thick cuticles of fully
425 expanded leaves of *E. globulus* (Johnson, 1926), since the cuticle limits BVOC diffusion from leaf
426 tissues to the atmosphere (Joensuu et al., 2016).

427 4.3 BVOC emissions from litter of deciduous tree species

428 Leaf litter from *Q. pubescens* showed a low emission rate ($0.61 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) with only non-
429 terpenic emissions, mostly low chain fatty acids derivatives (alkanes, alkenes). Contrastingly,
430 green leaves of *Q. pubescens* are a remarkable source of isoprene emissions with emissions of fully
431 expanded leaves ranging from 78.5 to $124.3 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ at 30°C (Genard-Zielinski et al., 2014;
432 Saunier et al., 2017a). According to the European inventory of BVOC emissions (Keenan et al.,

433 2009), *Q. pubescens* is the second major isoprene emitting species. Isoprene requires *de novo*
434 formation to be produced (Sharkey and Yeh, 2001) since there is no specific storage of isoprene in
435 leaves (probably due to its very high vapor pressure) explaining that it is not released by leaf litter.
436 However, isoprene can be synthesized from soil microorganisms (as shown in the data base mVOC
437 2.0; Lemfack et al., 2018) which probably explains why Warneke et al. (1999) showed isoprene
438 emission from oak, beech, spruce, grass litter (although the authors did not detail whether it was
439 fresh or decomposed litter). Despite the low non-terpenic emissions we reported for *Q. pubescens*,
440 this species grows in highly widespread deciduous forests where litter production is moderate (153
441 $\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$; Suppl. Table 3), suggesting that *Q. pubescens* litter could contribute to some extent to air
442 chemistry during the leaf abscission period (December-March in Mediterranean ecosystems).
443 *Q. pubescens* abundance is especially important in Southern France where it occupies more than 2
444 million ha only in the Sud-PACA (Saunier et al., 2017b).

445 Forests dominated by *A. monspessulanum* are probably a negligible source of BVOC
446 emissions independently of the ecological compartment considered, since litter released the lowest
447 emission rates ($< 0.01 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$, all of them being non-terpenic) and green leaves release only
448 negligible terpene emissions ($< 0.5 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Ormeño et al., 2020; Genard-Zielinski et al., 2014).

449 4.4 BVOC emissions from litter of shrub species

450 *T. vulgaris* released the highest terpene emission rates ($4.72 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) similar to green leaf
451 emissions ($3.6 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Parra et al., 2004). Accordingly, *T. vulgaris* contains remarkable
452 monoterpene concentrations in the glandular trichome of green leaves ($22.0 \text{mg}\cdot\text{g}_{\text{DM}}^{-1}$; Nielsen et
453 al., 2015). This major aromatic plant species largely widespread in Mediterranean garrigues, is thus
454 probably an important source of BVOC in these ecosystems.

455 Interestingly, BVOC emissions from litter or green leaves of *C. coggygria* have never been
456 studied despite *C. coggygria* leaves have been largely used for medicinal purposes own to its
457 essential oil (Matić et al., 2016). This species naturally coexists in the understory vegetation of
458 Mediterranean forests (Genard-Zielinski et al., 2014) and is highly used as ornamental plant in
459 urban vegetation. We observed, for the first time, the presence of a capitate sessile trichome, a type
460 glandular trichome, on the leaf abaxial (bottom) surface (Suppl. Fig 1) in agreement with other
461 Anacardiaceae species (Metcalf and Chalk, 1988). Despite the presence of these structures, *C.*
462 *coggygria* stores poor amounts of terpenes ($0.3 \text{mg}\cdot\text{g}_{\text{DM}}^{-1}$) both, in leaf litter (Santonja et al., 2018)
463 and green leaves (Romero et al., 2019).

464 *R. officinalis* appeared to be a low emitter ($0.71 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$) with numerous monoterpenes.
465 Green leaves of *R. officinalis* release low to moderate terpene emission rates ($2.2 - 5.0 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$;
466 Owen and Hewitt, 2000; Ormeño et al., 2007b) and store high amounts of terpenes ($6.0 \text{mg}\cdot\text{g}_{\text{DM}}^{-1}$;
467 Peñuelas and Llusà, 1997; Ormeño et al., 2007b). This result reveals that the presence of an
468 important storage of terpenes does not necessarily implies a high emission rate from leaf litter
469 although the underlying processes remain to be tested.

470 *C. albidus* and *C. salviifolius* showed low emission rates of both terpene and non-terpenic
471 compounds (0.30 and $0.21 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$ each). Accordingly, terpene emissions from green leaves of
472 *Cistus spp.* are low to moderate ($0.5-5.0 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$; Owen et al., 2001; Ormeño et al., 2007a), and
473 store low terpene amounts ($0.8 \text{mg}\cdot\text{g}_{\text{DM}}^{-1}$; Ormeño et al., 2007b). Epicuticular wax coverage (very
474 long chained and saturated wax components) in leaves of *C. albidus* is very high ($200 \mu\text{g}\cdot\text{cm}^{-2}$)
475 and represents about 3.5 % of the leaf dry weight (Hennig et al., 1988). Such coverage, which is a
476 mechanism to cope with the relative aridity that characterizes macchias and garigues, probably
477 limits leaf litter terpene emissions too. Although litter from *Cistus spp.* releases low emission rates,
478 its contribution to BVOC emissions in Mediterranean garrigue ecosystems could be compensated
479 by a high litter production ($673 \text{g}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$;

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>Q. agrifolia</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>Q. foliata</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>Q. pygmaea</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>Q. lobulus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>Q. coccifera</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Ormeno
<i>Q. ilex</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Owen & Ormeno & Llusia et al.
<i>Q. ilex</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>Q. officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>Q. agrifolia</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>Q. sessiliflora</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejeu (2014)
<i>Q. agrifolia</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>Q. coccifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Ormeno
<i>Q. agrifolia</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier (1999)
<i>Q. agrifolia</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier et al.
<i>Q. agrifolia</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio et al.
<i>Q. agrifolia</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

481 *Q. coccifera* also showed low emission rates (only non-terpenic emissions) but, again, its
482 important litter production (

483 Table 1) could compensate for such low emissions at the ecosystem scale. Finally, *U.*

Species	Collection	GPS Coordinates	Storage structure	Plant Functional Type	Leaf litter production		Green leaves emission		
					$\text{g}\cdot\text{m}^{-2}\cdot\text{y}^{-1}$	Reference	$\mu\text{g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$	Type	Reference
<i>Ulex</i>	April	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	673	Rodriguez-Ramirez et al. (2017)	0.1 - 4.5	MT>ST	Owen & Ormeno
<i>Lotus</i>	June	43°10'44.3"N 5°34'16.2"E	Glandular trichome	Shrub	Unknown		3.1	MT>ST	Owen et al.
<i>Pygmaea</i>	November	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	117	Unpublished data	Unknown	Unknown	
<i>Quercus</i>	November	43°04'13.5"N 5°51'12.3"E	Subdermal oil glands	Evergreen broad leaf tree	250 - 370	Arianoutsou (1989)	5.4 - 42.0	MT>ST	He et al. & Pio (2000)
<i>Quercus</i>	August	43°14'27.1"N 5°40'33.3"E	Secretory cavities	Evergreen needle leaf tree	Unknown		0.5 - 2.3	Iso>MT>ST	Owen & Hewitt
<i>Quercus</i>	August	43°56'06.0"N 5°42'37.7"E	Secretory cavities	Evergreen needle leaf tree	340 ^a	Arianoutsou & Radea (2000)	3.0 - 15.0	MT>ST	Owen & Llusia et al.
<i>Quercus</i>	August	43°17'22.1"N 5°24'35.2"E	Secretory cavities	Evergreen needle leaf tree	240 - 790	Arianoutsou & Radea (2000)	3.4 - 8.8	MT>ST	Street et al. & Hewitt
<i>Quercus officinalis</i>	August	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	385	Rodriguez-Ramirez et al. (2017)	2.2 - 5.0	MT>ST	Owen & Ormeno
<i>Quercus</i>	July	43°56'06.0"N 5°42'37.7"E	Glandular trichome	Shrub	Unknown		3.6	MT>ST	Parra et al.
<i>Quercus sessiliflora</i>	November	43°56'06.0"N 5°42'37.7"E	None	Deciduous tree	55 - 125	Unpublished data	< 0.3	Iso, MT	Genard-Dejeux (2014)
<i>Quercus</i>	June	43°10'44.3"N 5°34'16.2"E	None	Shrub	Unknown		19.0	Iso	Owen et al.
<i>Quercus coccifera</i>	September	43°56'06.0"N 5°42'37.7"E	None	Shrub	538	Arianoutsou (1989)	5.5 - 14.6	Iso>MT, ST	Owen & Staudt & Hewitt
<i>Quercus</i>	June	43°14'27.1"N 5°40'33.3"E	None	Evergreen broad leaf tree	180 - 250	Bello et al. (1992)	15.0 - 21.0	MT>ST	Kesselmeier (1999)
<i>Quercus</i>	January	43°10'44.3"N 5°34'16.2"E	None	Deciduous tree	166 ^b	Saunier et al. (2018)	90.0	Iso	Kesselmeier & Saunier et al.
<i>Quercus</i>	June	43°15'19.2"N 6°13'06.3"E	None	Evergreen broad leaf tree	349	Robert et al. (1996)	0.2 - 30.0	MT	Pio et al. & Osorio et al.
<i>Quercus</i>	June	43°21'54.9"N 5°25'30.9"E	None	Shrub	323	Rodriguez-Ramirez et al. (2017)	9.0 - 14.0	Iso	Owen et al.

484 *parviflorus* and the *E. arborea* showed negligible BVOC emissions ($<0.04 \mu\text{g}\cdot\text{g}_{\text{DM}}^{-1}\cdot\text{h}^{-1}$), while
 485 green leaves release high isoprene emissions (Owen and Hewitt, 2000). Since these species are not
 486 the most commonly found in garrigues their litter is unlikely to contribute significantly to the whole
 487 BVOC budget at the ecosystem scale.

488 4.5. Environmental implications of BVOC emissions from litter

489 According to our study, most emitted terpene compounds were limonene, β -caryophyllene,
490 α -pinene and α -humulene, in agreement with Leff and Fierer, (2008) and emissions from green
491 leaves (Owen et al., 2001). We note that highly volatile compounds such as methanol,
492 acetaldehyde, acetic acid, formic acid and acetone, are also commonly released by leaf litter alone
493 (Gray et al., 2010; Greenberg et al., 2012; Tang et al., 2019) or soil and litter together (Schade and
494 Goldstein, 2001) but these highly volatile compounds were not screened in our study.

495 The influence of BVOC emissions from soil (litter included) on secondary air pollution and
496 the oxidizing capacity of the atmosphere are, by far, less well understood than their emission rates.
497 Terpenes found in our study are known to take part of cloud condensation nuclei formation leading
498 to the production of SOA (Hartz et al., 2005). The SOA mass yield (%) varies according to the
499 species: β -caryophyllene, α -humulene and α -pinene have a high SOA mass yield (respectively
500 45 %, 45 % and 41 %), while monoterpenes such as β -pinene or myrcene have a smaller SOA mass
501 yield (17 % and 11 % respectively; Lee et al., 2006). It is important to highlight that BVOC from
502 litter extend the seasonal period where BVOC are emitted from the ecosystem and so participate
503 to SOA formation (Faiola et al., 2014) since the canopy of deciduous broadleaf species do not
504 contribute to BVOC emissions through the entire seasonal cycle.

505 Although litter is a source of BVOC, part of the emission is adsorbed or assimilated through
506 soil-related processes (Ramirez et al., 2010) and part readily reacts with atmospheric oxidants. For
507 example, most commonly emitted terpenes in this study (limonene, β -caryophyllene, α -pinene and
508 α -humulene) feature high reactivity with OH (Atkinson and Arey, 2003). Accordingly, studies on
509 OH reactivity in the forests highlight a missing OH sink own to incomplete characterization of
510 BVOC sources (Di Carlo et al., 2004) and several authors suggest the missing source is soil
511 (including litter) (Praplan et al., 2019 and citations therein).

512 It is worth to note that BVOC emissions from litter are not only important in atmospheric
513 chemistry but also in ecological-driven processes. For instance, Santonja et al. (2019) recently
514 showed that β -caryophyllene emissions from *P. halepensis* litter negatively affected the
515 development of neighboring herbaceous plants, eventually shaping the ecosystem diversity.
516 Moreover, terpenes impact the soil food web and slow down the litter decomposition process
517 (Chomel et al., 2016).

518 It is important to highlight, that soil acts as a sink of BVOC (highly volatile compounds,
519 isoprene and monoterpenes) through microbial uptake, dissolution in soil water and adsorption to
520 soil particles (Misra et al., 1996; Ramirez et al., 2010). Independently of the underlying processes,
521 several authors suggest that measuring BVOC emission from different compartments (litter alone,
522 soil+litter, bare soil) can greatly benefit further model parametrization (Tang et al., 2019).
523 Measuring BVOC emission rates from isolated leaf litter is thus necessary.

524 4.6. Potential participation of *P. halepensis* and *Q. pubescens* litter BVOC to air quality at regional 525 scale

526 The regional emissions of *P. halepensis* and *Q. pubescens* litter (8.9 and 3.2 kt.y⁻¹) could
527 contribute to global forest BVOC emissions in France, which are estimated to 457 kt.y⁻¹ of
528 isoprene, 350 kt.y⁻¹ of monoterpenes and 129 kt.y⁻¹ of other BVOC considering only the canopy
529 BVOC emissions (Simon et al., 2001). It should however be highlighted that our estimations are
530 preliminary since litter BVOC emission are likely to vary over the decomposition process as later
531 discussed.

532 Although the SOAP of *P. halepensis* and *Q. pubescens* were similar, it is likely that *P.*
533 *halepensis* litter contributes more than *Q. pubescens* to SOA formation at the regional scale since
534 the amount of BVOC emitted by *P. halepensis* was more than 2-times higher. Similar, SOAP of
535 the two tree species is explained by the emission chemical composition of their litter. While *P.*
536 *halepensis* mainly releases sesquiterpenes (~ 60 % of total emissions) which are considered to have
537 a high SOAP equal to 51.1, followed by monoterpenes with a SOAP equal to 18 (see Suppl Table
538 6), *Q. pubescens* features alkane emissions (77 % of total emissions) which have the highest SOAP
539 (54.5, Dominutti et al., 2019, Suppl Table 6). By contrast, litter emission of *P. halepensis* showed
540 a 2-times higher POCP than *Q. pubescens*. This difference is due to the moderate POCP of alkanes
541 (44.5, Suppl Table 6) which accounted for the major emissions of *Q. pubescens* (Dominutti et al.,
542 2019, Suppl Table 6) whereas the individual POCP of monoterpenes and sesquiterpenes released
543 by *P. halepensis* is much higher (109.2, Suppl Table 6). As described in material and methods, the
544 POCP of terpenes actually lacks in literature. As a result, the isoprene POCP was used in the present
545 study for all terpenes because of their similar chemical structure, thus further studies to calculate
546 terpenes POCP would be useful to improve our estimates. With a moderate emission rate from the
547 litter of *P. halepensis* and a high POCP, the litter of this species could contribute to the formation
548 of O₃ and SOA in a non-negligible way.

549 According to our results, species that mostly release sesquiterpenes such as *P. halepensis*,
550 *C. salviifolius* and *J. oxycedrus* (60 %, 74 % and 60 % of the total emission respectively, Figure 3)
551 could be of special importance in terms of SOA formation. Moreover, species that emit both,
552 monoterpenes and sesquiterpenes would have a higher potential of ozone formation assuming that
553 terpenes possess a POCP as high as isoprene. These preliminary estimations thereby require further
554 modelling exercises using models such as CHIMERE and ORCHIDE but also BVOC emission
555 algorithms able to calculate BVOC litter emissions.

556 **5. Conclusion**

557 The forest canopy is known to be the main source of BVOC emissions. This study shows that
558 litter emission of Mediterranean tree and shrub species should not be neglected in biosphere-
559 atmosphere interactions. We also suggest that leaf terpene content shown in literature is not
560 necessarily an indicator of the potential of litter to release BVOC. Of the three species known to
561 possess rich terpene concentrations (*T. vulgaris*, *R. officinalis*, *E. globulus*), only *T. vulgaris*
562 released high terpene amounts from litter. While litter of most species released low BVOC
563 emission rates with a few exceptions (*T. vulgaris*, *C. coggygia*, *P. halepensis*), literature shows
564 that litter is also a source of highly volatile compounds (methanol, ethanol, short-chained
565 aldehydes, acids, acetone), which were not investigated in this study (Gray et al., 2010; Tang et al.,
566 2019). Consequently, emission rates would have been higher if a more exhaustive BVOC
567 screening had been carried out including compounds below C₁₀.

568 We suggest future *in vitro* and *in situ* research tackles BVOC emission variability according
569 to biotic (microorganisms) and abiotic factors (climatic conditions, soil). Since a couple of studies
570 have reported that litter BVOC emissions vary with decomposition degree (Gray et al., 2010;
571 Isidorov et al., 2010), soil properties (Gray et al., 2014; Mancuso et al., 2015; Chaignaud et al.,
572 2018), and climatic conditions (Derendorp et al., 2011, Greenberg et al., 2012), future research
573 should take into account these factors. For these purposes, BVOC from litter that has undergone
574 decomposition in the field (using for example litterbags, Isidorov et al., 2010, Santonja et al., 2015)
575 could be collected over different decomposition times ranging from several months to years. BVOC
576 from these differently decomposed litter could be measured by combining *in vitro* measurements -
577 using a controlled environment (at different temperatures and humidity) - and *in situ* measurements
578 -using transparent dynamic enclosures equipped with temperature, humidity and light sensors as
579 described by Svendsen et al. (2016). Transparency of the enclosure system would allow to take

580 into account weather variability (e.g. sunny, cloudy and so light conditions). Additionally, for *in*
581 *vitro* measurements, BVOC emissions of litter free of microorganisms (using for example gamma
582 irradiation as sterilization technique) could be compared to natural litter in order to assess
583 microorganism-related BVOC and better estimate the contribution of litter itself. Characterization
584 of microbiota litter (Wilkinson et al., 2002; Isidorov et al., 2016), could also help to explain BVOC
585 emission variability over decomposition stages. These types of studies (combining decomposition
586 processes and contribution of microbiota to BVOC emissions) will be compared to BVOC
587 emissions from non-decomposed litter (our study). We also suggest that both, *in vitro* and *in situ*
588 measurements are performed using litter plus soil, litter alone and soil alone to gain insight into
589 soil capacity to act as a sink of BVOC as indicated by different authors (Isidorov et al., 1999;
590 Chaignaud et al., 2018; Rinnan and Albers, 2020). It is particularly important to perform such
591 studies with a focus on BVOC emissions from *P. halepensis* litter since it features a high litter
592 production, occupies a high forest surface in the Mediterranean area and shows the highest litter
593 BVOC emission rates among all tree species considered in this study. Results from this species
594 could serve as a model to other *Pinus* spp.

595 To conclude, although litter BVOC emissions are roughly lower compared to green leaf
596 emissions, litter accumulates in soil so BVOC emissions could occur all year round. This could be
597 particularly true in the Mediterranean region where litter production of numerous species is rather
598 high and decomposes slowly due to relative water scarcity during most part of the year (Quézel
599 and Médail, 2003). Litter could thereby importantly contribute to BVOC emissions, O₃ and SOA
600 formation in this region.

601 **Acknowledgements**

602 This study was performed in the frame of the CORVOLIM project (Inventaire d'émissions de
603 Composés ORganiques VOLatils par la Litière de la végétation Méditerranéenne : application à la
604 qualité de l'air), funded by the program BioDivMeX (BioDiversity of the Mediterranean
605 eXperiment) of the metaprogram MISTRALS (Mediterranean Integrated Studies at Regional And
606 Local Scales). The project LitterVOC from the ECCOREV program (ECosystèmes CONTinentaux
607 et Risques EnVironnementaux) and the ANR (Agence Nationale de Recherche) SecPrime² also
608 funded this study. We thank Alain Tonetto for support using SEM and Sylvie Dupouyet for litter
609 collection.

610 **Data availability :**

611 The data used in this study is available on demand.

612 **Bibliography**

- 613 Adams, R.P., 2007. Identification of essential oil components by gas chromatography/mass spectrometry. Allured
614 publishing corporation Carol Stream, Illinois.
- 615 Arianoutsou, M., 1989. Timing of litter production in a maquis ecosystem of north-eastern Greece. *Acta Oecoloecol.*
616 *Plant.* 10, 371–378.
- 617 Arianoutsou, M., Radea, C., 2000. Litter production and decomposition in *Pinus halepensis* forests. *Ecol. Biogeogr.*
618 *Manag. Pinus Halepensis Pinus Brutia For. Ecosyst. Mediterr. Basin Blackhuys Publ. Leiden Neth.* 183–190.
- 619 Asensio, D., Peñuelas, J., Ogaya, R., Llusà, J., 2007. Seasonal soil VOC exchange rates in a Mediterranean holm oak
620 forest and their responses to drought conditions. *Atmos. Environ.* 41, 2456–2466.
- 621 Atkinson, R., Arey, J., 2003. Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review.
622 *Atmos. Environ.* 37, 197–219.
- 623 Barik, A., Bhattacharya, B., Laskar, S., Banerjee, T.C., 2004. The determination of n-alkanes in the cuticular wax of
624 leaves of *Ludwigia adscendens* L. *Phytochem. Anal.* 15, 109–111.
- 625 Bellot, J., Sánchez, J.R., Lledó, M.J., Martínez, P., Escarré, A., 1992. Litterfall as a measure of primary production in
626 Mediterranean holm-oak forest, in: *Quercus Ilex L. Ecosystems: Function, Dynamics and Management,*
627 *Advances in Vegetation Science.* Springer Netherlands, pp. 69–76.
- 628 Bondada, B.R., Oosterhuis, D.M., Murphy, J.B., Kim, K.S., 1996. Effect of water stress on the epicuticular wax
629 composition and ultrastructure of cotton (*Gossypium hirsutum* L.) leaf, bract, and boll. *Environ. Exp. Bot.*
630 36, 61–69.
- 631 Bourtsoukidis, E., Williams, J., Kesselmeier, J., Jacobi, S., Bonn, B., 2014. From emissions to ambient mixing ratios:
632 online seasonal field measurements of volatile organic compounds over a Norway spruce-dominated forest
633 in central Germany. *Atmospheric Chem. Phys.* 14, 6495–6510.
- 634 Bruns, E.A., El Haddad, I., Keller, A., Klein, F., Kumar, N.K., Pieber, S.M., Corbin, J.C., Slowik, J.G., Brune, W.H.,
635 Baltensperger, U., Prévôt, A.S.H., 2015. Inter-comparison of laboratory smog chamber and flow reactor
636 systems on organic aerosol yield and composition. *Atmos. Meas. Tech. Discuss.* 8, 309–352.
- 637 Chaignaud, P., Morawe, M., Besaury, L., Kröber, E., Vuilleumier, S., Bringel, F., Kolb, S., 2018. Methanol
638 consumption drives the bacterial chloromethane sink in a forest soil. *ISME J.* 12, 2681.
- 639 Chomel, M., Guittonny-Larchevêque, M., Fernandez, C., Gallet, C., DesRochers, A., Paré, D., Jackson, B.G., Baldy,
640 V., 2016. Plant secondary metabolites: a key driver of litter decomposition and soil nutrient cycling. *J. Ecol.*
641 104, 1527–1541.
- 642 Derendorp, L., Holzinger, R., Wishkerman, A., Keppler, F., Röckmann, T., 2011. Methyl chloride and C₂–C₅
643 hydrocarbon emissions from dry leaf litter and their dependence on temperature. *Atmos. Environ.* 45, 3112–
644 3119.

645 Di Carlo, P., Brune, W.H., Martinez, M., Harder, H., Leshner, R., Ren, X., Thornberry, T., Carroll, M.A., Young, V.,
646 Shepson, P.B., Riemer, D., Apel, E., Campbell, C., 2004. Missing OH reactivity in a forest: evidence for
647 unknown reactive biogenic VOCs. *Science* 304, 722–725.

648 Derwent, R.G., Jenkin, M.E., Utembe, S.R., Shallcross, D.E., Murrells, T.P., Passant, N.R., 2010. Secondary organic
649 aerosol formation from a large number of reactive man-made organic compounds. *Sci. Total Environ.* 408,
650 3374–3381.

651 Dob, T., Dahmane, D., Chelghoum, C., 2006. Essential oil composition of *Juniperus oxycedrus*. Growing in Algeria.
652 *Pharm. Biol.* 44, 1–6.

653 Dominutti, P., Keita, S., Bahino, J., Colomb, A., Lioussé, C., Yoboué, V., Galy-Lacaux, C., Morris, E., Bouvier, L.,
654 Sauvage, S., Borbon, A., 2019. Anthropogenic VOCs in Abidjan, southern West Africa: From source
655 quantification to atmospheric impacts. *Atmospheric Chem. Phys.* 19, 11721–11741.

656 Faiola, C.L., VanderSchelden, G.S., Wen, M., Elloy, F.C., Cobos, D.R., Watts, R.J., Jobson, B.T., VanReken, T.M.,
657 2014. SOA formation potential of emissions from soil and leaf litter. *Environ. Sci. Technol.* 48, 938–946.

658 Fehsenfeld, F., Calvert, J., Fall, R., Goldan, P., Guenther, A.B., Hewitt, C.N., Lamb, B., Liu, S., Trainer, M., Westberg,
659 H., Zimmerman, P., 1992. Emissions of volatile organic compounds from vegetation and the implications for
660 atmospheric chemistry. *Glob. Biogeochem. Cycles* 6, 389–430.

661 Fernandez, C., Bousquet-Mélou, A., Prévosto, B., 2013. Répartition dans le bassin méditerranéen et en France, in: *Le*
662 *Pin d'Alep en France, Guide pratique.* pp. 23–27.

663 Genard-Zielinski, A.-C., Boissard, C., Fernandez, C., Kalogridis, C., Lathièrre, J., Gros, V., Bonnaire, N., Ormeño, E.,
664 2014. Variability of BVOC emissions from a Mediterranean mixed forest in southern France with a focus on
665 *Quercus pubescens*. *Atmospheric Chem. Phys. Discuss.* 14, 17225–17261.

666 Gray, C.M., Monson, R.K., Fierer, N., 2014. Biotic and abiotic controls on biogenic volatile organic compound fluxes
667 from a subalpine forest floor. *J. Geophys. Res. Biogeosciences* 119, 547–556.

668 Gray, C.M., Monson, R.K., Fierer, N., 2010. Emissions of volatile organic compounds during the decomposition of
669 plant litter. *J. Geophys. Res.* 115, G03015.

670 Greenberg, J.P., Asensio, D., Turnipseed, A., Guenther, A.B., Karl, T., Gochis, D., 2012. Contribution of leaf and
671 needle litter to whole ecosystem BVOC fluxes. *Atmos. Environ.* 59, 302–311.

672 Griffin, R.J., Cocker, D.R., Seinfeld, J.H., Dabdub, D., 1999. Estimate of global atmospheric organic aerosol from
673 oxidation of biogenic hydrocarbons. *Geophys. Res. Lett.* 26, 2721–2724.

674 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P.I., Geron, C., 2006. Estimates of global terrestrial
675 isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature). *Atmospheric*
676 *Chem. Phys.* 6, 3181–3210.

677 Hart, S.C., Firestone, M.K., Paul, E.A., 1992. Decomposition and nutrient dynamics of ponderosa pine needles in a
678 Mediterranean-type climate. *Can. J. For. Res.* 22, 306–314.

679 Hartz, K.E.H., Rosenørn, T., Ferchak, S.R., Raymond, T.M., Bilde, M., Donahue, N.M., Pandis, S.N., 2005. Cloud
680 condensation nuclei activation of monoterpene and sesquiterpene secondary organic aerosol. *J. Geophys. Res.*
681 *Atmospheres* 110, D14208.

682 He, C., Murray, F., Lyons, T., 2000. Monoterpene and isoprene emissions from 15 eucalyptus species in Australia.
683 *Atmos. Environ.* 34, 645–655.

684 Helmig, D., Ortega, J., Guenther, A., Herrick, J.D., Geron, C., 2006. Sesquiterpene emissions from loblolly pine and
685 their potential contribution to biogenic aerosol formation in the Southeastern US. *Atmos. Environ.* 40, 4150–
686 4157.

687 Hennig, S., Gülz, P.-G., Hängst, K., 1988. Organ specific composition of epicuticular waxes of *Cistus albidus* L.,
688 *Cistaceae*. *Z. Für Naturforschung C* 43, 806–812.

689 Hoffmann, T., Odum, J.R., Bowman, F., Collins, D., Klockow, D., Flagan, R.C., Seinfeld, J.H., 1997. Formation of
690 organic aerosols from the oxidation of biogenic hydrocarbons. *J. Atmos. Chem.* 26, 189–222.

691 Huang, G., Brook, R., Crippa, M., Janssens-Maenhout, G., Schieberle, C., Dore, C., Guizzardi, D., Muntean, M.,
692 Schaaf, E., Friedrich, R., 2017. Speciation of anthropogenic emissions of non-methane volatile organic
693 compounds: a global gridded data set for 1970–2012. *Atmospheric Chem. Phys.* 17, 7683–7701.

694 Isidorov, V., Povarov, V., Stepanov, A., 1999. Forest Soil Cover: VOC sink or source? *WIT Trans. Ecol. Environ.* 36.

695 Isidorov, V., Tyszkiewicz, Z., Pirożnikow, E., 2016. Fungal succession in relation to volatile organic compounds
696 emissions from Scots pine and Norway spruce leaf litter-decomposing fungi. *Atmos. Environ.* 131, 301–306.

697 Isidorov, V.A., Smolewska, M., Purzyńska-Pugaciewicz, A., Tyszkiewicz, Z., 2010. Chemical composition of volatile
698 and extractive compounds of pine and spruce leaf litter in the initial stages of decomposition. *Biogeosciences*
699 7, 2785–2794.

700 Isidorov, V.A., Vinogorova, V.T., Rafałowski, K., 2003. HS-SPME analysis of volatile organic compounds of

701 coniferous needle litter. *Atmos. Environ.* 37, 4645–4650.

702 Jaoui, M., Kamens, R.M., 2003. Gaseous and particulate oxidation products analysis of a mixture of α -pinene + β -
703 pinene/O₃/Air in the Absence of Light and α -pinene + β -pinene/NO_x/Air in the Presence of Natural Sunlight.
704 *J. Atmos. Chem.* 44, 259–297.

705 Joensuu, J., Altimir, N., Hakola, H., Rostás, M., Raivonen, M., Vestenius, M., Aaltonen, H., Riederer, M., Bäck, J.,
706 2016. Role of needle surface waxes in dynamic exchange of mono- and sesquiterpenes. *Atmospheric Chem.*
707 *Phys.* 16, 7813–7823.

708 Johnson, E.D., 1926. A comparison of the juvenile and adult leaves of *Eucalyptus globulus*. *New Phytol.* 25, 202–212.

709 Keenan, T., Niinemets, Ü., Sabate, S., Gracia, C., Peñuelas, J., 2009. Process based inventory of isoprenoid emissions
710 from European forests: model comparisons, current knowledge and uncertainties. *Atmospheric Chem. Phys.*
711 *Discuss.* 9, 6147–6206.

712 Kesselmeier, J., Staudt, M., 1999. Biogenic Volatile Organic Compounds (VOC): an overview on emission, physiology
713 and ecology. *J. Atmospheric Chem.* 33, 23–88.

714 Kramshøj, M., Vedel-Petersen, I., Schollert, M., Rinnan, Å., Nymand, J., Ro-Poulsen, H., Rinnan, R., 2016. Large
715 increases in Arctic biogenic volatile emissions are a direct effect of warming. *Nat. Geosci.* 9, 349–352.

716 Krishna, M.P., Mohan, M., 2017. Litter decomposition in forest ecosystems: a review. *Energy Ecol. Environ.* 2, 236–
717 249.

718 Kroll, J.H., Ng, N.L., Murphy, S.M., Flagan, R.C., Seinfeld, J.H., 2005. Secondary organic aerosol formation from
719 isoprene photooxidation under high-NO_x conditions. *Geophys. Res. Lett.* 32, L18808.

720 Lange, B.M., 2015. The evolution of plant secretory structures and emergence of terpenoid chemical diversity. *Annu.*
721 *Rev. Plant Biol.* 66, 139–159.

722 Lee, A., Goldstein, A.H., Kroll, J.H., Ng, N.L., Varutbangkul, V., Flagan, R.C., Seinfeld, J.H., 2006. Gas-phase
723 products and secondary aerosol yields from the photooxidation of 16 different terpenes. *J. Geophys. Res.*
724 *Atmospheres* 111, D17305.

725 Leff, J.W., Fierer, N., 2008. Volatile organic compound (VOC) emissions from soil and litter samples. *Soil Biol.*
726 *Biochem.* 40, 1629–1636.

727 Lemfack, M.C., Gohlke, B.-O., Toguem, S.M.T., Preissner, S., Piechulla, B., Preissner, R., 2018. mVOC 2.0: a
728 database of microbial volatiles. *Nucleic Acids Res.* 46, D1261–D1265.

729 Llusia, J., Penuelas, J., 2000. Seasonal patterns of terpene content and emission from seven mediterranean woody
730 species in field conditions. *Am. J. Bot.* 87, 133–140.

731 Maffei, M., 1994. Discriminant analysis of leaf wax alkanes in the Lamiaceae and four other plant families.
732 *Biochemical Systematics and Ecology* 22, 711–728.

733 Mäki, M., Heinonsalo, J., Hellén, H., Bäck, J., 2017. Contribution of understorey vegetation and soil processes to
734 boreal forest isoprenoid exchange. *Biogeosciences* 14, 1055–1073.

735 Mancuso, S., Taiti, C., Bazihizina, N., Costa, C., Menesatti, P., Giagnoni, L., Arenella, M., Nannipieri, P., Renella, G.,
736 2015. Soil volatile analysis by proton transfer reaction-time of flight mass spectrometry (PTR-TOF-MS).
737 *Appl. Soil Ecol.* 86, 182–191.

738 Matić, S., Stanić, S., Mihailović, M., Bogojević, D., 2016. *Cotinus coggygria* Scop.: An overview of its chemical
739 constituents, pharmacological and toxicological potential. *Saudi J. Biol. Sci.* 23, 452–461.

740 Messina, P., Lathiere, J., Sindelarova, K., Vuichard, N., Granier, C., Ghattas, J., Cozic, A., Hauglustaine, D.A., 2016.
741 Global biogenic volatile organic compound emissions in the ORCHIDEE and MEGAN models and sensitivity
742 to key parameters. *Atmospheric Chem. Phys.* 16, 14169–14202.

743 Metcalfe, C.R., Chalk, L., 1988. Anatomy of the dicotyledons. Volume I. Systematic anatomy of the leaf and stem,
744 with a brief history of the subject. *Anat. Dicotyledons Vol. Syst. Anat. Leaf Stem Brief Hist. Subj.*

745 Misra, G., Pavlostathis, S.G., Perdue, E.M., Araujo, R., 1996. Aerobic biodegradation of selected monoterpenes. *Appl.*
746 *Microbiol. Biotechnol.* 45, 831–838.

747 Navarro, J.C.A., Smolander, S., Struthers, H., Zorita, E., Ekman, A.M.L., Kaplan, J.O., Guenther, A., Arneth, A.,
748 Riipinen, I., 2014. Global emissions of terpenoid VOCs from terrestrial vegetation in the last millennium. *J.*
749 *Geophys. Res. Atmos.* 119, 6867–6885.

750 Nielsen, J.A., Frew, R.D., Whigam, P.A., Callaway, R.M., Dickinson, K.J.M., 2015. Germination and growth
751 responses of co-occurring grass species to soil from under invasive *Thymus vulgaris*. *Allelopathy J.* 35.

752 Nunes, T.V., Pio, C.A., 2001. Emission of volatile organic compounds from Portuguese eucalyptus forests.
753 *Chemosphere - Glob. Change Sci.* 3, 239–248.

754 Ormeño, E., Céspedes, B., Sánchez, I.A., Velasco-García, A., Moreno, J.M., Fernandez, C., Baldy, V., 2009. The
755 relationship between terpenes and flammability of leaf litter. *For. Ecol. Manag.* 257, 471–482.

756 Ormeño, E., Fernandez, C., Bousquet-Mélou, A., Greff, S., Morin, E., Robles, C., Vila, B., Bonin, G., 2007a.

757 Monoterpene and sesquiterpene emissions of three Mediterranean species through calcareous and siliceous
758 soils in natural conditions. *Atmos. Environ.* 41, 629–639.

759 Ormeño, E., Fernandez, C., Mévy, J.-P., 2007b. Plant coexistence alters terpene emission and content of Mediterranean
760 species. *Phytochemistry* 68, 840–852.

761 Ormeño, E., Viros, J., Mévy, J.-P., Tonetto, A., Saunier, A., Bousquet-Mélou, A., Fernandez, C., 2020. Exogenous
762 isoprene confers physiological benefits in a negligible isoprene emitter (*Acer monspessulanum* L.) under
763 water deficit. *Plants* 9, 159.

764 Owen, S., Boissard, C., Street, R.A., Duckham, S.C., Csiky, O., Hewitt, C.N., 1997. Screening of 18 Mediterranean
765 plant species for volatile organic compound emissions. *Atmos. Environ.* 31, 101–117.

766 Owen, S.M., Boissard, C., Hewitt, C.N., 2001. Volatile organic compounds (VOCs) emitted from 40 Mediterranean
767 plant species: VOC speciation and extrapolation to habitat scale. *Atmos. Environ.* 35, 5393–5409.

768 Owen, S.M., Hewitt, C.N., 2000. Extrapolating branch enclosure measurements to estimates of regional scale biogenic
769 VOC fluxes in the northwestern Mediterranean basin. *J. Geophys. Res. Atmospheres* 105, 11573–11583.

770 Parra, R., Gassó, S., Baldasano, J.M., 2004. Estimating the biogenic emissions of non-methane volatile organic
771 compounds from the North Western Mediterranean vegetation of Catalonia, Spain. *Sci. Total Environ.* 329,
772 241–259.

773 Peñuelas, J., Asensio, D., Tholl, D., Wenke, K., Rosenkranz, M., Piechulla, B., Schnitzler, J.P., 2014. Biogenic volatile
774 emissions from the soil. *Plant Cell Environ.* 37, 1866–1891.

775 Peñuelas, J., Llusia, J., 1997. Effects of Carbon Dioxide, Water Supply, and Seasonality on Terpene Content and
776 Emission by *Rosmarinus officinalis*. *J. Chem. Ecol.* 23, 979–993.

777 Peñuelas, J., Marino, G., Llusia, J., Morfopoulos, C., Farré-Armengol, G., Filella, I., 2013. Photochemical reflectance
778 index as an indirect estimator of foliar isoprenoid emissions at the ecosystem level. *Nat. Commun.* 4, 1–10.

779 Polade, S.D., Pierce, D.W., Cayan, D.R., Gershunov, A., Dettinger, M.D., 2014. The key role of dry days in changing
780 regional climate and precipitation regimes. *Sci. Rep.* 4, 1–8.

781 Praplan, A.P., Tykkä, T., Chen, D., Boy, M., Taipale, D., Vakkari, V., Zhou, P., Petäjä, T., Hellén, H., 2019. Long-
782 term total OH reactivity measurements in a boreal forest. *Atmospheric Chem. Phys.* 19, 14431–14453.

783 Préndez, M., Carvajal, V., Corada, K., Morales, J., Alarcón, F., Peralta, H., 2013. Biogenic volatile organic compounds
784 from the urban forest of the Metropolitan Region, Chile. *Environmental Pollution* 183, 143–150.

785 Pio, C.A., Silva, P.A., Cerqueira, M.A., Nunes, T.V., 2005. Diurnal and seasonal emissions of volatile organic
786 compounds from cork oak (*Quercus suber*) trees. *Atmos. Environ.* 39, 1817–1827.

787 Quézel, P., Médail, F., 2003. *Ecologie et biogéographie des forêts du bassin méditerranéen*. Elsevier, Paris.

788 Ramirez, K.S., Lauber, C.L., Fierer, N., 2010. Microbial consumption and production of volatile organic compounds
789 at the soil-litter interface. *Biogeochemistry* 99, 97–107.

790 R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical
791 Computing, Vienna, Austria. URL <https://www.R-project.org/>.

792 Rinnan, R., Albers, C.N., 2020. Soil uptake of volatile organic compounds: ubiquitous and underestimated? *J.*
793 *Geophys. Res. Biogeosciences* 125, e2020JG005773.

794 Robert, B., Caritat, A., Bertoni, G., Vilar, L., Molinas, M., 1996. Nutrient content and seasonal fluctuations in the leaf
795 component of coark-oak (*Quercus suber* L.) litterfall. *Vegetatio* 122, 29–35.

796 Rodriguez-Ramirez, N., Santonja, M., Baldy, V., Ballini, C., Montès, N., 2017. Shrub species richness decreases
797 negative impacts of drought in a Mediterranean ecosystem. *J. Veg. Sci.* 28, 985–996.

798 Romero, B., Fernandez, C., Lecareux, C., Ormeño, E., Ganteaume, A., 2019. How terpene content affects fuel
799 flammability of wildland–urban interface vegetation. *Int. J. Wildland Fire* 28, 614–627.

800 Sánchez-Osorio, I., López-Pantoja, G., Tapias, R., Pareja-Sánchez, E., Domínguez, L., 2019. Monoterpene emission
801 of *Quercus suber* L. highly infested by *Cerambyx welensii* Küster. *Ann. For. Sci.* 76, 98.

802 Santonja, M., Bousquet-Mélou, A., Greff, S., Ormeño, E., Fernandez, C., 2019. Allelopathic effects of volatile organic
803 compounds released from *Pinus halepensis* needles and roots. *Ecol. Evol.* 9, 8201–8213.

804 Santonja, M., Fernandez, C., Gauquelin, T., Baldy, V., 2015. Climate change effects on litter decomposition: intensive
805 drought leads to a strong decrease of litter mixture interactions. *Plant Soil* 393, 69–82.

806 Santonja, M., Foucault, Q., Rancon, A., Gauquelin, T., Fernandez, C., Baldy, V., Mirleau, P., 2018. Contrasting
807 responses of bacterial and fungal communities to plant litter diversity in a Mediterranean oak forest. *Soil Biol.*
808 *Biochem.* 125, 27–36.

809 Saunier, A., Ormeño, E., Boissard, C., Wortham, H., Temime-Roussel, B., Lecareux, C., Armengaud, A., Fernandez,
810 C., 2017a. Effect of mid-term drought on *Quercus pubescens* BVOCs’ emission seasonality and their
811 dependency on light and/or temperature. *Atmospheric Chem. Phys.* 17, 7555–7566.

812 Saunier, A., Ormeño, E., Havaux, M., Wortham, H., Ksas, B., Temime-Roussel, B., Blande, J.D., Lecareux, C., Mévy,

813 J.-P., Bousquet-Mélou, A., Gauquelin, T., Fernandez, C., 2018. Resistance of native oak to recurrent drought
814 conditions simulating predicted climatic changes in the Mediterranean region. *Plant Cell Environ.* 41, 2299–
815 2312.

816 Saunier, A., Ormeño, E., Wortham, H., Temime-Roussel, B., Lecareux, C., Boissard, C., Fernandez, C., 2017b. chronic
817 drought decreases anabolic and catabolic BVOC emissions of *Quercus pubescens* in a Mediterranean forest.
818 *Front. Plant Sci.* 8.

819 Schade, G.W., Goldstein, A.H., 2001. Fluxes of oxygenated volatile organic compounds from a ponderosa pine
820 plantation. *J. Geophys. Res. Atmospheres* 106, 3111–3123.

821 Sharkey, T.D., Yeh, S., 2001. Isoprene emission from plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52, 407–436.

822 Simon, V., Luchetta, L., Torres, L., 2001. Estimating the emission of volatile organic compounds (VOC) from the
823 French forest ecosystem. *Atmos. Environ., Selected Papers Presented at the Venice Conference* 35, S115–
824 S126.

825 Spittler, M., Barnes, I., Bejan, I., Brockmann, K.J., Benter, Th., Wirtz, K., 2006. Reactions of NO₃ radicals with
826 limonene and α -pinene: Product and SOA formation. *Atmos. Environ., Regional biogenic emissions of*
827 *reactive volatile organic compounds (BVOC) from forests: Process studies, modelling and validation*
828 *experiments (BEWA2000)* 40, 116–127.

829 Staudt, M., Lhoutellier, L., 2011. Monoterpene and sesquiterpene emissions from *Quercus coccifera* exhibit interacting
830 responses to light and temperature. *Biogeosciences* 8, 2757–2771.

831 Street, R.A., Owen, S., Duckham, S.C., Boissard, C., Hewitt, C.N., 1997. Effect of habitat and age on variations in
832 volatile organic compound (VOC) emissions from *Quercus ilex* and *Pinus pinea*. *Atmos. Environ.* 31, 89–
833 100.

834 Svendsen, S.H., Lindwall, F., Michelsen, A., Rinnan, R., 2016. Biogenic volatile organic compound emissions along
835 a high arctic soil moisture gradient. *Sci. Total Environ.* 573, 131–138.

836 Tang, J., Schurgers, G., Rinnan, R., 2019. Process Understanding of Soil BVOC Fluxes in Natural Ecosystems: A
837 Review. *Rev. Geophys.* 57, 966–986.

838 Tasoglou, A., Pandis, S.N., 2015. Formation and chemical aging of secondary organic aerosol during the β -
839 caryophyllene oxidation. *Atmospheric Chem. Phys.* 15, 6035–6046.

840 Vanhatalo, A., Ghirardo, A., Juurola, E., Schnitzler, J.-P., Zimmer, I., Hellén, H., Hakola, H., Bäck, J., 2018. Long-
841 term dynamics of monoterpene synthase activities, monoterpene storage pools and emissions in boreal Scots
842 pine. *Biogeosciences* 15, 5047–5060.

843 Vasic, P.S., Dubak, D.V., 2012. Anatomical analysis of red Juniper leaf (*Juniperus oxycedrus*) taken from Kopaonik
844 Mountain, Serbia. *Turk. J. Bot.* 36, 473–479.

845 Wagner, G.J., 1991. Secreting glandular trichomes: more than just hairs. *Plant Physiol.* 96, 675–679.

846 Warneke, C., Karl, T., Judmaier, H., Hansel, A., Jordan, A., Lindinger, W., Crutzen, P.J., 1999. Acetone, methanol,
847 and other partially oxidized volatile organic emissions from dead plant matter by abiological processes:
848 Significance for atmospheric HO_x chemistry. *Glob. Biogeochem. Cycles* 13, 9–17.

849 Wilkinson, S.C., Anderson, J.M., Scardelis, S.P., Tisiafouli, M., Taylor, A., Wolters, V., 2002. PLFA profiles of
850 microbial communities in decomposing conifer litters subject to moisture stress. *Soil Biol. Biochem.* 34, 189–
851 200.

852 Wilt, F.M., Miller, G.C., Everett, R.L., Hackett, M., 1993. Monoterpene concentrations in fresh, senescent, and
853 decaying foliage of single leaf pinyon (*Pinus monophylla* Torr. & Frem.: Pinaceae) from the western Great
854 Basin. *J. Chem. Ecol.* 19, 185–194.