

The decline of Fraxinus angustifolia Vahl in a Mediterranean salt meadow: Chlorophyll fluorescence measurements in long-term field experiment

Jean-Philippe Mevy, Frédéric Guibal, Caroline Lecareux, Franco Miglietta

▶ To cite this version:

Jean-Philippe Mevy, Frédéric Guibal, Caroline Lecareux, Franco Miglietta. The decline of Fraxinus angustifolia Vahl in a Mediterranean salt meadow: Chlorophyll fluorescence measurements in long-term field experiment. Estuarine, Coastal and Shelf Science, 2020, pp.107068. 10.1016/j.ecss.2020.107068 . hal-02980230

HAL Id: hal-02980230 https://amu.hal.science/hal-02980230

Submitted on 3 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

13004 Marseille, France Author for correspondence: Jean-Philippe Mevy Tel: +33 0413550766 Email; jean-philippe.mevy@imbe.fr

Abstract: 15

The ecological transition zones also known as ecotones are specific ecosystems but how global change 16 17 will affect their biodiversity and functioning is poorly understood. In the case of the areas at the 18 interface between riparian forest and salt meadows the biodiversity conservation remains a major 19 issue. The aim of this work was to evaluate the effects of a salty environment on Fraxinus angustifolia 20 a non-halophyte species. For this, the photochemistry of PSII was in situ explored during the summer 21 between 2007 and 2016. The predawn maximum quantum efficiency of PSII (Fv/Fm) was always 22 greater than 0.8 which indicates an absence of chronic photoinhibition induced by salinity. During these years, the light kinetic of the actual quantum yield of PSII (Φ_{PSII}) was studied. It appears that 23 this yield is generally higher in salty environments than in the riparian forest. Multivariate analysis, 24 25 showed that the dissipation regulated light energy faction (Φ_{NPO}), the photochemical quenching (qP) 26 and the Fd/Fs ratio constitute the main explanatory variables of Φ_{PSII} changes. The subsequent twoway-ANCOVA carried out demonstrated that Φ_{NPQ} was weaker in saline environments, suggesting a 27 28 better efficiency of photosystem collecting antennas. Meanwhile, the analysis of photosynthetic 29 pigments did not exhibited a salt effect except the significant accumulation of chlorophyll a in 2016. 30 In order to address the question of the salt-related toxicity, sodium ion contents in the leaves were found to be identical in the two sites, suggesting the establishment of an energy-dependent mechanism 31 32 of salt exclusion/retention in roots. The growth rings of *F.angustifolia* in saline areas highly follows the annual water deficit changes. Between 2007 and 2016, the biomass in the salt meadow was higher 33 with a peak in 2011 representing 60% of the biomass of control individuals. But, since 2011 a 34

The decline of Fraxinus angustifolia Vahl in a Mediterranean salt meadow: Chlorophyll 1 fluorescence measurements in long-term field experiment. 2

J.P. Mevy^{1*}, F. Guibal¹, C. Lecareux¹ and F. Miglietta^{2,3}. 3

1 Aix-Marseille University, Avignon University, CNRS, IRD, IMBE, Marseille, France 4

5 2 Institute of Biometeorology, National Research Council (CNR- IBIMET), Via Caproni 8,

50145 Firenze, Italy 6

7 3 IMèRA, Institut d'Etudes Avancées de l'Université Aix-Marseille, 2 Place Le Verrier,

- 8
- 9

10

11

12

13

35 continuous decline of growth was recorded. Water conditions seem to constitute a limiting factor even 36 if the photosynthetic apparatus generates the excess metabolic energy necessary to cope with the saline 37 toxicity. This study thus highlighted the plasticity of the photochemistry of the PSII depending to the 38 environmental conditions of the salt meadows. It also shows the vulnerability of an ecotone through 39 the decline in the productivity of a glycophyte with climate change. This augurs for changes in the 40 composition of plant communities which is in favor of strict halophytes.

41 Key words: *Fraxinus angustifolia*; salt stress; PSII photochemistry; salt meadow; water
42 deficit; ecophysiology.

43

44 **1. Introduction**

Global change projections of wetland ecosystems such as salt marshes include sea-level rise 45 and extreme climatic events (Thorne et al., 2012). Salt meadows spread over narrow 46 transitional areas between sea and terrestrial environments which are considered as high-risk 47 zones subject to climate change impacts (Thorne et al., 2012). In these ecosystems, several 48 taxonomic groups of plants coexist with salt requirements which are highly contrasted. They 49 are divided into euhalophytes and facultative halophytes but typically an halophyte is defined 50 as a plant that grows in soil conditions of more than 200 mM NaCl concentration (Flowers 51 and Colmer, 2008). Besides salinity conditions, biotic factors such as competition between 52 53 individuals are also to be considered in terms of saltmarshes functioning (Pennings et al., 2005). Overall, it appears that the hydro-period regime that depends on specific precipitation 54 and evaporation conditions is one of the main factors regulating the creation of niches for 55 wetland plant species (Foti et al., 2012). 56

57 The question of the spatial distribution of species between the seashore and the boundary of 58 non-halophytic terrestrial vegetation has been addressed by several authors (Pennings et al., 2005; Silvestri et al., 2005). For instance taking into account the elevation gradient, four 59 clusters reflecting the composition of plant community were shown (Bang et al., 2018). 60 However, the scarcity of studies focussing on the salt marsh-terrestrial boundaries was 61 underlined (Traut, 2005; Veldkornet et al., 2015). Indeed, the border between the salt marsh 62 and the adjacent upland (terrestrial vegetation) occupies a narrow surface also recognized as 63 ecotone. These habitats are submitted to particular abiotic conditions leading to a plant 64 community assembly that is specific to one or both of the adjacent ecosystems. In the 65 Mediterranean region and in the Bouches-du-Rhône in particular, salt meadows develop at the 66

67 confluence of rivers and the Etang de Berre. This is the case for the river Arc which exhibits a 68 steep fall of the presence of trees from the riparian forest in the upland. Among the riparian 69 woody species only few specimens of *Fraxinus angustifolia* Vahl (narrow-leaved ash) were 70 observed in the salt marsh upland transition zone possibly in the area so-called ecotone. 71 Hence, information on species with narrow biotope tolerance might best characterize the 72 effects of climate change (James and Zedler, 2000).

73

F. angustifolia belongs to the family Oleaceae and is widespread across Southern Europe. In 74 75 Mediterranean basin the plant is often encountered in riparian corridors. In Croatia for example, narrow-leaved ash is considered as one of the most important tree species of 76 77 lowland floodplain forests and its sensitivity to drought was shown (Drvodelic et al., 2016). But, how F. angustifolia will cope with salinity and drought expected with climate change is 78 poorly explored. More generally exploring the mechanisms of glycophytes adaptation to 79 salinity is becoming a great challenge because soil salinization is recognized as a global 80 problem resulting in soil degradation, arable land surface decrease and ecological 81 environmental disturbance (Hu et al., 2019). By comparing two Solanum species, one 82 glycophyte and the other salt-tolerant, it has been shown that ethylene induction plays a key 83 role in salinity tolerance through an improved stomatal conductance, an efficient osmotic 84 adjustment and the increase of Na⁺ accumulation in shoots (Gharbi et al., 2017). In roots of 85 the glycophyte Arabidopsis thaliana (L.) Heynh., salt stress triggered principally defence 86 proteins as those involved in ROS scavenging whereas for the xero-halophyte Zygophyllum 87 *xanthoxylum* Engl. Ex Dippel, Na⁺ uptake was improved in addition with metabolic shifts 88 towards de novo synthesis of glucose in order to sustain growth (Chai et al., 2019). Similarly, 89 90 retention of sodium cations in roots thus preventing their translocation to shoots is considered as a key adaptive mechanism of glycophytes to salt stress (Assaha et al., 2017; Hasegawa, 91 2013). Root sequestration of sodium cations not only improves the water balance by adequate 92 osmotic adjustment but also preserves the photosynthetic machinery from ionic toxicity. Thus, 93 from the xero-halophyte Nitraria retusa (Forssk.) Asch., PSII photochemistry and carotenoid 94 content were not affected by salinity whereas a significant reduction in pigment 95 96 concentrations and chlorophyll fluorescence parameters were observed from the glycophyte Medicago arborea L. (Boughalleb et al., 2009). This led to suggest that the decrease of the 97 photosynthetic activity is likely the result of damages in the photosynthetic apparatus rather 98 than factors driving stomatal closure. Other authors have also shown that the maximum 99 100 quantum efficiency of PSII is highly reduced for rice salt-sensitive cultivars consecutively to

salt stress application as compared to salt-tolerant cultivars. Similarly, the minimum 101 fluorescence intensity (F0), the actual quantum efficiency (PSII) and the None-Photochemical 102 Quenching (NPQ) were identified as the main fluorescence parameters that change depending 103 on the intensity of salt stress (Tsai et al., 2019). Besides the ionic toxicity induced by soil 104 salinity, water availability also constitutes a major constraint for glycophytes survival in 105 saltmarshes. Although, chlorophyll a fluorescence has been used as a signal for monitoring 106 plants response to drought in several reports (Dalberto et al., 2017; Vaz et al., 2016; Wang et 107 al., 2016) few works deal with both drought and salinity effects. Moreover, plant *in situ* often 108 109 experience multifactorial stress conditions.

Understanding how a glycophyte may stabilize its photosynthesis activity also maintaining its 110 ionic homeostasis under high soil salinity can help provide insight in terms of plant breeding 111 and ecological management. In this report we evaluate the ecological fitness of a glycophyte 112 113 F. angustifolia native to the saltmarsh/upland transition zone also termed ecotone upon a highly sensitive to global climatic change variables. Second, we hypothesized that the 114 predicted increase warming and summer drought in Mediterranean basin would lead to more 115 NaCl accumulation requiring additional metabolic energy to sustain growth and defence 116 molecules biosynthesis. Finally we considered that the global change would drive profound 117 changes in the assemblages of plant communities into the ecotones of the deltaic zones of the 118 Mediterranean rivers. 119

120 **2.** Material and methods

121 *2.1 The study site*

The area that was the subject of this study is located in the Arc delta ecotone and in the 122 riparian forest of the same river (Berre-l'Etang, France). The Arc delta occupies an area of 123 190 ha. It is associated with hydromorphic soils covered with a halophilic vegetation, which 124 extend about 700 m wide. Upstream the deltaic zone (1.4 km), the riparian forest is mainly 125 constituted by ash, poplar and white willow. The ecotone was characterized by the occurrence 126 127 of both euhalophytes and facultative halophytes as well as some specimens of *Fraxinus* angustifolia, namely: Sueda vera Forssk. ex J.F.Gmel., Limonium narbonense Mill., 128 Limonium cuspidatum (Delort) Erben, Chenopodium chenopodioides (L.) Aellen, Tamarix 129 gallica L., Obione portulacoides (=Atriplex portulacoides L.) and Plantago coronopus L. The 130 meteorological data were obtained from the nearest station namely Marignane which is 131

located at about 11 km away as the crow flies. Five to six trees (*F. angustifolia*) of similar
height and diameter were chosen in each zone during this study.

134 2.2 Modulated chlorophyll fluorescence parameters monitoring

Fluorescence data were obtained using Hansatech fluorometer (FMS2). Leaves (3 per tree) 135 were dark-adapted for 30 min using the dark-adaptation clip prior to the application of a low 136 pulsed amber light intensity to determine the minimum fluorescence (F_0) followed by the 137 application of a saturating pulse for the maximum fluorescence measurement (F_m) such that 138 all primary quinone electron acceptors in PSII are closed (Q_A fully reduced). These data allow 139 to calculate the maximum quantum efficiency through the ratio F_{v}/F_{m} where F_{v} $(F_{m}\mbox{-}F_{0})$ 140 corresponds to the variable fluorescence. This ratio was determined at predawn, dawn and in 141 diurnal period. 142

The diurnal course of chlorophyll parameters was monitored on 28th august 2007 from 9:00 h 143 to 18:00 h by using successively the dark adaptation clip and the PAR/temperature leaf-clip 144 which is designed for measurements under ambient light conditions. Thus additional variables 145 were recorded from light acclimated leaves: minimum (F'₀) maximum (F'_m) and the steady 146 state (F_s) values of fluorescence. Through these data as well as F_m and F_0 the following 147 fluorescence parameters were determined: The photochemical efficiency in the light adapted 148 state of photosystem II (Φ_{PSII}) calculated as (F'_m- F_s)/F'_m according to Genty et al. (1989), the 149 non-photochemical quenching coefficient of chlorophyll fluorescence (NPQ), calculated as 150 $(F_m - F'_m)/F'_m$ (Bilger and Björkman, 1990) which expresses the dissipation of excess energy 151 as heat and the photochemical quenching (qP)coefficient, which was calculated from (F'm -152 F_s /(F'_m - F'_0) that gives an indication of the proportion of PSII reaction centers that are open 153 (Maxwell and Johnson, 2000). An alternative non-photochemical quenching coefficient, QNP 154 =(Fm-F'm/Fm-F₀) was also determined (Schreiber et al., 1986). 155

Light response kinetic measurements were carried out at dawn by selecting four incident 156 photosynthetically active irradiance (PAR) intensities in order to simulate the half diurnal 157 light radiation change: 300, 600, 900 and 1200 µmol Photons m⁻²s⁻¹. The light saturating pulse 158 $(8.000 \ \mu mol Photons \ m^2s^{-1})$ was applied for each PAR intensity only when the steady-state 159 fluorescence intensity (F_s) was reached. This was checked prior to the design of a specific 160 light kinetic script. In addition to the above-mentioned fluorescence parameters, other 161 parameters were calculated for each actinic light application: the quantum efficiency of 162 regulated ΔpH and/or xanthophyll-dependent non-photochemical dissipation processes in the 163

antenna of PSII (Φ_{NPQ} = Fs/F'm – Fs/Fm). The quantum efficiency of the constitutive nonphotochemical fluorescence and energy dissipation ($\Phi_{f,D}$ =Fs/Fm) according to Hendrickson et al. (2005). The Chlorophyll fluorescence decrease ratio was calculated as: Fd/Fs= Fm-Fs/Fs (Lichtenthaler et al., 2005). Also, the rate of the linear electron transport was determined:

168 ETR = $0.5xPARx0.87x \Phi_{PSII}$ assuming that the photon flux are equally distributed between 169 the two photosystems.

170 2.3 Simultaneous gas exchange and fluorescence monitoring

The leaf gas exchange and fluorescence parameters were simultaneously determined using a 171 172 gas exchange system (LI-6400XT; LI-COR Inc., Lincoln, NE, USA) equipped with a fluorescence chamber head. CO_2 assimilation rate (A), internal CO_2 (Ci) and stomatal 173 conductance (g_s) were determined. All the measurements were carried out at 25°C and the 174 relative humidity of the chamber was set at 70%, while the amount of blue light was set to 175 10% of the photosynthetic photon flux density for optimized stomatal aperture. In light-176 adapted leaves, the steady state fluorescence yield (Fs) was recorded and the maximum 177 fluorescence (F'm) obtained after application of a saturating white light pulse (8.000 µmol 178 photons m⁻² s⁻¹, 0.8 s). The actinic light was then turned off and far-red illumination was 179 applied (2 µmol photons $m^{-2} s^{-1}$) to measure the light-adapted minimum fluorescence (F'0). 180 Using these parameters, Φ_{PSII} , qP and the ratio F'v/F'm = (F'm-F'o)/F'm which corresponds 181 to the efficiency of the energy harvesting by the oxidized PSII reaction centers in light 182 condition were determined. 183

184

185 2.4 Predawn leaf water potential determination

186 Water potential (Ψ) was measured using a pressure chamber of Scholander type (Scholander 187 et al., 1965), from mature fully expanded leaves of randomly selected plants. The 188 measurements were carried out from five trees per site at 05:00 h then finished before sunrise.

189 2.5 Soil and leaf sodium ions concentration

Five trees and five soil samples were considered for each condition: salt meadow and the control. Plant samples was composed by a pool of five leaves per tree. Soil and leaf samples were ash transformed in a muffle furnace. The extracted solutions were analysed for Na⁺ concentrations by using an Ion Chromatograph (Dionex Model DX 120, USA). The column

used was a CS16 with hydroxymethanesulfonic acid (26 mM) as eluent which a flow rate setat 1ml/min. Electrical conductivity was the mode of cations detection.

196

197

198 2.6 Photosynthetic pigments analysis

Leaves (a pool of five leaves per tree from five trees in each site) were collected at predawn then wrapped in aluminum foil to avoid for degradation of pigments by light and finally stored at -20° C until analysis in the laboratory. Chlorophylls and total carotenoids were extracted with acetone (80%) and filtered through Millipore membrane (0.2 µm pore size). Quantification was performed with a spectrophotometer (BioMateTM3) by measuring their absorbance at 663, 646 and 470 nm (Lichtenthaler and Wellburn, 1983).

205 2.7 Dendrochronology

The sampling process took place during the spring 2017. We selected six dominant and 206 207 mature ashes separated by at least 8 m from each other in the salt meadow; six other dominant ashes were selected as control trees. One core per tree at 1.3 m height was collected with a 208 Swedish increment borer. In the laboratory, cores were prepared according to standard 209 dendrochronological procedures (Fritts 1976). All cores were air-dried, glued onto groove 210 boards, sanded with progressively finer grade sandpaper to produce smoothed surfaces on 211 which the tree-ring boundaries were clearly visible and visually crossdated under a binocular 212 to ensure that each ring was dated to the correct calendar year. Then, ring-width series were 213 measured to the nearest 1/100 mm using a LINTAB6 measuring table (Rinntech®) and the 214 TSAP-Win software (Rinn 2003). All ring-width series were crossdated graphically and 215 crossdating was checked using the COFECHA software (Holmes, 1983) by assessing the 216 consistency of the different ring-width series among trees. 217

218

219 2.8 Statistical analysis

A Student test was performed to compare the two sites for each interval of time regarding tothe diurnal measurements of fluorescence parameters.

For the predawn physiological indicators (F_v/F_m , pigment content and Ψ) a two-way ANOVA was designed to assess the relationship between the two independent variables (Site x Year) on each dependent variable followed by Tukey's HSD (Honestly Significant Differences)

tests. The PSII photochemistry parameters obtained from light kinetics were first treated by a 225 multivariate analysis in order to determine the main explicative fluorescence parameters of 226 the actual quantum yield kinetics. This was performed using R statistical software (R 227 development Core Team, 2009) and the package "FactoMineR" Thus, covariance analyses 228 (ANCOVA), with light intensities fixed as covariate were used to examine the effect of site 229 and year on the pertinent explicative fluorescence variables. Curves fitting and statistical 230 analyses were performed using Statgraphics Centurion XV software and R statistical 231 software. 232

233

3. Results

235 *3.1 Annual changes in climatic conditions of the study site*

The study area was located near the village of Mauran at 11 km of the nearest meteorogical station which is that of Marignane (France). As shown in table 1, during this last decade, summer air temperature varied from 23 to 25°C. There was no significant difference between the years (one-way-ANOVA, F=2.12, P=0.065). However, the cumulated precipitation varied considerably (10-139 mm) resulting in similar changes in soil water deficit. For instance the highest water deficit was recorded in 2016 (-659 mm) while the lowest one was observed in 2011 (-4.18 mm).

243 3.2 Interannual radial growth

Analyzes of the cores collected from the individuals located at the two study sites allowed us to address their radial growth over the last ten years. The results clearly show that *F*. *angustifolia* grows better in salt meadows than the control site (Fig. 1). A peak of radial growth was recorded in 2014, where growth in the saline area was twice that of the riparian bank. However, since 2011, this growth has been falling to equal the radial growth of individuals in the control habitat.

250 *3.3 Leaf water and sodium ion status*

F. angustifolia environmental conditions in both salt meadow and riparian forest bank were first characterized at the beginning of august 2007. Soil water availability was characterized by the measurement of leaf water potential in the absence of any transpiration flux as well as leaves and soil sodium chloride contents. Predawn water potential was higher in control stand

compared to salt marsh individuals. Leaves Na^+ content were similar while soil Na^+ content in salt meadow site was about 13-fold that of the riparian forest bank (Table 2).

257

258

259 *3.4 The diurnal variation of chlorophyll fluorescence parameters*

From 9:00 h to 18:00 h several chlorophyll fluorescence data were recorded (Fig. 2). The 260 maximum quantum efficiency of PSII (Fv/Fm) was significantly higher for samples of salt 261 meadow site compared to those of the riparian forest only from 9:00 h to 13:00 h. For 262 instance, at 13:00 h the ratios Fv/Fm were 0.739 and 0.614, respectively (Fig. 2A). When 263 leaves were light acclimated, the actual efficiency of PSII photochemistry (Φ_{PSII}) decreases by 264 late morning to 16:00 h then rises until 6:00 h pm in both cases (Fig. 2B). However at 9:00h 265 Φ_{PSII} of F. angustifolia located in salty soils was about 2-fold that of none salty grown 266 specimens. The exitonic energy used for the reduction of the plastoquinone pool, determined 267 268 by the photochemical quenching coefficient (qP) exhibited similar patterns for both sites but with a significant difference obtained also only at 9:00 h am (Fig. 2C). Alternatively, the 269 270 excess of absorbed excitation energy may be dissipated through non-photochemichal processes referred as NPQ. No significant difference was found between the two sites in the 271 morning (Fig. 2D). 272

Hence, it seems that *F. angustifolia* quantum yield in salty soil condition is higher than the control during the morning and this may be explained by both photochemical quenching mechanisms and PSII antenna efficiency. This led to further investigation regarding to the predawn maximum quantum efficiency and the plant response to light kinetic taking into account the interannual variability.

278 3.5 Predawn Fv/Fm and steady-state light response of chlorophyll fluorescence parameters in
279 field conditions

280 *3.5.1 Predawn Fv/Fm*

The analysis of the maximum quantum efficiency carried out between 2007 and 2016 showed that the factors site and year interact together (F=5.31; p<0.001). Hence, the predawn ratios Fv/Fm of the sites were compared year by year. As a result (Fig. 3) *F. angustifolia* exhibited significant differences between sites in 2009 and 2016. For the former date, a lower ratio was obtained in salt meadow compare to the control; 0.819 and 0.886, respectively. For the latter,

an opposite response was shown because salt meadow Fv/Fm was about 0.05 units higher than the data recorded from control samples. The predawn ratio was also tested site by site for all the years which revealed that the ratio Fv/Fm tends to be lower in control with greater variability throughout the years (Fig. 4).

290

291 3.5.2 Dawn light kinetic fluorescence parameters measurements

Solar course during half a day was mimed by the implementation of a script where the 292 intensity of the light increases from 300 to 1200 μ mol. photon.m⁻².s⁻¹ in 4 steps. Fluorescence 293 data were recorded during 6 years then first examined through the quantum yield kinetic. 294 Two-way ANCOVA with light intensity as covariate was carried out. A significant effect of 295 the factors year (F=61.54 p<0.001) and site (F=9.38 p<0.001) was shown and no interaction 296 was found between them. Thus, the site effect on the dependent variable Φ_{PSII} was tested year 297 by year. As a result, the photochemical quantum yield decreases exponentially with light 298 intensity (Fig. 5) with significant differences between the two sites except in 2014. The 299 photosynthetic performance of individuals in salt meadows was higher than that of the 300 controls. It is thus possible to summarize the functioning of the photosynthetic machinery in 2 301 types of response. The first one is characterized by large variations in the actual quantum 302 yield at low light intensities as in 2007, 2008 and 2016. The second type of response is 303 manifested by an increasing difference as light intensities increase. This is the case in 2015 304 305 and 2009 where the quantum yields were about 0.8. It appears that soil salinity contributes to increase F. angustifolia PSII quantum yield through physiological mechanisms that change 306 307 depending to the year. This arises the question of the identification of the fluorescence parameters that explain these variations of the effective quantum yield. A multivariate 308 309 analysis was performed in order to prioritize the fluorescence parameters involved. The factor map of the variables denotes a total variance of 89.46 % (Fig.6). Interestingly, Φ_{PSII} is 310 strongly correlated to axis 1 (correlation coefficient: 0.99) followed by Φ_{NPO} (correlation 311 coefficient: 0.96) and qP (correlation coefficient: -0.87). As for axis 2, the main explanatory 312 variable is Fd/Fs (correlation coefficient: 0.87). In summary the variations of Φ_{PSII} are mainly 313 explained by non-photochemical quenching mechanisms followed by the ratio Fd/Fs. For the 314 former, the two-way ANCOVA performed showed a significant site (F=11.69, p<0.001) and 315 year (F=50.35, p<0.001) effects, no interaction was found between the two factors. In these 316 conditions, Φ_{NPO} was compared between sites, year by year. The results showed that Φ_{NPO} in 317 the salty areas was lower than that of the control except in 2014, 2015 (Fig. 7). 318

319

320 *3.6 Gas exchange and fluorescence measurements*

Simultaneous measurements of gas exchanges and fluorescence data were carried out in 2009 321 with a Licor 6400 equipped with the fluorescence chamber. The responses of F. angustifolia 322 to salt meadow environmental conditions were examined through the measurements of leaf 323 net photosynthesis (A), leaf internal CO₂ concentration (Ci), PSII antenna efficiency 324 (Fv'/Fm'), the photochemical quenching qP and the quantum yield of PSII. As shown in table 325 3 salt meadow plants exhibited a slight increase of the net photosynthesis (3.17 µmol. CO₂.m⁻ 326 2 .s⁻¹) compared to the controls (2.93 µmol. CO₂.m⁻².s⁻¹). Similar result was also obtained for 327 leaf internal CO₂ content. However, no differences were found for the quantum yield and 328 related fluorescence data. 329

330

331 *3.7 Photosynthetic pigments analysis*

The concentrations of chlorophylls and total carotenoids contents were measured from the 332 two sites throughout six years. Based on the data obtained for chlorophyll a there was an 333 interaction between the factors site and year (F=2.43; p<0.05). Sites were thus compared year 334 by year but no significant difference was found except in 2016 where chlorophyll a contents 335 from salt meadow samples were 3-fold higher compare to the controls (Fig. 8A). For the other 336 pigments no interaction was ascertained between the two factors and significant changes were 337 obtained only over the years. Hence, the highest contents of chlorophyll b were recorded in 338 2008 whereas the lowest values were obtained in 2007 (Fig. 8B). Total carotenoids 339 concentration (Fig. 8C) in 2008 was about 75 μ g/g of DW which represents the highest values 340 341 recorded during the years investigated while the lowest data were obtained in 2015 (27 μ g/g of DW). 342

343

4. Discussion

In this report we showed for the first time a long term survey of the *in situ* adaptation of a glycophyte to salt march conditions based on chlorophyll fluorescence signals. The model used was *F. angustifolia* which occurs naturally in several Mediterranean deltaic zones unlike to most of riparian forest species.

During the last decade, the tree-rings of individuals growing in the salt marsh showed clearly 349 two effects: on one hand, the biomass is greater than that of the individuals in the control area 350 (Fig. 1) and on the other hand, this biomass varies according to the fluctuations of water 351 deficit (Table 1). In particular, from the peak of 2011 the biomass has steadily decreased to 352 reach that of the control individuals in 2017. A higher biomass could be the result of intense 353 photosynthetic activity at the level of both light and enzymatic reactions but also the fact of a 354 greater availability of nutrients. Indeed, river mouths are recognized as areas with high 355 concentrations of mineral elements because of the importance of alluvial deposits. This is true 356 for halophytes but glycophytes have to cope with salinity: 13-fold higher as found in this 357 study (Table 2) suggesting the need of additional metabolic energy. Thus, the light-dependent 358 reactions of photosynthesis were followed through several parameters based on the 359 fluorescence of chlorophyll a. Measurements of the maximum quantum efficiency before 360 361 sunrise did not show any particular stress since the values obtained were all greater than 0.75 (Björkman and Demmig, 1987). This also indicates the lack of chronic photoinhibition or salt 362 363 induced toxicity to PSII (Yan et al., 2015). However, for the year 2016, which was characterized by the greatest water deficit (P=10 mm), the ratio Fv/Fm measured in the saline 364 365 zone was significantly higher than of the control zone. Similarly, the levels of chlorophyll a were three times higher compared to individuals in the control zone (Fig.7A). This is an 366 unexpected result because the ratio Fv/Fm trends to decrease with stress intensity as well as 367 chlorophyll a contents (Manaa et al., 2019; Masondo et al., 2019). Nevertheless, the increase 368 of chlorophyll a content under salinity was observed from individuals of Phragmites karka 369 (Retz.) Trin ex Sternd (Shoukat et al., 2019) as such, the increase of chlorophyll content has 370 been considered to be a marker of resistance to salinity (Acosta-Motos et al., 2017). This 371 would mean that the increase in the Fv/Fm ratio corresponds to an adaptive response of F. 372 angustifolia to both drought and salinity. Indeed chlorophyll a is present as a dimer in the 373 374 reaction center of the photosystems, an increase in its concentration could result in an increase in the number of photosystems and consequently a better photosynthetic capacity of 375 individuals to mitigate the environmental constraints. 376

Given the absence of chronic photoinhibition, fluorescence measurements were conducted in order to understand the dynamics of photosynthesis light reactions. Initially, the measurements made between 9:00 h and 18:00 h clearly showed that the actual photochemical yield (Φ_{PSII}) of individuals in salty environment was higher than those in the control area (Fig. 2B). Subsequently, kinetics measurements were carried out by simulation of light intensity variations for half a day. During the six years of measurement the kinetic curves show that

 Φ_{PSII} in saline medium is always higher than the control except during 2014. The water 383 conditions alone do not explain this difference between the years because the water deficits 384 between the years 2014 and 2015 were roughly the same (138 and 139). It seems that other 385 environmental factors contributed to buffer Φ_{PSII} variations in 2014. Nevertheless, the 386 adaptive capacity of an invasive plant in salt-meadows, Spartina patens (Aiton) Muhl. was 387 related to maintaining a high level of Φ_{PSII} despite the increase in salinity (Duarte et al., 388 2015). We can therefore consider that maintaining a high level of Φ_{PSII} of F. angustifolia in a 389 390 salty environment is an adaptive trait.

391

In order to highlight the mechanisms involved in the regulation of Φ_{PSII} , a multivariate 392 analysis was conducted. PCA analysis revealed that the main explanatory variables were the 393 fraction of absorbed energy that is dissipated thermally by the ΔpH - and the xanthophyll 394 cycle-dependent quenching in PSII (Φ_{NPO}), qP and Fd/Fs ratio. The one-way ANCOVA 395 performed on Φ_{NPO} confirms significant differences between salty and control environments 396 (Fig. 6) with the exception of the years 2014 and 2015. In particular, the high yields of Φ_{PSII} 397 in saline environments may be explained by a better efficiency in the conversion of the light 398 399 energy absorbed into reducing power since Φ_{NPQ} is low. However, in another survey it has been reported that low Φ_{NPO} values indicate a lower photoprotection capacity in the salt marsh 400 environmental condition (Arena et al., 2013). Photoinhibition mitigation was checked through 401 the analysis of carotenoid contents but this did not show any site effect suggesting the need of 402 further studies on individual xanthophylls characterization. Because no difference was found 403 in leaves Na⁺ content it is likely that ash individuals in the meadow have evolved specific 404 mechanisms of salt retention/exclusion at the root level. Indeed, several reports have shown 405 that glycophytes may adapt to salt stress by sequestering sodium ions in their root 406 compartment (Assaha et al., 2017; Chen et al., 2018). Among the mechanisms involved, the 407 apoplastic barrier through the casparian strip seems to play a critical role in controlling Na⁺ 408 uptake and transport into the shoots. This is supported by the identification of suberin-forming 409 410 key genes as *SbKCS11* (Chen et al., 2018).

Given the exclusion of salt from foliar cytoplasm, it appears that the major adaptive constraint of ash is water availability since the water deficit follows the growth pattern. Precisely, the greater the water deficit is the more saline concentration increases thus making the water less accessible to glycophytes. The concomitant fluorescence and gas exchange measurements carried out in 2009 showed that the net assimilation of CO_2 was the same regardless of the difference of salinity between the two sites. It is likely that the higher values of Φ_{PSII} obtained

in saline areas corresponds to a supplementary energy demand to sustain growth and 417 adaptation to water/salt stress. When considering the radial growth of F. angustifolia that 418 declines since 2011, both ATP and the anabolic coenzyme NADPH from photosynthesis 419 light-dependent reactions and the carbon assimilated are rather allocated to defence 420 processes. The tradeoffs between lignin biosynthesis and defense still remain challenging 421 although jasmonic acid and phytochrome B signalling pathway was identified as a hub that 422 coordinate plants growth and defense (Xie et al., 2018). Interestingly, differential analysis of 423 genes expressed in response to salt stress in Fraxinus velutina showed that the up-regulated 424 genes mainly concerned those of the biosynthetic pathway of alkaloids which are also known 425 as defense components (Yan et al., 2019). 426

427 **5.** Conclusion

In conclusion, this work, carried out for 6 years between 2007 and 2016, highlighted some of 428 the adaptive mechanisms of a glycophyte in salt meadows environmental condition. Since 429 2011, a decline has been observed in the production of ash biomass in saline environments. 430 431 This decline seems mainly related to recurrent water deficit. In the perspective of the climate change announced in the Mediterranean region, one could expect an extinction of ash in salt 432 433 meadows. F. angustifolia resistance was shown by the establishment of an efficient photochemical yield both by a reduction of the fraction of the absorbed radiation which is 434 dissipated as heat and also by increasing the concentration of chlorophyll a. However, non-435 native salt marsh species with a high tolerance to salinity and drought will be least affected by 436 437 the sea-level increase and drought associated to global change. Further studies are needed to establish the metabolic network involved in the adaptive strategies of glycophytes in saline 438 environments. 439

440

441 Acknowledgements: The authors gratefully acknowledge Mrs Christine Desouches and
442 Météo-France for providing the meteorological data. Dr Julien Ruffault is also acknowledged
443 for graph fitting with R software.

444





the control riparian forest from 2007-2017. n=6, means \pm SE)





Fig. 2 : Diurnal course of summer fluorescence measurements from *Fraxinus angustifolia* in salt meadow and riparian forest sites: The ratio Fv/Fm (A), the actual quenching yield (B), the photochemical quenching qP (C), and the Non-Photochemical Quenching NPQ (D). A student test was applied to compare the sites at each interval of time.*- P<0.05, **- P<0.01, ***-P<0.001. n = 6, means \pm SE



ns

470

457

471 Fig.3: Predawn Fv/Fm ratio of *Fraxinus angustifolia* from salt meadow and the control

- 472 riparian forest between 2007 and 2016. The test was performed year by year (one-way-
- 473 ANOVA, *- p < 0.05, ns- not significant, n=18, means \pm SD).

475 Mevy et al



476

477 Fig. 4: Predawn Fv/Fm ratio of *Fraxinus angustifolia* from salt meadow and the control

riparian forest between 2007 and 2016. The test was performed site by site (one-way-

479 ANOVA, **-p<0.001, ns- not significant, n=18, means ± SD).

481 Mevy et al



484 Fig. 5: Light response of PSII actual quantum yield of *Fraxinus angustifolia* in salt meadow
485 (red dotted line) and riparian forest sites (blue line). one-way-ANCOVA. *- P<0.05, ** -
486 P<0.01, ***- P<0.001.





491

Fig. 6: Principal Component Analysis (PCA) of the fluorescence variables map on axes 1 and 492 493 2 plus the correlation circle: The photochemical efficiency in the light adapted state of photosystem II (Φ_{PSII}), the non-photochemical quenching coefficient (NPQ), the 494 photochemical quenching coefficient (qP), the regulated ΔpH and/or xanthophyll-dependent 495 non-photochemical dissipation processes (Φ_{NPQ}), the constitutive non-photochemical 496 fluorescence and energy dissipation ($\Phi_{f,D}$), the ratio Fd/Fs, the Photosynthetic Photon Flux 497 498 Density (PPFD), the Electron Transport Rate (ETR) and the alternative non-photochemical 499 quenching coefficient, QNP.

501 Mevy et al.





Fig.7: Light response of PSII Non-Photochemical Quenching yield of *Fraxinus angustifolia* in
salt meadow (red dotted line) and riparian forest sites (blue line). one-way-ANCOVA.







- 511 Sites are compared year-by-year, one-way ANOVA, * *P*<0.05. B; Chlorophyll b content (no
- site effect, P>0.05). C; Carotenoids content (no site effect, P>0.05).

Journal Pre-proof					
Sampling date	Air temperature (°C)	Precipitations (P;mm)	Potential evapotranspiration (PET ; mm)	Water deficit (mm)	
Summer 2007	23.53±1.10	36.8	635.7	-598.9	
Summer 2008	23.87±2.28	27.6	649.1	-621.5	
Summer 2009	24.88±1.79	109.8	691.3	-581.5	
Summer 2011	22.86±1.81	191.2	609.6	-418.4	
Summer 2014	23.53±0.99	138.5	646.0	-507.5	
Summer 2015	25.14±1.66	139.0	675.8	-536.8	
Summer 2016	24.42±1.80	10.0	669.8	-659.8	

513

Table 1: Annual variations and climatic variables of Marignane (France) station close to the

515 study area. Climatic data for 3 months in summer before each sampling date. Average air 516 temperature (\pm SD) of decadal data in each year (n = 9). Cumulative values (P and PET) are

calculated from daily data for each summer (n = 90). Water deficit: P-PET. Air temperature

518 test: One-way-ANOVA, *F*=2.12, *P*=0.065).

	Predawn leaf water potential (MPa)	Leaf cation content (g/g of as	Na ⁺ Soil Na ⁺ content (g/ h) of soil solution)
Salt meadow	-4.61 (0.24)	0.74 (0.23)	20.74 (8.50)
Riparian forest	-4.13 (0.21)	0.76 (0.23)	1.63 (0.09)
			<u> </u>

Table 2: Predawn leaf water potential, leaf and soil Na⁺ contents from Salt meadow and 520

riparian forest of Fraxinus angustifolia in august 2007. 521 5. hidition

522

Mevy et al.					
Site	Net photosynthesis (A; μ mol CO ₂ . m ⁻² .s ⁻¹)	$\Phi_{\rm PSII}$	Fv'/Fm'	qP	Ci (µmol CO ₂ mol ⁻¹)
Control	2.93	0.24	0.44	0.54	261.70
	(0.53)	(0.01)	(0.01)	(0.03)	(25.82)
Salt meadow	3.17	0.24	0.42	0.55	273.98
	(0.84)	(0.03)	(0.01)	(0.06)	(26.58)

Journal Pre-proof

Table 3: Gas exchange and fluorescence measurements from leaves of Fraxinus angustifolia

in salt meadow and riparian forest stands. (n=12, means \pm SD).

References 529 Acosta-Motos, J.R., Ortuño, M.F., Bernal-Vicente, A., Diaz-Vivancos, P., Sanchez-Blanco, 530 M.J., Hernandez, J.A., 2017. Plant Responses to Salt Stress: Adaptive Mechanisms. 531 Agronomy 7, 18. https://doi.org/10.3390/agronomy7010018 532 Arena, C., De Micco, V., De Maio, A., Mistretta, C., Aronne, G., Vitale, L., 2013. Winter and 533 summer leaves of Cistus incanus: differences in leaf morphofunctional traits, 534 photosynthetic energy partitioning, and poly(ADP-ribose) polymerase (PARP) 535 activity. Botany 91, 805-813. https://doi.org/10.1139/cjb-2013-0121 536 Assaha, D.V.M., Mekawy, A.M.M., Liu, L., Noori, M.S., Kokulan, K.S., Ueda, A., Nagaoka, 537 T., Saneoka, H., 2017. Na+ Retention in the Root is a Key Adaptive Mechanism to 538 Low and High Salinity in the Glycophyte, Talinum paniculatum (Jacq.) Gaertn. 539 (Portulacaceae). J Agro Crop Sci 203, 56-67. https://doi.org/10.1111/jac.12184 540 Bang, J.H., Bae, M.-J., Lee, E.J., 2018. Plant distribution along an elevational gradient in a 541 542 macrotidal salt marsh on the west coast of Korea. Aquatic Botany 147, 52-60. https://doi.org/10.1016/j.aquabot.2018.03.005 543 Bilger, W., Björkman, O., 1990. Role of the xanthophyll cycle in photoprotection elucidated 544 545 by measurements of light-induced absorbance changes, fluorescence and 546 photosynthesis in leaves of Hedera canariensis. Photosynth Res 25, 173–185. 547 https://doi.org/10.1007/BF00033159 Björkman, O., Demmig, B., 1987. Photon yield of O2 evolution and chlorophyll fluorescence 548 characteristics at 77 K among vascular plants of diverse origins. Planta 170, 489-504. 549 https://doi.org/10.1007/BF00402983 550 551 Boughalleb, F., Denden, M., Tiba, B.B., 2009. Photosystem II photochemistry and physiological parameters of three fodder shrubs, Nitraria retusa, Atriplex halimus and 552 Medicago arborea under salt stress. Acta Physiol Plant 31, 463–476. 553 554 https://doi.org/10.1007/s11738-008-0254-3 Chai, W.-W., Wang, W.-Y., Ma, O., Yin, H.-J., Hepworth, S.R., Wang, S.-M., 2019. 555 Comparative transcriptome analysis reveals unique genetic adaptations conferring salt 556 tolerance in a xerohalophyte. Funct. Plant Biol. 46, 670–683. 557 https://doi.org/10.1071/FP18295 558 Chen, M., Yang, Z., Liu, J., Zhu, T., Wei, X., Fan, H., Wang, B., 2018. Adaptation 559 Mechanism of Salt Excluders under Saline Conditions and Its Applications. Int. J. 560 561 Mol. Sci. 19, 3668. https://doi.org/10.3390/ijms19113668 Dalberto, D.S., Martinazzo, E.G., Bacarin, M.A., 2017. Chlorophyll a fluorescence reveals 562 adaptation strategies in drought stress in Ricinus communis. Braz. J. Bot. 40, 861-870. 563 564 https://doi.org/10.1007/s40415-017-0412-1 Drvodelic, D., Ugarkovic, D., Orsanic, M., Paulic, V., 2016. The Impact of Drought, Normal 565 Watering and Substrate Saturation on the Morphological and Physiological Condition 566 567 of Container Seedlings of Narrow-Leaved Ash (Fraxinus angustifolia Vahl). SEEFOR-South-East Eur. For. 7, 135–142. https://doi.org/10.15177/seefor.16-11 568 569 Duarte, B., Santos, D., Marques, J.C., Caçador, I., 2015. Ecophysiological constraints of two invasive plant species under a saline gradient: Halophytes versus glycophytes. 570 Estuarine, Coastal and Shelf Science, Coastal systems under change: tuning 571 assessment and management tools 167, 154–165. 572 573 https://doi.org/10.1016/j.ecss.2015.04.007 Flowers, T.J., Colmer, T.D., 2008. Salinity tolerance in halophytes. New Phytol. 179, 945-574 963. https://doi.org/10.1111/j.1469-8137.2008.02531.x 575 Foti, R., Jesus, M. del, Rinaldo, A., Rodriguez-Iturbe, I., 2012. Hydroperiod regime controls 576 the organization of plant species in wetlands. PNAS 109, 19596-19600. 577 https://doi.org/10.1073/pnas.1218056109 578

Journal Pre-proof

579	Genty, B., Briantais, JM., Baker, N.R., 1989. The relationship between the quantum yield of
580	photosynthetic electron transport and quenching of chlorophyll fluorescence.
581	Biochimica et Biophysica Acta (BBA) - General Subjects 990, 87–92.
582	https://doi.org/10.1016/S0304-4165(89)80016-9
583	Gharbi, E., Martínez, JP., Benahmed, H., Lepoint, G., Vanpee, B., Quinet, M., Lutts, S.,
584	2017. Inhibition of ethylene synthesis reduces salt-tolerance in tomato wild relative
585	species Solanum chilense. Journal of Plant Physiology 210, 24–37.
586	https://doi.org/10.1016/j.jplph.2016.12.001
587	Hasegawa, P.M., 2013. Sodium (Na+) homeostasis and salt tolerance of plants. Environ. Exp.
588	Bot. 92, 19–31. https://doi.org/10.1016/j.envexpbot.2013.03.001
589	Hendrickson, L., Förster, B., Pogson, B.J., Chow, W.S., 2005. A simple chlorophyll
590	fluorescence parameter that correlates with the rate coefficient of photoinactivation of
591	Photosystem II. Photosynth Res 84, 43–49. https://doi.org/10.1007/s11120-004-6430-
592	4
593	Holmes, R.L., 1983. Computer-Assisted Quality Control in Tree-Ring Dating and
594	Measurement.
595	Hu, J., Peng, J., Zhou, Y., Xu, D., Zhao, R., Jiang, Q., Fu, T., Wang, F., Shi, Z., 2019.
596	Quantitative Estimation of Soil Salinity Using UAV-Borne Hyperspectral and Satellite
597	Multispectral Images. Remote Sens. 11, 736. https://doi.org/10.3390/rs11070736
598	James, M.L., Zedler, J.B., 2000. Dynamics of Wetland and Upland Subshrubs at the Salt
599	Marsh-Coastal Sage Scrub Ecotone. The American Midland Naturalist 143, 298–311.
600	Lichtenthaler, H.K., Buschmann, C., Knapp, M., 2005. How to correctly determine the
601	different chlorophyll fluorescence parameters and the chlorophyll fluorescence
602	decrease ratio RFd of leaves with the PAM fluorometer. Photosynthetica 43, 379–393.
603	https://doi.org/10.1007/s11099-005-0062-6
604	Lichtenthaler, H.K., Wellburn, A.R., 1983. Determinations of total carotenoids and
605	chlorophylls a and b of leaf extracts in different solvents. Biochem Soc Trans 11, 591–
606	592. https://doi.org/10.1042/bst0110591
607	Manaa, A., Goussi, R., Derbali, W., Cantamessa, S., Abdelly, C., Barbato, R., 2019. Salinity
608	tolerance of quinoa (Chenopodium quinoa Willd) as assessed by chloroplast
609	ultrastructure and photosynthetic performance. Environmental and Experimental
610	Botany 162, 103–114. https://doi.org/10.1016/j.envexpbot.2019.02.012
611	Masondo, N.A., Aremu, A.O., Kulkarni, M.G., Petřík, I., Plačková, L., Šubrtová, M., Novák,
612	O., Grúz, J., Doležal, K., Strnad, M., Finnie, J.F., Van Staden, J., 2019. How Do
613	Different Watering Regimes Affect the Growth, Chlorophyll Fluorescence,
614	Phytohormone, and Phenolic Acid Content of Greenhouse-Grown Ceratotheca triloba?
615	J Plant Growth Regul 38, 385–399. https://doi.org/10.1007/s00344-018-9848-1
616	Maxwell, K., Johnson, G.N., 2000. Chlorophyll fluorescence—a practical guide. J Exp Bot
617	51, 659–668. https://doi.org/10.1093/jexbot/51.345.659
618	Pennings, S.C., Grant, MB., Bertness, M.D., 2005. Plant zonation in low-latitude salt
619	marshes: disentangling the roles of flooding, salinity and competition. Journal of
620	Ecology 93, 159–167. https://doi.org/10.1111/j.1365-2745.2004.00959.x
621	Scholander, P.F., Bradstreet, E.D., Hemmingsen, E.A., Hammel, H.T., 1965. Sap Pressure in
622	Vascular Plants: Negative hydrostatic pressure can be measured in plants. Science
623	148, 339–346. https://doi.org/10.1126/science.148.3668.339
624	Schreiber, U., Schliwa, U., Bilger, W., 1986. Continuous recording of photochemical and
625	non-photochemical chlorophyll fluorescence quenching with a new type of modulation
626	fluorometer. Photosyn. Res. 10, 51–62. https://doi.org/10.1007/BF00024185
627	Shoukat, E., Abideen, Z., Ahmed, M.Z., Gulzar, S., Nielsen, B.L., 2019. Changes in growth
628	and photosynthesis linked with intensity and duration of salinity in Phragmites karka.

630 https://doi.org/10.1016/j.enveyphot.2010.03.024	
050 https://doi.org/10.1010/j.envexp001.2017.05.024	
631 Silvestri, S., Defina, A., Marani, M., 2005. Tidal regime, salinity and salt marsh p	lant
zonation. Estuarine, Coastal and Shelf Science 62, 119–130.	
633 https://doi.org/10.1016/j.ecss.2004.08.010	
Thorne, K.M., Takekawa, J.Y., Elliott-Fisk, D.L., 2012. Ecological Effects of Clin	nate Change
635 on Salt Marsh Wildlife: A Case Study from a Highly Urbanized Estuary. J	ournal of
636 Coastal Research 1477–1487. https://doi.org/10.2112/JCOASTRES-D-11-	00136.1
637 Traut, B.H., 2005. The role of coastal ecotones: a case study of the salt marsh/upla	and
638 transition zone in California. Journal of Ecology 93, 279–290.	
639 https://doi.org/10.1111/j.1365-2745.2005.00969.x	
640 Tsai, YC., Chen, KC., Cheng, TS., Lee, C., Lin, SH., Tung, CW., 2019. Ch	nlorophyll
641 fluorescence analysis in diverse rice varieties reveals the positive correlation	on between
642 the seedlings salt tolerance and photosynthetic efficiency. BMC Plant Biol	. 19, 403.
643 https://doi.org/10.1186/s128/0-019-1983-8	
Vaz, M., Coelho, R., Rato, A., Samara-Lima, R., Silva, L.L., Campostrini, E., Mo	ta, J.B.,
645 2016. Adaptive strategies of two Mediterranean grapevines varieties (Arag	gonez syn.
646 Tempranilio and Trincadeira) face drought: physiological and structural re	sponses.
647 Theor. Exp. Plant Physiol. 28, 205–220. https://doi.org/10.100//s40626-01	6-00/4-6
648 Veldkornet, D.A., Adams, J.B., Potts, A.J., 2015. Where do you draw the line? De	etermining
649 the transition thresholds between estuarine sait marsnes and terrestrial veg	etation.
bill South African Journal of Bolany, Biome Boundaries in South Africa 101,	135–139.
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	iona 7
652 Walls, 1., All, 1., 1u, J., Zhou, Z., He, S., Ku, W., Cul, D., Zhang, 1., Hall, K., L 652 2016 Different responses of photosystem II and antioxidants to drought st	race in two
654 contrasting populations of Sour jujube from the Loess Plateau China Eco	1 Res 31
761_{775} bttps://doi.org/10.1007/s11284_016_1384_5	I. ICS. 51,
656 Xie M Zhang I Tschanlinski T I Tuskan G A Chen I-G Muchero W 20	118
657 Regulation of Lignin Biosynthesis and Its Role in Growth-Defense Tradeo	ffs Front
658 Plant Sci 9, 1427 https://doi.org/10.3389/fpls.2018.01427	115. 1 10110.
659 Yan, K., Wu, C., Zhang, L., Chen, X., 2015, Contrasting photosynthesis and photo	oinhibition
660 in tetraploid and its autodiploid honeysuckle (Lonicera japonica Thunb.) u	nder salt
661 stress. Front Plant Sci 6. https://doi.org/10.3389/fpls.2015.00227	
662 Yan, L., Liu, C., Wang, Y., Wang, K., Ren, F., Yao, J., Wu, D., 2019. De novo tra	inscriptome
663 analysis of Fraxinus velutina Torr in response to NaCl stress. Tree Genetic	s &
664 Genomes 15, 56. https://doi.org/10.1007/s11295-019-1340-y	

Highlights

Adaptation of a glycophyte as *Fraxinus angustifolia* to salt meadow is determined by:

- An increase of the actual quantum yield of PSII photochemistry -
- _ An increase of chlorophyll a contents
- A lower fraction of absorbed energy that is dissipated thermally by the ΔpH and the xanthophyll cycle-dependent quenching in PSII (Φ_{NPO})
- _ A higher photochemical quenching (qP) and the ratio Fd/Fs

Fraxinus angustifolia in salt meadow is more sensitive to global climate change. Since 2011 a decrease of the tree-rings growth was shown suggesting changes in the specific composition in these areas.

inges in

Declaration of interests

 \boxtimes The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: