



Improving the antioxidant properties of *Calophyllum inophyllum* L. seed oil from French Polynesia: development and biological applications of ethanol-soluble resin extracts

Mathieu Cassien, Anne Mercier, Sophie Thétiot-Laurent, Marcel Culcasi,
Alice Asteian, Emilie Ricquebourg, Gaëtan Herbette, Jean-Pierre Bianchini,
Phila Raharivelomanana, Sylvia Pietri

► To cite this version:

Mathieu Cassien, Anne Mercier, Sophie Thétiot-Laurent, Marcel Culcasi, Alice Asteian, et al.. Improving the antioxidant properties of *Calophyllum inophyllum* L. seed oil from French Polynesia: development and biological applications of ethanol-soluble resin extracts. 2020. hal-03019052

HAL Id: hal-03019052

<https://amu.hal.science/hal-03019052>

Preprint submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improving the antioxidant properties of *Calophyllum inophyllum* L. seed oil from French Polynesia: development and biological applications of ethanol-soluble resin extracts.

Mathieu Cassien^{1,2,3}, Anne Mercier¹, Sophie Thétiot-Laurent¹, Marcel Culcasi¹, Alice Asteian¹, Emilie Ricquebourg¹, Gaëtan Herbette⁴, Jean-Pierre Bianchini², Phila Raharivelomanana² and Sylvia Pietri^{1,2*}

1-Aix Marseille University, CNRS, ICR UMR 7273, SMBSO, Marseille, France

2- EIO UMR241, Université de la Polynésie Française, Faa'a, Tahiti, French Polynesia

3-YELEN Analytics, Ensùès la-Redonne, France

3-Aix Marseille Univ, CNRS, Spectropole, FSCM, Marseille, France

Corresponding Author

S. Pietri, Ph.D.

Aix Marseille University, CNRS, ICR UMR 7273, Institut de Chimie Radicalaire

Sondes Moléculaires en Biologie et Stress Oxydant, Service 522

Faculté des Sciences de Saint-Jérôme, Avenue Escadrille Normandie-Niemen

13397 Marseille Cedex 20 - France

Tel: 33 (0)4 91 28 85 79 - Fax: 33 (0)4 91 28 87 58

E-Mail: sylvia.pietri@univ-amu.fr

Abstract: Tamanu oil (TO) from *Calophyllum inophyllum* L. has long been used for its numerous properties in traditional medicine. With the aim to investigate the intrinsic antioxidant and pharmacological activities of the complex pool of constituents contained in the resin part of TO, extraction with methanol and ethanol were undertaken to give MeTO and EtTO extracts, this latter being further partitioned into neutral (NTR) and acidic (ATR)

fractions. Further LPLC/HPLC separations led to sixteen pure metabolites. For two of them, calanolide D and 12-oxocalanolide A, we report here their first identification from a natural source. Extracts, subfractions and metabolites were evaluated for their antioxidant, anti-inflammatory, antimicrobial (*Staphylococcus aureus*) and antimycobacterial (*Mycobacterium tuberculosis H37RV*) properties, and their protective effect on *t*BuOOH-exposed cells and UV-exposed rat skin slices. Data showed that NTR and ATR resin fractions gathering the main amounts of bioactive compounds exhibited enhanced antioxidant and anti-inflammatory activities as compared to the lipid phase of TO. Antimicrobial activity was highlighted for NTR, while ATR was more cytoprotective in cells and rat tissues. All together, data suggest that ethanol-soluble fractions enriched in phenolics, pyranocoumarin derivatives and flavonoids could be of interest for further development of new and better bioavailable medicinal preparation.

Keywords: *Calophyllum inophyllum*, Tamanu oil, ethanol soluble extracts, lipophilic antioxidants, phenolics, pyranocoumarins, antimicrobial activity, anti-inflammatory activity, UV protection.

Table of contents:

¹ H NMR spectra of isolated or identified phytoconstituents of Tamanu resins:	2
Figure S1: ¹ H NMR spectra of Inocalophyllin B	2
Figure S2: ¹ H NMR spectra of Inophyllum C	3
Figure S3: ¹ H NMR spectra of Inophyllum E	3
Figure S4: ¹ H NMR spectra of Inophyllum P	4
Figure S5: ¹ H NMR spectra of Calophyllolide	4
Figure S6: ¹ H NMR spectra of 12-oxocalanolide A	5
Figure S7: ¹ H NMR spectra of Calanolide A	5
Figure S8: ¹ H NMR spectra of Calanolide D	6
Figure S9: ¹ H NMR spectra of Calanolide GUT70	6
Figure S10: ¹ H NMR spectra of Tamanolide	7

Figure S11: ^1H NMR spectra of Tamanolide D

7

Figure S12: ^1H NMR spectra of Fraction D

8

Figure S13: ^1H NMR spectra of Fraction P

8

^1H NMR spectra of isolated or identified phytoconstituents of Tamanu resins

Figure S1: ^1H NMR spectra of Inocalophyllin B




Figure S2: ^1H NMR spectra of Inophyllum C




Figure S3: ^1H NMR spectra of Inophyllum E




Figure S4: ^1H NMR spectra of Inophyllum P




Figure S5: ^1H NMR spectra of Calophyllolide




Figure S6: ^1H NMR spectra of 12-oxocalanolide A




Figure S7: ^1H NMR spectra of Calanolide A




Figure S8: ^1H NMR spectra of Calanolide D




Figure S9: ^1H NMR spectra of Calanolide GUT70




Figure S10: ^1H NMR spectra of Tamanolide




Figure S11: ^1H NMR spectra of Tamanolide D




Figure S12: ^1H NMR spectra of Fraction D




Figure S13: ^1H NMR spectra of Fraction P

