

Description of Gracilibacillus phocaeensis sp. nov., a new halophilic bacterium isolated from Senegalian human stool

I. I. Ngom, I. Hasni, Bruno Senghor, C. I. Lo, N. Armstrong, C. Sokhna, D. Raoult, P. -E. Fournier, J. -C. Lagier

► To cite this version:

I. I. Ngom, I. Hasni, Bruno Senghor, C. I. Lo, N. Armstrong, et al.. Description of Gracilibacillus phocaeensis sp. nov., a new halophilic bacterium isolated from Senegalian human stool. NEW MICROBES AND NEW INFECTIONS, 2020, 38, 10.1016/j.nmni.2020.100799 . hal-03149259

HAL Id: hal-03149259 https://amu.hal.science/hal-03149259

Submitted on 15 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Description of *Gracilibacillus phocaeensis* sp. nov., a new halophilic bacterium isolated from a Senegalian human stool.

Issa Isaac NGOM^{1,2}, Issam HASNI^{1,2,4}, Bruno SENGHOR^{1,2}, Cheikh Ibrahima LO^{2,3}, Nicholas ARMSTRONG^{1,2}, Cheikh SOKHNA^{2,3}, Didier RAOULT^{1,2}, Pierre-Edouard FOURNIER^{2,3}, Jean-Christophe LAGIER^{1,2*}.

¹ Aix Marseille Univ, IRD, AP-HM, MEPHI, Marseille, France

² IHU-Méditerranée Infection, Marseille, France

³ Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

⁴ Amoeba R&D Department, Chassieu, France.

* Corresponding author : Jean-Christophe LAGIER, Institut Hospitalo-Universitaire
Méditerranée-Infection, 19-21 Boulevard Jean Moulin, 13385, Marseille cedex 05, France ;
tel +33 413 732 401 ; fax : +33 413 732 402 ; e-mail: jean-christophe.lagier@ap-hm.fr

Running title: Gracilibacillus phocaeensis sp. nov.

Keywords: Gracilibacillus phocaeensis sp. nov. ; genome; taxonogenomics; culturomics; gut

Abbreviations :

CSUR: Collection de Souches de l'Unité des Rickettsies MALDI-TOF: Matrix-Assisted Laser-Desorption/Ionization Time-Of-Flight FAME: Fatty Acid Methyl Ester GC/MS: Gaz Chromatography/Mass Spectrometry ASP: Antibiotic susceptibility profile

1 Abstract

Using Taxonogenomics method, we described here *Gracilibacillus phocaeensis* strain Marseille-P380, a new species previously isolated from a salty stool of a 20-year-old man from N'Diop. It is Gram-positive, aerobic and motile bacilli. The major fatty acids are C_{15:0} anteiso (59 %), C_{16:0} (16 %) and C_{17:0 anteiso} (11 %). Strain Marseille-P3801 exhibits a 98.45% sequence similarity with *Gracilibacillus thailandensis* TP2-8, the phylogenetically closest species. Its genome is 4.66 Mb with 39.6 mol% G+C content.

9 Introduction

The genus Gracilibacillus has been described as moderately halophilic, motile endospore-10 forming bacteria [1]. Moderately halophilic bacteria have been found in a variety of 11 fermented foods and one of the most important food preservation methods in history has been 12 the use of salt (NaCl) [2]. Salt is the main source of sodium in our diet. Some gut bacteria, 13 such as lactobacilli, are very sensitive to salt [3]. However, it has been demonstrated to favor 14 the emergence and growth of others, mainly halophilic bacteria, including *Gracilibacillus* [4]. 15 Halophilic and halotolerant bacteria are most commonly isolated from the human gut 16 microbiota [5,6]. The culturomics approach, based on the multiplication of culture conditions 17 18 (variation of media, temperature, and atmosphere) with a more rapid bacterial identification by Matrix-Assisted Laser-Desorption/Ionization Time-Of-Flight Mass Spectrometry 19 (MALDI-TOF MS) [7] was used to explore the human gut halophilic microbiota. Thanks to 20 culturomics, we used high salt-containing culture media, which allowed us to isolate a new 21 moderately halophilic bacterial strain Marseille-P3801^T, which belongs to the genus 22 Gracilibacillus [8]. The genus Gracilibacillus currently compromises 13 species validly 23 standing in nomenclature [9]. Gracilibacillus species were isolated from diverse salty 24 environmental samples, including sea water, salty lakes [4,10,11], soil [12,13] and/or food 25 [1,14,15] and gut microbiota [6]. 26

Various parameters, including phenotypic and genotypic characteristics, such as DNA-DNA hybridization, were used to define a new species, but they present certain limitations [16]. In our study, using a taxonogenomic approach that includes phenotypic characteristics, proteomic information obtained by MALDI-TOF MS and the analysis of the complete genome sequence, we were able to achieve a complete description of the new halophilic species called *G. phocaeensis*; the strain Marseille-P3801T (= CSUR P3801) is the type strain of *Gracilibacillus phocaeensis* sp. nov.

34 Materials and Methods

35 Bacterial strains and growth conditions

Strain Marseille-P3801^T was isolated from stool samples of a 20-year-old man from N'Diop. 36 The study was approved by the ethics committee of the Institut Hospitalo-Universitaire 37 Méditerranée Infection under number 2016-011, and the patient provided a signed informed 38 consent. The percentage of NaCl in the stool sample was determined using a salinity 39 refractometer (Thermo Scientific, Villebon-sur-Yvette, France) by diluting 1 g in 10 mL of 40 distilled water and centrifuging it for 10 minutes at 5000g. In a second time, 100 µL of 41 supernatant was deposited in the refractometer; the result was in a straight line, displayed on 42 43 the screen in per mille and then reported in percentage of NaCl. To culture the bacteria from stool samples, we used an aerobic blood culture bottle (Becton Dickinson, Le Pont-de-Claix, 44 France) containing a halophilic medium prepared in a modified Columbia broth (Sigma-45 Aldrich, Saint-Quentin-Fallavier, France), by adding (per litre: 1% (w/v) MgSO4, 0.1% (w/v) 46 MgCl2, 0.4% (w/v) KCl, 0.1% (w/v) CaCl2, 0.05% (w/v) NaHCO3, 0.2% (w/v) of glucose, 47 0.5% (w/v) of yeast extract (Becton Dickinson), and from 10 to 15% (w/v) NaCl according to 48 the required salinity with a pH adjusted to 7.5 and incubated for 3 days in an aerobic 49 atmosphere at 37°C [8]. All strains were first isolated in a halophilic culture medium with 50 15% (w/v) NaCl. The initial growth of colonies on agar were obtained after 24 hours of 51 incubation at 37°C under aerobic conditions. The oxygen requirement was evaluated by 52 incubating strain Marseille-P3801^T under aerobic, microaerophilic, and anaerobic conditions 53 using AnaeroGenTM (Atmosphere Generation Systems, Dardily, France) at 37°C. The 54 colonies isolated were identified using MALDI-TOF MS, as previously described [17]. As for 55 unidentified colonies, the 16S rRNA gene was sequenced and the obtained sequence was 56 matched against the NCBI database using the BLAST algorithm [18]. 57

58 16S rRNA gene sequencing and phylogenetic analysis

The 16S rRNA gene sequence of the strain was determined for subsequent phylogenetic 59 analysis. The genomic DNA of the strain was amplified by PCR using the primers pair fD1 60 and rP2 (Eurogentec, Angers, France) [19] and sequenced with the MiSeq Technology 61 (Illumina Inc, San Diego, CA, USA) as previously described [20]. The 16S rRNA nucleotide 62 assembled and corrected using CodonCode Aligner 63 sequences were software (http://www.codoncode.com). A BLAST research (Basic Local Alignment Search Tool) was 64 further performed against the GenBank nucleotide collection 65 (http://blast.ncbi.nlm.nih.gov/Blast.cgi). If the 16S rRNA sequence similarity value was lower 66 than 98.65% with the most closely related species with standing in nomenclature, as proposed 67 by Stackebrandt [21], the strain was proposed as belonging to a new species [22]. 68

69 Phenotypic and biochemical characteristics

70 The morphology of strain Marseille-P3801^T was revealed by a negative staining observed on

a Hitachi SU5000 scanning electron microscope (Hitachi Group, Krefeld, Germany) and

72 Gram staining observed on photonic microscope Leica DM2500 (Leica Microsystems,

73 Nanterre, France) with a 100X oil-immersion objective. Sporulation, motility catalase and

oxidase were tested, as previously reported [23,24]. To determine the optimal growth

conditions, the strain Marseille-P3801 was cultivated in Mueller Hinton agar (Sigma-Aldrich)

by varying the NaCl concentrations from 5 to 20% (w/v) as well as the pH values (5, 5.5, 6,

6.5, 7, 7.5 and 8). It was also seeded on 5% sheep blood-enriched Columbia agar

78 (bioMérieux, Marcy l'Etoile, France) and incubated under different temperatures (25, 28, 37,

45 and 55°C). API 50CH, ZYM and 20 NE test strips (bioMérieux, Marcy-l'Etoile, France)

80 were used to study the carbohydrate metabolism, enzymes activities and biochemical criteria

81 of strain Marseille-P3801, according to the manufacturer's instructions.

82 Cellular fatty acid methyl ester (FAME) analysis was performed by GC/MS. Two samples

83 were prepared with approximately 90 mg of bacterial biomass per tube, harvested from

several culture plates. Fatty acid methyl esters were prepared as described by Sasser [25].
GC/MS analyses were carried out as previously described [26]. Briefly, fatty acid methyl
esters were separated using an Elite 5-MS column and monitored by mass spectrometry
(Clarus 500 - SQ 8 S, Perkin Elmer, Courtaboeuf, France). Spectral database search was
performed using MS Search 2.0 operated with the Standard Reference Database 1A (NIST,
Gaithersburg, USA) and the FAMEs mass spectral database (Wiley, Chichester, UK).

90 Genome: extraction, sequencing and assembly

Genomic DNA (gDNA) of Gracilibacillus phocaeensis was extracted in two steps : a 91 mechanical treatment was first performed by glass beads acid washed (G4649-500g Sigma) 92 using a FastPrep BIO 101 instrument (Qbiogene, Strasbourg, France) at maximum speed 93 (6.5m/s) for 90 seconds. After 30 minutes incubation of the lysozyme at 37°C, the DNA was 94 extracted on the EZ1 biorobot (Qiagen) with the EZ1 DNA tissue kit. The elution volume is 95 96 50µL. Genomic DNA was evaluated with a Qubit test (Life technologies, Carlsbad, CA, USA). The mate pair library was prepared with 1.5 µg of genomic DNA using the Nextera 97 mate pair Illumina guide. The DNA sample was simultaneously fragmented and tagged with a 98 mate pair junction adapter. The pattern of fragmentation was validated on an Agilent 2100 99 BioAnalyzer (Agilent Technologies Inc, Santa Clara, CA, USA) with a DNA 7500 labchip 100 kit. The DNA fragments size ranged from 1.5 kb up to 11 kb, with an optimal size at 8.10 kb. 101 No size selection was performed and 600 ng of tagmented fragments were circularized. The 102 circularized DNA was mechanically sheared to small fragments with an optimal at 1086 bp on 103 the Covaris device S2 in T6 tubes (Covaris, Woburn, MA, USA). The library profile was 104 105 visualized on a High Sensitivity Bioanalyzer LabChip kit (Agilent Technologies Inc) and the final concentration library was measured at 31.31 nmol/l. The libraries were normalized at 106 2nM and pooled. After a denaturation step and dilution at 15 pM, the pool of libraries was 107 loaded onto the reagent cartridge and then onto the instrument along with the flow cell. 108

Automated cluster generation and sequencing run were performed in a single 39-hour run in a 2x251-bp. Total information of 8.2 Gb was obtained from a 932 K/mm2 cluster density with a cluster passing quality control filters of 91 %. Within this run, the index representation for *Gracilibacillus phocaeensis* was determined at 13.20 %. The 2 141 870 paired end reads were filtered according to the read qualities.

The assembly was performed with a pipeline incorporating different softwares, including Velvet[27], Spades [28] and Soap Denovo [29] on trimmed (Trimmomatic) [30] or raw data. To reduce assembly gap, GapCloser software was used. The scaffolds inferior to 800 bp and scaffolds with a depth value < 25% of the mean depth were discarded. The best assembly was selected using different criteria (number of scaffolds, N50, number of N).

119 Genome annotation and comparisons

Open reading frames (ORFs) were predicted using the Prodigal tool (http://prodigal.ornl.gov) 120 121 with defaults parameters. tRNAs and rRNAs were detected using tRNAscan-SE v.1.2129 and RNAmmer v.1.230, respectively[31, 32]. The protein sequence annotation was performed on 122 NCBI GenBank non-redundant protein sequence database (nr) using BLAST protein with an 123 e-value of 1e-03 as significance thresholds [33]. Then, we obtained the functional 124 classification of gene families (COG ID and letters) using EggNOG against the COG database 125 [34]. The genome of *Gracilibacillus phocaeensis* strain Marseille-P3801^T (EMBL EBI 126 accession number UZBG0000000) was compared with that of Gracilibacillus 127 boraciitolerans strain JCM 21714 (BAVS0000000), Gracilibacillus massiliensis strain Awa-128 1^T (CZRP00000000), Gracilibacillus lacisalsi DSM 19029 (ARIY00000000), Gracilibacillus 129 ureilyticus CGMCC (FOGL0000000), Gracilibacillus dipsosauri (QGTD00000000) and 130 Halobacillus karajensis DSM 14948 (FNWW0000000) using OrthoANI software [35]. 131

132 **Results**

133 Strain identification and phylogenetic analysis

Strain Marseille-P3801^T formed yellow colonies after 1 day of culture on agar with horse 134 blood ranging from 2 to 20% (w/v) NaCl (optimum at 7.5 (w/v) NaCl) at 37°C. The spectrum 135 resulting from the 8 pure colonies of strain Marseille-P3801 deposited on MALDI-TOF target 136 plate did not allow the identification of this bacterium because there was no spectrum match 137 with those in the Bruker database (Supplementary Fig. S1). Using 16S rRNA sequence of G. 138 phocaeensis (LT934503.1), the phylogenetic analysis revealed that strain Marseille-P3801 139 exhibited a sequence similarity of 98.39% with Gracilibacillus thailandensis strain TP2-8 140 (GenBank accession no. NR_116568.1), the phylogenetically closest species with standing in 141 nomenclature (Fig. 1). Therefore, we classified this strain as a member of a new species 142 143 within the genus Gracilibacillus, family Bacillaceae and phylum Firmicutes. This value was lower than the 98.7% 16S rRNA gene sequence, threshold advised by Meier-Kolthoff et al. 144 [36] to delineate a new species without carrying out DNA-DNA hybridization. Classification 145 146 and general features of strain Marseille-P3801 are summurazed in Table S1.

147 Physiological and biochemical characteristics

G. phocaeensis sp. nov. strain Marseille-P3801^T (= CSUR P3801) is Gram positive. Colonies
are circular, yellow with a mean diameter of 2 mm after 2-3 days of growth on 5% sheep
blood-enriched Columbia agar medium (BioMérieux, Marcy l'Etoile, France). Bacterial cells
of strain Marseille-P3801 were motile, rod shaped and polymorphic (Fig. 2).

The major fatty acids were saturated structures mainly branched: 12-methyl-tetradecanoic acid (59 %), hexadecanoic acid (16 %) and 14-methyl-hexadecanoic acid (11 %). Other saturated and branched fatty acids were also described. 7-Hexadecenoic acid was the only one unsaturated structure detected (**Supplementary Table S2**). Catalase and oxidase were positive. Using API ZYM trips, positive reaction were detected for lipases (C4, C8 and C14), leucine arylamidase, acid phosphatase, naphthol-as-bi-phosphohydrolase, β -galactosidase, α glucosidase, β -glucosidase and N-acetyl- β -glucosaminidase, but no reaction was observed for

alkaline phosphatase, valine and cysteine arylamidase, α -chymotrypsin, α -galactosidase, 159 160 trypsin, β -glucuronidase, α -mannosidase and α -fucosidase. API 20NE strip exhibited positive reactions of fermentation of glucose, urease activity, and metabolism of L-arginine and 161 esculin. In contrast, negative reactions were observed for nitrate and indole production as well 162 metabolism of D-glucose, L-arabinose, D-mannose, D-maltose, D-mannitol, as 163 N-acetyl-glucosamine, potassium gluconate, capric acid, malic acid and phenylacetic acid. 164 Using API 50CH strip, strain Marseille-P3801^T exhibited esculin hydrolysis and negative 165 reactions for D-galactose, D-lactose, D-maltose, D-ribose, D-saccharose, D-lyxose, 166 D-mannose, L-sorbose, D-tagatose, D-turanose, D-xylose, L-xylose, D-arabinose, 167 L-arabinose, D-sorbitol, D-cellobiose, D-melezitose, D-melibiose, D-trehalose, D-raffinose, 168 D-arabitol, L-arabitol, D-glucose, D-fructose, D-fucose, L-rhamnose, D-adonitol, D-mannitol, 169 L-fucose, amygdalin, arbutin, erythritol, dulcitol, gentiobiose, glycerol, glycogen, inositol, 170 inulin, salicin, αD-glucopyranoside, methyl-βD-xylopyranoside, 171 starch, xylitol, methyl-αD-mannopyranoside, potassium gluconate and N-acetylglucosamine. The differential 172 characteristics of Gracilibacillus phocaeensis with respect to other bacteria related to the 173 genus Gracilibacillus are outlined in Table 1. Phenotypic characterization of Gracilibacillus 174 175 phocaeensis sp. nov., based on analytical profile index (API) tests was summurized on Table 2. 176

177 Genome properties

The genome length of *Gracilibacillus phocaeensis* strain Marseille-P3801^T is 4.66 megabases encompassing 12 scaffolds (11 contigs). The guanine-cytosine (GC) content is 39.6 mol%. Among the 4,390 predicted genes, 4,255 were protein-coding genes, 67 were RNAs (11 rRNA, 52 tRNAs and 4 non-coding RNAs) and 68 pseudogenes. The BLASTp annotation of *G. phocaeensis* strain Marseille-P3801 had assigned a putative function to 3,606 genes and 649 genes annotated as hypothetical proteins. To further insight on the gene function, a 184 comparison of *G. phocaeensis* protein sequences with COG database was realized. Out of

- 185 4,390 protein encoding genes, 3,774 were assigned to a COG function (86%) distributed in 20
- 186 COG categories (Supplementary Fig. S2).

187 Genome comparison

To explore the genomic similarity of G. phoceeanensis with closely related bacteria, we 188 performed an OrthoANI analysis. Among closely related species, we found OrthoANI values 189 ranging from 66.64% between Gracilibacillus phocaeensis strain Marseille-P3801 and 190 Halobacillus karajensis DSM 14948, to 78.39 % between Gracilibacillus boraciitolerans 191 strain JCM 21714 and Gracilibacillus massiliensis strain Awa-1. When Gracilibacillus 192 phocaeensis strain Marseille-P3801 was compared with these closely related species, we 193 found values ranging from 66.64% with Halobacillus karajensis DSM 14948, to 72.42% with 194 Gracilibacillus lacisalsi strain DSM 19029 (Fig. 3). The representations of strain Marseille-195 196 P3801 genome and its genes repartition into functional categories are illustrated in Fig. 4.

197 Description of Gracilibacillus phocaeensis

Gracilibacillus phocaeensis (pho.ca.een'sis, N.L. masc. adj., from phocaeensis, related to the 198 Phocaeans, the founders of Marseille, France where the type strain was isolated and 199 characterized like many others species). It is Gram positive, motile and aerobic bacterium. 200 The colonies are yellow, circular with a mean diameter of 2 mm. Bacterial cells were rod-201 shaped and polymorphic. Strain Marseille-P3801 grows at an optimal temperature of 37°C, 202 pH 7 with 7.5% (w/v) of NaCl. It is catalase and oxidase positive. Positive reactions were 203 observed for esterase (C4), esterase lipase (C8), acid phosphatase, naphthol-as-bi-204 phosphohydrolase, leucine arylamidase, β -galactosidase, α -glucosidase, β -glucosidase and N-205 acetyl-β-glucosaminidase. G. phocaeensis strain Marseille-P3801^T was isolated from a stool 206 sample of a 20-year-old man from N'Diop, Senegal. This strain exhibited a G+C content of 207 39.6 mol%. Its 16S rRNA sequence was deposited in GenBank under accession number 208

LT934503, and the whole genome shotgun sequence was deposited in GenBank under accession number UZBG00000000.

211 Discussion and conclusion

The concept of "microbial culturomics", based on the variation of physicochemical 212 parameters of the culture conditions, allows the exploration of microbial diversity of different 213 ecosystems such as gut microbiota [7]. Microbial culturomics provides culture conditions that 214 simulate, reproduce or mimic all the selective constraints that have shaped the natural 215 microbiota for millions of years. Many new bacterial species have been discovered, 216 particularly those belonging to the Bacillales order, which is one of the most represented 217 bacterial orders [37]. To explore the halophilic microbiota of the human gut, the use of culture 218 media with a high salt content, allowed us to isolate a new moderately halophilic bacterial 219 strain, Marseille-P3801^T which belongs to the genus *Gracilibacillus* [8]. To the best of our 220 221 knowledge, this is the second Gracilibacillus species described in the human gut. Based on its phenotypic, phylogenetic and genomic characteristics, this strain is proposed to represent a 222 novel species in the genus Gracilibacillus, for which the name Gracilibacillus phocaeensis 223 sp. nov. is proposed, with Marseille-P3801^T as the type strain. 224

References

- Oh YJ, Lee H-W, Lim SK, Kwon M-S, Lee J, Jang J-Y, et al. *Gracilibacillus kimchii* sp. nov., a halophilic bacterium isolated from kimchi. J Microbiol Seoul Korea. 2016;54:588–93.
- Gibtan A, Park K, Woo M, Shin J-K, Lee D-W, Sohn JH, et al. Diversity of Extremely Halophilic Archaeal and Bacterial Communities from Commercial Salts. Front Microbiol [Internet]. 2017 [cited 2019 Aug 19];8. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5423978/
- 3. Miranda PM, De Palma G, Serkis V, Lu J, Louis-Auguste MP, McCarville JL, et al. High salt diet exacerbates colitis in mice by decreasing *Lactobacillus* levels and butyrate production. Microbiome. 2018; 6:57.
- 4. Wainø M, Tindall BJ, Schumann P, Ingvorsen K. *Gracilibacillus* gen. nov., with description of *Gracilibacillus halotolerans* gen. nov., sp. nov.; transfer of *Bacillus dipsosauri* to *Gracilibacillus dipsosauri* comb. nov., and *Bacillus salexigens* to the genus *Salibacillus* gen. nov., as *Salibacillus salexigens* comb. nov. Int J Syst Bacteriol. 1999; 49 Pt 2:821–31.
- Khelaifia S, Lagier J-C, Bibi F, Azhar EI, Croce O, Padmanabhan R, et al. Microbial Culturomics to Map Halophilic Bacterium in Human Gut: Genome Sequence and Description of *Oceanobacillus jeddahense* sp. nov. OMICS J Integr Biol. 2016; 20:248–58.
- Diop A, Seck E hadji, Dubourg G, Armstrong N, Blanc-Tailleur C, Raoult D, et al. Genome sequence and description of *Gracilibacillus timonensis* sp. nov. strain Marseille-P2481T, a moderate halophilic bacterium isolated from the human gut microflora. MicrobiologyOpen [Internet]. 2018 [cited 2019 Aug 8]; 8. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6485780/
- 7. Lagier J-C, Armougom F, Million M, Hugon P, Pagnier I, Robert C, et al. Microbial culturomics: paradigm shift in the human gut microbiome study. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis. 2012;18:1185–93.
- Senghor B, Seck EH, Khelaifia S, Bassène H, Sokhna C, Fournier P-E, et al. Description of 'Bacillus dakarensis' sp. nov., 'Bacillus sinesaloumensis' sp. nov., 'Gracilibacillus timonensis' sp. nov., 'Halobacillus massiliensis' sp. nov., 'Lentibacillus massiliensis' sp. nov., 'Oceanobacillus senegalensis' sp. nov., 'Oceanobacillus timonensis' sp. nov., 'Virgibacillus dakarensis' sp. nov. and 'Virgibacillus marseillensis' sp. nov., nine halophilic new species isolated from human stool. New Microbes New Infect. 2017; 17:45–51.
- 9. Parte AC. LPSN—list of prokaryotic names with standing in nomenclature. Nucleic Acids Res. 2014;42:D613–6.
- Gao M, Liu Z-Z, Zhou Y-G, Liu H-C, Ma Y-C, Wang L, et al. *Gracilibacillus kekensis* sp. nov., a moderate halophile isolated from Keke Salt Lake. Int J Syst Evol Microbiol. 2012;62:1032–6.
- 11. Jeon CO, Lim J-M, Jang HH, Park D-J, Xu L-H, Jiang C-L, et al. *Gracilibacillus lacisalsi* sp. nov., a halophilic Gram-positive bacterium from a salt lake in China. Int J Syst Evol Microbiol. 2008;58:2282–6.

- Huo Y-Y, Xu X-W, Cui H-L, Wu M. *Gracilibacillus ureilyticus* sp. nov., a halotolerant bacterium from a saline–alkaline soil. Int J Syst Evol Microbiol. 2010;60:1383–6.
- 13. Chen Y-G, Cui X-L, Zhang Y-Q, Li W-J, Wang Y-X, Xu L-H, et al. *Gracilibacillus halophilus* sp. nov., a moderately halophilic bacterium isolated from saline soil. Int J Syst Evol Microbiol. 2008;58:2403–8.
- Chamroensaksri N, Tanasupawat S, Akaracharanya A, Visessanguan W, Kudo T, Itoh T. *Gracilibacillus thailandensis* sp. nov., from fermented fish (pla-ra). Int J Syst Evol Microbiol. 2010;60:944–8.
- 15. Diop A, Khelaifia S, Armstrong N, Labas N, Fournier P-E, Raoult D, et al. Microbial culturomics unravels the halophilic microbiota repertoire of table salt: description of *Gracilibacillus massiliensis* sp. nov. Microb Ecol Health Dis [Internet]. 2016 [cited 2019 Aug 18];27. Available from:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5071648/

- 16. Vandamme P, Pot B, Gillis M, de Vos P, Kersters K, Swings J. Polyphasic taxonomy, a consensus approach to bacterial systematics. Microbiol Rev. 1996;60:407–38.
- Lo CI, Fall B, Sambe-Ba B, Diawara S, Gueye MW, Mediannikov O, Sokhna C et al. MALDI-TOF Mass Spectrometry: A Powerful Tool for Clinical Microbiology at Hôpital Principal de Dakar, Senegal (West Africa). PLoS One 2015; 10(12):e0145889. doi: 10.1371/journal.pone.0145889.
- Drancourt M, Bollet C, Carlioz A, Martelin R, Gayral J-P, Raoult D. 16S Ribosomal DNA Sequence Analysis of a Large Collection of Environmental and Clinical Unidentifiable Bacterial Isolates. J Clin Microbiol. 2000;38:3623–30.
- 19. Weisburg WG, Barns SM, Pelletier DA, Lane DJ. 16S ribosomal DNA amplification for phylogenetic study. J Bacteriol. 1991;173:697–703.
- 20. Morel A-S, Dubourg G, Prudent E, Edouard S, Gouriet F, Casalta J-P, et al. Complementarity between targeted real-time specific PCR and conventional broadrange 16S rDNA PCR in the syndrome-driven diagnosis of infectious diseases. Eur J Clin Microbiol Infect Dis Off Publ Eur Soc Clin Microbiol. 2015; 34:561–70.
- 21. Stackebrandt E. Taxonomic parameters revisited : tarnished gold standards. Microbiol Today. 2006;33:152–5.
- 22. Sentausa E, Fournier P-E. Advantages and limitations of genomics in prokaryotic taxonomy. Clin Microbiol Infect. 2013; 19:790–5.
- 23. Diop K, Diop A, Khelaifia S, Robert C, Pinto FD, Delerce J, Raoult D, Fournier PE, Bretelle F, Fenollar F. Characterization of a novel Gram-stain-positive anaerobic coccus isolated from the female genital tract: Genome sequence and description of *Murdochiella vaginalis* sp. nov. Microbiologyopen. 2018; 7(3):e00570. doi: 10.1002/mbo3.570.
- 24. Bilen M, Mbogning Fonkou MD, Nguyen TT, Richez M, Daoud Z, Fournier PE, Raoult D, Cadoret F. *Miniphocibacter massiliensis* gen. nov., sp. nov., a new species isolated from the human gut and its taxono-genomics description. Microbiologyopen. 2019; 8(5):e00735. doi: 10.1002/mbo3.735.
- 25. Sasser M. Bacterial Identification by Gas Chromatographic Analysis of Fatty Acids Methyl Esters (GC-FAME). :6.

- 26. Dione N, Sankar SA, Lagier JC, Khelaifia S, Michele C, Armstrong N, et al. Genome sequence and description of *Anaerosalibacter massiliensis* sp. nov. New Microbes New Infect. 2016; 10:66–76.
- 27. Zerbino DR, Birney E. Velvet: Algorithms for de novo short read assembly using de Bruijn graphs. Genome Res. 2008; 18:821–9.
- 28. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol. 2012;19:455–77.
- 29. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, et al. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. GigaScience. 2012; 1:18.
- 30. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014; 30:2114–20.
- 31. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Nucleic Acids Res. 1997; 25:955–64.
- Lagesen K, Hallin P, Rødland EA, Staerfeldt H-H, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids Res. 2007; 35:3100–8.
- 33. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990; 215:403–10.
- 34. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, et al. Fast Genome-Wide Functional Annotation through Orthology Assignment by eggNOG-Mapper. Mol Biol Evol. 2017; 34:2115–22.
- Lee I, Ouk Kim Y, Park S-C, Chun J. OrthoANI: An improved algorithm and software for calculating average nucleotide identity. Int J Syst Evol Microbiol. 2016; 66:1100– 3.
- 36. Meier-Kolthoff JP, Göker M, Spröer C, Klenk H-P. When should a DDH experiment be mandatory in microbial taxonomy? Arch Microbiol. 2013; 195:413–8.
- Lagier J-C, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, et al. Culture of previously uncultured members of the human gut microbiota by culturomics. Nat Microbiol. 2016; 1:16203.
- 38. Hirota K, Hanaoka Y, Nodasaka Y, Yumoto I. *Gracilibacillus alcaliphilus* sp. nov., a facultative alkaliphile isolated from indigo fermentation liquor for dyeing. Int J Syst Evol Microbiol. 2014 Sep;64(Pt 9):3174-3180. doi: 10.1099/ijs.0.060871-0.
- 39. Tang SK, Wang Y, Lou K, Mao PH, Jin X, Jiang CL, Xu LH, Li WJ. *Gracilibacillus saliphilus* sp. nov., a moderately halophilic bacterium isolated from a salt lake. Int J Syst Evol Microbiol. 2009 Jul;59(Pt 7):1620-4. doi: 10.1099/ijs.0.006569-0.
- 40. Carrasco IJ, Márquez MC, Yanfen X, Ma Y, Cowan DA, Jones BE, Grant WD, Ventosa A. *Gracilibacillus orientalis* sp. nov., a novel moderately halophilic bacterium isolated from a salt lake in Inner Mongolia, China. Int J Syst Evol Microbiol. 2006 Mar;56(Pt 3):599-604. doi: 10.1099/ijs.0.63971-0. PMID: 16514034.



0.0050

Fig. 1: Phylogenetic tree highlighting the phylogenetic position of *Gracilibacillus phocaeensis* strain Marseille-P3801T relative to other phylogenetically close members of the family *Bacillaceae*. Sequences were aligned using CLUSTALW, and phylogenetic inferences were obtained using the maximum likelihood method within the MEGA7 software. Numbers at the nodes are percentages of bootstrap values obtained by repeating the analysis 500 times to generate a majority consensus tree.



(a) (b) **Fig. 2:** (a) Gram staining of *Gracilibacillus phocaeensis* strain Marseille-P3801T. (b) Morphological structure of *G. phocaeensis* strain Marseille-P3801T obtained with Hitachi SU5000 scanning electron microscope. Scale bar and acquisition settings are shown of figure.



- 1 2 Fig. 3: Heatmap generated with OrthoANI values calculated using the OAT software between
- 3 Gracilibacillus phocaeensis sp. nov. strain Marseille-P3801. nov. and other closely related species with standing in nomenclature.



7

Fig. 4: Distribution of functional classes of predicted genes according to the clusters of orthologous groups of proteins of *Gracilibacillus phocaeensis* strain Marseille-P3801

1 Table 1: Differential characteristics of 1, Gracilibacillus phocaeensis strain Marseille-P3801 compared to other close bacteria of the genus Gracilibacillus: 2,

2 *G. timonensis* strain Marseille-P2481[6]; **3**, *G. massiliensis* strain Marseille-P1441[15]; **4**, *G. alcaliphilus* strain SG103 [38]; **5**, *G. saliphilus* strain YIM 91119

3	[39]; 6 , <i>G</i> .	orientalis strain	XH-63 [40]	and 7, G. h	halophilus strain	YIM-C55.5	[13]
---	-----------------------------	-------------------	------------	-------------	-------------------	-----------	------

Properties	1	2	3	4	5	6	7
Cell diameter (µm)	0.3-0.6	0.5-0.8 Creamy	0.3-1.8	0.5-0.7	0.7-0.9 Creamy	0.7-0.9	0.3-0.5
Pigmentation	Yellow	orange	White	Creamy white	white	Creamy	White
Oxygen requirement	Aerobic	Aerobic	Aerobic	Aerobic	Aerobic	Aerobic	Aerobic
Gram stain	+	+	+	+	+	+	+
Salt requirement	+	+	+	+	+	+	+
Motility	+	+	+	+	+	+	+
Sporulation	+	+	-	+	+	+	+
Indole	-	-	-	-	_	_	-
Production of :							
Alkaline phosphate	-	-	-	-	+	NA	+
Catalase	+	+	+	+	+	+	+
Oxidase	+	_	-	_	+	_	+
Nitrate reductase	-	_	-	+	+	_	+
Urease	+	+	+	+	+	_	_
β-galactosidase	+	+	-	NA	+	NA	+
α-galactosidase	-	-	+	-	_	NA	-
N-acetyl-glucosamine	-	-	-	+	+	NA	-
Acid from							
L-Arabinose	-	_	-	+	+	+	_
D-mannose	-	_	-	_	+	_	_
D-mannitol	-	_	-	+	+	+	+
D-glucose	-	_	-	+	+	+	+
D-fructose	-	_	-	+	+	+	+
D-maltose	-	_	-	+	+	+	_
D-lactose	_	_	_	+	+	+	_

	DNA G+C	content (mol%)	39.6	39.8	36.05	41.3	40.1	37.1	42.3	
	Habitat		Human gut	Human gut	Cooking salt	Fermentation liquor	Salt lake	Salt lake	Salt soil	
4	+,	positive	reaction;	-,	negative	reaction;	NA,	Not	available	data.

- 1 Table 2: Phenotypic characterization of *Gracilibacillus phocaeensis* sp. nov., strain Marseille-P3801
- 2 based on analytical profile index (API) tests.

Tests	Characteristics	Results Tests	Characteristics	Results
	Alkaline phosphatase	-	Glycerol	-
	Esterase (C4)	+	Erythritol	-
	Esterase Lipase (C8)	+	D-arabinose	-
	Lipase (C14)	+	L-arabinose	-
	Leucine arylamidase	+	D-ribose	-
	Valine arylamidase	-	D-xylose	-
	Cystine arylamidase	-	L-xylose	-
	Trypsin	-	D-Adonitol	-
AI	α-chymotrypsin	-	Methyl βD-xylopyranoside	-
ZIG	Acid phosphatase	+	D-galactose	-
IX	Naphthol-AS-BI-			
A	phosphohydrolase	+	D-glucose	-
	α-galactosidase	-	D-fructose	-
	β-galactosidase	+	D-mannose	-
	β-glucuronidase	-	L-sorbose	-
	α-glucosidase	+	L-rhamnose	-
	β-glucosidase	+	Dulcitol	-
	N-acetyl-β-glucosaminidase	+	Inositol	-
	α-mannosidase	- AP	D-mannitol	-
	α-fucosidase		D-sorbitol	-
	Nituatas to nituitas	C C	Methyl αD-	
	Indules to muntes	- 표	Mathul aD alugaramanasida	-
	Indole Character formation	-	Metnyl aD-glucopyranoside	-
	Glucose fermentation	+	N-acetyi-giucosamine	-
	Arginine dinydrolase	-		-
	Urease 8 shassidase	+	Arbutin Econtin formic citrate	-
	p-giucosidase	-		+
		-		-
AP	p-galactosidase	-	D-cellobiose	-
I 2(Glucose assimilation	-	D-maltose	-
Ň	Arabinose	-	D-lactose	-
L.	Mannose	-	D-melibiose	-
	Mannitol	-	Sucrose	-
	N-acetyl-glucosamine	-	D-trehalose	-
	Maltose	-	Inulin	-
	Potassium gluconate	-	D-melezitose	-
	Capric acid	-	D-raffinose	-
	Adipic acid	-	Starch	-
	Malate	-	Glycogen	-
	Trisodium Citrate	-	Xylitol	-

Phenylacetic acid -	Gentiobiose	-
	D-turanose	-
	D-lyxose	-
	D-tagalose	-
	D-fucose	-
	L-fucose	-
	D-arabitol	-
	L-arabitol	-
	Potassium gluconate	-
	Potassium 2-ketogluconate	-
	Potassium 5-ketogluconate	-

3 +, positive reaction; -, negative reaction.