

Vitamin D deficiency and the COVID-19 pandemic

Patrick Zemb, Peter Bergman, Carlos A Camargo, Etienne Cavalier, Catherine Cormier, Marie Courbebaisse, Bruce Hollis, Fabrice Joulia, Salvatore Minisola, Stefan Pilz, et al.

► To cite this version:

Patrick Zemb, Peter Bergman, Carlos A Camargo, Etienne Cavalier, Catherine Cormier, et al.. Vitamin D deficiency and the COVID-19 pandemic. Journal of Global Antimicrobial Resistance, 2020, 22, pp.133-134. 10.1016/j.jgar.2020.05.006 . hal-03150330

HAL Id: hal-03150330 https://amu.hal.science/hal-03150330

Submitted on 23 Feb 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Contents lists available at ScienceDirect



Journal of Global Antimicrobial Resistance

journal homepage: www.elsevier.com/locate/jgar



Short Communication

Vitamin D deficiency and the COVID-19 pandemic

Patrick Zemb^{a,*}, Peter Bergman^b, Carlos A. Camargo Jr^c, Etienne Cavalier^d, Catherine Cormier^e, Marie Courbebaisse^f, Bruce Hollis^g, Fabrice Jouliaⁿ, Salvatore Minisola^h, Stefan Pilzⁱ, Pawel Pludowski^j, François Schmitt^k, Mihnea Zdrenghea^l, Jean-Claude Souberbielle^m

^a Gynecology Department, Centre hospitalier de Lorient, Lorient, France

^b Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden

^c Department of Emergency Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

^d Department of Clinical Chemistry, University of Liège, CHU de Liège, Liège, Belgium

^e Department of Rheumatology, Cochin Hospital, AP-HP, Paris, France

^f Department of Physiology, Georges Pompidou European Hospital, AP-HP, Paris, France

^g Medical University of South Carolina, Charleston, SC, USA

h Department of Clinical, Internal, Anaesthesiologic and Cardiovascular Sciences, 'Sapienza', Rome University, Viale del Policlinico 155, Rome, 00161 Italy

ⁱ Division of Endocrinology and Diabetology, Department of Internal Medicine, Medical University of Graz, Graz, Austria

^j Department of Biochemistry, Radioimmunology and Experimental Medicine, The Children's Memorial Health Institute, Warsaw, Poland

^k Department of Laboratory Medicine, Centre hospitalier de Lorient, Lorient, France

¹Iuliu Hatieganu University of Medicine and Pharmacy, Cluj-Napoca, Romania

^m Department of Physiology, Necker-Enfants malades hospital, AP-HP, Paris, France

ⁿ Center of Research in Cardiovascular and Nutrition (C2VN), Aix-Marseille University & INSERM 1260, France

ARTICLE INFO

Article history: Received 24 April 2020 Received in revised form 5 May 2020 Accepted 19 May 2020 Available online 29 May 2020

Keywords: Vitamin D coronavirus respiratory infections Covid-19

The COVID-19 (coronavirus disease 2019) pandemic is having a colossal human, societal and economic cost that is encouraging individuals, physicians, collectives and authorities to re-evaluate the benefit/cost ratio of the various resources that could help us to better address the crisis—and limit the risk of its recurrence. In this context, we cannot ignore evidence in a recent 'individual patient data' meta-analysis that vitamin D supplementation protects against acute respiratory infections (ARIs) [1]. In this meta-analysis of 25 randomised controlled trials (RCTs) for which the full data of almost 11 000 individual patients were available, subgroup analyses revealed that the benefit was observed in those who received frequent (e.g. daily) doses of vitamin D but not in those who received bolus doses, and that the effect was largest when

vitamin D was given to individuals with vitamin D deficiency. Since then, several new RCTs have been published with mixed results, with some studies showing beneficial effects of vitamin D on ARIs while other did not. It is known that vitamin D stimulates innate immunity and modulates acquired immunity [2], explaining at least in part how vitamin D may fight against ARIs. Given that vitamin D deficiency is very common [3], especially during the 'cold' season owing to a lack of sunlight exposure, and given that containment at home will prevent sunlight exposure for many people worldwide, even in spring/summer, we believe that vitamin D supplementation should be encouraged, at least in any individual with risk factors of vitamin D deficiency (e.g. obesity, old age, dark skin, wearing covering clothes, no sunshine exposure). We acknowledge that while COVID-19, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is a type of ARI, we have no data enabling us to assert that vitamin D supplementation will reduce the incidence of COVID-19 infection.

http://dx.doi.org/10.1016/j.jgar.2020.05.006

2213-7165/© 2020 The Author(s). Published by Elsevier Ltd on behalf of International Society for Antimicrobial Chemotherapy. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).



^{*} Corresponding author. E-mail address: patrick.zemb@orange.fr (P. Zemb).

However, except for certain rare situations of vitamin D hypersensitivity (e.g. mutation in the CYP24A1 gene or sarcoidosis), daily supplementation with moderate doses of vitamin D_3 is safe. Specifically, there was no increased occurrence of renal stones in recent mega-trials that tested vitamin D_3 doses of 2000 IU/day [4] or 4000 IU/day [5]. In conclusion, despite a lack of direct evidence of an effect of vitamin D status on COVID-19 infection, we believe that vitamin D deficiency is an easily modifiable risk factor of ARIs and should be actively corrected through inexpensive, safe and readily-available vitamin D supplements. Even a small decrease in COVID-19 infections would easily justify this intervention. Nevertheless, we acknowledge that specific high-quality data are needed to demonstrate the efficacy of vitamin D supplementation in the prevention of COVID-19.

Funding

None.

Competing interests

None declared.

Ethical approval

Not required.

References

- [1] Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. BMJ 2017;356:i6583, doi:http://dx.doi.org/10.1136/bmj.i6583.
- Brighenti S, Bergman P, Martineau AR. Vitamin D and tuberculosis: where next? J Intern Med 2018;284:145–62, doi:http://dx.doi.org/10.1111/joim.12777.
 Cashman KD, Dowling KG, Škrabáková Z, Gonzalez-Gross M, Valtueña J, De
- [3] Cashman KD, Dowling KG, Škrabáková Z, Gonzalez-Gross M, Valtueña J, De Henauw S, et al. Vitamin D deficiency in Europe: pandemic? Am J Clin Nutr 2016;103:1033-44, doi:http://dx.doi.org/10.3945/ajcn.115.120873.
- [4] Manson JE, Cook NR, Lee IM, Christen W, Bassuk SS, Mora S, et al. Vitamin D supplements and prevention of cancer and cardiovascular disease. N Engl J Med 2019;380:33–44, doi:http://dx.doi.org/10.1056/NEJMoa1809944.
- [5] Pittas AG, Dawson-Hughes B, Sheehan P, Ware JH, Knowler WC, Aroda VR, et al. Vitamin D supplementation and prevention of type 2 diabetes. N Engl J Med 2019;381:520–30, doi:http://dx.doi.org/10.1056/NEJMoa1900906.