



HAL
open science

Taxonogenomics description of *Bacillus dakarensis* sp. nov., *Bacillus sinesaloumensis* sp. nov. and *Bacillus massiliogabonensis* sp. nov., three new species isolated from human stools

M. Sarr, C.I. Lo, M.L. Tall, A. Fadlane, Bruno Senghor, C. Sokhna, D. Raoult, M. Million, F. Fenollar

► To cite this version:

M. Sarr, C.I. Lo, M.L. Tall, A. Fadlane, Bruno Senghor, et al.. Taxonogenomics description of *Bacillus dakarensis* sp. nov., *Bacillus sinesaloumensis* sp. nov. and *Bacillus massiliogabonensis* sp. nov., three new species isolated from human stools. *New Microbes and New Infections*, 2020, 37, pp.100718. 10.1016/j.nmni.2020.100718 . hal-03159650

HAL Id: hal-03159650

<https://amu.hal.science/hal-03159650v1>

Submitted on 14 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Taxono-genomics description of *Bacillus dakarensis* sp. nov., *Bacillus sinesaloumensis* sp. nov., and *Bacillus massiliogabonensis* sp. nov., three new species isolated from human stools.

Marièma SARR^{1,4}, Cheikh Ibrahima LO^{2,3}, Mamadou Lamine TALL^{1,2}, Amael FADLANE^{1,2},
Bruno SENHOR^{1,4}, Cheikh SOKHNA^{1,4}, Didier RAOULT^{1,2}, Matthieu MILLION^{1,2} and
Florence FENOLLAR^{2,3*}

¹Aix Marseille Univ, IRD, AP-HM, MEΦI, Marseille, France

²IHU-Méditerranée Infection, Marseille, France

³Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

⁴Campus Commun UCAD-IRD of Hann, Dakar, Senegal

* Corresponding author : Florence FENOLLAR, Institut Hospitalo-Universitaire
Méditerranée-Infection, 19-21 Boulevard Jean Moulin, 13385, Marseille cedex 05, France ;
tel +33 413 732 401 ; fax : +33 413 732 402 ; e-mail: florence.fenollar@univ-amu.fr.

Running title: *B. dakarensis* sp. nov., *B. sinesaloumensis* sp. nov., and *B. massiliogabonensis* sp. nov.

Keywords: *Bacillus* sp.; Taxono-genomics; Culturomics; Human stool; Africa.

1 **Abstract:**

2 By microbial culturomics, three *Bacillus* strains were isolated, identified and characterized
3 following the taxono-genomics strategy. *Bacillus dakarensis* strain Marseille-P3515^T
4 (=CSURP3515), *Bacillus sinesaloumensis* strain Marseille-P3516^T (=CSURP3516), and
5 *Bacillus massiliogabonensis* strain Marseille-P2639^T (=CSURP2639) were isolated from
6 human stool samples. The phylogenic analysis, phenotypic characteristics, and genotypic data
7 presented here prove that these three bacteria are different from previously known bacterial
8 species with standing in nomenclature and represent new *Bacillus* species.

9 **Introduction**

10 Genus *Bacillus* was created in 1872 by Ferdinand Julius Cohn [1]. To date, there are 379
11 species and seven subspecies with validly published names. Most of *Bacillus* species are
12 environmental bacteria which are found in food, soil, fresh water and sea. Others *Bacillus*
13 species may be saprophytic [2] or endophytic plants [3]. Otherwise, there are two particularly
14 redoubtable species in public health namely *Bacillus cereus* (associated with food poisoning)
15 and *Bacillus anthracis* (responsible for anthrax) [4, 5].

16 Several bacteria involved in normal physiological functions and predisposition to human
17 diseases must be studied for a better understanding [6]. The culturomics method that isolates
18 bacteria under different culture conditions is complemented in our laboratory by the
19 systematic sequencing of the 16S rRNA gene, which allows us to explore the microbial
20 diversity of the human gut [7-9]. The new species, for which we reported here, were described
21 using a combination of genotypic and phenotypic characteristics, following a taxono-
22 genomics strategy previously described [10, 11].

23 Herein, we present the details of the isolation and taxono-genomics characterization of strain
24 Marseille-P3515^T, strain Marseille-P3516^T, and strain Marseille-P2639^T as type strains of
25 *Bacillus dakarensis* sp. nov., *Bacillus sinesaloumensis* sp. nov., and *Bacillus*
26 *massiliogabonensis* sp. nov., respectively.

27 **Isolation and growth conditions**

28 In 2016, three stool samples were collected aiming to study halophilic bacteria involved in
29 human gut functioning [12]. Marseille-P3515^T and Marseille-P3516^T were isolated from the
30 stool samples of a 17-year-old adolescent boy and 10-year-old girl, respectively and both
31 living in Ndiop, a rural area in Senegal. But strain Marseille-P2639^T was isolated from the
32 stool sample of a 16-year-old boy living in Gabon. After numerous attempts at identification
33 by MALDI-TOF mass spectrometry, no reliable recognition was obtained for the three strains.

34 The screening was performed on a Microflex LT spectrometer (Bruker, Daltonics, Bremen,
35 Germany) as previously described [13]. The obtained spectra (Figure 1) were imported into
36 MALDI Biotyper 3.0 software and analyzed against the Bruker database that was permanently
37 improving with MEPHI database (<https://www.mediterranee-infection.com/urms-data-base>).
38 The strains (Marseille-P3515^T, Marseille-P3516^T, and Marseille-P2639^T) were first isolated
39 after 1 to 2 days of pre-incubation of stool samples at aerobic condition in blood-culture bottle
40 enriched with 5% rumen fluid sterilized by filtration at 0.2 µm and seeded on 5% sheep-blood
41 Columbia agar (bioMérieux) under aerobic condition at 37°C.

42 **Strain identification**

43 After missing identification with MALDI-TOF instrument, the 16S rRNA genes for each
44 strain was sequenced in order to classify these bacteria. These genes were amplified by using
45 universal primer pairs fD1 and rP2 (Eurogentec, Angers, France) and sequenced with the Big
46 Dye® Terminator v1.1 Cycle Sequencing Kit and 3500xL Genetic Analyzer capillary
47 sequencer (ThermoFisher, Saint-Aubin, France), as previously described [14]. CodonCode
48 Aligner software was used to assembly and to correct the 16S rRNA nucleotide sequences
49 (<http://www.codoncode.com>). The sequences of each strain were submitted for BLAST in
50 NCBI database aiming to determine the phylogenetically closest species with standing in
51 nomenclature. It's in this context that we found that strain Marseille-P3515^T exhibited a
52 97.96% sequence identity with *Bacillus circulans* strain NBRC 13626 (Genbank accession
53 number: AY724690), strain Marseille-P3516^T showed a 98.51% sequence identity with
54 *Bacillus humi* strain LMG 22167 (AJ627210), and strain Marseille-P2639^T displayed 98.41%
55 sequence identity with *Bacillus ciccensis* strain 105-2 (KP965576). These values are below
56 the threshold value recommended (<98.7% sequence similarity of the 16S rRNA gene) by
57 authors to delineate new bacterial species within a genus without performing DNA-DNA
58 hybridization [15, 16]. Based on this observation, we declare that these strains are new

59 members of the genus *Bacillus* belonging to the family *Bacillaceae* within the phylum
60 *Firmicutes* (Figure 2).

61 **Phenotypic characteristics**

62 The strains from Senegal were grown easily in aerobic atmosphere. Apparent colonies are
63 obtained after 24h of incubation at 37°C on 5% sheep's blood–Columbia agar medium
64 (bioMérieux, Marcy l'Etoile, France). Strains Marseille-P3515^T and Marseille-P3516^T were
65 recovered from human stool sample from Senegalese village named N'diop. Their colonies
66 appear rounded, beige and shiny with a mean diameter of 1.2 mm. Cells were Gram-positive
67 bacteria, rod shaped and catalase positive. In addition, the oxidase reaction test was positive
68 for these two strains which were mobile and spore forming. These strains were cultured on a
69 halophilic medium with varied NaCl concentration of 50, 75,100 and 150 g/L of NaCl. In
70 parallel, the growth of bacteria was tested on media at different pH as 6, 6.5, 7, 7.5 and 8.
71 Tests have shown that the two strains grow better in 48 hours at pH 7.5, at 75 g/L of NaCl and
72 37 ° C. Consequently, these data prove that strains Marseille-P3515^T and Marseille-P3516^T
73 are halophilic bacteria.

74 Strain Marseille-P2639^T is endospore forming and motile. It was a Gram-negative bacterium
75 that exhibited catalase activity. Bacterial cells do not have an oxidase reaction. They measure
76 3.7 µm in length and 0.8 µm in diameter. Strain Marseille-P2639^T is aerobic bacterium that
77 can grow between 23°C and 45°C in less one day of incubation. Colonies of strain Marseille-
78 P2639^T are white with a mean diameter of 3 mm on 5% sheep's blood-enriched Columbia
79 agar. The strain Marseille-P2639^T is a bacterium that weakly tolerates salt concentrations
80 above 50g/L of NaCl but is able to grow on media with a pH ranging from 6 to 10. The
81 optimal growth temperature is 37° C under aerobic conditions.

82 The shape of these bacteria was highlighted with the Tecnai G20 transmission electron
83 microscope (FEI Company) (Figure 3). The biochemical characteristics of these strains were

84 tested using the API ZYM and API 50 CH strips (bioMérieux) and presented in Table 1. A
85 comparative study of the differential characteristics of these strains with other closely related
86 species is displayed in Table 2.

87 **Genome sequencing**

88 Genomic DNA was extracted using the EZ1 biorobot (Qiagen, Courtaboeuf, France) with the
89 EZ1 DNA tissue Kit and then sequenced on the MiSeq technology (Illumina, San Diego, CA,
90 USA) with the Nextera Mate Pair sample prep kit and Nextera XT Paired end (Illumina), as
91 previously described [17]. The assembly was performed with a pipeline incorporating
92 different softwares (Velvet [18], Spades [19] and Soap Denovo [20]), and trimmed (MiSeq
93 and Trimmomatic [21] softwares) or untrimmed data (only MiSeq software). GapCloser was
94 used to reduce assembly gaps. Scaffolds < 800 base pairs (bp) and scaffolds with a depth
95 value lower than 25% of the mean depth were removed. The best assembly was selected by
96 using different criteria (number of scaffolds, N50, number of N). The degree of genomic
97 similarity of these three strains with closely related species was estimated using the OrthoANI
98 software [22]. *OrthoANI values among Bacillus species (Figure 4) ranged from 68.03%*
99 *between *B. acidicola* and *B. humi* to 88.91% between *B. ciccensis* and *B. massiliogabonensis*.*
100 *These values were lower than 95%, the threshold value suggested delineating new bacterial*
101 *species doing OrthoANI comparison [23].*

102

103 **Genome properties**

104 The genome of strain Marseille-P3515^T is 5,482,351 bp long with 38.6 mol% G+C content
105 (Table 3). It is composed of 11 scaffolds (composed of 62 contigs). Of the 5,282 predicted
106 genes, 5,040 were protein-coding genes and 242 were RNAs (15 genes are 5S rRNA, 7 genes
107 are 16S rRNA, 9 genes are 23S rRNA, 211 genes are tRNA genes). A total of 3,600 genes
108 (71.43%) were assigned as putative function (by cogs or by NR blast). 206 genes were

109 identified as ORFans (4.09%). The remaining genes were annotated as hypothetical proteins
110 (1,014 genes => 20.12%).

111 Strain Marseille-P3516^T has a genome size of 4,556,426 bp long with 37.9 mol% G+C
112 content (Table 3). It is composed of 3 scaffolds which 26 contigs. Of the 4,466 predicted
113 genes, 4 323 were protein-coding genes and 143 were RNAs. These RNA genes are divided
114 into 12 5S rRNA genes, 13 16S rRNA genes, 13 23S rRNA genes and 105 tRNA genes.
115 Strain Marseille-P3516^T possesses 3,095 genes (71.59%) as putative function (by cogs or by
116 NR blast) and 130 genes as ORFans (3.01%). 868 genes (20.08%) were annotated as
117 hypothetical proteins.

118 The genome of strain Marseille-P2639^T counts in all 5,224,786 bp with 37.9 mol% G+C
119 content and is composed of 9 scaffolds which 69 contigs (Table 3). Overall in 5,234 predicted
120 genes, 5,045 genes encode proteins and 189 genes are RNAs (12 genes are 5S rRNA, 18
121 genes are 16S rRNA, 16 genes are 23S rRNA, 143 genes are tRNA genes). A total of 3,476
122 genes (68.9%) were assigned as putative function (by the cogs or by NR blast) while 179
123 genes were recognized as ORFans (3.55%) and 1,151 other genes that remained (22.81%)
124 were annotated as hypothetical proteins.

125 The digital DNA-DNA hybridization (dDDH) between the genomes of three new *Bacillus* and
126 other available genomes of the phylogenetically closest species, was calculated using GGDC
127 online calculator with formula 2 (Table 4). Indeed, 70% threshold is fixed par Meier-Kolthoff
128 *et al.*, to differentiate two distinct species [24]. While, DDH values ranged from 21.5%
129 between *Bacillus ciccensis* and *Bacillus firmus* to 34.9% between *Bacillus sinesaloumensis*
130 and *Bacillus acidicola*, thus supporting the previous data that suggest the classification of
131 these as a new bacterial species (Table 4). Likewise, repartition of genes into the 25 general
132 COG categories was illustrated in Table 5 and Figure 5. The average percentage nucleotide
133 identity calculated using the Average Genomic Identity of Orthologous Gene Sequences

134 (AGIOS) in-house software [10] and number of orthologous genes of *B. dakarensis*, *B.*
135 *sinesaloumensis* and *B. massiliogabonensis* shared with others species are displayed in Table
136 6.

137 **Conclusion**

138 Basing on the results from unique phenotypic characteristics, including API galleries tests,
139 MALDI-TOF spectrum, and phylogenetic and genomic analysis such as 16S rRNA sequence
140 similarity lower than 98.7% and OrthoANI value lower than 95% with the phylogenetically
141 closest species with standing in nomenclature, we consequently proposed strains Marseille-
142 P3515^T, Marseille-P3516^T, and Marseille-P2639^T, respectively as being the type strains of
143 *Bacillus dakarensis* sp. nov., *Bacillus sinesaloumensis* sp. nov., and *Bacillus*
144 *massiliogabonensis* sp. nov., which are new species in the genus *Bacillus*.

145 **Description of *Bacillus dakarensis* sp. nov.**

146 *Bacillus dakarensis* (da.ka.ren'sis, N.L. masc. adj. *dakarensis* of Dakar, the name of the
147 capital of Senegal where the stool sample was collected). The colonies of the strain appear
148 beige and circular on blood agar with a mean diameter of 1.2 mm. The cells are mobile and
149 spores forming. They are Gram-positive bacilli and present positive oxidase and positive
150 catalase activities. The draft genome size of strain Marseille-P3515^T is about 5.33 Mb with a
151 38.6 mol% of G+C content. The 16S rRNA gene sequence and whole-genome shotgun
152 sequence of *Bacillus dakarensis* strain Marseille-P3515^T were deposited in GenBank under
153 accession numbers LT671589 and FTOZ00000000, respectively. The type strain Marseille-
154 P3515^T (=CSURP3515) was isolated from stool sample of 17-year-old-boy living in Senegal.

155 **Description of *Bacillus sinesaloumensis* sp. nov.**

156 *Bacillus sinesaloumensis* (si.ne.sa.lou.men'sis, N.L. masc. adj. *sinesaloumensis* of Sine-
157 Saloum, a former administrative region of Senegal where the village of Ndiop is currently
158 located from which this strain was sampled). Colonies grow on 5% sheep blood Colombia

159 agar plate after 24 hours of incubation under aerobic condition. They are shine and beige with
160 2 mm of diameter. The cells are Gram-positive bacteria, mobile and spores forming. Oxidase
161 and catalase activities were positive. The DNA G+C content of the type strain is 37.9 mol% in
162 a genome sequence long of 4.52 Mb. Its type strain Marseille-P3516^T (= CSURP3516^T) was
163 isolated from a 10-year-old girl from N'diop, a rural area in Senegal.

164 **Description of *Bacillus massiliogabonensis* sp. nov.**

165 *Bacillus massiliogabonensis* (mas.si.li.ga.bo.nen'sis: NL. masc. adj. a composed name
166 designating Marseille and Gabon, the city and the country where the strain and the stool
167 specimen was characterized and collected, respectively). The strain grows at temperatures
168 ranging from 23°C to 45°C in aerobic conditions at an optimum temperature at 37°C.
169 Colonies with a white aspect had a mean diameter of 3 mm on blood agar medium. The strain
170 Marseille-P2639^T is a Gram-negative bacterium and exhibits positive catalase and negative
171 oxidase. The genome of the Marseille-P2639^T strain was 5.13 Mb with 37.9 mol% of G+C
172 content. The potential pathogenicity of the type strain Marseille-P2639^T (= CSURP2639)
173 presently is unknown. It was isolated from the stool sample of a healthy 16-year-old
174 Gabonese boy.

175

176 **Nucleotide sequence accession number.** The following Table shows the 16S rRNA gene and
177 genome sequences accession numbers deposited in Genbank for these three new bacterial
178 species:

Species	Strain Numbers	16S rRNA Numbers	Genome Accession Numbers
<i>Bacillus dakarensis</i>	Marseille-P3515	LT671589	FTOZ00000000
<i>Bacillus sinesaloumensis</i>	Marseille-P3516	LT671591	FTOX00000000
<i>Bacillus massiliogabonensis</i>	Marseille-P2639	LT598571	FZRJ00000000

179

180 **Conflict of interest:**

181 None to declare.

182 **Acknowledgments:**

183 The authors thank Catherine Robert for sequencing the genome, Aurelia Caputo for
184 submitting the genomic sequence to GenBank and Carine Couderc for doing MALDI-TOF
185 reference spectrum.

186 **Funding sources:**

187 This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée
188 Infection, the National Research Agency under the program « Investissements d'avenir »,
189 reference ANR-10-IAHU-03, the Région Provence Alpes Côte d'Azur and European funding
190 FEDER PRIM1.

191 **Ethics and consent**

192 The study and consent procedures were approved by the ethics committee of the Institut
193 Hospitalo-Universitaire Méditerranée Infection (N°2011-11), the National ethics committee
194 of Gabon (N° 0023 / 2013 / SG / CNE) and IFR48 of Marseille France (N° 09-022). The
195 volunteers gave a written consent.

196

- 198 1. Cohn F. Untersuchungen über Bakterien. Beitrage zur Biologie der Pflanzen Heft.
199 1872; 1:127-224.
- 200 2. Lopez MS, Hodde MK, Chamakura KR, Kutty Everett GF. Complete Genome of
201 *Bacillus megaterium* Podophage Page. Genome Announce 2014; 2(2). pii: e00332-14.
202 doi: 10.1128/genomeA.00332-14.
- 203 3. Zhang YZ, Chen WF, Li M, Sui XH, Liu HC, Zhang XX, *et al.* *Bacillus endoradicis*
204 sp. nov., an endophytic bacterium isolated from soybean root. Int J Syst Evol
205 Microbiol 2012; 62:359-63. dx.doi.org/10.1099/ijs.0.028936-0.
- 206 4. Jernigan JA, Stephens DS, Ashford DA, Omenaca C, Topiel MS, Galbraith M *et al.*
207 Bioterrorism-related inhalational anthrax: the first 10 cases reported in the United
208 States. Emerg Infect Dis 2001; 7:933-44. doi.org/10.3201/eid0706.010604.
- 209 5. Bottone EJ. *Bacillus cereus*, a volatile human pathogen. Clin Microbiol Rev 2010;
210 23:382-98. doi.org/10.1128/CMR.00073-09.
- 211 6. Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R and Gordon JI. The
212 human Microbiome Project. Nature 2007; 449: 804-810.
213 <https://www.nature.com/articles/nature06244>.
- 214 7. Lagier JC, Armougom F, Million M, Hugon P, Pagnier I, Robert C, *et al.* Microbial
215 culturomics: paradigm shift in the human gut microbiome study. Clin Microbiol Infect
216 2012; 18:1185–93. doi:10.1111/1469-0691.12023.
- 217 8. Lagier JC, Hugon P, Khelaifia S, Fournier PE, La Scola B, Raoult D. The rebirth of
218 culture in microbiology through the example of culturomics to study human gut
219 microbiota. Clin Microbiol Rev 2015; 28:237–64. doi:10.1128/CMR.00014-14.
- 220 9. Lagier JC, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, *et al.* Culture of
221 previously uncultured members of the human gut microbiota by culturomics. Nat
222 Microbiol 2016; 1:16203. doi:10.1038/nmicrobiol.2016.203.
- 223 10. Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi M, Sentausa E, *et al.* A
224 polyphasic strategy incorporating genomic data for the taxonomic description of novel
225 bacterial species. Int J Syst Evol Microbiol 2014; 64:384–91.
226 doi:10.1099/ijs.0.057091-0.
- 227 11. Fournier PE, Lagier JC, Dubourg G, Raoult D. From culturomics to
228 taxonomogenomics: A need to change the taxonomy of prokaryotes in clinical
229 microbiology. Anaerobe 2015; 36:73-8. doi: 10.1016/j.anaerobe.2015.10.011.
- 230 12. Seck EH, Diop A, Armstrong N, Delerce J, Fournier PE, Raoult D, *et al.* Microbial
231 culturomics to isolate halophilic bacteria from table salt: genome sequence and
232 description of the moderately halophilic bacterium *Bacillus salis* sp. nov. New
233 Microbes New Infect 2018; 23:28-38. doi: 10.1016/j.nmni.2017.12.006.
- 234 13. Lo CI, Fall B, Sambe-Ba B, Diawara S, Gueye MW, Mediannikov O, Sokhna C *et al.*
235 MALDI-TOF Mass Spectrometry: A Powerful Tool for Clinical Microbiology at
236 Hôpital Principal de Dakar, Senegal (West Africa). PLoS One 2015; 10(12):e0145889.
237 doi: 10.1371/journal.pone.0145889.
- 238 14. Morel AS, Dubourg G, Prudent E, Edouard S, Gouriet F, Casalta JP, *et al.*
239 Complementarity between targeted real-time specific PCR and conventional broad-

- 240 range 16S rDNA PCR in the syndrome-driven diagnosis of infectious diseases. Eur J
241 Clin Microbiol Infect Dis 2015; 34:561-70. doi:10.1007/s10096-014-2263-z.
- 242 15. Meier-Kolthoff JP, Göker M, Spröer C, Klenk HP. When should a DDH experiment
243 be mandatory in microbial taxonomy? Arch Microbiol 2013; 195(6):413-8.
244 doi.org/10.1007/s00203-013-0888-4.
- 245 16. Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards.
246 Microbiol Today 2006; 33:152-5.
- 247 17. Lo CI, Sankar SA, Fall B, Ba BS, Diawara S, Gueye MW, *et al.* High-quality draft
248 genome sequence and description of *Haemophilus massiliensis* sp. nov. Standards in
249 Genomic Sciences 2016; 11:31. doi: 10.1186/s40793-016-0150-1.
- 250 18. Zerbino DR, Birney E. Velvet: Algorithms for de novo short read assembly using de
251 Bruijn graphs. Genome Res 2008; 18:821–9. doi:10.1101/gr.074492.107.
- 252 19. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, *et al.*
253 SPAdes: a new genome assembly algorithm and its applications to single-cell
254 sequencing. J Comput Biol 2012; 19:455–77. doi:10.1089/cmb.2012.0021.
- 255 20. Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, *et al.* SOAPdenovo2: an empirically
256 improved memory-efficient short-read de novo assembler. Gigascience. 2012; 1:18.
257 doi:10.1186/2047-217X-1-18.
- 258 21. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
259 sequence data. Bioinformatics 2014; 30:2114–20. doi:10.1093/bioinformatics/btu170.
- 260 22. Lee I, Ouk Kim Y, Park SC, Chun J. OrthoANI: An improved algorithm and software
261 for calculating average nucleotide identity. Int J Syst Evol Microbiol 2016; 66:1100–3.
262 doi:10.1099/ijsem.0.000760.
- 263 23. Richter M, Rosselló-Móra R. Shifting the genomic gold standard for the prokaryotic
264 species definition. Proc Natl Acad Sci U S A. 2009; 106(45):19126-19131.
265 doi:10.1073/pnas.0906412106
- 266 24. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. Genome sequence-based species
267 delimitation with confidence intervals and improved distance functions. BMC
268 Bioinformatics. 2013; 14: 60. doi:10.1186/1471-2105-14-60

Table 1. Phenotypic characterization of *Bacillus dakarensis* sp. nov., strain Marseille-P3515^T, *Bacillus sinesaloumensis* sp. nov., strain Marseille-P3516^T and *Bacillus massiliogabonensis* sp. nov., strain Marseille-P2639^T sp. nov., based on analytical profile index (API) tests.

Tests	Number	Characteristics	Marseille-P3515 ^T	Marseille-P3516 ^T	Marseille-P2639 ^T
API ZYM	2	Alkaline phosphatase	-	-	+
	3	Esterase (C4)	+	+	+
	4	Esterase Lipase (C8)	-	-	-
	5	Lipase (C14)	-	-	-
	6	Leucine arylamidase	-	-	-
	7	Valine arylamidase	-	-	-
	8	Cystine arylamidase	-	-	-
	9	Trypsin	-	-	-
	10	α -chymotrypsin	-	-	+
	11	Acid phosphatase	+	-	+
	12	Naphthol-AS-BI-phosphohydrolase	-	-	-
	13	α -galactosidase	-	+	-
	14	β -galactosidase	-	+	-
	15	β -glucuronidase	-	+	-
	16	α -glucosidase	-	+	-
	17	β -glucosidase	-	+	-
	18	N-acetyl- β -glucosaminidase	-	+	-
	19	α -mannosidase	-	-	-
	20	α -fucosidase	-	-	-
	API 50CH	1	Glycerol	-	-
2		Erythritol	-	-	-
3		D-arabinose	+	-	-
4		L-arabinose	+	-	-
5		D-ribose	-	-	-
6		D-xylose	+	-	-
7		L-xylose	-	-	-
8		D-Adonitol	+	-	-
9		Methyl β D-xylopyranoside	-	-	-
10		D-galactose	-	-	+
11		D-glucose	-	-	-
12		D-fructose	-	-	-

13	D-mannose	-	-	-
14	L-sorbose	-	-	-
15	L-rhamnose	+	-	-
16	Dulcitol	-	-	-
17	Inositol	+	-	-
18	D-mannitol	-	-	-
19	D-sorbitol	-	-	-
20	Methyl α D-mannopyranoside	-	-	-
21	Methyl α D-glucopyranoside	-	-	+
22	N-acetyl-glucosamine	-	-	+
23	Amygdalin	-	-	+
24	Arbutin	-	-	+
25	Esculin ferric citrate	+	+	+
26	Salicin	-	-	+
27	D-cellobiose	-	-	-
28	D-maltose	-	-	-
29	D-lactose	-	-	-
30	D-melibiose	+	-	-
31	Sucrose	-	-	-
32	D-trehalose	-	-	+
33	Inulin	-	-	+
34	D-melezitose	-	-	+
35	D-raffinose	-	-	+
36	Starch	-	-	+
37	Glycogen	-	-	+
38	Xylitol	+	-	-
39	Gentiobiose	-	-	-
40	D-turanose	-	-	-
41	D-lyxose	+	-	-
42	D-tagalose	-	-	-
43	D-fucose	+	-	-
44	L-fucose	-	-	-
45	D-arabitol	-	-	-
46	L-arabitol	-	-	+

47	Potassium gluconate	-	-	-
48	Potassium 2-ketogluconate	-	-	-
49	Potassium 5-ketogluconate	-	-	-

Table 2. Differential characteristics of *Bacillus dakarensis* strain Marseille-P3515^T, *Bacillus sinesaloumensis* strain Marseille-P3516^T, *Bacillus massiliogabonensis* strain Marseille-P2639^T compared with *Bacillus ndiopicus* strain FF3^T and *Bacillus dielmoensis* strain FF4^T.

Property	P3515 ^T	P3516 ^T	P2639 ^T	FF3 ^T	FF4 ^T
Cell diameter (µm)	0.5-1	1-1.9	0.7-1	0.8-1.6	0.5-0.8
Oxygen requirement	Aerobic	Aerobic	Aerobic	Aerobic	Aerobic
Gram stain	+	+	-	+	+
Motility	+	+	-	+	+
Endospore formation	+	+	+	+	-
Production of:					
Alkaline phosphatase	-	-	+	+	+
Acid phosphatase	+	-	+	-	+
Catalase	+	+	+	+	+
Oxidase	+	+	-	-	-
β-Galactosidase	-	+	-	-	+
α-Glucosidase	-	+	-	-	+
Esterase	+	+	+	+	+
Esterase lipase	-	-	-	+	+
Naphthol-AS-BI-phosphohydrolase	-	-	-	-	+
N-acetyl-β-glucosaminidase	-	+	-	-	-
Utilization of:					
Potassium 5-ketogluconate	-	-	-	-	-
D-Xylose	+	-	-	-	-
D-Fructose	-	-	-	-	-
D-Glucose	-	-	-	-	-
D-Mannose	-	-	-	-	-
Habitat	Human sample	Human sample	Stool sample	Human skin	Human skin

+, positive result; -, negative result; NA, data not available.

Table 3. Nucleotide content and gene count levels of genomes of *Bacillus dakarensis* strain Marseille-P3515^T, *Bacillus sinesaloumensis* strain Marseille-P3516^T and *Bacillus massiliogabonensis* strain Marseille-P2639^T

Attribute	<i>B. dakarensis</i>		<i>B. sinesaloumensis</i>		<i>B. massiliogabonensis</i>	
	Value	% of total ^a	Value	% of total ^a	Value	% of total ^a
Size (bp)	5,482,351	100	4,556,426	100	5,224,786	100
G+C content (bp)	2,057,335	38.6	1,711,281	37.9	1,944,649	37.9
Total of genes	5,282	100	4,466	100	5,234	100
RNA genes	242	4.5	143	3.2	189	3.6
Coding sequence size (bp)	4,474,355	81.6	3,886,129	85.3	4,337,909	83.0
Protein-coding genes	5,04	100	4,323	100	5,045	100
Protein assigned to COGs	3,246	64.4	2,741	63.4	3,063	60.7
Genes with peptid signals	595	11.8	478	11.0	561	11.1
Genes with transmembrane helices	1,189	23.6	1,183	27.3	1,278	25.3
Genes associated to mobilome	2,147	42.6	1,714	39.6	2,068	40.9
Genes associated to virulence	951	18.8	790	18.2	948	18.7

Table 4. Genome comparison between three new bacterial strains and closely related species using GGDC and formula 2 (dDDH estimates based on identities over HSP length), upper right. The inherent uncertainty in assigning dDDH values from intergenomic distances is presented in the form of confidence intervals.

	Bdak	Bmas	Bsin	Baci	Bcic	Bcir	Bfir	Bnov	Btim
Bdak	100%	24.3% ± 4.8	27.5% ± 4.9	28.4% ± 4.9	24.5% ± 4.8	28.5% ± 4.9	22.7% ± 4.7	24.3% ± 4.8	23.2% ± 4.8
Bmas		100%	31.5% ± 4.9	32.1% ± 4.9	37.6% ± 5	30.3% ± 4.9	22.0% ± 4.7	26.3% ± 4.8	25.9% ± 4.8
Bsin			100%	34.9% ± 4.9	27.8% ± 4.8	27.1% ± 4.8	26.4% ± 4.9	32.1% ± 5	23.8% ± 4.7
Baci				100%	32.7% ± 4.9	33.9% ± 4.9	27.3% ± 4.9	29.3% ± 4.9	29.3% ± 4.9
Bcic					100%	28.9% ± 4.8	21.5% ± 4.7	25.6% ± 4.8	25.0% ± 4.8
Bcir						100%	23.4% ± 4.8	32.6% ± 4.9	25.0% ± 4.8
Bfir							100%	22.2% ± 4.7	22.6% ± 4.7
Bnov								100%	26.2% ± 4.8
Btim									100%

Bacillus dakarensis strain Marseille-P3515T (**Bdak**), *Bacillus massiliogabonensis* strain Marseille-P2639T (**Bmas**), *Bacillus sinesaloumensis* strain Marseille-P3516T (**Bsin**), *Bacillus acidicola* strain DSM 14745T (**Baci**), *Bacillus ciccensis* strain KCTC 33663T (**Bcic**), *Bacillus circulans* strain NBRC 13626 (**Bcir**), *Bacillus firmus* strain NBRC 15306 (**Bfir**), *Bacillus novalis* strain NBRC 102450 (**Bnov**) and *Bacillus timonensis* strain Marseille-P162 (**Btim**).

Table 5. Number of genes associated with 25 general COGs functional categories




Code	<i>Bacillus sinesaloumensis</i>		<i>Bacillus dakarensis</i>		<i>Bacillus massiliogabonensis</i>		Description
	Value	% of total	Value	% of total	Value	% of total	
[J]	236	5.45	237	4.70	250	4.95	Translation
[A]	0	0	0	0	0	0	Rna processing and modification
[K]	219	5.06	217	4.30	255	5.05	Transcription
[L]	104	2.40	111	2.20	126	2.49	Replication, recombination and repair
[B]	1	0.02	1	0.01	1	0.01	Chromatin structure and dynamics
[D]	52	1.20	56	1.11	58	1.14	Cell cycle control, mitosis and meiosis
[Y]	0	0	0	0	0	0	Nuclear structure
[V]	79	1.82	58	1.15	98	1.94	Defense mechanisms
[T]	151	3.49	197	3.90	198	3.92	Signal transduction mechanisms
[M]	122	2.82	132	2.61	139	2.75	Cell wall/membrane biogenesis
[N]	53	1.22	69	1.36	72	1.42	Cell motility
[Z]	0	0	0	0	1	0.01	Cytoskeleton
[W]	8	0.18	9	0.17	8	0.15	Extracellular structures
[U]	29	0.67	40	0.79	34	0.67	Intracellular trafficking and secretion
[O]	113	2.61	128	2.53	149	2.95	Posttranslational modification, protein turnover, chaperones
[X]	26	0.60	32	0.63	66	1.30	Mobilome: prophages, transposons
[C]	167	3.86	254	5.03	208	4.12	Energy production and conversion
[G]	294	6.80	245	4.86	201	3.98	Carbohydrate transport and metabolism
[E]	291	6.73	368	7.30	329	6.52	Amino acid transport and metabolism
[F]	100	2.31	94	1.86	111	2.20	Nucleotide transport and metabolism
[H]	156	3.60	187	3.71	178	3.52	Coenzyme transport and metabolism
[I]	158	3.65	345	6.84	160	3.17	Lipid transport and metabolism
[P]	172	3.97	237	4.70	227	4.49	Inorganic ion transport and metabolism
[Q]	81	1.87	164	3.25	98	1.94	Secondary metabolites biosynthesis, transport and catabolism
[R]	278	6.43	363	7.20	300	5.94	General function prediction only
[S]	230	5.32	220	4.36	239	4.73	Function unknown
—	1582	36.59	1794	35.59	1982	39.28	Not in COGs

1 **Table 6.** Numbers of orthologous proteins shared between genomes (upper right) ^a

Genomes	<i>Bacillus circulans</i>	NCIMB8773	Marseille-P3515	105-2	WCC4585	IAM12464
<i>Bacillus circulans</i>	4950	1829	1857	1842	2031	2037
<i>Bacillus lentus</i> strain NCIMB8773	60.91	4088	1754	1698	1830	1808
<i>Bacillus dakarensis</i> strain Marseille-P3515	61.84	67.91	5040	1816	2099	2244
<i>Bacillus acidicola</i> strain 105-2	60.54	67.96	67.56	4876	1839	1976
<i>Bacillus gottheilii</i> strain WCC4585	61.69	67.52	70.24	67.66	4450	2285
<i>Bacillus firmus</i> strain IAM12464	59.95	58.64	60.36	59.11	61.28	4922
	PB1NCIMB	LMG21833	AM31D	LMG22167	Marseille-P3516	IFO15566
<i>Bacillus methanolicus</i> strain PB1NCIMB	3410	1704	1524	1627	1628	1715
<i>Bacillus bataviensis</i> strain LMG21833	61.99	5207	1754	2168	2086	2449
<i>Bacillus krulwichiae</i> strain AM31D	53.99	55.98	4596	1774	1793	1880
<i>Bacillus humi</i> strain LMG22167	55.03	58.02	66.45	4842	2316	2198
<i>Bacillus sinosaloumensis</i> strain Marseille-P3516	55.00	57.56	66.84	81.22	4323	2161
<i>Bacillus niacini</i> strain IFO15566	55.76	60.61	66.02	68.48	68.34	5952
	Marseille-P2639	LMG21831	LMG21837	LMG21838	A1-2	DSM13966
<i>Bacillus massiliogabonensis</i> strain Marseille-P2639	5045	2208	2364	2338	2369	1904
<i>Bacillus drentensis</i> strain LMG21831	70.62	5043	2707	2796	2107	1902
<i>Bacillus novalis</i> strain LMG21837	70.47	77.50	5425	3014	2246	1958
<i>Bacillus soli</i> strain LMG21838	70.39	79.26	80.82	5340	2244	1944
<i>Bacillus eiseniae</i> strain A1-2	72.08	69.53	69.15	69.22	5468	1870
<i>Bacillus subterraneus</i> strain DSM13966	60.44	60.27	60.17	60.27	60.07	3465

2 ^aAverage percentage similarity of nucleotides corresponding to orthologous protein shared between genomes (green) and numbers of proteins per genome (blue).

3



Legend:

Bacillus dakarensis strain Marseille-P3515^T

Bacillus sinesaloumensis strain Marseille-P3516^T


Bacillus massiliogabonensis strain Marseille-P2639^T

4

5 **Figure 1:** MALDI-TOF MS reference spectrum of the three new species described above.




6 The reference spectra were generated by comparison of spectra from 12 individual colonies

7 for each species.



9

10 **Figure 2:** Phylogenetic tree highlighting the position of three new bacterial species relative to
 11 its most closely related type strains and validly published. Genbank accession numbers of 16S
 12 rRNA are indicated in parentheses. Sequences were aligned using MUSCLE with default
 13 parameters, phylogenetic inference were obtained using the Maximum likelihood method and
 14 the MEGA 7 software. Numbers at the nodes are percentages of bootstrap values obtained by
 15 repeating the analysis 1000 times to generate a majority consensus tree. The scale bar
 16 indicates a 5% nucleotide sequence divergence.



Legend:

Bacillus dakarensis strain Marseille-P3515^T

Bacillus sinesaloumensis strain Marseille-P3516^T


Bacillus massiliogabonensis strain Marseille-P2639^T

17

18 **Figure 3:** Transmission electron microscopy of three new *Bacillus*. Cells are observed on

19 Tecnai G20 transmission electron microscope operated at 200 keV. Scales are displayed on

20 figures.



21

22 **Figure 4:** Heatmap generated with OrthoANI values calculated using the OAT software for of


23 *Bacillus dakarensis* strain Marseille-P3515^T, *Bacillus sinesaloumensis* strain Marseille-

24 P3516^T and *Bacillus massiliogabonensis* strain Marseille-P2639^T among other related

25 *Bacillus* species with standing in nomenclature.

26

27



29

30 **Figure 5:** Distribution of functional classes of predicted genes according to the COG of
 31 proteins of *Bacillus dakarensis* strain Marseille-P3515^T, *Bacillus sinesaloumensis* strain
 32 Marseille-P3516^T and *Bacillus massiliogabonensis* strain Marseille-P2639^T among other
 33 related *Bacillus* species.

34