



How useful is serology for COVID-19?

Didier Raoult

► To cite this version:

Didier Raoult. How useful is serology for COVID-19?. International Journal of Infectious Diseases, 2021, 102, pp.170-171. 10.1016/j.ijid.2020.10.058 . hal-03216115

HAL Id: hal-03216115

<https://amu.hal.science/hal-03216115>

Submitted on 15 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

How useful is serology for COVID-19?

Didier Raoult¹⁻²

Affiliations :

¹ IHU-Méditerranée Infection, Marseille, France

² Aix Marseille University, IRD, AP-HM, MEPHI, Marseille, France

Corresponding author: Didier Raoult, IHU - Méditerranée Infection, 19-21 boulevard Jean
Moulin, 13005 Marseille, France. Tel.: +33 413 732 401, Fax: +33 413 732 402; email:
didier.raoult@gmail.com

Keywords: COVID-19, seroepidemiology, prevalence

A paper in this journal, reports on the different sero-epidemiological studies carried out around the world for SARS-CoV-2¹. This study helps to put into perspective the potential interest of serology in SARS-CoV-2 infection. The first question was, according to serological techniques, is it possible to identify positive serologies in patients sampled in 2019, before the epidemic? The answer is yes⁸. Depending on the different serological techniques used, cross-reactions may be observed, particularly in children, which could explain why SARS-CoV-2 is to date the only respiratory viral disease that does not affect children^{2;3}. Indeed, in most cases, children are the most affected by respiratory viral infections, this situation is reversed for SARS-CoV-2 for reasons that are not clear at this time. The second major interest that these serological studies could have, would be to confirm *a posteriori* that patients who were included in studies based on purely clinical criteria were actually infected with SARS-CoV-2. Indeed, several studies, including the Recovery study in England and that of Skipper et al.^{4;5}, have included in their evaluation patients who had not been tested, or tested negative, at the time of inclusion. Performing serology retrospectively, as in one recent study for example⁶, ensures that the patients included were indeed affected with SARS-CoV-2. A recent study conducted in France, shows that the predictive value of the clinical diagnosis of SARS-CoV-2 in patients who were subsequently serologically tested for SARS-CoV-2 is less than 50%⁷. This means that studies in which there was no diagnostic test to confirm the diagnosis included at least 50% of people who were not infected with SARS-CoV-2. This also depends on the sensitivity of the serological test. This makes series without a diagnostic test hard to interpret. Seroprevalence studies help to evaluate the extension of epidemics. Incidences based on samples tested by PCR from one area to another have not provided a complete epidemiological picture of the spread and lethality of the disease. In Marseilles, we have systematically tested all symptomatic and non-symptomatic people since February 2020⁸ and have tested 10% of the whole population. We were therefore able to

evaluate the percentage of positive people in the initial epidemic (incidence), at 8%. We had also samples of peoples from three other regions in France: the Ile de France (which includes Paris), the Grand Est, and the New Aquitaine regions. The incidence rates among patients tested in these regions by PCR (unpublished data) surprisingly give a result exactly superimposable to that of seroprevalence in the whole population of these areas, with 8% for Marseille area (Etablissement Français du Sang, France, unpublished data), 10% for Ile de France, 9% for Grand Est and 3% for Aquitaine³. The extent of population seroprevalence in a region or country will provide a sense of the importance of social control measures such as lockdown, the use of masks, or generalized testing. Moreover, it may allow clarifying the lethality in our area independently of the qPCR testing capacity during the outbreak. In the work carried out in this field, particularly in Italy ⁹ and Spain ¹⁰, it has been shown that persons confined in house, when not tested for putative contagiousity, have more often been positive than working unconfined persons. This is counter-intuitive, as the consequence that SARS-CoV-2 carriers have not been detected. This raises the question of the value of the lockdown of families when asymptomatic carriers are not detected. Seroprevalence is also a very good evaluation of the prevention measures of the health care staff. In general, the measurement of seroprevalence in Spain ¹⁰ and Italy ⁹ showed much higher prevalence of the health care personnel, indicating a weakness in prevention measures as well as their high risk. In Marseilles, the percentage of positive test results among health care staff in direct contact remained very low in may 2020, at 3.5% ¹¹, due to the importance of the protective measures deployed. As a matter of fact, we had to import protective artefacts directly from China, as no stock was available in France. All in all, this work opens the door to a more general reflection on the conduct of epidemiological studies in order to refine the consequences of social or protective measures in the spread of the virus in a given site.

Conflict of interest: I declare that I have no competing interests

Funding source : None

Ethical approval : Not applicable

Reference List

- (1) Chih-Cheng Lai, Jui-Hsiang Wang, Po Ren Hsueh. Population-based seroprevalence surveys of anti-SARS-CoV-2 antibody: An up-to-date review. *Int J Antimicrobial Agents* 2020;Submitted.
- (2) Colson P, Tissot-Dupont H, Morand A et al. Children account for a small proportion of diagnoses of SARS-CoV-2 infection and do not exhibit greater viral loads than adults. *Eur J Clin Microbiol Infect Dis* 2020;39:1983-1987.
- (3) Maltezou HC, Magaziotou I, Dedoukou X, Eleftheriou E, Raftopoulos V et al. Children and Adolescents With SARS-CoV-2 Infection: Epidemiology, Clinical Course and Viral Loads. *Pediatr Infect Dis J* 2020;In press.
- (4) <http://www.recoverytrial.net/>. Recovery. Randomised Evaluation of COVID-19 Therapy. 2020. University of Oxford.
Ref Type: Report
- (5) Skipper CP, Pastick KA, Engen NW et al. Hydroxychloroquine in Nonhospitalized Adults With Early COVID-19: A Randomized Trial. *Ann Intern Med* 2020.
- (6) Scholz M, Derwand R, Zelenko V. COVID-19 Outpatients – Early Risk-Stratified Treatment with Zinc Plus Low Dose Hydroxychloroquine and Azithromycin: A Retrospective Case Series Study. *Medicine and Pharmacology* 2020.
- (7) Carrat F, De Lamballerie X, Rahib D, Blanche H. Seroprevalence of SARS-CoV-2 among adults in three regions of France following the lockdown and associated risk factors: a multicohort study. *MedRxiv* 2020;<https://doi.org/10.1101/2020.09.16.20195693>.
- (8) Lagier JC, Million M, Gautret P et al. Outcomes of 3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in Marseille, France: A retrospective analysis. *Travel Med Infect Dis* 2020;36:101791.
- (9) ISTAT. Primi risultati dell'indagine di sieroprevalenza sul SARS-CoV-2. 3-8-2020.
Ref Type: Report
- (10) Pollán M, Pérez-Gómez B, Pastor-Barriuso R et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet* 2020;396:535-544.

97 (11) Edouard S, Colson P, Melenotte C et al. Evaluating the serological status of COVID-19 patients
98 using an indirect immunofluorescent assay, France. *J Clin Microbiol* 2020;Submitted.
99
100