



Updating the repertoire of cultured bacteria from the human being

Ami Diakite, Grégory Dubourg, Didier Raoult

► To cite this version:

Ami Diakite, Grégory Dubourg, Didier Raoult. Updating the repertoire of cultured bacteria from the human being. *Microbial Pathogenesis*, 2021, 150, pp.104698. 10.1016/j.micpath.2020.104698 . hal-03216125

HAL Id: hal-03216125

<https://amu.hal.science/hal-03216125>

Submitted on 2 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Updating the repertoire of cultured bacteria from the human being

Ami Diakite^{1, 2}, Grégory Dubourg^{1, 2}, Didier Raoult^{1, 2*}

¹*Aix Marseille Université, IRD, AP-HM, MEPHI, Marseille, France*

²*IHU Méditerranée Infection, Marseille, France*

***Corresponding author :** Didier Raoult¹, Aix Marseille Université, IHU Méditerranée Infection, Marseille, France, E-mail : didier.raoult@gmail.com

Phone number: +33 4 13 73 24 01

Fax number: +33 4 13 73 24 02

Abstract

The recent renewal of cultural approach has substantially enriched knowledge of the human microbiota, notably through the discovery of new taxa from various anatomical sites. As an increasing number of these recent species are currently considered beneficial or harmful for human health, a constant updating of the repertoire of bacteria and archaea isolated from humans by culture is essential. Herein, we show that the number of cultured bacterial species associated with human beings increased, from 2776 in 2018, to 3253 in 2020, representing a 17% increase in 2 years by adding 477 species, of which 64% are new species (N=307). A wide majority of the species added (i.e., 63%) were isolated using the culturomics approach, while 16% were cultured as part of clinical microbiology laboratories. Human microbiota studies would benefit from the completeness of the repertoire of bacteria associated with human beings, which would require continued efforts to culture microbes from human specimens.

Keywords: Human repertoire; Bacteria: Culturomics

1. Introduction

The Human microbiota is defined as all the bacteria, parasites, viruses, fungi and archaea found in the human body. In terms of biomass, the number of bacteria colonizing the human body has been estimated to be equal to the number of human cells [1]. However, the bacterial diversity associated with the human host remains so far unknown. Indeed, the tools dedicated to the elucidation of the human microbiota suffers from various biases that were extensively discussed [2]. There is however a growing evidence of the influence of the human microbiota in a variety of diseases, including obesity, diabetes and cancer [3–6].

In addition, the advent of bacteriotherapy and the growing use of the fecal microbiota transplantation (FMT) demonstrate the clinical importance of these microorganisms [7,8].

While most of this data is derived from high-throughput sequencing methods by screening for taxa of potential medical interest, biological objects remain essential. Unattributed sequences of potential anticancer prokaryotes have been shown to have already been isolated [9].

Various cultivation approaches have indeed recently contributed to the renewal of culture.

Among these, culturomics is defined as a high-throughput culture technique combining variations in culture conditions and rapid identification of microorganisms using mass

spectrometry type Matrix Assisted Laser Desorption Ionization - Time of Flight (MALDI-

TOF MS) [10]. But there are many other high-throughput cultural approaches currently

developed to reduce the metagenomics dark matter [11,12] resulting in an increasing number of new bacterial species discovered from human being. Reference databases used for

taxonomic assignments currently do not include the entirety of sequences derived from these novel taxa. We therefore believe that updating the previous repertoire is crucial to improve the

identification of sequences generated by high-throughput sequencing approaches. We have

already provided such repertoires in 2015 and 2018 and spontaneously propose herein to

update the repertoire of human bacteria isolated at least once in humans [13,14].

2. Bibliographical methods

A manual query was conducted on three major databases, including Google Scholar, PubMed and Web of science. We started by defining terms and text words referring to Bacteria (Q1), Archaea (Q2), Man (Q3) and Culture Technology (Q4). Then, we combined these terms in order to search the different databases. We build different requests for bacteria and archaea (*Supplementary Data 1*). Considering the period covered by the preceding repertoire update, we focused our literature search on data published from 01/01/2018 to 25/02/2020. Using request for bacteria, we retrieved 5950, 27472 and 44562 papers on Google Scholar, PubMed and Web of science databases, respectively. Using request for archaea, we retrieved 92, 642 and 2,368 papers on Google Scholar, PubMed and Web of science databases, respectively. These data were exported to the EndNote version X9 platform (<https://endnote.com>) in order to sort and analyze the publications. Following removal of duplicates, we obtained 66,622 publications from bacterial request and 2,731 publications from archaeal request (*Figure 1A, 1B*). After analysis of the titles and abstracts, we excluded 56590 studies that did not involve humans, did not use culture-dependent methods and that did not focus on bacteria. Then, we analyzed the full paper and supplementary data of 10,032 publications and excluded those that did not add more bacterial species when compared to the dataset from Bilen et al. in 2018 [14]. We checked the name of each species, filled in their taxonomic information, the origin of their first isolation, whether they were found in pathological situations and whether they were isolated for diagnostic or research purposes. These information were collected from the websites www.ncbi.nlm.nih.gov/taxonomy, www.bacdiv.dsmz.de and the original papers relating to these bacteria. The oxygen tolerance of bacterial species was determined using the database <https://www.mediterranee-infection.com/acces-ressources/base-de-donnees/list-of-prokaryotes-according-to-their-aerotolerant-or-obligate-anaerobic-metabolism/> www.mediterranee-infection.com.

3. Number of species added and description of their characteristics.

Based on our bibliographical research, we were able to add 475 bacterial species and 2 archaea to the last update of the human bacterial repertoire in 2018 by Bilen *et al* [14]. This represents a 17% increase of the repertoire in 2 years (*Figure 2, Supplementary Table 1*). The 477 species added to this repertoire represent 9 phyla with a majority of *Firmicutes* (N=238; 50%), followed by *Actinobacteria* (N=109; 23%), *Proteobacteria* (N=73; 15%) and *Bacteroidetes* (N=46; 10%). Phyla such as *Euryarchaeota* (N=2), *Fusobacteria* (N=2), *Synergistetes* (N=1), *Deferribacteres* (N=1) and *Chlamydiae* (N=1) each represented less than 1% of the total (*Supplementary Figure 1A*). The unique bacterial species isolated in each of these three phyla are *Anaerobaculum hydrogeniformans* (isolated from gut), *Simkania negevensis* (isolated from oral cavity) and *Mucispirillum schaedleri* (isolated from breast milk), respectively. For this latter, it represents the second member of the *Deferribacteres* phylum ever isolated from the human being. This distribution of phylum is consistent with what was demonstrated in the previous human repertoire by Bilen *et al.* in 2018[14]. The 9 phyla are divided into 240 bacterial genera, the majority of which belong to the *Clostridium* (N=25), *Bifidobacterium* (N=20) and *Bacillus* (N=18) genera (*Supplementary Figure 1B*). Anaerobic species constitute half of the species added (N=239; 50%). Majority of the species added, 64% (N=307), are new species discovered from humans samples. The remaining 36% (N=170) species are bacterial species first isolated from the environment (N=75; 16%), animals (N=72; 15%) and food (N=23; 5%) (*Supplementary Figure 2*).

4. Contribution of Culturomics to the expansion of the human bacterial repertoire.

Culturomics is defined as a high-throughput culture method that has so far led to the discovery of hundreds of new taxa associated with humans [10]. In this work, we qualified as culturomics all studies using a high-throughput culture method with or without reference to

the term culturomics and evaluated whether the species included in this study were isolated by this approach or not [11,12]. As a result, 301 bacterial species were isolated as a part of culturomics studies, representing a contribution of 63% of the total number of species added by this work (N=301/477) (*Figure 2, Supplementary Table 1*). Most of these species are strictly anaerobic (64%; N=194/301) and are new taxa (73%; N=220/301). Gut is the most represented anatomical site as it is the source of 73% (N=160/220) of these new taxa, followed by vagina (9%; N=19/220), skin (5%; N=11/220) and less than 5% in other sites such as oral cavity (N=9), urine (N=7), respiratory tract (N=6), breast milk (N=4), blood (N=3) and nasopharyngeal swabs (N=1). These new species are mostly anaerobes (N=150/220; 68%). The remaining species (27%; N=81/301) isolated from humans through culturomics are species of non-human origin known previously. They have been found at 99% (N=80/81) in the gut and 1% (N=1/81) in breast milk with an anaerobic proportion estimated at 54% (N=44/81).

5. The role of clinical microbiology in updating the repertoire of human microbes

Microbial culture in the diagnostic laboratory has also been the subject of several innovations that make it faster and more efficient. The introduction of mass spectrometry type Matrix Assisted Laser Desorption Ionization - Time of Flight (MALDI-TOF MS) in the identification of bacterial species in the routine laboratory has revolutionized bacterial identification by progressively abandoning morphological and biochemical methods [15]. MALDI-TOF MS therefore saved a considerable amount of time and increased the colony identification rate because the database used for spectrum comparison is versatile. It currently represents a powerful screening tool. Indeed, it has probably led microbiologists to test an increasing number of colonies cultivated from clinical specimens that have contributed to expand the human bacterial repertoire. In this work, we tried to evaluate the percentage of species added using the microbial culture method in the diagnostic laboratory. As a result, 16% (N=76/477)

of the species added to the repertoire were isolated from clinical specimen (*Figure 2, Supplementary Table 1*). Most of these species are tolerant to oxygen (i.e., 88%, 67/76) and are new taxa (64/76; 84%). The clinical specimens were respiratory samples (25%; N=16/64), blood cultures (23%; N=15/64), wounds (11%, N=7/64) and other sites with low percentages. The species that are not new (N=12/76; 16%) are species already known from the environment, animals and food.

6. Updating the human archaeal repertoire

Archaea are prokaryotic organisms that were first discovered in extreme environments and were gradually isolated from humans, in particular from the intestinal microbiota and, more recently, from the breast milk [16,17]. Importantly, archaeal species were isolated from clinical specimen, including urine [18]. The most important discovery was the detection of *Methanobrevibacter smithii* in the blood of febrile patients, including subjects with endocarditis. These findings were screened by PCR and confirmed by various assays, including culture. Regarding the repertoire, only 10 of 715 species of archaea known according to NCBI taxonomy have so far been found by culture in humans in 2018[14]. The present study enables to add two species of archaea, including "*Candidatus Methanomethylophilus Alvus*" and "*Candidatus Methanomassiliicoccus intestinalis*", representing a new order of methanogens related to *Thermoplasmata*. Both species were isolated from the human intestinal microbiota, thus increasing the number of archaea isolated by culture from humans to a total of 12 species [19,20].

7. Description of the current human bacterial repertoire

The aim of this work was to update the latest repertoire of bacteria and archaea cultured at least once in humans carried out in 2018 by Bilen. et al [14]. Species identified in this work were isolated from humans according to our data between 1982 and 2020 with a peak in 2019

(N=239; 50%) (*Supplementary Figure 3*). A significant number of species (N=113; 24%) were reported in 2018, the year of the last update where data collection ceased in April 2018. The year 2020 represents 9% (N=41) of the added species, while 18% of the species added (N=84) were found between the years 1980 and 2017, which are periods normally covered by the previous repertoires. This underlines the exhaustiveness of the methodology used herein, that was mostly performed and verified manually. As a result, 477 were added, an increase of 17% in two years. The number of bacteria and archaea isolated from humans thereby increases from N=2776 to N=3253, with the proportion of new species increasing from 19% to 26% (N=831) (*Figure 3, Supplementary Table 2*). Most species are *Firmicutes* 37%, *Proteobacteria* 25%, *Actinobacteria* 25% and *Bacteroidetes* (*Figure 4*). A total of 711 bacterial genera compose the human repertoire to date with the addition of 108 genera yet unknown in humans. The genera *Mycobacterium* (N=163), *Clostridium* (N=133), *Bacillus* (N=108), *Corynebacterium* (N=89) and *Streptococcus* (N=78) are the most represented in terms of number of species. Despite the diversification of human anatomical sites subjected to high-throughput culture analysis, the intestinal tract remains the site where the most species are isolated. Indeed, 52% (N=247) of the total species added by this study (N=477) came from the gut of which 67% (N=166) are new taxa. IHU Méditerranée-Infection contributed up to 20% (N= 639/3253) of the total human bacteria and archaea repertoire, mostly through the discovery of new species (*Figure 5*). The proportion of anaerobic species increased globally from 24% of the previous repertoire (N=662) to 28% in this new repertoire (N=901) with the addition of the 239 anaerobic species, most of which are new species. These findings show that the anaerobes constitute a substantial fraction of the microbial dark matter associated with humans.

8. Conclusion

189 This present study makes it possible to significantly increase the repertoire of human-
190 associated bacteria in a very short period of time. We are currently witnessing the revival of
191 culture, by culturomics, but not only, as many works have reported the isolation of new taxa
192 through innovative cultural strategies [11,12,21,22]. Update of such repertoire is thus crucial
193 as the taxa recently discovered are implied in human health and disease[9,23]. Infectious
194 diseases are also concerned as new species discovered thanks to culturomics were further
195 detected in clinical specimen [24]. We believe that such repertoire could also help in the
196 interpretation of datasets from high sequencing approaches. Indeed, with the current increase
197 of the sequencing depth, the number of sequences amplified is exponentially growing. It can
198 therefore be tricky to determine the relevance relevance of the identification of certain taxa as
199 part of human microbiome studies. Some sequences could be due to the presence of
200 contaminants, or transient microbes that are not commensal to the studied anatomical site.
201 This is also supported by the fact that not all microbes detected are viable in a human
202 specimen [25]. This repertoire, along with other repertoires dedicated to specific human
203 ecological niches [13,26–29], may at least partly help to resolve these interpretation issues.
204 Improving culture approaches to isolate of microbes that are considered "unculturable" is
205 therefore crucial, as shown by Kaboré et al for the culture of *Gemmata* genus. Known for
206 being tedious to cultivate, it was isolated using a culture medium including the skeleton of
207 marine sponges and the aqueous heat filtrate of the skeleton sponge [30]. Belkacemi et al., for
208 the same purpose, developed a successful culture technique based on passive filtration for the
209 culture of *Treponema* species [31]. Thanks to metagenomic studies, we know that the number
210 of species of archaea in humans is much higher compared to the number cultivated [32]. To
211 date, we have listed several 12 species of archaea cultivated at least once in humans. It is
212 therefore imperative to develop culture techniques to be able to isolate them and explore their
213 role as they are often found to be associated with several bacterial infections, such as

periodontitis, chronic sinusitis, muscular or cerebral abscesses. *Methanobrevibacter smithii* was detected in blood cultures from febrile patients, thus suggesting that these microbes should not be longer neglected. Guindo et al. had, for example, developed a technique for cultivating methanogens archaea through the production of hydrogen necessary for their growth by a chemical technique based on the use of iron filings and acetic acid [33]. In parallel, we observed that most of the newly isolated species were anaerobic species, hence the need to develop anaerobic culture strategies. The fact that most of the new taxa isolated from clinical specimen are, on the contrary, tolerant to oxygen, reflects the gradual abandonment of anaerobic culture in clinical microbiology. Yet, Diakit   et al., in analyzing the cost-effectiveness of the culture conditions used in culturomic studies, showed that anaerobic culture conditions allow the isolation of 5 times more bacteria than aerobic culture conditions [34]. Efforts must continue to cultivate bacterial species that do not tolerate oxygen, including extremely oxygen-sensitive bacteria (EOS). To this end, special attention should be paid to maintaining anaerobiosis from specimen collection to incubation [35]. The important contribution of culturomics technique initiated 8 years ago (2012) to the human repertoire demonstrates its effectiveness in highlighting and understanding the diversity of the human microbiota. These innovative approaches to culture techniques are to be encouraged in order to continue isolating bacteria and filling the dark matter of high-throughput sequencing approaches.

References

- [1] R. Sender, S. Fuchs, R. Milo, Revised Estimates for the Number of Human and Bacteria Cells in the Body, PLoS Biol. 14 (2016) e1002533. <https://doi.org/10.1371/journal.pbio.1002533>.
- [2] J.-C. Lagier, G. Dubourg, M. Million, F. Cadoret, M. Bilen, F. Fenollar, A. Levasseur, J.-M. Rolain, P.-E. Fournier, D. Raoult, Culturing the human microbiota and culturomics, Nat. Rev. Microbiol. 16 (2018) 540–550. <https://doi.org/10.1038/s41579-018-0041-0>.

- 240 [3] I. Sekirov, S.L. Russell, L.C.M. Antunes, B.B. Finlay, Gut Microbiota in Health and
241 Disease, *Physiological Reviews*. 90 (2010) 859–904.
242 <https://doi.org/10.1152/physrev.00045.2009>.
- 243 [4] L. Zhao, The gut microbiota and obesity: from correlation to causality, *Nature*
244 *Reviews Microbiology*. 11 (2013) 639–647. <https://doi.org/10.1038/nrmicro3089>.
- 245 [5] N. Larsen, F.K. Vogensen, F.W.J. van den Berg, D.S. Nielsen, A.S. Andreasen, B.K.
246 Pedersen, W.A. Al-Soud, S.J. Sørensen, L.H. Hansen, M. Jakobsen, Gut microbiota in human
247 adults with type 2 diabetes differs from non-diabetic adults, *PLoS ONE*. 5 (2010) e9085.
248 <https://doi.org/10.1371/journal.pone.0009085>.
- 249 [6] L. Zitvogel, L. Galluzzi, S. Viaud, M. Vétizou, R. Daillère, M. Merad, G. Kroemer,
250 Cancer and the gut microbiota: An unexpected link, *Science Translational Medicine*. 7 (2015)
251 271ps1-271ps1. <https://doi.org/10.1126/scitranslmed.3010473>.
- 252 [7] N. Kobyliak, C. Conte, G. Cammarota, A.P. Haley, I. Styriak, L. Gaspar, J. Fusek, L.
253 Rodrigo, P. Kruzliak, Probiotics in prevention and treatment of obesity: a critical view, *Nutr*
254 *Metab (Lond)*. 13 (2016) 14. <https://doi.org/10.1186/s12986-016-0067-0>.
- 255 [8] M. Hocquart, J.-C. Lagier, N. Cassir, N. Saidani, C. Eldin, J. Kerbaj, M. Delord, C.
256 Valles, P. Brouqui, D. Raoult, M. Million, Early Fecal Microbiota Transplantation Improves
257 Survival in Severe *Clostridium difficile* Infections, *Clin Infect Dis*. 66 (2018) 645–650.
258 <https://doi.org/10.1093/cid/cix762>.
- 259 [9] J.-C. Lagier, M. Million, A.H. Togo, S. Khelaifia, D. Raoult, Culturomics provides
260 critical prokaryote strains for anti-*Listeria* and anti-cancer probiotics, *Int. J. Antimicrob.*
261 *Agents*. 54 (2019) 407–409. <https://doi.org/10.1016/j.ijantimicag.2019.05.017>.
- 262 [10] J.-C. Lagier, P. Hugon, S. Khelaifia, P.-E. Fournier, B.L. Scola, D. Raoult, The
263 Rebirth of Culture in Microbiology through the Example of Culturomics To Study Human
264 Gut Microbiota, *Clinical Microbiology Reviews*. 28 (2015) 237–264.
265 <https://doi.org/10.1128/CMR.00014-14>.
- 266 [11] Y. Zou, W. Xue, G. Luo, Z. Deng, P. Qin, R. Guo, H. Sun, Y. Xia, S. Liang, Y. Dai,
267 D. Wan, R. Jiang, L. Su, Q. Feng, Z. Jie, T. Guo, Z. Xia, C. Liu, J. Yu, Y. Lin, S. Tang, G.
268 Huo, X. Xu, Y. Hou, X. Liu, J. Wang, H. Yang, K. Kristiansen, J. Li, H. Jia, L. Xiao, 1,520
269 reference genomes from cultivated human gut bacteria enable functional microbiome
270 analyses, *Nat. Biotechnol*. 37 (2019) 179–185. <https://doi.org/10.1038/s41587-018-0008-8>.
- 271 [12] H.P. Browne, S.C. Forster, B.O. Anonye, N. Kumar, B.A. Neville, M.D. Stares, D.
272 Goulding, T.D. Lawley, Culturing of “unculturable” human microbiota reveals novel taxa and
273 extensive sporulation, *Nature*. 533 (2016) 543–546. <https://doi.org/10.1038/nature17645>.
- 274 [13] P. Hugon, J.-C. Lagier, P. Colson, F. Bittar, D. Raoult, Repertoire of human gut
275 microbes, *Microb. Pathog*. 106 (2017) 103–112.
276 <https://doi.org/10.1016/j.micpath.2016.06.020>.

277 [14] M. Bilen, J.-C. Dufour, J.-C. Lagier, F. Cadoret, Z. Daoud, G. Dubourg, D. Raoult,
 278 The contribution of culturomics to the repertoire of isolated human bacterial and archaeal
 279 species, *Microbiome*. 6 (2018) 94. <https://doi.org/10.1186/s40168-018-0485-5>.

280 [15] P. Seng, M. Drancourt, F. Gouriet, B. La Scola, P.-E. Fournier, J.M. Rolain, D. Raoult,
 281 Ongoing Revolution in Bacteriology: Routine Identification of Bacteria by Matrix-Assisted
 282 Laser Desorption Ionization Time-of-Flight Mass Spectrometry, *Clin Infect Dis*. 49 (2009)
 283 543–551. <https://doi.org/10.1086/600885>.

284 [16] N. Gaci, G. Borrel, W. Tottey, P.W. O'Toole, J.-F. Brugère, Archaea and the human
 285 gut: New beginning of an old story, *World J Gastroenterol*. 20 (2014) 16062–16078.
 286 <https://doi.org/10.3748/wjg.v20.i43.16062>.

287 [17] A.H. Togo, G. Grine, S. Khelaifia, C. des Robert, V. Brevaut, A. Caputo, E. Baptiste,
 288 M. Bonnet, A. Levasseur, M. Drancourt, M. Million, D. Raoult, Culture of Methanogenic
 289 Archaea from Human Colostrum and Milk, *Scientific Reports*. 9 (2019) 18653.
 290 <https://doi.org/10.1038/s41598-019-54759-x>.

291 [18] G. Grine, R. Lotte, D. Chirio, A. Chevalier, D. Raoult, M. Drancourt, R. Ruimy, Co-
 292 culture of *Methanobrevibacter smithii* with enterobacteria during urinary infection,
 293 *EBioMedicine*. 43 (2019) 333–337. <https://doi.org/10.1016/j.ebiom.2019.04.037>.

294 [19] G. Borrel, H.M.B. Harris, W. Tottey, A. Mihajlovski, N. Parisot, E. Peyretailade, P.
 295 Peyret, S. Gribaldo, P.W. O'Toole, J.-F. Brugère, Genome sequence of “*Candidatus*
 296 *Methanomethylophilus alvus*” Mx1201, a methanogenic archaeon from the human gut
 297 belonging to a seventh order of methanogens, *J. Bacteriol*. 194 (2012) 6944–6945.
 298 <https://doi.org/10.1128/JB.01867-12>.

299 [20] G. Borrel, H.M.B. Harris, N. Parisot, N. Gaci, W. Tottey, A. Mihajlovski, J. Deane, S.
 300 Gribaldo, O. Bardot, E. Peyretailade, P. Peyret, P.W. O'Toole, J.-F. Brugère, Genome
 301 Sequence of “*Candidatus Methanomassiliicoccus intestinalis*” Issoire-Mx1, a Third
 302 Thermoplasmatales-Related Methanogenic Archaeon from Human Feces, *Genome Announc*.
 303 1 (2013). <https://doi.org/10.1128/genomeA.00453-13>.

304 [21] S. Ghimire, C. Roy, S. Wongkuna, L. Antony, A. Maji, M.C. Keena, A. Foley, J.
 305 Scaria, Identification of *Clostridioides difficile*-Inhibiting Gut Commensals Using
 306 Culturomics, Phenotyping, and Combinatorial Community Assembly, *MSystems*. 5 (2020).
 307 <https://doi.org/10.1128/mSystems.00620-19>.

308 [22] Y. Chang, F. Hou, Z. Pan, Z. Huang, N. Han, L. Bin, H. Deng, Z. Li, L. Ding, H. Gao,
 309 F. Zhi, R. Yang, Y. Bi, Optimization of Culturomics Strategy in Human Fecal Samples, *Front*
 310 *Microbiol*. 10 (2019) 2891. <https://doi.org/10.3389/fmicb.2019.02891>.

311 [23] C. Gómez-Gallego, S. Pohl, S. Salminen, W.M. De Vos, W. Kneifel, *Akkermansia*
 312 *muciniphila*: a novel functional microbe with probiotic properties, *Benef Microbes*. 7 (2016)
 313 571–584. <https://doi.org/10.3920/BM2016.0009>.

314 [24] G. Dubourg, S. Baron, F. Cadoret, C. Couderc, P.-E. Fournier, J.-C. Lagier, D. Raoult,
315 From Culturomics to Clinical Microbiology and Forward, *Emerging Infect. Dis.* 24 (2018)
316 1683–1690. <https://doi.org/10.3201/eid2409.170995>.

317 [25] S. Bellali, J.-C. Lagier, D. Raoult, J. Bou Khalil, Among Live and Dead Bacteria, the
318 Optimization of Sample Collection and Processing Remains Essential in Recovering Gut
319 Microbiota Components., *Front Microbiol.* 10 (2019) 1606.
320 <https://doi.org/10.3389/fmicb.2019.01606>.

321 [26] K. Diop, J.-C. Dufour, A. Levasseur, F. Fenollar, Exhaustive repertoire of human
322 vaginal microbiota, *Human Microbiome Journal.* (2019).

323 [27] A. Morand, F. Cornu, J.-C. Dufour, M. Tsimaratos, J.-C. Lagier, D. Raoult, Human
324 Bacterial Repertoire of the Urinary Tract: a Potential Paradigm Shift., *J Clin Microbiol.* 57
325 (2019). <https://doi.org/10.1128/JCM.00675-18>.

326 [28] A. Togo, J.-C. Dufour, J.-C. Lagier, G. Dubourg, D. Raoult, M. Million, Repertoire of
327 human breast and milk microbiota: a systematic review., *Future Microbiol.* 14 (2019) 623–
328 641. <https://doi.org/10.2217/fmb-2018-0317>.

329 [29] M.D. Fonkou, J.-C. Dufour, G. Dubourg, D. Raoult, Repertoire of bacterial species
330 cultured from the human oral cavity and respiratory tract. *Future Microbiol.* 13 (2018) 1611–
331 1624. <https://doi.org/10.2217/fmb-2018-0181>.

332 [30] O.D. Kaboré, S. Godreuil, M. Drancourt, Improved culture of fastidious Gemmata
333 spp. bacteria using marine sponge skeletons, *Sci Rep.* 9 (2019) 11707.
334 <https://doi.org/10.1038/s41598-019-48293-z>.

335 [31] S. Belkacemi, J. Bou Khalil, Y. Ominami, A. Hisada, A. Fontanini, A. Caputo, A.
336 Levasseur, J.-C. Lagier, S. Khelaifia, D. Raoult, Passive Filtration, Rapid Scanning Electron
337 Microscopy, and Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass
338 Spectrometry for *Treponema* Culture and Identification from the Oral Cavity, *J. Clin.*
339 *Microbiol.* 57 (2019). <https://doi.org/10.1128/JCM.00517-19>.

340 [32] H.T.T. Huynh, M. Pignoly, V.D. Nkamga, M. Drancourt, G. Aboudharam, The
341 repertoire of archaea cultivated from severe periodontitis, *PLoS ONE.* 10 (2015) e0121565.
342 <https://doi.org/10.1371/journal.pone.0121565>.

343 [33] C.O. Guindo, E. Terrer, E. Chabrière, G. Aboudharam, M. Drancourt, G. Grine,
344 Culture of salivary methanogens assisted by chemically produced hydrogen, *Anaerobe.* 61
345 (2020) 102128. <https://doi.org/10.1016/j.anaerobe.2019.102128>.

346 [34] A. Diakite, G. Dubourg, N. Dione, P. Afouda, S. Bellali, I.I. Ngom, C. Valles, M.L.
347 Tall, J.-C. Lagier, D. Raoult, Optimization and standardization of the culturomics technique
348 for human microbiome exploration, *Sci Rep.* 10 (2020) 9674. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-020-66738-8)
349 020-66738-8.

[35] E. Guilhot, S. Khelaifia, B. La Scola, D. Raoult, G. Dubourg, Methods for culturing anaerobes from human specimen, *Future Microbiol.* 13 (2018) 369–381. <https://doi.org/10.2217/fmb-2017-0170>.

Fig. 1: Flow chart of bibliographical bacterial (A) and archaea (B) request.

Fig. 2: All 477 added species isolated at least once from humans by culture. Using the online tool wordle (www.wordle.net), the size of the name of each species is proportional to the number of times it occurs in the repertoire. The circles realized thanks to the cytoscape software (www.cytoscape.org) represent the partition of the culturomics technique, clinical microbiology and other culture techniques in this update of the repertoire. Red edges represent the contribution of IHU Méditerranée infection and green edges represent that of other laboratories. Orange dots at species level represent new taxa and the grey dots represent already known species other than humans.

Fig. 3: Evolution of the number of known bacterial and archaea species in humans through culture in 2015, 2018 and 2020. Percentage added in relation to the previous repertoire is represented by arrow. Orange proportion represents evolution of new species isolated from 2015 to nowadays.

Fig. 4: Distribution in terms of phylum and genera of the 3253 species of the current human bacterial and archaea repertoire. In blue the distribution of phylum from the last update of the repertoire in 2018 by Bilen et al, and in orange the distribution of phylum added by this study.

Fig. 5: All 3253 species of the current human bacterial isolated at least once from humans using culture. Using the online tool wordart (<https://wordart.com/>), the size of the name of each species is proportional to the number of times it occurs in the database. Circle in dark

- 375 blue represent IHU Méditerranée infection contribution in the total repertoire and circle in
- 376 light blue represent species firstly isolated by IHU Méditerranée infection in environment.