



**HAL**  
open science

## **Pedological characteristics of artificialized soils: A snapshot**

Sophie S. Cornu, Catherine Keller, Béatrice Béchet, Cécile Delolme,  
Christophe Schwartz, Laure Vidal Beaudet

► **To cite this version:**

Sophie S. Cornu, Catherine Keller, Béatrice Béchet, Cécile Delolme, Christophe Schwartz, et al..  
Pedological characteristics of artificialized soils: A snapshot. *Geoderma*, 2021, 401, pp.115321.  
10.1016/j.geoderma.2021.115321 . hal-03293308

**HAL Id: hal-03293308**

**<https://amu.hal.science/hal-03293308>**

Submitted on 2 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Pedological characteristics of artificialized soils: a snapshot

2

3 Sophie Cornu<sup>1\*</sup>, Catherine Keller<sup>1\*</sup>, Béatrice Béchet<sup>2</sup>, Cécile Delolme<sup>3</sup>, Christophe

4 Schwartz<sup>4</sup>, Laure Vidal-Beaudet<sup>5</sup>,

5 1- Aix Marseille Univ., CNRS, IRD, INRAE, Coll France, CEREGE Aix-en-Provence,  
6 France

7 2- Université Gustave Eiffel, GERS, LEE, F-44344 Bouguenais, France

8 3- Université de Lyon, INSA Lyon, DEEP, F-69621 Villeurbanne Cedex, France

9 4- Université de Lorraine, INRAE, LSE, F-54000 Nancy, France

10 5- Institut Agro, Agrocampus ouest, EPHor, F-49045, Angers, France

11 \* Corresponding authors

12

13 **Abstract**

14

15 The onset of industrialization entailed the increase in soil artificialization worldwide.

16 Soils from artificialized zones (SAZ) are the result of urban, industrial, traffic, mining or

17 recreational land use. Knowledge of their characteristics is a prerequisite for evaluating

18 the ability of SAZ to perform key functions and provide soil ecosystem services. Based

19 on a selection of 102 papers published between 1991 and 2017, this article aims at

20 providing a quantitative vision of the pedological characteristics of the different

21 categories of SAZs as a function of the land use and soil type (natural profile, soil

22 developed on allochthonous material). We show that, overall, the inherent

23 characteristics of SAZ topsoil generally depend on the soil type - soil of allochthonous

24 material being more stony and poorer in the < 2 µm fraction - while manageable

25 characteristics are a function of both the soil type and the land use, with the exception of

26 pH that is mainly driven by the land use. However, a good characterization of SAZs

27 requires a more systematic description, sampling and analysis to obtain comprehensive

28 knowledge of their characteristics as several features are still too rarely measured such

29 as cationic exchange capacity, electrical conductivity, soil depth, and nutrient  
30 concentrations. Sealed soils are still too rarely considered despite their large surface  
31 area. Such a characterization would allow a better management of the ecosystem  
32 services that SAZs fulfill.

33

34 **Keywords:** urban soils, mining soils, industrial soils, recreational soils, traffic soils,  
35 artificialization, land use, soil properties, anthropogenic impact

36

37

38 Introduction

39

40 As Vitousek et al. (1997) highlighted several years ago, man has profoundly modified  
41 terrestrial ecosystems, including the soil compartment. This change is particularly  
42 extreme in the case of artificialized environments. In these environments, the question  
43 arises of the persistence of the soil functions as defined by e.g. Robinson et al., (2009)  
44 and Dominati et al. (2010). These functions depend on the soil characteristics. Some of  
45 them, considered as inherent, are therefore little impacted by human activities that  
46 initially focused on agriculture (Karlen et al., (1997); Robinson et al., (2009); Dominati et  
47 al., (2010)). However, for artificialized soils such as Technosols for example, the  
48 hypothesis of no human impact on the so-called inherent characteristics is no longer  
49 valid (Barles et al., 1999). More generally, the actual effect of artificialization on the  
50 pedological profile and associated soil characteristics is still insufficiently documented,  
51 with in particular the absence of synthesis except for organic matter, nutrients or  
52 contamination (e.g. Joimel, 2016). It is essential, however, to determine the soil  
53 characteristics in order to assess the functions they provide, such as 1) water flow and

54 retention, 2) retention and cycling of nutrients, 3) physical stability and support, 4)  
55 maintenance of biodiversity and habitat, and 5) preservation of the natural soil heritage  
56 (Commission of the European Communities, 2006; Bone et al., 2010). Knowledge of both  
57 their inherent and manageable (Dominati et al., 2010) characteristics is therefore a  
58 prerequisite for evaluating the ability of soils in artificialized areas (SAZ) to provide  
59 services in order to propose appropriate management.

60

61 Artificialized areas are generally identified according to their land-use as defined by the  
62 European Corine Land Cover inventory (CLC) that differentiates (i) urban fabric; (ii)  
63 industrial, commercial and transport units; (iii) mine, landfill; and (iv) artificial non-  
64 agricultural. The soils of these artificialized zones were categorized by the International  
65 Union of Soil Sciences (IUSS) in 1998 with land use as the determining factor (the  
66 SUITMA categorization for Soils of Urban, Industrial, Traffic, Mining and Military Areas).

67 It is postulated that a given land use implies a certain degree of soil artificialization  
68 understood as degradation as shown by Vasenev et al. (2012) who point out the  
69 "increasing human load" when moving from recreational areas to residential and  
70 industrial areas. It is also implicitly postulated that the more "artificialized" the soils are,  
71 the more degraded they are (Kurucu and Chiristina, 2008; Vasenev et al., 2012; Peng et  
72 al. 2013; Morel et al., 2015) and that they will be unable to perform their functions,  
73 leading eventually to the abandonment of artificial soils considered as sterile.

74 Attempts at characterizing the impact of artificialization by comparing SAZ  
75 characteristics with those of soils under natural vegetation or agriculture are rare,  
76 except within the study by Joimel et al. (2016) for French SUITMAs. They showed that  
77 SUITMAs exhibit a greater variability of characteristics than that recorded for forested  
78 or agricultural soils. The reasons for this variability may lie in that:

- 79 1- all the SUITMAs were grouped together while specificities might occur among the  
80 different land uses;
- 81 2- each SUITMA category contains soils with contrasted levels of artificialization  
82 and thus different pedological profiles, ranging from (pseudo)natural soils to  
83 Technosols (Morel et al., 2005; Rossiter, 2007; IUSS Working Group World  
84 Reference Base, 2015).

85 This paper aims at (i) providing a quantitative vision of the pedological characteristics of  
86 the different categories of soils of artificialized zones (SAZ) in relation to a soil profile  
87 categorization that we developed based on the descriptions available in the literature  
88 (called soil types hereafter), and in comparison to those of other environments (natural  
89 vegetation and cultivated soils), and (ii) identifying some of the drivers of their  
90 variability (land use or soil type). The level of contamination is beyond the scope of this  
91 paper.

92

## 93 2. Methodology

94

### 95 2.1. Building the corpus of scientific papers

96

97 In France, in 2017, a pluridisciplinary and collective scientific state-of-the-art on the drivers  
98 and the environmental, social and economic effects – positive and adverse- of soil  
99 artificialisation was published (Béchet et al., 2017). Among a global corpus of about 2500  
100 scientific references a corpus of 970 articles on SUITMAs was constituted from an  
101 automatic query on the Web of Science (WoS) on the basis of the script presented in SI-  
102 1. From this corpus, only papers dealing with pedological characteristics were kept for

103 the present review. It was further enriched with additional research on the WoS  
104 including the following key words: urban or mining or industrial and soil and pH or  
105 cation exchange capacity (CEC) or electrical conductivity (EC) or gravel or stone (coarse  
106 elements) or carbonates, or organic matter or compost or C cycles. Articles focusing on  
107 agricultural soils and agricultural activities were excluded as agriculture was not  
108 considered as artificialized soils. Only book chapters and papers in referenced journals  
109 were selected.

110 Out of this corpus a database was built with all the studies that fulfilled the following  
111 two conditions:

112 (i) the soils studied could be assigned with the provided information to a given land  
113 use;

114 (ii) the article contained pedological information and data for surface soil horizons (0  
115 to 0.3 m maximum) and/or a deep horizon (> 0.4 m).

116 The resulting database contains data gathered from 102 articles published from 1991 to  
117 2017.

118 In the database, the soils of the considered studies were assigned to a land use category.  
119 Following the CLC classification, we classified the artificialized surfaces into urban,  
120 industrial, mining, traffic and recreational, the latter referring to parks and urban  
121 forests. Since private gardens were referred to as “residential” in most papers, they were  
122 included in the “urban” land-use. The urban class may also include soils from industrial,  
123 recreational or traffic areas when no further information was provided than “*soils*  
124 *sampled in the city of...*”, resulting in a relatively heterogeneous category. In addition, in  
125 order to provide a vision of the pedological characteristics of the SAZs with reference to  
126 soils in other environments (forest, cultivated land...), “control” or “natural” soils, which  
127 were explicitly designated as such in the articles, were also entered in the database.

128 They were categorized as soils with “natural vegetation”, “crop cover” or “unspecified  
129 cover”.

130 The soils entered in the database consisted either of a single soil profile or a topsoil  
131 sample; the latter consisted either of a single sample or of the average value of a set of  
132 samples (counted as one, as few authors gave individual data). Based on the information  
133 at hand, and when a minimal description of soil profiles was provided, we were able to  
134 categorize the corresponding soil profiles into four rough classes, namely “natural soil  
135 profile”, “natural truncated soil profile”, “soil profile developed on allochthonous  
136 materials (of natural or technogenic origin)”, and “sealed profile”. When the description  
137 was reduced to “mining soil or urban soils” without any further specification, the soil  
138 was categorized in a fifth class called unspecified unsealed soil profile.

139 Both manageable and inherent characteristics were entered in the database. Manageable  
140 characteristics are soil water pH, soil organic carbon (SOC) content, cation exchange  
141 capacity (CEC), electrical conductivity (EC) and bulk density. Nutrient concentrations  
142 were either too seldom reported or were measured with methods that were too  
143 different to be compared. Thus, N, P and K were not included in the database. The  
144 inherent characteristics recorded were stoniness (coarse element content) and < 2  $\mu\text{m}$   
145 fraction content. For each study, we also recorded the morphology of the soil profiles (i.e  
146 description of profile), the type of soil used as a reference, as well as the occurrence and  
147 nature of artefacts (here defined as by-products derived from industrial or artisanal  
148 activities).

149 Methods used to measure the different parameters varied for the different studies. Only  
150 data obtained with methods considered as comparable were entered in the database.  
151 Nevertheless, a variability due to the various methods used cannot be excluded, that is  
152 the reason why the parameters were presented as classes (see below). Soil water pH

153 was measured with the soil/water ratio varying from 1/1 (Chichester and Hauser, 1991;  
154 Johnson and Skousen 1995; Hearing et al., 2004; Howard et al., 2013) to 1/5 (Nikiforova  
155 et al. 2014); soil organic carbon (SOC) content was measured by wet (e.g. Jim, 1998a) or  
156 dry (e.g. Abel et al., 2015a) combustion, CNS analyser (e.g. Hiller 2000; Norra et al.,  
157 2008), dichromate oxidation (e.g. Wang et al., 2007; Oliveira et al., 2014; Ahirval and  
158 Maiti, 2016; Pellegrini et al., 2016); cation exchange capacity (CEC) was measured after  
159 ammonium acetate extraction (e.g. Johnson and Skousen, 1995; Jim, 1998a; Fu et al.,  
160 2011; Ahirval and Maiti, 2016; Pellegrini et al., 2016) or cobalt hexamine extraction (e.g.  
161 Capilla et al., 2006), electrical conductivity (EC) was measured with the soil/water ratio  
162 varying from 1/2 (Johnson and Skousen, 1995) to 1/2.5 (Greinert, 2015); bulk density  
163 was measured on clods (Jim, 1998a) or by the cylinder methods (e.g. Oliveira et al.,  
164 2014; Ahirval and Maiti, 2016); particle size fractionation was performed by sieving and  
165 Stocks law (e.g. Jim, 1998a; Schleuß et al., 1998; Hiller, 2000; Oliveira et al., 2014;  
166 Greinert, 2015) or sieving and the pipette method (e.g. Capilla et al., 2006; Ahirval and  
167 Maiti, 2016; Pellegrini et al., 2016). Coarse elements, when measured, were generally  
168 defined as > 2 mm without specification of an upper boundary size limit (e. g. Hiller,  
169 2000; Beyer et al., 2001; Golubiewski et al., 2006). Schleuß et al. (1998) defined them as  
170 gravels and stones that were less than 0.20 m in size according to the IUSS Working  
171 Group WRB (2015). Jim (1998a) also mentioned boulders.

172

## 173 2.2. Database structure

174

175 For each parameter considered, the data were organized into classes. Consequently, the  
176 database does not contain the absolute value of each soil parameter considered for each



177 soil, but the number of soil samples belonging to each class of the parameter considered.  
178 The classes were defined according to international references as specified for each  
179 parameter in the figure captions (www.fao.org; FAO, 1993; Jim, 1998a, b; Baize, 2000;  
180 Schoeneberger et al., 2002; Vrščaj et al., 2008, among others...).

181 Two depths were considered: topsoils and subsoils. Topsoil thickness was usually 0.3 m,  
182 but ranged in some cases from 0.05, 0.10, 0.15, 0.20 to 0.30 m or was sometimes not  
183 even specified (Lottermoser et al., 1999; Flynn et al., 2003; Chung et al., 2005; Wahsha et  
184 al., 2016). This may have induced a bias, notably for SOC. When several layers were  
185 analyzed within the 0.3 m, a mean value was calculated. The so-called subsoil layer  
186 consisted in the soil horizon ideally below 0.4 m, but the limit could be between 0.4 and  
187 0.6 m. When soil sampling was more superficial than 0.4 m, no subsoil was recorded.

188

### 189 2.3. Data representation and statistical treatments

190 Tables 1 and 2 present the number of topsoil and subsoil samples respectively for each  
191 soil category within each SAZ type and for each parameter. In addition, Table SI-2 gives,  
192 for each of the papers included in the database, the type of land use, the existence of a  
193 reference soil (where relevant), the type of sample (surface or profile), and the  
194 pedological characteristics measured. Data were further represented in Figures SI-1  
195 and 2 as radar plots for references and SAZs topsoils and subsoils with more than eight  
196 occurrences. Correspondence analysis (CA) was performed for each parameter and for  
197 the soil types, SAZs and references with more than eight occurrences. In order to  
198 identify the similarity among the different soil types/SAZs and references, an  
199 agglomerative hierarchical cluster analysis (AHC) was performed on the two first factors  
200 of the CA.

201

## 202 3. Results and discussion

### 203 3.1 Location and publication date of the articles contained in the database

204

205 The earliest papers contained in our database were published in the early 90s and  
206 reported studies related to mining soils, following the decommissioning of many mines  
207 (Figure 1). Among the articles collected, studies dealing with soils associated to traffic,  
208 urban and recreational areas only appeared from the mid-90s. Lastly, soils associated to  
209 industry appeared after 2000. The total number of studies related to mining,  
210 recreational and urban soils in the database is about the same, around 35 to 40, while  
211 those focusing on traffic and industrial soils are less numerous with only 15 and 10  
212 studies respectively (Figure 1).

213 Studies were mainly performed for European sites (accounting for 45 to 65 %  
214 depending on the land use considered, Figure 2). Studies in North America (mainly the  
215 USA) and East-Asia are also well represented in the database (5 to 40 %), while we  
216 found only few studies for the rest of the world, except for mine soils. Lastly, only a third  
217 of the studies provided a reference soil. For the latter, the land use was described only in  
218 60 % of the cases, most of it being covered by natural vegetation (Figure 3).

219

### 220 3.2. Morphological characteristics of soils of artificialized zones

221

222 **SAZ profiles are generally poorly described.** When described, the profiles are not  
223 classified according to any pedological classification. Only a few studies identify them as  
224 Anthrosols or Technosols according to the WRB (IUSS Working Group WRB, 2015). For  
225 one third of the urban soils, and 70 % of the industrial soils, no description at all was  
226 provided, with the notable exception of soils in mining areas. Indeed, for mining soil, the

227 use of the soil as “dump” or “tailing” allowed an easy categorization as soil profile  
228 developed on allochthonous materials, resulting in a particularly low proportion of  
229 unspecified profiles (10 %, Figure 4).

230 **Soils developed on allochthonous materials of natural or technogenic origin are**  
231 **the most common SAZ profiles specified.** They represent more than 50 % of cases in  
232 urban and mining contexts and 10 % to 40 % of cases for the other SAZs (Figures 4 and  
233 5). However, for these soils, the substrate was rarely explicitly described, despite its  
234 potential diversity and its importance in soil profile development. Natural soil profiles  
235 represented a maximum of 20 % of all the studied profiles and were mainly encountered  
236 in recreational, industrial and urban contexts. When encountered in a mining context  
237 (less than 10 % of the cases), they consisted of soils "not physically disturbed" but  
238 considered as receivers of airborne particles from mining or associated smelters or  
239 impacted by acid mine drainage. Truncated natural soil profiles were only encountered  
240 in mining contexts and sealed soils in traffic areas, as expected. Both were nevertheless  
241 poorly represented in the database (13 and 5 cases respectively): truncated natural soil  
242 profiles were thus merged with natural soil profiles in the rest of this analysis and sealed  
243 soils were not further considered.

244 **The vertical differentiation of the SAZs is strongly dependent on the nature of the**  
245 **soil profile.** While the soils with a natural or pseudo-natural pedon (possibly truncated  
246 at the surface) comprise horizon types and sequences that depended on the nature of  
247 the native soils present in the area, the soils developed on allochthonous materials are  
248 characterized by clear limits between layers generally strongly marked by their  
249 formation history (e.g. Jim, 1998a; Abel et al., 2015a, Figures 5a, b, e, f, g). The nature of  
250 the parent materials is generally markedly different from native materials with a strong  
251 vertical contrast with depth (Figures 5c, e). Pedogenetic development is often quite

252 limited with an A over C profile-type (Murolo et al., 2005, Figures 5d, f) as these soils are  
253 rather young as compared to natural soils.

254 Lastly, **SAZs are often characterized by a complex pattern of spatial variability**  
255 resulting from the superimposition of the complexity of their recent history on the  
256 natural heterogeneity. In some cases, the spatial variability becomes metric to  
257 centimetric with discontinuities in the soil cover as in the extreme case of soil  
258 development in pavement joints (Burghardt and von Bertrab, 2016, Figure 5h).

259

### 260 3.3. Inherent pedological characteristics of soils of artificialized zones

261

262 Inherent soil characteristics are considered as stable over a time scale of a few years  
263 (Dominati et al., 2010) in contrast to manageable soil characteristics, which are  
264 considered as more dynamic. The inherent soil characteristics encompass mainly  
265 texture, depth, structure at depth, horizonation and the nature of the mineral phases  
266 (Dominati et al., 2010).

267

#### 268 *3.3.1. SAZ stoniness and fine (< 2 µm) fraction content is a function of land use and soil type*

269 Stoniness is usually considered when assessing agronomic performances because the  
270 presence of coarse materials modulates the surface available for water infiltration,  
271 water storage and root development (Yilmaz et al., 2018). The non-consideration of  
272 these coarse elements may result in an overestimation of the C and N stocks (Rytter,  
273 2012). Stoniness may also create fissures at the interface with fine backfill materials  
274 (Cousin et al., 2003; Wang et al., 2018), thus allowing for preferential flows. However,  
275 the abundance of coarse elements was recorded in only nine percent of the articles,  
276 mainly from soils developed in allochthonous materials (Tables 1 and 2), resulting in 58  
277 and 42 surface and deep horizons respectively (including reference soils, Tables 1 and  
278 2). This low amount of data on soil stoniness is not surprising since Bünemann et al.  
279 (2018) concluded in their literature review that stoniness was not considered as an  
280 important soil characteristic to define soil quality. The two first factors of the CA  
281 performed for the topsoils and subsoils explain 95 and 96 % of the total variance  
282 respectively (Figures 6a and 7a). In both case, three poles are identified by the AHC  
283 made on the two CA factors, one very rich in coarse elements, one with less than 1 %

284 coarse element and one with 1 to 25 % of coarse elements (Figures 6a and 7a).  
285 Nevertheless, these results showed that the reference soils in the database contained  
286 less than 25 % of coarse elements (whatever the land use considered, agriculture or  
287 natural vegetation, Figure 6a), which is consistent with results of the LUCAS European  
288 soil database showing that only Mediterranean and mountain soils had higher coarse  
289 element contents (European Commission – JRC - ESDAC, 2020). When considering SAZs,  
290 the amount of coarse elements is higher than in natural soils, especially at depth  
291 (Figures 7a). This global trend masks a more complex reality. Artificialized natural soil  
292 profiles were poorly represented and did not differ from the reference soils in terms of  
293 stoniness. Most of the soils developed on allochthonous materials in urban and  
294 recreational zones contained 5 to 25 % of coarse elements as seen for the reference  
295 soils, with a small proportion (10 %) containing more than 25%, while mining soils on  
296 allochthonous materials contained generally between 50 and 80 %, of coarse elements  
297 in the topsoils (Figure 6a). Industrial soils exhibit a more complex distribution pattern  
298 and traffic soils on allochthonous materials have a low amounts of coarse elements in  
299 the topsoils (Figure 6a). At depth, all SAZs contained more coarse elements than the  
300 reference soils, mining soils remaining the ones with the largest amounts of stones and  
301 gravels (Figures 7a). Indeed, a layer of “arable” soil is often applied to the surface  
302 notably in recreational soil profiles developed on allochthonous material (e. g. Jim,  
303 1998a; Abel et al., 2015a).

304

305 Conversely, the content of the  $< 2 \mu\text{m}$  fraction was considered since it controls most of  
306 the soil parameters such as cation exchange capacity (CEC), structural stability, and  
307 water holding capacity. The results were measured in fine earth ( $< 2 \text{ mm}$ ). The  
308 distribution of the reference soils in the different classes of  $< 2 \mu\text{m}$  percentage is similar

309 to that provided at the global scale by Shangguan et al. (2014) and Dai et al. (2019). The  
310 two first factors of the CA performed for the topsoils and subsoils explain 98 and 93 % of  
311 the total variance respectively (Figures 6b and 7b). In both cases, three poles are  
312 identified by the AHC made on the two first CA factors, one with concentrations in the <  
313 2  $\mu\text{m}$  fraction lower than 15 %, one with a contribution of soils having concentrations in  
314 the < 2  $\mu\text{m}$  fraction higher than 35 % and, one with a large contribution of soils having a  
315 < 2  $\mu\text{m}$  fraction ranging from 15 to 35 % (Figures 6b and 7b). In general, SAZ topsoils  
316 have a < 2  $\mu\text{m}$  fraction lower than 15 % (Figure 6b), unlike reference topsoils, which  
317 have a < 2  $\mu\text{m}$  fraction content between 35 and 60 %. However, topsoils of urban natural  
318 soil profiles, mining soils on allochthonous materials and unspecified recreational soils  
319 may exhibit concentrations in the < 2  $\mu\text{m}$  fraction ranging from 15 to 35 % as some  
320 topsoils of the reference soils (Figure 6b). At depth, the situation is more contrasted  
321 with natural profiles that are clearly differentiated from SAZs. Recreational soils on  
322 allochthonous materials have the lowest concentration in the < 2  $\mu\text{m}$  fraction (Figure  
323 7b). Since the same SAZs are also richer in coarse elements than the reference soils, the  
324 < 2  $\mu\text{m}$  fraction in the whole soil matrix is even lower compared to the reference soils.

325

326 This analysis clearly shows that, while as expected SAZ with a natural profile do not  
327 differ much in terms of stoniness and < 2  $\mu\text{m}$  content from reference soils, SAZ on  
328 allochthonous materials are generally stonier with a lower < 2  $\mu\text{m}$  fraction.

329

### 330 *3.3.2. SAZs are characterized by the presence of artefacts*

331 Beside the occurrence of the minerals classically encountered in soils, a characteristic of  
332 SAZs is their content in artefacts of different sizes, from coarse elements, which  
333 constitute a large fraction of the allochthonous soil profiles, to micro artefacts such as

334 ash deposition also encountered in SAZs with natural profiles (Fu et al., 2011; Howard  
335 and Olriki, 2016).

336 The nature of these materials depends on the type of artificialized zone. Due to their  
337 generally long and complex history, urban soils on allochthonous material contain a  
338 wide variety of technogenic materials such as building/demolition materials (brick  
339 (Nehls et al., 2013), mortar, nails, glass, wood, plaster (Schleuß et al., 1998; Howard et  
340 al., 2013; Greinert, 2015)), ashes, coals, slags, asphalt and ferrous residues (Howard et  
341 al., 2013), industrial deposits (Greinert, 2015), municipal waste (Schleuß et al., 1998;  
342 Greinert, 2015), and stormwater sediments (El-Mufleh et al., 2014a,b). On the contrary,  
343 other SAZ contain more specific materials related to their land use: (i) disturbed  
344 recreational soils may contain demolition materials (Abel et al., 2015a) or filling with  
345 materials of natural origin (Jim, 1998a); (ii) soils from traffic areas contain building  
346 materials, ashes, coals, slag, ballast (Hiller, 2000), natural embankment (Kida and  
347 Kawahigashi, 2015) or dredged sediments for soils near waterways (Capilla et al., 2006);  
348 (iii) soils from quarries contain rock debris with a large particle size distribution (Fu et  
349 al., 2011), while metalliferous mining soils contain sludge, mine wastes (Pellegrini et al.,  
350 2016) or even foundry wastes when associated with smelters, and soils from coal and  
351 lignite mining contain various types of spoils (Shankar et al., 1993; Kundu and Ghose ,  
352 1994; Pedrol et al., 2010; Adeli et al., 2013; Clouard et al., 2014; Ahirwal and Maiti, 2016;  
353 Kolodziej et al., 2016), with the presence of fossil carbon. To a much less extent, airborne  
354 technogenic materials of the same nature are also encountered in SAZ with natural  
355 profiles (Fu et al., 2011; Howard and Olriki, 2016).

356

357 For all the inherent characteristics, the main differences are observed between the  
358 reference soils and soils on allochthonous materials. The latter are modified to such an



359 extent that the distinction between inherent and manageable properties might not be  
360 meaningful anymore. In addition, it remains to be clarified whether allochthonous  
361 materials will follow the same line of evolution as natural parental material.

362

### 363 3.4. Soil manageable pedological characteristics in soils from artificialized zones

364

365 Soil characteristics are considered as dynamic or manageable (Dominati et al., 2010)  
366 when they can be impacted by land use or management decisions (Karlen et al., 2003).  
367 In the following, we address the physical manageable properties (bulk density,  
368 hydrodynamic properties, aggregation), organic matter concentration and physico-  
369 chemical manageable characteristics (pH, CEC, electrical conductivity).

370

#### 371 *3.4.1. SAZs have a large range of bulk densities depending on their land use and* 372 *soil profile type*

373

374 The two first factors of the CA performed for the topsoils and subsoils explain 79 and 75  
375 % of the total variance respectively (Figures 6c and 7c).

376 The topsoil bulk densities of reference soils are lower than  $1.4 \text{ g cm}^{-3}$  under natural  
377 vegetation with a mode at  $1\text{-}1.2 \text{ g cm}^{-3}$  (Figure 6c). Cultivated soils have a bulk density  
378 mode at  $1.4\text{-}1.55 \text{ g cm}^{-3}$  with nearly 10 % of the soil having a bulk density greater than  
379  $1.55 \text{ g cm}^{-3}$  (Figure 6c), which agrees with results reported by Kern (1995). SAZs have an  
380 even bulk density distribution from low densities ( $< 1 \text{ g cm}^{-3}$ ) to densities of  $1.7 \text{ g cm}^{-3}$ ,  
381 with a small percentage of very high bulk densities ( $> 1.8 \text{ g cm}^{-3}$ , Figure 6c). This  
382 distribution indicates a higher variability linked both to the land use and the soil profile  
383 type. Recreational soils with natural profiles exhibit bulk densities comparable to those  
384 measured for reference soils, while natural soils at mining sites have a higher proportion

385 of bulk densities larger than  $1.55 \text{ g cm}^{-3}$  (35 %) since their topsoil has often been  
386 stripped (truncated) and compacted (Figure 6c). SAZs on allochthonous materials  
387 exhibit bulk densities comparable to those encountered in cultivated soils (Figure 6c).  
388 Edmondson et al. (2011) and Pouyat et al. (2007) showed that the urban soils for the  
389 City of Leicester (UK) (136 sites with a median bulk density class of  $0.81\text{-}1 \text{ g cm}^{-3}$ ) and  
390 Baltimore (USA) ( $0.71$  to  $1.74 \text{ g cm}^{-3}$ , with a median value of  $1.18 \text{ g cm}^{-3}$ ) were less  
391 compact than those of the surrounding agricultural area, while Jim's study of the city of  
392 Hong Kong (1998b) showed that in two-thirds of the cases, urban soils of alignment  
393 trees were often very compacted ( $1.4$  to  $2.2 \text{ g cm}^{-3}$ ), with bulk densities greater than  $1.6$   
394  $\text{g cm}^{-3}$ , which is the threshold for root development (Hanks and Lewandowski, 2003).  
395 Scharenbroch et al. (2005) also showed that the bulk density was a function of the age of  
396 the residential area with bulk densities around  $1.73 \text{ g cm}^{-3}$  in new residential areas  
397 against  $1.41 \text{ g cm}^{-3}$  in old residential areas. In recreational areas, studies showed an  
398 extreme spatial variability of the topsoil bulk density with locally very high bulk density  
399 values linked to the spatial variability of trampling (Jim, 1998a; Cakir et al., 2010).  
400 Traffic topsoils on allochthonous materials have very low bulk density (Figure 6c). At  
401 depth, SAZs generally exhibit higher bulk density than reference soils, especially mine  
402 soils developed on allochthonous materials (Figure 7c). This is however not the case for  
403 urban soils on allochthonous materials or recreational soils (Figure 7c).  
404 The strong compaction of certain types of SAZ results in (i) a reduced total porosity, (ii)  
405 a few coarse pores and a large percentage of fine pores (Baumgartl, 1998; Yang and  
406 Zhang, 2011), which induces (iii) low aeration, (iv) a poorly effective drainage (Jim,  
407 1998b) and increases (v) the risks of runoff and erosion (Gregory et al., 2006).  
408

409 *3.4.2. SAZs have a highly variable organic carbon content*

410

411 Soil organic carbon (SOC) content is one of the most frequently reported soil parameters  
412 in the database with 63 references resulting in 410 topsoil and 100 subsoil SAZ SOC data  
413 respectively (Tables 1 and 2). In France, a country considered as having a very high  
414 pedovariability (Minasny et al., 2010), Joimel et al. (2016) observed that the median soil  
415 organic carbon (SOC) concentration of the topsoil (0-0.3 m) was 9 g kg<sup>-1</sup> for vineyards  
416 and orchards, 14.9 g kg<sup>-1</sup> for cultivated soils, 24.8 g kg<sup>-1</sup> for grassland, 26.2 g kg<sup>-1</sup> for  
417 gardens, 26.8 g kg<sup>-1</sup> for forests and 37.3 g kg<sup>-1</sup> for SAZs grouped in one category (all  
418 SUITMAs). The latter seem thus to contain higher SOC concentrations than vineyards  
419 and orchards, agricultural and grassland soils, while the difference is less marked with  
420 forest soils. On the other hand, the authors showed that the variability of the SOC  
421 concentrations in SAZs was higher than that observed for other land uses (as also seen  
422 on Figure 6d). This result probably derives from the high variability of the soil types  
423 grouped under the term “SAZ” as observed in our database.

424 The two first factors of the CA performed for the topsoils and subsoils explain 74 and  
425 64 % of the total variance respectively. The AHC made on the two first CA factors  
426 identified three types of profiles for the topsoils and four at depth (Figures 6d and 7d).

427 While urban and industrial topsoils have a comparable level of SOC to natural soils  
428 whatever the soil profile considered, 40% of traffic and 50% of mining topsoils on  
429 allochthonous materials are very poor in SOC (Figure 6d). Recreational natural profiles  
430 also exhibit a large amount of very poor SOC topsoils, notably in natural profiles (Figure  
431 6d). Soil organic carbon concentrations in the top 0.30 m of urban zones are often higher  
432 than in agricultural soils (Pouyat et al., 2006; Lorenz and Lal 2009; Edmondson et al.,  
433 2014; Vasenev et al., 2014; Cambou et al., 2018; Figures 6d) and sometimes higher than

434 those of some forest soils, in the case of soils in residential neighbourhoods (Pouyat et  
435 al., 2002; Cambou et al., 2018). It may be one of the factors responsible for the lower  
436 bulk density measured in urban topsoils (Edmondson et al., 2011; 2014). In addition,  
437 SOC concentrations in urban soils vary widely. For example, in the city of Stuttgart,  
438 Germany, SOC concentrations in soils ranged from 5 to 207 g C kg<sup>-1</sup> (Lorenz et al., 2006).  
439 The soil cover for a given land use may be responsible for this variability. As an example,  
440 tree-planted gardens had higher SOC concentrations than non-planted community  
441 gardens or non-domestic spaces planted with trees or lawns (Edmondson et al., 2014).  
442 However, the difficulty of this analysis lies in the very variable depth of sampling of the  
443 topsoil horizons, which ranges from 0.05 to 0.30 m depending on the authors. This  
444 variability probably induces a bias as SOC decreases exponentially with depth over the  
445 first 0.50 m of soil.

446 At depth, the differences are more marked among SAZs, especially soils on  
447 allochthonous materials, with some mining soils having lower concentrations than both  
448 the reference soils and urban soils (Figure 7d). Conversely, some urban and industrial  
449 soils have a higher SOC concentration at depth than the reference soils (Figures 7d) as  
450 also shown by Cambou et al. (2018) for Paris and New York. In some remediated soils in  
451 New York City, for instance, the concentrations reached 160 g C kg<sup>-1</sup> at 0.40 m depth and  
452 were still 80 g C kg<sup>-1</sup> at 1 m depth (Huot et al., 2017). Soil organic carbon concentration  
453 along the whole soil profile is also investigated to assess the soil potential for climate  
454 change mitigation through the calculation of carbon stocks. The depth of SOC  
455 accumulation seems to be related to the history of use, which determines the dynamic of  
456 the SOC stock changes (Bae and Ryu, 2015; Cambou et al., 2018) but is still too seldom  
457 measured to give reliable trends (Cambou et al., 2018).

458 However, some authors have shown that urbanization, through land-use change and the

459 removal of vegetation cover, creates in some areas a critical SOC stock reduction, e. g. a  
460 30 % reduction in 25 years for an exponentially growing city such as Changzhou  
461 (increase in surface area by a factor of 10 in 20 years from 1980 to 2000; Tao et al.,  
462 2015). Obviously, organic carbon stock decreases or increases in SAZs need to be  
463 investigated in greater detail.

464

#### 465 *3.4.3. The pH of SAZs is mainly basic*

466 The water pH is the other most frequently reported soil parameter in the database with  
467 64 references resulting in 416 topsoil and 115 subsoil SAZ pH data (Tables 1 and 2).

468 The two first factors of the CA performed for the surface and deep horizons explain 70  
469 and 66 % of the total variance respectively. The AHC made on the two first CA factors  
470 identified three types of profile for the topsoils and four at depth (Figures 6e and 7e).

471 pH of all SAZs is higher than that of the reference soils (Figures 6e and 7e). This is in  
472 agreement with Joimel et al. (2016) who found that the pH of SAZs was mainly basic  
473 while the pH of agricultural and natural soils varied over a very wide range with rather  
474 acidic soils in forest and grassland, neutral in agriculture and rather basic for orchards  
475 and vines.

476 pH values above 7 observed for urban, recreational, industrial and traffic infrastructure  
477 at both depths (Figures 6e and 7e) are due to the frequent presence of 1) technogenic  
478 carbonated materials, notably due to rubble for profiles on allochthonous materials (Jim,  
479 1998b; Abel et al., 2015b); 2) carbonate, siliceous and fossil carbon microparticles for all  
480 profile types (Howard and Orlicki, 2015), whose pH ranges mostly from 8 to 12; 3)  
481 and/or the use of road deicing products in winter, which may result in the formation of  
482 salty alkaline soils as observed in a cold urban context (Nikiforova et al., 2014).

483 Mining soils and sometimes industrial soils can exhibit very acid pH values (Figures 6e

484 and 7e), with low pH (< 5) accounting for 40 to 50% of the cases depending on the soil  
485 type considered. In the case of soils from mining areas, the acidic pH are usually  
486 restricted to the metalliferous areas, since soils from coal and lignite mine areas are  
487 predominantly neutral to alkaline (Ogala et al., 2012). Some very low pH (less than 4.5)  
488 are observed in nearly 30 % of the allochthonous soils from metalliferous areas (Figure  
489 7e) that can notably be due to the presence of pyrite in the mine spoils (Schaaf and Hüttl,  
490 2005; Rivas-Pérez et al., 2016).

491

#### 492 *3.4.4. Electrical conductivity is very high in some SAZ types*

493 The electrical conductivity (EC) data are scarce (Tables 1 and 2) as this parameter was  
494 mainly measured in soils where high EC values were expected (Table SI-3f). In most of  
495 the SAZ types, the EC range was not different from that of the reference soils and was  
496 always lower than 2 mS cm<sup>-1</sup> which is the limit for non-saline soils (Baize, 2000;  
497 Schoeneberger et al., 2002). With EC above this threshold, saline soils represent 3.4 % of  
498 soils worldwide according to the FAO ([www.fao.org](http://www.fao.org)). However, in 20 % of urban and  
499 mining soils, EC higher than 2 mS cm<sup>-1</sup> were recorded and EC values exceeding 4 mS cm<sup>-1</sup>  
500 were encountered in about 15 % of the cases. Some soils reached extremely high EC  
501 ( $\geq 16$  mS cm<sup>-1</sup>), in relation with the salinization observed in Northern cities like Chicago  
502 (Scharenbroch and Catania, 2012) or Moscow (Nikiforova et al., 2014) due to the use of  
503 salt for deicing roads and the concentrated drainage observed at some metal mine sites  
504 (Pellegrini et al., 2016). At depth, high EC was observed only for mining soils (Table SI-  
505 3f).

506

#### 507 *3.4.5. CEC is rarely measured in SAZs*

508 CEC was measured in only 12% of the studies (Tables 1 and 2), resulting in the absence

509 of data for some SAZs or some soil types, notably for deep horizons (Table 2). If one  
510 considers the soil with CEC larger than  $6 \text{ cmol}^+ \text{ kg}^{-1}$  (which is a low value if fertility is  
511 considered), there is no difference between the SAZs and the reference soils with the  
512 exception of the 10 % of traffic soils on allochthonous materials and 10 to 20 % of the  
513 mining soils on allochthonous materials or undefined. However, these CECs were  
514 measured for the fraction less than 2 mm, which in the case of soils on allochthonous  
515 materials may represent a rather small fraction of the whole soil because of their high  
516 stoniness as mentioned above.

517

#### 518 4. Synthesis and recommendations

519 Among the different SAZs, more data on pedological characteristics are available in the  
520 literature on urban, mining or recreational soils than on industrial and traffic soils. The reason  
521 behind this lower representation of industrial and traffic soils is probably their partial  
522 inclusion in urban soils as very often only the mention “soil sampled in an urban area” is  
523 reported in the articles. Care should be taken to better describe the land use in SAZ studies. In  
524 addition, despite their surficial abundance (70 % of the artificial surfaces) only a few studies  
525 have focused on sealed soils, which were therefore not considered in this synthesis. However,  
526 the few existing studies showed that sealed soils can store a substantial amount of SOC  
527 (Raciti et al., 2012), despite the decrease in the SOC stock in the first years after sealing (Jim  
528 et al., 1998b; Wei et al., 2014). The amount of organic carbon stored in sealed soils, and the  
529 potential importance of its role, have not been sufficiently investigated to draw general  
530 conclusions and more studies dedicated to this soil type should be performed.

531 For unsealed soils, we showed that different soil types are encountered, ranging from only  
532 slightly disturbed soils to Technosols on allochthonous material, the latter being the most

533 abundant in the literature. Studies often focus on disturbed areas, but it remains necessary to  
534 assess urban land surfaces with natural or little disturbed profiles for functions such as  
535 supporting urban food production.

536 We also showed that both the soil profile type and the land use determine the  
537 pedological characteristics of SAZs. While inherent characteristics are strongly linked to  
538 the type of soil profile, with allochthonous soils being richer in coarse elements and  
539 poorer in  $< 2 \mu\text{m}$  fractions, manageable characteristics are linked either to land use (the  
540 case of pH), or to both land use and soil type (the case of organic carbon or bulk density).  
541 Here again, the frequent absence of descriptions of soil profiles prevents a detailed  
542 understanding of the determinants of the acquisition of their pedological characteristics.  
543 In addition, the characterization SAZs is mainly limited to their surface horizons  
544 (topsoils) and not enough studies have been conducted on subsoils. Drawing  
545 conclusions from our set of data for deep horizons is thus more difficult. This highlights  
546 the need for a more systematic study of soil profiles. The articles reporting information  
547 at depth point to the importance of subsoils in fulfilling some ecosystem services such as  
548 carbon storage as already observed in natural soils (Balesdent et al., 2018) or, on the  
549 contrary, preventing water infiltration.

550 Lastly, not all soil characteristics are investigated systematically. As observed from the  
551 102 papers in the database, some parameters were either too seldom reported (e.g. total  
552 soil depth) or with too various methods (e.g. nutrients, pH other than water pH) to be  
553 included for discussion in this analysis. Some parameters were discussed but in a limited  
554 number of studies (e.g. % coarse elements, EC and CEC).

555 To take advantage of future data to be produced on SAZs and thus improve their  
556 characterization, we suggest that the following points should be systematically



557 addressed:

558 - the history of the soil formation by systematically associating data on land use,  
559 historical background, demographic growth, economic growth, and typology of uses, to  
560 the studied profiles;

561 - the whole soil profile must be considered when characterizing the soil physicochemical  
562 properties on well-defined horizons;

563 - the use of a common set of sampling and analytical methods and a common typology of  
564 the technogenic materials present in soils;

565 - assessment of the representativeness and the spatial extension of the reported points  
566 or profiles, in the same way as it is usually performed for agricultural or forest soils.

567

## 568 **Acknowledgements**

569 We thank Sophie Le Perchec, INRAE, for providing us with the initial database on  
570 artificialized soils, Maylis Desrousseaux and Bertrand Schmitt (INRAE), coordinators of  
571 the project, and Yves Le Bissonnais ( INRAE, LISAH) and Anne Ruas (Univ. Eiffel, COSYS),  
572 the scientific co-pilots together with Béatrice Béchet. This work is also part of the MUSE  
573 and SUPRA projects (funding by ADEME).

574

## 575 **References**

576 Abel, S., Nehls, T., Mekiffer, B., Mathes, M., Thieme, J., Wessolek, G., 2015a. Pools of sulfur  
577 in urban rubble soils. *Journal of Soils and Sediments*, 15 (3): 532-540.  
578 <http://dx.doi.org/10.1007/s11368-014-1014-1>

579 Abel, S., Nehls, T., Mekiffer, B., Wessolek, G., 2015b. Heavy metals and benzo a pyrene in

580 soils from construction and demolition rubble. *Journal of Soils and Sediments*, 15 (8):  
581 1771-1780. <http://dx.doi.org/10.1007/s11368-014-0959-4>

582 Adeli, A., McLaughlin, M.R., Brooks, J.P., Read, J.J., Willers, J.L., Lang, D.J., McGrew, R.,  
583 2013. Age Chronosequence Effects on Restoration Quality of Reclaimed Coal Mine Soils  
584 in Mississippi Agroecosystems. *Soil Science*, 178 (7): 335-343.  
585 <http://dx.doi.org/10.1097/SS.0b013e3182a79e37>

586 Ahirwal, J., Maiti, S.K., 2016. Assessment of soil properties of different land uses  
587 generated due to surface coal mining activities in tropical Sal (*Shorea robusta*) forest,  
588 India. *Catena*, 140: 155-163. <http://dx.doi.org/10.1016/j.catena.2016.01.028>

589 Bae, J., Ryu, Y., 2015. Land use and land cover changes explain spatial and temporal  
590 variations of the soil organic carbon stocks in a constructed urban park. *Landscape and*  
591 *Urban Planning*, 136: 57-67. <http://dx.doi.org/10.1016/j.landurbplan.2014.11.015>

592 Baize, D., 2000. Teneurs totales en métaux lourds dans les sols français, résultats  
593 généraux du programme ASPITET. *Courrier de l'environnement de l'INRA*, 39: 39-54.

594 Balesdent J., Basile-Doelsch I., Chadoeuf J., Cornu S., Derrien D., Fekiacova Z. & Hatté C.,  
595 2018. Atmosphere–soil carbon transfer as a function of soil depth. *Nature*, 559, 599–602.

596 Barles S., Breysse D., Guillerme A., Leyval C., coord. (1999). *Le sol urbain*. Collection  
597 *Villes, Anthropos*, 278p.

598 Baumgartl, T., 1998. Physical soil properties in specific fields of application especially in  
599 anthropogenic soils. *Soil & Tillage Research*, 47 (1-2): 51-59.  
600 [http://dx.doi.org/10.1016/s0167-1987\(98\)00072-5](http://dx.doi.org/10.1016/s0167-1987(98)00072-5)

601 Béchet B. (coord.), Le Bissonnais Y. (coord.), Ruas A. (coord.), Aguilera A., André M.,  
602 Andrieu H., Ay J.-S., Baumont C., Barbe E., Beaudet-Vidal L., Belton-Chevallier L., Berthier

603 E., Billet Ph., Bonin O., Cavailhès J., Chancibault K., Cohen M., Coisson T., Colas R., Cornu  
604 S., Cortet J., Dablanc L., Darly S., Delolme C., Fack G., Fromin N., Gadal S., Gauvreau B.,  
605 Géniaux G., Gilli F., Guelton S., Guérois M., Hedde M., Houet T., Humbertclaude S. (expert  
606 technique), Jolivet L., Keller C., Le Berre I., Madec P. (expert technique), Mallet C., Marty  
607 P., Mering C., Musy M., Oueslati W., Paty S., Polèse M., Pumain D., Puissant A., Riou S.,  
608 Rodriguez F., Ruban V., Salanié J., Schwartz C., Sotura A., Thébert M., Thévenin T., Thisse  
609 J., Vergnès A., Weber C., Werey C., Desrousseaux M, 2017. Sols artificialisés et processus  
610 d'artificialisation des sols, Déterminants, impacts et leviers d'action. INRA (France),  
611 609p.

612 Beyer, L.; Cordsen, E.; Blume, H.P.; Schleuss, U.; Vogt, B.; Wu, Q., 1996. Soil organic matter  
613 composition in urbic anthrosols in the city of Kiel, NW-Germany, as revealed by wet  
614 chemistry and CPMAS C-13-NMR spectroscopy of whole soil samples. *Soil Technology*, 9  
615 (3): 121-132. [http://dx.doi.org/10.1016/s0933-3630\(96\)00007-4](http://dx.doi.org/10.1016/s0933-3630(96)00007-4)

616 Bone J., Head M., Barraclough D., Archer M., Scheib C., Flight D. et Voulvoulis N., 2010 -  
617 Soil quality assessment under emerging regulatory requirements. *Environment*  
618 *International*, 36 (6): 609–622.

619 Bünemann E.K., Bongiorno G., Bai Z., Creamer R.E., De Deyn G., de Goede R., Fleskens L.,  
620 Geissen V., Kuyper T.W., Mäder P., Pulleman M., Sukkel W., van Groenigen J.W.,  
621 Brussaard L., 2018. Soil quality – A critical review. *Soil Biology and Biochemistry*, 120:  
622 105-125. [doi.org/10.1016/j.soilbio.2018.01.030](https://doi.org/10.1016/j.soilbio.2018.01.030)

623 Burghardt, W.; von Bertrab, M., 2016. Dialeimmasol, urban soil of pavements. *Journal of*  
624 *Soils and Sediments*, 16 (11): 2500-2513. <http://dx.doi.org/10.1007/s11368-016-1526->  
625 *y*

626 Cakir, M., Makineci, E., Kumbasli, M., 2010. Comparative study on soil properties in a

627 picnic and undisturbed area of Belgrad forest, Istanbul. *Journal of Environmental*  
628 *Biology*, 31 (1-2): 125-128.

629 Cambou A., Shaw R.K., Huot H., Vidal-Beaudet L., Hunault G., Cannavo P., Nold F.,  
630 Schwartz C. 2018. Estimation of soil organic carbon stocks of two cities, New York City  
631 and Paris. *Sciences of the total environment*, 644 : 452-464.  
632 [doi.org/10.1016/j.scitotenv.2018.06.322](https://doi.org/10.1016/j.scitotenv.2018.06.322)

633 Capilla, X., Schwartz, C., Bedell, J.P., Sterckeman, T., Perrodin, Y., Morel, J.L., 2006.  
634 Physicochemical and biological characterisation of different dredged sediment deposit  
635 sites in France. *Environmental Pollution*, 143 (1): 106-116.  
636 <http://dx.doi.org/10.1016/j.envpol.2005.11.007>

637 Chichester, F.W., Hauser, V.L., 1991. Change in chemical-properties of constructed  
638 minesoils developing under forage grass management. *Soil Science Society of America*  
639 *Journal*, 55 (2): 451-459.  
640 <http://dx.doi.org/10.2136/sssaj1991.03615995005500020026x>

641 Chung, E., Lee, J.S., Chon, H.T., Sager, M., 2005. Environmental contamination and  
642 bioaccessibility of arsenic and metals around the Dongjeong Au-Ag-Cu mine, Korea.  
643 *Geochemistry-Exploration Environment Analysis*, 5: 69-74.  
644 <http://dx.doi.org/10.1144/1467-7873/03-060>

645 Clouard, M., Criquet, S., Borschneck, D., Ziarelli, F., Marzaioli, F., Balesdent, J., Keller, C.,  
646 2014. Impact of lignite on pedogenetic processes and microbial functions in  
647 Mediterranean soils. *Geoderma*, 232-234: 257-269.  
648 <http://dx.doi.org/10.1016/j.geoderma.2014.05.00>

649 CEC - Commission of the European Communities, 2006 - Communication from the  
650 Commission to the Council, the European Parliament, the European Economic and Social

651 Committee and the Committee of the Regions - Thematic Strategy for Soil Protection,  
652 COM(2006)231 final, 22.9.2006.

653 Cousin, I., Nicoullaud, B., Coutadeur, C., 2003. Influence of rock fragments on the water  
654 retention and water percolation in a calcareous soil. *Catena* 53(2), 97–114.  
655 doi:10.1016/S0341-8162(03)00037-7

656 Dai Y., Shangguan W., Wei N., Xin Q., Yuan H., Zhang S., Liu S., Lu X., Wang D., Yan F., 2019.  
657 A review of the global soil property maps for Earth system models. *Soil* 5: 137–158,  
658 2019. doi.org/10.5194/soil-5-137-2019 SOIL

659 Dominati, E., Patterson, M., Mackay, A., 2010. A framework for classifying and  
660 quantifying the natural capital and ecosystem services of soils. *Ecological Economics*, 69  
661 (9): 1858-1868. <http://dx.doi.org/10.1016/j.ecolecon.2010.05.002>

662 Edmondson, J.L., Davies, Z.G., Gaston, K.J., Leake, J.R., 2014. Urban cultivation in  
663 allotments maintains soil qualities adversely affected by conventional agriculture.  
664 *Journal of Applied Ecology*, 51 (4): 880-889. [http://dx.doi.org/10.1111/1365-](http://dx.doi.org/10.1111/1365-2664.12254)  
665 [2664.12254](http://dx.doi.org/10.1111/1365-2664.12254)

666 Edmondson, J.L., Davies, Z.G., McCormack, S.A., Gaston, K.J., Leake, J.R., 2011. Are soils in  
667 urban ecosystems compacted? A citywide analysis. *Biology Letters*, 7 (5): 771-774.  
668 <http://dx.doi.org/10.1098/rsbl.2011.0260>

669 El-Mufleh, A., Bechet, B., Basile-Doelsch, I., Geffroy-Rodier, C., Gaudin, A., Ruban, V.,  
670 2014a. Distribution of PAHs and trace metals in urban stormwater sediments:  
671 combination of density fractionation, mineralogy and microanalysis. *Environmental*  
672 *Science and Pollution Research*, 21 (16): 9764-9776.  
673 <http://dx.doi.org/10.1007/s11356-014-2850-7>

674 El-Mufleh, A., Bechet, B., Ruban, V., Legret, M., Clozel, B., Barraud, S., Gonzalez-Merchan,  
675 C., Bedell, J.P., Delolme, C., 2014b. Review on physical and chemical characterizations of  
676 contaminated sediments from urban stormwater infiltration basins within the  
677 framework of the French observatory for urban hydrology (SOERE URBIS).  
678 Environmental Science and Pollution Research, 21 (8): 5329-5346.  
679 <http://dx.doi.org/10.1007/s11356-013-2490-3>

680 European Commission - JRC - ESDAC: LUCAS 2015 Topsoil Data.  
681 <https://esdac.jrc.ec.europa.eu/>

682 FAO. [www.fao.org](http://www.fao.org). seen in April 2020.

683 FAO, 1993. FESLM: An international framework for evaluating sustainable land  
684 management. Land and water development division. Rome, Italy.

685 Flynn, H.C., Meharg, A.A., Bowyer, P.K., Paton, G.I., 2003. Antimony bioavailability in mine  
686 soils. Environmental Pollution, 124 (1): 93-100. [http://dx.doi.org/10.1016/s0269-](http://dx.doi.org/10.1016/s0269-7491(02)00411-6)  
687 [7491\(02\)00411-6](http://dx.doi.org/10.1016/s0269-7491(02)00411-6)

688 Fu, S.S., Li, P.J., Feng, Q.A., Li, X.J., Li, P., Sun, Y.B., Chen, Y., 2011. Soil Quality Degradation  
689 in a Magnesite Mining Area. Pedosphere, 21 (1): 98-106.  
690 [http://dx.doi.org/10.1016/s1002-0160\(10\)60084-7](http://dx.doi.org/10.1016/s1002-0160(10)60084-7)

691 Golubiewski, N.E., 2006. Urbanization increases grassland carbon pools: Effects of  
692 landscaping in Colorado's front range. Ecological Applications, 16 (2): 555-571.  
693 [http://dx.doi.org/10.1890/1051-0761\(2006\)016\[0555:uigcpe\]2.0.co;2](http://dx.doi.org/10.1890/1051-0761(2006)016[0555:uigcpe]2.0.co;2)

694 Gregory, J.H., Dukes, M.D., Jones, P.H., Miller, G.L., 2006. Effect of urban soil compaction  
695 on infiltration rate. Journal of Soil and Water Conservation, 61 (3): 117-124.  
696 <http://www.jsowonline.org/content/61/3/117.short>

697 Greinert, A., 2015. The heterogeneity of urban soils in the light of their properties.  
698 Journal of Soils and Sediments, 15 (8): 1725-1737. <http://dx.doi.org/10.1007/s11368->  
699 014-1054-6

700 Hanks, D., Lewandowski, A., 2003. Protecting urban soil quality: examples for landscape  
701 codes and specifications. Washington, DC: USDA-NRCS, 20 p.

702 Haering, K.C.; Daniels, W.L.; Galbraith, J.M., 2004. Appalachian mine soil morphology and  
703 properties: Effects of weathering and mining method. Soil Science Society of America  
704 Journal, 68 (4): 1315-1325. <http://dx.doi.org/10.2136/sssaj2004.1315>

705 Hiller, D.A., 2000. Properties of Urbic Anthrosols from an abandoned shunting yard in  
706 the Ruhr area, Germany. Catena, 39 (4): 245-266. <http://dx.doi.org/10.1016/s0341->  
707 8162(00)00081-3

708 Howard, J.L., Dubay, B.R., Daniels, W.L., 2013. Artifact weathering, anthropogenic  
709 microparticles and lead contamination in urban soils at former demolition sites, Detroit,  
710 Michigan. Environmental Pollution, 179: 1-12.  
711 <http://dx.doi.org/10.1016/j.envpol.2013.03.053>

712 Howard, J.L., Orlicki, K.M., 2015. Effects of Anthropogenic Particles on the Chemical and  
713 Geophysical Properties of Urban Soils, Detroit, Michigan. Soil Science, 180 (4-5): 154-  
714 166. <http://dx.doi.org/10.1097/ss.0000000000000122>

715 Howard, J.L., Orlicki, K.M., 2016. Composition, micromorphology and distribution of  
716 microartefacts in anthropogenic soils, Detroit, Michigan, USA. Catena, 138: 103-116.

717 Huot, H., Joyner, J., Córdoba, A., Shaw, R.K., Wilson, M.A., Walker, R., Muth, T.R., Cheng, Z.,  
718 2017. Characterizing urban soils in New York City: profile properties and bacterial  
719 communities. Journal of Soils and Sediments, 17 (2): 393-407.

720 <http://dx.doi.org/10.1007/s11368-016-1552-9>

721 IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 2014.  
722 International soil classification system for naming soils and creating legends for soil  
723 maps: update 2015. Roma: FAO, World Soil Resources Reports, (n°106), 192 p.  
724 [www.fao.org/3/a-i3794e.pdf](http://www.fao.org/3/a-i3794e.pdf)

725 Jim, C.Y., 1998a. Soil characteristics and management in an urban park in Hong Kong.  
726 *Environmental Management*, 22 (5): 683-695.  
727 <http://dx.doi.org/10.1007/s002679900139>

728 Jim, C.Y., 1998b. Urban soil characteristics and limitations for landscape planting in Hong  
729 Kong. *Landscape and Urban Planning*, 40 (4): 235-249.  
730 [http://dx.doi.org/10.1016/S0169-2046\(97\)00117-5](http://dx.doi.org/10.1016/S0169-2046(97)00117-5)

731 Johnson, C.D., Skousen, J.G., 1995. Minesoil properties of 15 abandoned mine land sites in  
732 West-Virginia. *Journal of Environmental Quality*, 24 (4): 635-643.  
733 <http://dx.doi.org/10.2134/jeq1995.00472425002400040014x>

734 Joimel, S., Cortet, J., Jolivet, C.C., Saby, N.P., Chenot, E.D., Branchu, P., Consales, J.N., Lefort,  
735 C., Morel, J.L., Schwartz, C., 2016. Physico-chemical characteristics of topsoil for  
736 contrasted forest, agricultural, urban and industrial land uses in France. *Science of the*  
737 *Total Environment*, 545-546: 40-47. <http://dx.doi.org/10.1016/j.scitotenv.2015.12.035>

738 Karlen D.L., Mausbach M.J., Doran J.W., Cline R.G., Harris R.F. et Schuman G.E., 1997. Soil  
739 quality: A concept, definition, and framework for evaluation. *Soil Science Society of*  
740 *America Journal*, 61 (1): 4-10.

741 Karlen D.L., Ditzler C.A., Andrews S.S., 2003. Soil quality: Why and how? *Geoderma*, 114:  
742 145-156.



743 Kern, J.S., 1995. Geographic patterns of soil water-holding capacity in the contiguous  
744 USA. *Soil Science Society of America Journal*, 59:1126-1133.

745 Kida, K., Kawahigashi, M., 2015. Influence of asphalt pavement construction processes  
746 on urban soil formation in Tokyo. *Soil Science and Plant Nutrition*, 61: 135-146.  
747 <http://dx.doi.org/10.1080/00380768.2015.1048182>

748 Kolodziej, B., Bryk, M., Slowinska-Jurkiewicz, A., Otremba, K., Gilewska, M., 2016. Soil  
749 physical properties of agriculturally reclaimed area after lignite mine: A case study from  
750 central Poland. *Soil & Tillage Research*, 163: 54-63.  
751 <http://dx.doi.org/10.1016/j.still.2016.05.001>

752 Kundu, N.K., Ghose, M.K., 1994. Studies on the topsoil of an underground coal-mining  
753 project. *Environmental Conservation*, 21 (2): 126-132.  
754 <http://dx.doi.org/10.1017/S0376892900024553>

755 Kurucu, Y., Chiristina, N.K., 2008. Monitoring the impacts of urbanization and  
756 industrialization on the agricultural land and environment of the Torbali, Izmir region,  
757 Turkey. *Environmental Monitoring and Assessment*, 136 (1-3): 289-297.  
758 <http://dx.doi.org/10.1007/s10661-007-9684-4>

759 Lorenz, K., Lal, R., 2009. Biogeochemical C and N cycles in urban soils. *Environment*  
760 *International*, 35 (1): 1-8. <http://dx.doi.org/10.1016/j.envint.2008.05.006>

761 Lorenz, K., Lal, R., 2015. Managing soil carbon stocks to enhance the resilience of urban  
762 ecosystems. *Carbon Management*, 6 (1-2): 35-50.  
763 <http://dx.doi.org/10.1080/17583004.2015.1071182>

764 Lorenz, K., Preston, C.M., Kandeler, E., 2006. Soil organic matter in urban soils:  
765 Estimation of elemental carbon by thermal oxidation and characterization of organic

766 matter by solid-state C-13 nuclear magnetic resonance (NMR) spectroscopy. *Geoderma*,  
767 130 (3-4): 312-323. <http://dx.doi.org/10.1016/j.geoderma.2005.02.004>

768 Lottermoser, B.G., Ashley, P.M., Lawie, D.C., 1999. Environmental geochemistry of the  
769 Gulf Creek copper mine area, north-eastern New South Wales, Australia. *Environmental*  
770 *Geology*, 39 (1): 61-74. <http://dx.doi.org/10.1007/s002540050437>

771 Minasny, B., McBratney, A., Hartemink, A., 2010. Global pedodiversity, taxonomic  
772 distance, and the World Reference Base. *Geoderma* 155: 132-139.  
773 <https://doi.org/10.1016/j.geoderma.2009.04.024>

774 Morel, J.L., Chenu, C., Lorenz, K., 2015. Ecosystem services provided by soils of urban,  
775 industrial, traffic, mining, and military areas (SUITMAs). *Journal of Soils and Sediments*,  
776 15 (8): 1659-1666. <http://dx.doi.org/10.1007/s11368-014-0926-0>

777 Morel, J.L., Schwartz, C., Florentin, L., de Kimpe, C., 2005. Urban soils. In D. Hillel (ed.)  
778 *Encyclopedia of Soils in the Environment*, Elsevier Ltd., Academic Press, London, 202-  
779 208

780 Murolo, M., Pugliano, M.L., Ermice, A., 2005. Landfill and natural soils on the Somma-  
781 Vesuvius volcanic complex, Italy: Differences and similarities in soil morphology and  
782 properties. *Soil Science*, 170 (8): 652-668.  
783 <http://dx.doi.org/10.1097/01.ss.0000178202.51170.2a>

784 Nehls, T., Rokia, S., Mekiffer, B., Schwartz, C., Wessolek, G., 2013. Contribution of bricks to  
785 urban soil properties. *Journal of Soils and Sediments*, 13 (3): 575 - 584

786 Nikiforova, E.M., Kasimov, N.S., Kosheleva, N.E., 2014. Long-term dynamics of the  
787 anthropogenic salinization of soils in Moscow (by the example of the Eastern district).  
788 *Eurasian Soil Science*, 47 (3): 203-215. <http://dx.doi.org/10.1134/s1064229314030041>

789 Norra, S.; Fjer, N.; Li, F.; Chu, X.; Xie, X.; Stuben, D., 2008. The influence of different land  
790 uses on mineralogical and chemical composition and horizonation of urban soil profiles  
791 in Qingdao, China. *Journal of Soils and Sediments*, 8 (1): 4-16.  
792 <http://dx.doi.org/10.1065/jss2007.08.250>

793 Ogala, J., Siavalas, G., Christanis, K., 2012. Coal petrography, mineralogy and  
794 geochemistry of lignite samples from the Ogwashi–Asaba Formation, Nigeria. *Journal of*  
795 *African Earth Sciences*, 66-67 (Supplement C): 35-45.  
796 <http://dx.doi.org/10.1016/j.jafrearsci.2012.03.003>

797 Oliveira, V.; Makeschin, F.; Sano, E.; Lorz, C., 2014. Physical and chemical analyses of bare  
798 soil sites in Western Central Brazil: a case study. *Environmental Earth Sciences*, 72 (12):  
799 4863-4871. <http://dx.doi.org/10.1007/s12665-014-3103-2>

800 Pedrol, N., Puig, C.G., Souza, P., Forján, R., Vega, F.A., Asensio, V., González, L., Cerqueira,  
801 B., Covelo, E.F., Andrade, L., 2010. Soil fertility and spontaneous revegetation in lignite  
802 spoil banks under different amendments. *Soil & Tillage Research*, 110 (1): 134-142.  
803 <http://dx.doi.org/10.1016/j.still.2010.07.005>

804 Pellegrini, S., Garcia, G., Penas-Castejon, J.M., Vignozzi, N., Costantini, E.A.C., 2016.  
805 Pedogenesis in mine tails affects macroporosity, hydrological properties, and pollutant  
806 flow. *Catena*, 136: 3-16. <http://dx.doi.org/10.1016/j.catena.2015.07.027>

807 Peng, C., Ouyang, Z.Y., Wang, M.E., Chen, W.P., Li, X.M., Crittenden, J.C., 2013. Assessing  
808 the combined risks of PAHs and metals in urban soils by urbanization indicators.  
809 *Environmental Pollution*, 178: 426-432.  
810 <http://dx.doi.org/10.1016/j.envpol.2013.03.058>

811 Pouyat, R., Groffman, P., Yesilonis, I., Hernandez, L., 2002. Soil carbon pools and fluxes in  
812 urban ecosystems. *Environmental Pollution*, 116: S107-S118.

813 [http://dx.doi.org/10.1016/s0269-7491\(01\)00263-9](http://dx.doi.org/10.1016/s0269-7491(01)00263-9)

814 Pouyat, R.V., Yesilonis, I.D., Nowak, D.J., 2006. Carbon storage by urban soils in the  
815 United States. *Journal of Environmental Quality*, 35 (4): 1566-1575.  
816 <http://dx.doi.org/10.2134/jeq2005.0215>

817 Pouyat, R.V., Yesilonis, I.D., Russell-Anelli, J., Neerchal, N.K., 2007. Soil Chemical and  
818 Physical Properties That Differentiate Urban Land-Use and Cover Types. *Soil Science  
819 Society of America Journal*, 71 (3): 10 p. <http://dx.doi.org/10.2136/sssaj2006.0>

820 Raciti, S.M., Hutyra, L.R., Finzi, A.C., 2012. Depleted soil carbon and nitrogen pools  
821 beneath impervious surfaces. *Environmental Pollution*, 164: 248-251.  
822 <http://dx.doi.org/10.1016/j.envpol.2012.01.046>

823 Rivas-Pérez, I.M., Fernández-Sanjurjo, M.J., Núñez-Delgado, A., Monterroso, C., Macías, F.,  
824 Álvarez-Rodríguez, E., 2016. Evolution of Chemical Characteristics of Technosols in an  
825 Afforested Coal Mine Dump over a 20-year Period. *Land Degradation & Development*, 27  
826 (6): 1640-1649. <http://dx.doi.org/10.1002/ldr.2472>

827 Robinson, D.A., Lebron, I., Vereecken, H., 2009. On the Definition of the Natural Capital of  
828 Soils: A Framework for Description, Evaluation, and Monitoring. *Soil Science Society of  
829 America Journal*, 73 (6): 1904-1911. <http://dx.doi.org/10.2136/sssaj2008.0332>

830 Rossiter, D.G., 2007. Classification of urban and industrial soils in the world reference  
831 base for soil resources. *J Soils Sediments*, 7(2):96-100

832 Rytter, R.M., 2012. Stone and gravel contents of arable soils influence estimates of C and  
833 N stocks. *Catena*, 95: 153-159.

834 Schaaf, W., Hüttl, R.F., 2005. Soil chemistry and tree nutrition of post-lignite-mining sites.  
835 *Journal of Plant Nutrition and Soil Science*, 168 (4): 483-488.

836 <http://dx.doi.org/10.1002/jpln.200421692>

837 Scharenbroch B.C., Catania M., 2012. Soil quality attributes as indicators of urban tree  
838 performance. *Arboriculture & Urban Forestry*, 38 (5): 214-228.

839 Scharenbroch, B.C., Lloyd, J.E., Johnson-Maynard, J.L., 2005. Distinguishing urban soils  
840 with physical, chemical, and biological properties. *Pedobiologia*, 49 (4): 283-296.  
841 <http://dx.doi.org/10.1016/j.pedobi.2004.12.002>

842 Schleuß, U., Wu, Q.L., Blume, H.P., 1998. Variability of soils in urban and periurban areas  
843 in Northern Germany. *Catena*, 33 (3-4): 255-270. <http://dx.doi.org/10.1016/s0341->  
844 [8162\(98\)00070-8](http://dx.doi.org/10.1016/s0341-8162(98)00070-8)

845 Schoeneberger, P.J., Wysocki, D.A., Benham, E.C., Broderson, W.D., 2002. Field book for  
846 describing and sampling soils, v. 2.0. Natural Resources Conservation Service, National  
847 Soil Survey Center, U.S. Department of Agriculture, Lincoln, NE.

848 Shankar, U., Boral, L., Pandey, H.N., Tripathi, R.S., 1993. Degradation of land due to coal-  
849 mining and its natural recovery pattern. *Current Science*, 65 (9): 680-687.

850 Tao, Y., Li, F., Liu, X.S., Zhao, D., Sun, X., Xu, L.F., 2015. Variation in ecosystem services  
851 across an urbanization gradient: A study of terrestrial carbon stocks from Changzhou,  
852 China. *Ecological Modelling*, 318: 210-216.  
853 <http://dx.doi.org/10.1016/j.ecolmodel.2015.04.027>

854 Vasenev, V.I., Ananyeva, N.D., Makarov, O.A., 2012. Specific features of the ecological  
855 functioning of urban soils in Moscow and Moscow oblast. *Eurasian Soil Science*, 45 (2):  
856 194-205. <http://dx.doi.org/10.1134/s1064229312020147>

857 Vasenev, V.I., Stoorvogel, J.J., Vasenev, II, Valentini, R., 2014. How to map soil organic  
858 carbon stocks in highly urbanized regions? *Geoderma*, 226: 103-115.

859 <http://dx.doi.org/10.1016/j.geoderma.2014.03.007>

860 Vrščaj B., Poggio L., Ajmone-Marsan F., 2008. A method for soil environmental quality  
861 evaluation for management and planning in urban areas. *Landscape and Urban Planning*,  
862 88 (2–4): 81–94. <http://dx.doi.org/10.1016/j.landurbplan.2008.08.005>

863 Wahsha, M., Nadimi-Goki, M., Fornasier, F., Al-Jawasreh, R., Hussein, E.I., Bini, C., 2016.  
864 Microbial enzymes as an early warning management tool for monitoring mining site  
865 soils. *Catena*, 148: 40-45. <http://dx.doi.org/10.1016/j.catena.2016.02.021>

866 Wang, X.S.; Qin, Y.; Chen, Y.K., 2007. Leaching characteristics of arsenic and heavy metals  
867 in urban roadside soils using a simple bioavailability extraction test. *Environmental*  
868 *Monitoring and Assessment*, 129 (1-3): 221-226. [http://dx.doi.org/10.1007/s10661-](http://dx.doi.org/10.1007/s10661-006-9355-x)  
869 [006-9355-x](http://dx.doi.org/10.1007/s10661-006-9355-x)

870 Wang, P., Zheng, H., Ren, Z., Zhang, D., Zhai, C., Mao, Z., Tang, Z., He, X., 2018. Effects of  
871 Urbanization, Soil Property and Vegetation Configuration on Soil Infiltration of Urban  
872 Forest in Changchun, Northeast China. *CHINESE Geogr. Sci.* 28(3), 482–494.  
873 [doi:10.1007/s11769-018-0953-7](https://doi.org/10.1007/s11769-018-0953-7)

874 Wei, Z.Q., Wu, S.H., Zhou, S.L., Li, J.T., Zhao, Q.G., 2014. Soil Organic Carbon  
875 Transformation and Related Properties in Urban Soil Under Impervious Surfaces.  
876 *Pedosphere*, 24 (1): 56-64. [http://dx.doi.org/10.1016/S1002-0160\(13\)60080-6](http://dx.doi.org/10.1016/S1002-0160(13)60080-6)

877 Yang, J.L., Zhang, G.L., 2011. Water infiltration in urban soils and its effects on the  
878 quantity and quality of runoff. *J. Soils Sediments* 11(5), 751–761.

879 Yilmaz, D., Cannavo, P., Séré, G., Vidal-Beaudet, L., Legret, M., Damas, O., Peyneau, P.E.,  
880 2018. Physical properties of structural soils containing waste materials to achieve urban  
881 greening. *J. Soils Sediments* 18 (2), 442–455. <https://doi.org/10.1007/s11368-016->

882 1524-0.

883

884 **List of figure captions**

885 Figure 1: Evolution over time of the number of studies included in the database for the  
886 different SUTIMAs. Papers including several SUTIMAs are counted once per SUTIMA,  
887 which explains the discrepancy between the number of papers included in the database  
888 (n=102) and the number of studies recorded for the different SUTIMAs (n=138).

889

890 Figure 2: Origin of the studies of the database per SUTIMA. Papers including several  
891 SUTIMAs are counted once per SUTIMA, which explains the discrepancy between the  
892 number of papers included in the database (n=102) and the number of studies recorded  
893 for the different SUTIMAs (n=138).

894

895 Figure 3: Land-cover of the reference soils in the database (n=83)

896

897 Figure 4: Distribution of the soil classification of the soils of artificialized zones  
898 contained in the database (n=561)

899

900 Figure 5: Examples of artificialized soil profiles: a) Soil from a mine tailing: earthy  
901 material on spoil (TC, M); b) succession of gravel and earthy layers on concrete (TC, T);  
902 earth material on top of a natural profile (AT, U/R); c) Superposition of earthy material  
903 on a technogenic material (TC, U/R); d) multiple layers of alternate earthy and  
904 technogenic material (TC, U); e) succession of backfilling material, earthy material and  
905 granular material embedding a pipe (TC, U/T); f) Soil from a mine tailing: earthy  
906 material on spoil (TC, M); g) earthy material on red mud (TC, I); h) Joint soil and  
907 partially sealed soil (TC, U).



908

909 Figure 6: Two first factors of the CA performed on the different soil types of the different  
910 SAZs and reference soils for the different parameters: a- coarse elements (CE); b- < 2  $\mu$ m  
911 fraction; c- bulk density; d- soil organic carbon; and e- water pH for the topsoil (0-0.1, 0-  
912 0.2 or 0-0.3 m depending on the study). Ellipses represent the AHC performed on the  
913 two first factors of the CA.

914

915 Figure 7: Two first factors of the CA performed on the different soil types of the different  
916 SAZs and reference soils for the different parameters: a- coarse elements (CE); b- < 2  $\mu$ m  
917 fraction; c- bulk density; d- soil organic carbon; and e- water pH for a deep horizon  
918 selected around or under 0.4 m depth. Ellipses represent the AHC performed on the two  
919 first factors of the CA.

920

### 921 **List of table captions**

922 Table 1 : Number of topsoils (0-30 cm) in the data base for each considered soil  
923 property, each soil type and each land use (Tot, stands for total; Urb, for urban; Recre,  
924 for recreational; Indust, for industrial; Mine, for mining; Nat., for natural vegetation and  
925 Unsp., for unspecified vegetation).

926

927 Table 2 : Number of subsoils (> 0.4 m) in the data base for each considered soil property,  
928 each soil type and each land use (Tot, stands for total; Urb, for urban; Recre, for  
929 recreational; Indust, for industrial; Mine, for mining; Nat., for natural vegetation and  
930 Unsp., for unspecified vegetation).

931




Figure 1: Evolution over time of the number of studies included in the database for the different SUTMAs. Papers including several SUTMAs are counted once per SUTMA, which explains the discrepancy between the number of papers included in the database (n=102) and the number of studies recorded for the different SUTMAs (n=138).




Figure 2: Origin of the studies of the database per SUTMA. Papers including several SUTMAs are counted once per SUTMA, which explains the discrepancy between the number of papers included in the database (n=102) and the number of studies recorded for the different SUTMAs (n=138).




Figure 3: Land-cover of the reference soils in the database (n=83)




Figure 4: Distribution of the soil classification of the soils of artificialized zones contained in the database (n=561)




Figure 5: Examples of artificialized soil profiles: a) Soil from a mine tailing: earthy material on spoil (TC, M); b) succession of gravel and earthy layers on concrete (TC, T); earth material on top of a natural profile (AT, U/R); c) Superposition of earthy material on a technogenic material (TC, U/R); d) multiple layers of alternate earthy and technogenic material (TC, U); e) succession of backfilling material, earthy material and granular material embedding a pipe (TC, U/T); f) Soil from a mine tailing: earthy material on spoil (TC, M); g) earthy material on red mud (TC, I); h) Joint soil and partially sealed soil (TC, U).

Figure 6: Two first factors of the CA performed on the different soil types of the different SAZs and reference soils for the different parameters: a- coarse elements (CE); b- < 2 μm fraction; c- bulk density; d- soil organic carbon; and e- water pH for the topsoil (0-0.1, 0-0.2 or 0-0.3 m depending on the study). Ellipses represent the AHC performed on the two first factors of the CA.




Figure 7: Two first factors of the CA performed on the different soil types of the different SAZs and reference soils for the different parameters: a- coarse elements (CE); b- < 2 μm fraction; c- bulk density; d- soil organic carbon; and e- water pH for a deep horizon selected around or under 0.4 m depth. Ellipses represent the AHC performed on the two first factors of the CA.






Table 1 : Number of topsoils (0-30 cm) in the data base for each considered soil property, each soil type and each land use (Tot, stands for total; Urb, for urban; Recre, for recreational; Indust, for industrial; Mine, for mining; Nat., for natural vegetation and Unsp., for unspecified vegetation).

|                                 | SAZ        |            |            |           |           |            | References |      |      |       |
|---------------------------------|------------|------------|------------|-----------|-----------|------------|------------|------|------|-------|
|                                 | Tot        | Urb        | Recre      | Indust    | Traffic   | Mine       | Tot        | Nat. | Crop | Unsp. |
| <b>pH</b>                       |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>5</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>69</b>  | 14         | 16         | 10        | 0         | 29         | <b>64</b>  | 20   | 11   | 33    |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>184</b> | 52         | 18         | 6         | 13        | 95         |            |      |      |       |
| Unspecified profile             | <b>158</b> | 47         | 35         | 36        | 15        | 25         |            |      |      |       |
| <b>Total</b>                    | <b>416</b> | <b>113</b> | <b>69</b>  | <b>52</b> | <b>28</b> | <b>149</b> |            |      |      |       |
| <b>SOC</b>                      |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>1</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>55</b>  | 12         | 21         | 2         | 0         | 20         | <b>63</b>  | 27   | 13   | 23    |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>183</b> | 46         | 26         | 22        | 22        | 67         |            |      |      |       |
| Unspecified profile             | <b>171</b> | 56         | 53         | 30        | 18        | 14         |            |      |      |       |
| <b>Total</b>                    | <b>410</b> | <b>114</b> | <b>100</b> | <b>54</b> | <b>40</b> | <b>101</b> |            |      |      |       |
| <b>Bulk density</b>             |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>2</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>28</b>  | 5          | 8          | 0         | 0         | 15         | <b>31</b>  | 7    | 10   | 14    |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>113</b> | 20         | 6          | 2         | 15        | 70         |            |      |      |       |
| Unspecified profile             | <b>72</b>  | 41         | 24         | 2         | 5         | 0          |            |      |      |       |
| <b>Total</b>                    | <b>215</b> | <b>66</b>  | <b>38</b>  | <b>4</b>  | <b>20</b> | <b>85</b>  |            |      |      |       |
| <b>&lt; 2 µm</b>                |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>2</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>29</b>  | 13         | 6          | 0         | 0         | 10         | <b>39</b>  | 10   | 11   | 18    |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>123</b> | 31         | 26         | 0         | 3         | 63         |            |      |      |       |
| Unspecified profile             | <b>66</b>  | 25         | 18         | 4         | 14        | 5          |            |      |      |       |
| <b>Total</b>                    | <b>220</b> | <b>69</b>  | <b>50</b>  | <b>4</b>  | <b>17</b> | <b>78</b>  |            |      |      |       |
| <b>Coarse elements</b>          |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>1</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>2</b>   | 2          | 0          | 0         | 0         | 0          | <b>5</b>   | 2    | 3    | 0     |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>49</b>  | 8          | 9          | 0         | 15        | 17         |            |      |      |       |
| Unspecified profile             | <b>2</b>   | 1          | 1          | 0         | 0         | 0          |            |      |      |       |
| <b>Total</b>                    | <b>54</b>  | <b>11</b>  | <b>10</b>  | <b>0</b>  | <b>15</b> | <b>17</b>  |            |      |      |       |
| <b>Cation Exchange Capacity</b> |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>0</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>9</b>   | 0          | 4          | 0         | 0         | 5          | <b>8</b>   | 2    | 3    | 3     |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>66</b>  | 10         | 9          | 0         | 10        | 37         |            |      |      |       |
| Unspecified profile             | <b>40</b>  | 28         | 2          | 0         | 0         | 10         |            |      |      |       |
| <b>Total</b>                    | <b>115</b> | <b>38</b>  | <b>15</b>  | <b>0</b>  | <b>10</b> | <b>52</b>  |            |      |      |       |
| <b>Electrical conductivity</b>  |            |            |            |           |           |            |            |      |      |       |
| sealed                          | <b>1</b>   |            |            |           |           |            |            |      |      |       |
| Natural profile                 | <b>16</b>  | 6          | 9          | 0         | 0         | 1          | <b>13</b>  | 2    | 1    | 10    |
| Profile on                      |            |            |            |           |           |            |            |      |      |       |
| allochthonous materials         | <b>53</b>  | 14         | 1          | 0         | 0         | 38         |            |      |      |       |
| Unspecified profile             | <b>43</b>  | 10         | 3          | 19        | 1         | 10         |            |      |      |       |
| <b>Total</b>                    | <b>113</b> | <b>30</b>  | <b>13</b>  | <b>19</b> | <b>1</b>  | <b>49</b>  |            |      |      |       |

Table 2 : Number of subsoils (> 0.4 m) in the data base for each considered soil property, each soil type and each land use (Tot, stands for total; Urb, for urban; Recre, for recreational; Indust, for industrial; Mine, for mining; Nat., for natural vegetation and Unsp., for unspecified vegetation).

|                                 | <b>Tot</b> | <i>Urb</i> | <i>Recre</i> | <b>SAZ</b>    |                |             | <b>Tot</b> | <b>References</b> |             |              |
|---------------------------------|------------|------------|--------------|---------------|----------------|-------------|------------|-------------------|-------------|--------------|
|                                 |            |            |              | <i>Indust</i> | <i>Traffic</i> | <i>Mine</i> |            | <i>Nat.</i>       | <i>Crop</i> | <i>Unsp.</i> |
| <b>pH</b>                       |            |            |              |               |                |             |            |                   |             |              |
| sealed                          |            |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>18</b>  | 2          | 4            | 8             | 0              | 4           | <b>37</b>  | 10                | 9           | 18           |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>75</b>  | 11         | 9            | 1             | 2              | 52          |            |                   |             |              |
| Unspecified profile             | <b>22</b>  | 7          | 3            | 1             | 4              | 7           |            |                   |             |              |
| <b>Total</b>                    | <b>115</b> | <b>20</b>  | <b>16</b>    | <b>10</b>     | <b>6</b>       | <b>63</b>   |            |                   |             |              |
| <b>SOC</b>                      |            |            |              |               |                |             |            |                   |             |              |
| sealed                          |            |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>14</b>  | 1          | 4            | 0             | 0              | 9           | <b>41</b>  | 15                | 10          | 16           |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>72</b>  | 15         | 10           | 11            | 2              | 34          |            |                   |             |              |
| Unspecified profile             | <b>14</b>  | 5          | 6            | 0             | 3              | 0           |            |                   |             |              |
| <b>Total</b>                    | <b>100</b> | <b>21</b>  | <b>20</b>    | <b>11</b>     | <b>5</b>       | <b>43</b>   |            |                   |             |              |
| <b>Bulk density</b>             |            |            |              |               |                |             |            |                   |             |              |
| sealed                          | <b>0</b>   |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>3</b>   | 2          | 0            | 0             | 0              | 1           | <b>19</b>  | 3                 | 3           | 13           |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>44</b>  | 11         | 6            | 3             | 5              | 19          |            |                   |             |              |
| Unspecified profile             | <b>12</b>  | 2          | 8            | 0             | 2              | 0           |            |                   |             |              |
| <b>Total</b>                    | <b>59</b>  | <b>15</b>  | <b>14</b>    | <b>3</b>      | <b>7</b>       | <b>20</b>   |            |                   |             |              |
| <b>&lt; 2 µm</b>                |            |            |              |               |                |             |            |                   |             |              |
| sealed                          | <b>0</b>   |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>6</b>   | 1          | 0            | 0             | 0              | 5           | <b>24</b>  | 4                 | 8           | 12           |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>67</b>  | 18         | 9            | 0             | 0              | 40          |            |                   |             |              |
| Unspecified profile             | <b>12</b>  | 1          | 3            | 1             | 2              | 5           |            |                   |             |              |
| <b>Total</b>                    | <b>85</b>  | <b>20</b>  | <b>12</b>    | <b>1</b>      | <b>2</b>       | <b>50</b>   |            |                   |             |              |
| <b>Coarse elements</b>          |            |            |              |               |                |             |            |                   |             |              |
| sealed                          | <b>0</b>   |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>0</b>   | 0          | 0            | 0             | 0              | 0           | <b>5</b>   | 2                 | 3           | 0            |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>37</b>  | 8          | 9            | 0             | 5              | 15          |            |                   |             |              |
| Unspecified profile             | <b>0</b>   | 0          | 0            | 0             | 0              | 0           |            |                   |             |              |
| <b>Total</b>                    | <b>37</b>  | <b>8</b>   | <b>9</b>     | <b>0</b>      | <b>5</b>       | <b>15</b>   |            |                   |             |              |
| <b>Cation Exchange Capacity</b> |            |            |              |               |                |             |            |                   |             |              |
| sealed                          | <b>0</b>   |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>7</b>   | 0          | 4            | 0             | 0              | 3           | <b>7</b>   | 2                 | 3           | 2            |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>41</b>  | 10         | 9            | 0             | 0              | 22          |            |                   |             |              |
| Unspecified profile             | <b>5</b>   | 0          | 1            | 0             | 0              | 4           |            |                   |             |              |
| <b>Total</b>                    | <b>53</b>  | <b>10</b>  | <b>14</b>    | <b>0</b>      | <b>0</b>       | <b>29</b>   |            |                   |             |              |
| <b>Electrical conductivity</b>  |            |            |              |               |                |             |            |                   |             |              |
| sealed                          | <b>0</b>   |            |              |               |                |             |            |                   |             |              |
| Natural profile                 | <b>1</b>   | 0          | 0            | 0             | 0              | 1           | <b>4</b>   | 1                 | 1           | 2            |
| Profile on                      |            |            |              |               |                |             |            |                   |             |              |
| allochthonous materials         | <b>36</b>  | 3          | 1            | 0             | 0              | 32          |            |                   |             |              |
| Unspecified profile             | <b>14</b>  | 1          | 0            | 9             | 0              | 4           |            |                   |             |              |
| <b>Total</b>                    | <b>51</b>  | <b>4</b>   | <b>1</b>     | <b>9</b>      | <b>0</b>       | <b>37</b>   |            |                   |             |              |