



HAL
open science

Management of an invasive plant in a mediterranean protected area: the experience of *Senecio deltoideus* in Italy

Luigi Minuto, Gabriele Casazza, Davide Dagnino, Maria Guerrina, Carmelo Macrì, Giorgio Mariotti Mauro

► To cite this version:

Luigi Minuto, Gabriele Casazza, Davide Dagnino, Maria Guerrina, Carmelo Macrì, et al.. Management of an invasive plant in a mediterranean protected area: the experience of *Senecio deltoideus* in Italy. *Annali di Botanica*, 2020, 11, pp.1-12. 10.13133/2239-3129/16852 . hal-03314556

HAL Id: hal-03314556

<https://amu.hal.science/hal-03314556v1>

Submitted on 5 Aug 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MANAGEMENT OF AN INVASIVE PLANT IN A MEDITERRANEAN PROTECTED AREA: THE EXPERIENCE OF *SENECIO DELTOIDEUS* IN ITALY

MINUTO L.^{1,3,*}, CASAZZA G.², DAGNINO D.¹, GUERRINA M.¹, MACRÌ C.¹, MARIOTTI M.G.^{1,3}

¹ *Dipartimento di Scienze della terra, Ambiente e Vita, Università di Genova, Corso Europa 26, 16132 Genova, Italy.*

² *Institut Méditerranéen de Biodiversité et d'Écologie marine et continentale (IMBE), Aix Marseille Université, Avignon Université, CNRS, IRD, Technopole de l'Arbois-Méditerranée, BP 80, 13545 Aix-en-Provence Cedex 4, France.*

³ *Giardini Botanici Hanbury, Università di Genova, Corso Montecarlo 43, I-18039 Ventimiglia, Italy.*

*Corresponding author; telephone: +390102099361; email: luigi.minuto@unige.it

(RECEIVED 30 JUNE 2020; RECEIVED IN REVISED FORM 27 OCTOBER 2020; ACCEPTED 2 NOVEMBER 2020)

ABSTRACT - Biological invasions are one of the most serious threats to global biodiversity and ecosystem integrity. The problem is growing year by year and a large number of protected areas worldwide are today invaded by at least one exotic species. In this study, we tested the eco-friendly and cost-effective weeding control of *Senecio deltoideus* in a Regional Protected Area in the North Mediterranean region. During a two years experiment, four techniques compatible with the local laws on protected areas (natural-herbicide, flame-weeding, mulching and mowing) were applied five times a season on sixty plots, compared with fifteen untreated controls. All techniques were effective in reducing *Senecio* covering: after the first year the maximum covering was limited to 37.93% (flame weeding) with a mean value of 10%; after second year the covering was further reduced (17.02% max; 2% mean). Interestingly, during the second year all plots submitted to a one-year treatment showed an enduring control of *S. deltoideus* covering (40.73% max; 20% mean). Taking into account feasibility and their impact on the environment, the weeding control recommendable for *S. deltoideus* is mowing. In a long-time management strategy, the selected treatment could be applied every two years with a drastic reduction in costs for the manager of the protected area.

KEYWORDS: ALIEN SPECIES; ECO-FRIENDLY AND COST; EFFECTIVE; MANAGEMENT; NATURAL ENVIRONMENT.

INTRODUCTION

Biological invasions are one of the most serious threats to global biodiversity and ecosystem integrity (Mack et al., 2000; Essl et al., 2020; Shackleton et al., 2020). Alien species may cause huge ecological (Hulme, 2003), economical (Pimentel et al., 2000; Oerke, 2006), social (García-Llorente et al., 2011) and sanitary impacts (McNeely, 2001). Invasive alien species are characterized by rapid growth rates, extensive dispersal capabilities, large and rapid reproductive output and broad environmental tolerance (Geesing et al., 2000). It is estimated that as many as 50% of invasive species in general can be classified as ecologically harmful, (Richardson et al., 2000) forming alien-dominated plant communities (Viciani et al., 2020). Despite invasive

species mainly occur in vegetational disturbed areas, their prevalence has gone nowadays beyond these areas and they invade natural protected areas across the world (DePoorter et al., 2007). Invasions into natural habitats can produce detrimental consequences in native communities (e.g. Bais et al., 2003; Miller & Gorchov, 2004; Wearne & Morgan, 2004) and represent one of the main threats in protected areas (Ervin, 2003), becoming a threat to endangered or threatened plant species around the world (Pimentel et al., 2005). The problem is growing year by year and today at least one exotic species has invaded a large number of the protected areas in the world (Shackleton et al., 2020). For example, in Spain, more than 90% of invasive species are found in at least

one protected area (Andreau et al., 2009). For this reason, invasive plant control is today a major challenge for natural resource management (Kettenring & Adams, 2011). Despite effort to understand the characteristics of invasive plants (e.g., Rejmánek & Richardson, 1996; Bossdorf et al., 2005; Ellstrand & Schierenbeck, 2006) and what makes some areas more susceptible to invasion than others (e.g. Stohlgren et al., 1999; Alpert et al., 2000), the control of invasive plants in the context of ecosystem conservation have had mixed success. In particular, few studies aimed to assess the eradication and control of invasive plant species in protected areas (Child et al., 1998; Shaw, 2003; Dehnen-Schumutz et al., 2004; Nielsen et al., 2005; Křivánek, 2006). Most of them applied mechanical methods in substitution to the chemical control, forbidden within protected areas of many countries (Villaverde et al., 2014). Mechanical control allows the exhaustion of reserves, provoking withering and depriving the plant of any capacity to regenerate (Chicouene, 2007). Mowing (Milakovic et al., 2014), uprooting (Chicouene, 2007), grazing (Erskine Ogden & Rejmánek, 2005; Bandarra et al., 2012;), and burning (Kettenring & Adams, 2011) are the main controls applied in protected areas today but new weeding techniques aiming at lower environmental impact and cost are now available. However, their efficiency and applicability with invasive plants still need to be tested in protected areas.

Senecio deltoideus Less. (Asteraceae - Canary creeper) is one out of the 791 non-native naturalized plant species detected in Italy (Galasso et al., 2018). It is an extremely invasive exotic plant introduced in the Mediterranean area since the beginning of the XIX century for ornamental purposes (Zappa & Campodonico, 2005). Today the species is a casual neophyte in SE France (Fried, 2010) and Spain, but it is an invasive plant in NW Italy (Galasso et al., 2018).

In this study, we tested the best eco-friendly and cost-effective weeding control of *S. deltoideus* in a Regional Protected Area near Ventimiglia (Italy - close to the French border). More specifically, in a Mediterranean climate area, the study aimed at: a) assessing the re-growth of *S. deltoideus* in a non-native area; b) verifying the effect of different weeding techniques for the control of *S. deltoideus* invasion; c) testing the enduring effect of each technique on *S. deltoideus* after one-year treatment.

MATERIALS AND METHODS

Study species

Senecio deltoideus Less. is a vigorous perennial climbing herb native from Tropical Africa, producing masses of bright

yellow flower heads in early winter. In native range, it is a species of open terrain, such as bracken and ericoid scrub or bush clumps in montane grassland, where it often forms masses on the ground and it climbs on lower scrub (Hyde et al., 2020). In non-native range, the species grows mainly in open vegetation, in Mediterranean grassland and shrubs, frequently becoming invasive. The plant blooms in December and later it shows a vigorous vegetative growth favoured by its many basal gems from which many shoots sprout, or resprout in summer when the plant is cut.

Study site

We performed the study at Capo Mortola [Ventimiglia (IM), North West Italy - 43.781576° N; 7.554797°E] inside the Regional Protected Area Giardini Botanici Hanbury. This territory is part of the Special Area of Conservation (IT13116118 “Capo Mortola”), protected by the Habitats European Directive 1992/43/EEC, aimed to conserve natural habitats and wild fauna and flora in Europe. The territory has a Mediterranean climate, with an arid period in July (annual average rainfall 663 mm, mean annual temperature: 12.9 °C). In 1867 part of the SAC (18 ha) was bought by British merchant Thomas Hanbury (Campodonico, 2010), who transformed the land in an acclimatization garden for exotic plants (Campodonico & Zappa, 2006). *Senecio deltoideus* entered the SAC in 1902 (Zappa & Campodonico, 2005). Currently there is an extension of about 10 hectares invaded by the African plant, rapidly covering fields and shrubby places with its covering and suffocating shoots (Galasso et al., 2018; personal observations).

Experimental design

The experiment was carried out in a homogenous plane (10 m a.s.l., exp. SE, incl. 2 %, 100 m distant from seaside) within an area of about one square hectare (43.781576° N; 7.554797°E). The selected area was never previously cultivated and located in a Aleppo-pine wood (*Pinus halepensis* Mill.) in which occurs some manna ash (*Fraxinus ornus* L.) and holm oak (*Quercus ilex* L.) individuals with a sparse underwood vegetation of Mediterranean scrubs (*Rhamnus alaternus* L., *Pistacia lentiscus* L., *Coronilla valentina* L.) and some grassland parcels. In those areas infested by *S. deltoideus* with a covering area of at least 90%, we randomly placed seventy-five 1 x 1 m plots. In January 2018 and 2019 (at the beginning of vegetative growth), we mowed all plots to have the same starting covering percentages and to measure the colonization capability of *S. deltoideus*. This preliminary operation corresponds to the usual management control of an alien invasive plant performed in a protected area management (Braun et al.,

2016; Weidlich et al., 2020). Fifteen plots were untreated and used as control (hereafter CTR) while the other sixty plots were treated five times during the first years with four eco-friendly and cost-effective weeding techniques (fifteen plots for each treatment). Then in the second year, we repeated six times the treatments on thirty-six plots (nine for each technique) with a timetable similar to the first year. To test the enduring effect of each technique, we did not treat any more the remnant twenty-four plots (1-year treatment), and we compared them with those treated for a second year (2-years treatment). In particular, we used the following techniques and specific application methods:

1. organic herbicide - performed with pelargonic acid (FINALSAN Professional, W. Neudorff GmbH KG, Germany) 20 L/hl per ha, applied manually with shoulder sprayer five time a year with a regular interval between each treatment (5 day/ha manpower per year). Pelargonic acid (hereafter PEL) is a contact herbicide killing the sprayed parts of the plant, which does not translocate the active substance (Muñoz et al., 2020). Thus, PEL has a local effect like fire and does not leave any polluting traces in the environment;
2. flame-weeding (hereafter FLW) - performed with a flame-weeder Pirotrolley15 Maito, fuelled by liquefied petroleum gas (LPG), 1 kg/ha, five time a year with a regular interval between each treatment (5 day/ha manpower per year). The operator made a subjective use (up to visual plant desiccation, about 3-5 sec/m²) in order to kill herbs but not affecting woody plants and not generating a fire. Plants are exposed to ultra-high temperatures producing cellular death due to the initial thermal disruption of the cellular membranes rapidly followed by dehydration of the affected tissue (Ellwanger et al., 1973);
3. mulching (hereafter MUL) - performed with chopped wood locally produced with the management of the protected area (pruning, chopped wood, etc.). A layer of 20 cm height of chopped wood was scattered on the ground after the preliminary mowing made in January (3 day/ha manpower per year);
4. mowing (hereafter MOW), performed with a handy machine trimmer, five time a year with a regular interval between each treatment (5 day/ha manpower per year).

Data collection and analysis

Before any new treatment (MUL included), we took a high definition photograph of each plot to record *S. deltoideus* re-growth. The photographs were made with a Canon EOS 350D equipped with an EFS 18-25 mm lens, fixed on a tripod at a distance of 120 cm from the ground. Specifically,

each year we took 450 photos in six sessions (a total of 900 photos in 12 sessions during the whole study): the first session was aimed to detect the starting covering, the others to quantify the re-growth. At the end of the study we uploaded: 90 photographs of CTR, PEL, FLW, MUL, MOW each belonging to 2018; 90 photos of CTR belonging to 2019; 216 of PEL, FLW, MUL, MOW after 2 years treatment belonging to 2019; 144 of PEL, FLW, MUL, MOW after 1-years treatment belonging to 2019.

The images were elaborated with ImageJ 1.52a (Wayne Rasband, National Institute of Health, USA. <http://imagej.nih.gov/ij>) producing a final matrix of the covering percentage of *S. deltoideus* only per plot per weeding technique. The recolonization of other plants was detected but not analysed in this study.

The repeated measures of covering proportion were analysed using a generalized linear mixed effect model by using lme4 package (Bates et al., 2015) implemented in R (R Core Team, 2020). To determine whether there was a significant effect of techniques, we set them as a fixed effect, and plot as random factor. To determine difference between CTR and 1-year treatment plots we set them as a fixed effect and plot as random factor. Then, to determine for each weeding technique any difference between 1-year treatment and 2-years treatment plots, we set them as a fixed effect and plot as random factor. The data were analysed with a negative binomial distribution because the response variable is proportion data. We performed post hoc analysis using the emmeans function in the “emmeans” package in R (Lenth, 2020).

RESULTS

The CTR plots had a high re-growth after the shared mowing performed at the beginning of each study year (Tables 1 and 2). *S. deltoideus* covering grows constantly up to the mean value of 96.4% (sd 4.77) and 88.68% (sd 9.39) of the total surface at the end of 2018 and 2019 respectively.

The re-growth was very fast in spring and early summer, a period favoured by a sensible amount of precipitations (Figure 1). The summer aridity, on the contrary, limited the vegetative growth of the plant, with a fast restart of the vegetation colonization during the autumn (Figure 1).

During the first year of experiment, the four weeding techniques reduced *S. deltoideus* covering (Tables 1) but they had significantly different effect (Table 3). In particular, FLW was significantly different from all other treatments, reaching the highest covering value of 37.93% (sd 23.98) in late autumn (Table 1). The other techniques had a better

Table 1. Percentage of surface and standard deviation (in brackets) covered by *S. deltoideus* during the first year in the four weeding techniques. The total number of plots performed (N) is reported. The columns 1-6 correspond to the treatment applied: 1 = winter mowing as pre-treatment; 2-6 the five subsequent treatment from early spring to late autumn.

Treatments	N	1	2	3	4	5	6
MOW	15	0.28 (0.28)	7.31 (3.21)	7.31 (3.56)	3.10 (2.87)	5.17 (6.72)	4.72 (4.88)
FLW	15	0.29 (0.15)	15.50 (12.16)	22.57 (15.54)	23.13 (19.70)	24.19 (14.83)	37.93 (23.98)
MUL	15	0.20 (0.30)	0.16 (0.25)	0.40 (0.83)	2.34 (4.48)	0.22 (8.38)	5.12 (11.15)
PEL	15	0.31 (0.33)	1.32 (0.98)	4.29 (3.15)	0.59 (0.86)	0.16 (0.19)	0.22 (0.30)
CTR	15	0.23 (0.15)	19.83 (12.45)	51.46 (25.59)	90.49 (9.54)	89.99 (10.73)	96.40 (4.77)

Table 2. Percentage of surface and standard deviation (in brackets) covered by *S. deltoideus* during the second year in the four weeding techniques. Values are reported for 2-years treated plots (y-t = 2; grey stipes) on which the technique was repeated and 1-year treated plots (y-t = 1; white stripes) on which the technique was not repeated. The total number (N) of plots used is reported. The columns 1-6 correspond to the treatment applied: 1 = winter mowing as pre-treatment; 2-6 the five subsequent treatment from early spring to late autumn.

Technique	y-t	N	1	2	3	4	5	6	7
MOW	1	6	1.18 (0.73)	8.34 (4.78)	20.07 (9.48)	4.08 (3.82)	2.59 (2.11)	11.97 (13.34)	40.73 (31.15)
	2	9	0.61 (0.29)	1.54 (1.39)	2.06 (2.45)	0.05 (0.07)	0.01 (0.03)	0.05 (0.11)	0.38 (0.83)
FLW	1	6	0.75 (0.98)	8.05 (5.57)	23.84 (18.20)	8.18 (7.66)	3.61 (2.88)	9.51 (12.06)	25.32 (16.68)
	2	9	0.57 (0.43)	17.02 (11.36)	6.64 (9.51)	0.94 (0.86)	0.21 (0.34)	0.91 (1.80)	1.44 (2.15)
MUL	1	6	1.34 (0.96)	5.92 (3.75)	12.56 (11.28)	6.13 (5.77)	3.67 (3.43)	11.01 (15.20)	16.34 (23.77)
	2	9	1.25 (1.03)	1.29 (2.52)	1.47 (1.64)	0.09 (0.23)	0.05 (0.16)	0.38 (0.51)	0.70 (1.98)
PEL	1	6	0.38 (0.27)	0.39 (0.31)	13.06 (10.19)	0.07 (0.07)	0.03 (0.04)	0.03 (0.03)	0.01 (0.02)
	2	9	0.86 (0.71)	0.23 (0.43)	0.18 (0.13)	0.02 (0.02)	0.00 (0.01)	0.10 (0.24)	0.00 (0.00)
CTR	0	15	0.18 (0.13)	47.19 (19.00)	83.03 (10.45)	84.41 (10.42)	71.88 (20.85)	54.48 (26.81)	88.68 (9.39)

performance reaching a maximum covering of 7.31% (sd 3.56), 5.12% (sd 11.15) and 4.29% (sd 3.15) in MOW, MUL and PEL respectively (Table 1 and Fig. 2a).

Differently, during the second year, the effect of the four techniques was similar and not significant differences were detected (Table 2 and 3). In 2-years treatment plots the maximum covering surface was reached in FLW again (17.02%, sd 11.36), while the other techniques showed a maximum covering of 2.06% (sd 2.45), 1.47% (sd 1.64)

and 0.23% (sd 0.43) in MOW, MUL and PEL respectively (Table 2 and Fig. 2b).

The 1-year treatment plots were significantly lower and different to CTRCTR plots in the proportion of *Senecio* covering (Table 4 and Fig. 3). The highest maximum covering surface was reached in MOW (40.73%, sd 31.15), while the other treatments showed maximum covering values of 25.32% (sd 16.68), 16.34% (sd 23.77) and 13.06% (sd 10.19) for FLW, MUL and PEL respectively (Table 2 and Fig. 3).




Figure 1. Natural re-growth of *S. deltoideus* in the CTR plots in 2018 (black continuous line) and 2019 (yellow dotted line), after the preliminary mowing operation corresponding to the usual management control of an alien invasive plant performed in a protected area management. Proportion indicates the mean percentage of covering area of *S. deltoideus* in the study plots used as control and never treated with any weeding techniques.




Figure 2. Comparison in covering surface (%) of *S. deltoideus* after one year and two years treatment with the four weeding techniques (MOW = mowing; FLW = flame weeding; MUL = mulching; PEL = pelargonium acid).

Table 3. Post hoc test for the difference over time of the proportion of *Senecio* covering detected in 2018 and 2019, according to the different weeding techniques (FLW, MOW, MUL and PEL) and the control (CTR). Fix terms is the interaction treatment with days after first treatment (treatment x days).

Year	Contrast	Estimate	SE	z ratio	p value
2018	CTR vs FLW	1.0362	0.2486	4.1684	0.0003
	CTR vs MOW	2.4970	0.4547	5.4914	<0.0001
	CTR vs MUL	3.3646	0.6768	4.9714	<0.0001
	CTR vs PEL	3.9226	0.9172	4.2766	0.0002
	FLW vs MOW	1.4608	0.5025	2.9071	0.0300
	FLW vs MUL	2.3284	0.7099	3.2801	0.0092
	FLW vs PEL	2.8864	0.9418	3.0649	0.0185
	MOW vs MUL	0.8676	0.8065	1.0758	0.8191
	MOW vs PEL	1.4256	1.0166	1.4023	0.6262
MUL vs PEL	0.5580	1.1335	0.4923	0.9881	
2019	CTR vs FLW	2.7412	0.6138	4.466	0.0001
	CTR vs MOW	4.5139	1.0357	4.3582	0.0001
	CTR vs MUL	4.4093	2.1447	2.0559	0.2395
	CTR vs PEL	5.7303	1.2766	4.4886	0.0001
	FLW vs MOW	1.7727	1.1968	1.4812	0.5747
	FLW vs MUL	1.6681	2.2164	0.7526	0.9439
	FLW vs PEL	2.9891	1.4136	2.1146	0.2137
	MOW vs MUL	-0.1046	2.3729	-0.0441	0.9999
	MOW vs PEL	1.2164	1.6422	0.7407	0.9469
MUL vs PEL	1.3210	2.4918	0.5301	0.9843	

Table 4. Post hoc test for the difference over time of the proportion of *S. deltoideus* covering in the 1-year treated plots with different weeding techniques (FLW, MOW, MUL and PEL) and the control (CTR). Fix terms is the interaction treatment with days after first treatment (treatment:days).

Contrast	Estimate	SE	z.ratio	p.value
CTR - FLW	1.6907	0.4693	3.6028	0.0029
CTR - MOW	1.4656	0.3513	4.1721	0.0003
CTR - MUL	1.6445	0.4602	3.5734	0.0032
CTR - PEL	3.4266	0.9313	3.6795	0.0022
FLW - MOW	-0.2251	0.5606	-0.4015	0.9945
FLW - MUL	-0.0462	0.6344	-0.0728	1.0000
FLW - PEL	1.7358	1.0322	1.6817	0.4453
MOW - MUL	0.1789	0.5529	0.3235	0.9976
MOW - PEL	1.9609	0.9842	1.9924	0.2696
MUL - PEL	1.7820	1.0281	1.7334	0.4133




Figure 3. Natural re-growth of *S. deltoideus* in the CTR plots in 2018 (black continuous line) compared with the natural re-growth in the 1-year treated plots, verifying the enduring effect of one-year treatment (MOW = mowing; FLW = flame weeding; MUL = mulching; PEL = pelargonium acid).

The proportion of *Senecio* covering was partially different between 1-year treatment and 2 years treatment plots. In particular, MOW and MUL were significantly different between 1-year treatment and 2 years treatment plots, while in FLW and PEL no significant differences were detected (Table 2 and 5 and Fig. 4).

Table 5. Difference in proportion of *S. deltoideus* covering between 1-year treated and 2-years treated plots with the different weeding techniques (FLW, MOW, MUL and PEL), assessed using generalized linear mixed effect model with negative binomial distribution.

	χ^2	df	P value
FLW	1.9873	1	0.1586
MOW	7.3703	1	0.0066
MUL	4.3117	1	0.0378
PEL	0.9088	1	0.3404




Figure 4. Comparison between 1-year (black continuous line) and 2-years (yellow dotted line) treated plots detected in 2019. The first are those no more treated during the second year, the second are those on which the 4 weeding techniques were applied a second time (MOW = mowing; FLW = flame weeding; MUL = mulching; PEL = pelargonium acid).

DISCUSSION

In this study, we tested different eco-friendly and cost-effective weeding techniques on *S. deltoideus*, in order to define the easiest and most convenient strategy in managing this invasive plant in a protected area of the northern Mediterranean coasts. The untreated plots quantified for the first time the capability in vegetative growth of the species in a non-native area and its invasive potential toward natural vegetation: the species showed a constant, rapid and complete re-colonization of the plot after the start of the mowing operation at the beginning of the vegetative seasons (Table 1 and 2). When stimulated by a single control cutting a year (corresponding to a basic invasive control made in protected areas), the plant shows a vigorous gemmation activity at the base (personal observations) as observed in *Senecio pterophorus* DC. (alien invasive species in Italy - Barberis et al., 1998) in the same geographical area of this study. The spring and autumn precipitations of the Mediterranean climate support the plant growth and the seed germination, as already observed in the other co-generic *S. pterophorus* and *S. inaequidens* DC., alien species to Spain and Italy (Caño et al., 2007). In particular, *S. deltoideus* seems to be sensible to the summer aridity typical of the Mediterranean climate (Fig. 1) when it temporarily stops its fast and vigorous vegetative growth, similarly to other congeneric species (Caño et al., 2007).

The multiple application of all four eco-friendly and cost-effective weeding techniques during the same year resulted in a high reduction of *S. deltoideus* covering (Table 1 and 2), evidencing that to obtain effective control of perennial alien plants, more frequently repeated treatments are required (Rask et al., 2013). According to literature, a single weeding control may have a negative effect by promoting the invasion of an alien invasive species and decreasing natural vegetation competitiveness (Trtikova, 2009). On the contrary, as observed in *Senecio jacobaea* and other alien perennial plant species, consecutive cuts are necessary to achieve maximum control (Suter et al., 2007; Roos et al., 2011), depleting the reserves in the plants by repeated killing of the leaves (Raghu et al., 2006), interrupting seedling growth and causing seed bank depletion (Sebastian et al., 2017).

All four weeding techniques, applied for two consecutive years, showed a progressively and drastic reduction of *S. deltoideus* covering (Tables 1 and 2), but they had significantly different effect. During the first-year treatments, FLW was less effective but in the second-year treatment the same ineffectiveness was not detected. Among the four weeding techniques applied, PEL showed the most consistent and constant covering reduction along years. MUL and MOW showed a similar efficiency with a covering reduction during the first year treatments but definitely lower in the second year treatment.

Flame weeding (FLW) is the most common thermal weed control method used in agriculture in both organic and conventional systems (Ascard, 1995), as an alternative to herbicide applications (Peruzzi et al., 2007; Datta & Knezevic 2013; Fontanelli et al., 2013), and few times in the management of natural areas. One of the limits of FLW is that the operator has the subjective responsibility to define the sufficient quantity of heat to cause the killing of weeds. In addition, this personal assessment is also combined with the environmental conditions under which the treatment is carried out. The weather conditions of the day (wind, rain, etc.), the trend of the season (more or less dry), the characteristics of the habitat (amount of litter and undergrowth, etc.) can lead to different results in the application of FLW (Ulloa et al., 2012). For these reasons, FLW resulted in a lower effectiveness in our experiments along the first year treatments and in the inconsistency between the first and second year. Moreover, the use of FLW in natural environment is still debated and controversial: on the one hand, there is growing evidence that these management projects in wide scale in natural vegetation alter ecosystem structure in ways that promote alien plant invasion (Keeley, 2006). On the other hand, when used in altered environments, fire is likely to promote persistence of aliens (Brooks et al., 2004), and being difficult to apply to a targeted species, it may favour other alien vigorous re-sprouting plants (Keeley, 2001). However, the use of FLW to control invasive herbaceous plants in open natural spaces may be advantageous because it avoids the spreading of chemical residues in the environment and the herbicide carryover to the following season, it is effective on a very wide spectrum of weeds (Ascard, 1995; Mojžiš, 2002; Fontanelli et al., 2013). The other two physical eradication techniques used to control *S. deltoideus* (i.e., mulching, MUL and mowing, MOW) showed an effectiveness similar to FLW but they were more constant between years. However, these techniques have two common drawbacks: they can be expensive because of the manpower involved and they disrupt the soil, creating disturbed sites prone to new invasion (DiTomaso, 2000), as also supported by the re-growth observed at the end of the second year in 1-year treatment plots. In general, MUL is most effective on controlling small seeded species and it is marginally effective on established re-sprouting perennials (Hayes et al., 2019), but in this study we recorded satisfactory results on *S. deltoideus* too (Table 1 and 2). This method has the advantage that the operator work is limited in time (the mulch layer is laid once a season) and it is made with the sustainable use of locally produced recycled material. On the other side, MUL increases the risk of summer fire in the natural environment for the huge accumulation of highly flammable material. Finally, MUL may be also hazardous in terms of introduction, promotion and establishment of further alien species (Kruse et al., 2004; Keeley, 2006). MOW is often used to control annuals but can occasionally reduce

seed production and can provide suppression of biennials and perennials (Biazzo & Milbrath, 2019), if used repeatedly as we did. Timing is critical to the success of MOW and the repetition of several cuts in the year leads to a complete consumption of the plant's reserves (DiTomaso et al., 2013). The advantages of MOW are its quick and easy execution and its applicability on large surfaces.

The highest effectiveness and consistency between years was achieved using PEL, a chemical control never performed on *Senecio* before. Recent studies report the use of synthetic systemic herbicides (2,4-D, Asulam and MCPA) to control some invasive species of *Senecio* (Roberts & Pullin, 2004). They effectively reduced population density (e.g. *S. jacobaea* and *S. aquaticus*), but there are legitimate concerns over the use of herbicides in terms of potential environmental impacts. Although newer herbicides are less toxic, have shorter residence times, and are more specific, concerns over detrimental environmental impacts remain (Borokini & Babalola, 2012), especially if applied in natural protected areas. Differently, the pelargonic acid is a natural extract with a contact herbicide effect, and the plant does not translocate it (Muñoz et al., 2020). Thus, it behaves similarly to fire, and if application is timed correctly, it may be used in a targeted way on specific alien plants (Ciriminna et al., 2019). The main limitation is that the product is expensive: the suggested dosage is 170 litres of product per ha, resulting in a cost of about 1,300 euros per hectare (Muñoz et al., 2020).

The interruption of treatments on some plots during the second year showed a potential long lasting effect of the weeding techniques on *S. deltoideus*, keeping its covering reduced over time (Table 4 and Fig. 3). In particular, FLW and PEL did not show any significantly difference in the comparison between 1-year and 2-year treated plots, demonstrating an enduring effect as a control technique. This result opens interesting future perspective for *S. deltoideus* management: the control might be limited to biannual cycles of treatment, maintaining the invasion degree at a low level and reducing the cost.

CONCLUSION

Evaluating both advantages and disadvantages of the four weeding techniques and their implementation costs, mowing is the recommendable control method for *S. deltoideus* (Table 6) because it fulfils the best compromise among a good covering decrement, a lasting effect and a moderate demand of manpower, without harmful consequences for the environment. Despite sometimes more effective according to our results, the other treatments bring some important contraindications (Table 6): they are either much more expensive (PEL), or very dangerous in creating potential wood fires (FLW and MUL).

However, the selection of the best strategy to control *S. deltoideus* is difficult to define. As suggested by many studies in literature, long-term and repeated control methods would be needed to effectively manage invasive plants, but it is always necessary to consider the costs and benefits of each methodology when implementing control programmes (Matarczyk et al., 2002; Wootton et al., 2005; MacDonald et al., 2007). If an exotic weed is already widespread, a species-specific organic control may be the only long-term effective method able to suppress its abundance over large areas (Mack & Lonsdale, 2002). In this context, our second-year results for no-treated plots revealed that in a long-time management strategy the selected treatment could be applied every two years with a drastic reduction in costs for the manager of the protected area.

Table 6. Comparison among the weeding techniques on their eco-friendly and cost-effective performances during an experimental year. Efficacy = mean covering resulted after 1-year treatment. Manpower = expressed as days per hectare per year for the application of the technique. Cost = referred to the material used applying the technique. Risk = potential risk applying the technique. Enduring = mean covering resulted in 1-year treated plots if untreated during the second year.

	Efficacy (%)	Manpower (day)	Cost	Risk	Enduring (%)
MOW	2-7	5	cheap	-	0-2.5
FLW	17-38	5	cheap	Wood fire	1-17
MUL	1.5-5	2	cheap	Wood fire	0-2.5
PEL	0.2-4	5	expensive	Veg. desicc.	0-13

ACKNOWLEDGEMENTS

We would like to thank Francesco Ghilino for supporting the experimental set-up of the pots experiment. We are grateful to Oscar Marchelli for supporting photos sampling during 2019. This study was performed in frame of the 2014-2020 Interreg Maritime Program, Project "ALIEM - Action pour Limiter les risques de diffusion des espèces Introduites Envahissantes en Méditerranée". Gabriele Casazza gratefully acknowledges financial support from the European Union's Horizon 2020 Research and Innovation Programme [Grant Agreement No. 793226].

REFERENCES

- Alpert P., Bone E., Holzapfel C., 2000. Invasiveness, invasibility and the role of environmental stress in the spread of non-native plants. *Perspectives in Plant Ecology Evolution and Systematics* 3, 52-66.
- Andreau J., Vilà M., Hulme P.E., 2009. An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management* 43, 1244-1255.
- Ascard J., 1995. Effects of flame weeding on weed species at different developmental stages. *Weed Research* 35(5), 397-411.
- Bais H.P., Vepachedu R., Gilroy S., Callaway R.M., Vivanco J.M., 2003. Allelopathy and exotic plant invasion: from molecules and genes to species interactions. *Science* 301, 1377-1380.
- Bandarra P.M., de Oliveira L.G., Cabrera Dalto A., Marques Boabaid F., Juffo G., Riet-Correa F., Driemeier D., Farias da Cruz C.E., 2012. Sheep production as a *Senecio* spp. control tool. *Pesquisa Veterinária Brasileira* 32(10), 1017-1022.
- Barberis G., Minuto L., Peccenini S., 1998. *Senecio grisebachii* Baker (Compositae), new to the Italian flora. *Webbia* 52, 201-206.
- Bates D., Maechler M., Bolker B., Walker S., 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67(1), 1-48.
- Biazzo J., Milbrath L.R., 2019. Response of pale swallowwort (*Vincetoxicum rossicum*) to multiple years of mowing. *Invasive Plant Science and Management* 12, 169-175.
- Borokini T.I., Babalola F.D., 2012. Management of invasive plant species in Nigeria through economic exploitation: lessons from other countries. *Management of Biological Invasions* 3(1), 45-55.
- Bossdorf O., Auge H., Lafuma L., Rogers W.E., Siemann E., Prati D., 2005. Phenotypic and genetic differentiation between native and introduced plant populations. *Oecologia* 144, 1-11.
- Braun M., Schindler S., Essl F., 2016. Distribution and management of invasive alien plant species in protected areas in Central Europe. *Journal for Nature Conservation* 33, 48-57.
- Brooks M.L., D'Antonio C.M., Richardson D.M., Grace J.B., Keeley J.E., DiTomaso J.M., Hobbs R.J., Pellant M., Pyke D., 2004. Effects of invasive alien plants on fire regimes. *BioScience* 54, 677-688.
- Campodonico P.G., 2010. Thomas Hanbury and his Garden. *Bollettino dei Musei e degli Istituti Biologici dell'Università di Genova* 72, 12-14.
- Campodonico P.G., Zappa E., 2006. Introduction of ornamental species of Labiatae at Hanbury Botanic Gardens – La Mortola (Italy). *Proceedings of the First International Symposium on the Labiatae. Acta Horticulture* 723, 105-109.
- Caño L., Escarré J., Sans F.X., 2007. Factors affecting the invasion success of *Senecio inaequidens* and *S. pterophorus* in Mediterranean plant communities. *Journal of Vegetation Science* 18, 281-288.
- Chicouene D., 2007. Mechanical destruction of weeds. A review. *Agronomy for Sustainable Development* 27, 19-27.
- Child L.E., Wade M., Wagner M., 1998. Cost effective control of *Fallopia japonica* using combination treatments. In *Plant invasions: ecological mechanisms and human responses* (eds. Starfinger U., Edwards K., Kowarik I., Williamson M.), pp. 143-154. Backhuys Publishers, Leiden, The Netherlands.
- Ciriminna R., Fidalgo A., Ilharco L.M., Pagliaro M., 2019. Herbicides based on pelargonic acid: Herbicides of the bioeconomy. *Biofuels, Bioproducts and Biorefining* 13(6), 1476-1482.
- Datta A., Knezevic S.Z., 2013. Flaming as an alternative weed control method for conventional and organic agronomic crop production systems: A review. In *Advances in Agronomy* (ed. Sparks D.), 118, 399-428. Elsevier Academic Press, San Diego, CA.
- Dehnen-Schumutz K., Perrings C., Williamson M., 2004. Controlling *Rhododendron ponticum* in the British Isles: an economic analysis. *Journal of Environmental Management* 70, 323-332.
- DePoorter M., Pagad S., Ullah M.I., 2007. Invasive alien species and protected areas: a scoping report. The World Bank, Washington, DC. Available at: www.issg.org/Animal%20Imports%20Webpage/Electronic%20References/IASinPAS.pdf. (last accessed 02.03.2020)
- DiTomaso J.M., 2000. Invasive weeds in rangelands: species, impacts, and management. *Weed Science* 48, 255-265.
- DiTomaso J.M., Kyser G.B., Oneto S.R., Wilson R.G., Orloff S.B., Anderson L.W., Wright S.D., Roncoroni J.A., Miller T.L., Prather T.S., Ransom C., Beck K.G., Duncan C., Wilson K.A., Mann J.J., 2013. Weed control in natural areas in the western United States. *Weed Research and Information Center, University of California*, 544.
- Ellstrand N., Schierenbeck K., 2006. Hybridization as a stimulus for the evolution of invasiveness in plants? *Euphytica* 148, 35-46.

- Ellwanger T.C., Bingham S.W., Chappell W.E., Tolin S.A., 1973. Cytological effects of ultra-high temperatures on corn. *Weed science* 21, 299-303.
- Erskine Ogden J.A., Rejmánek M., 2005. Recovery of native plant communities after the control of a dominant invasive plant species, *Foeniculum vulgare*: implications for management. *Biological Conservation* 125, 427-439.
- Ervin J., 2003. Rapid assessment of protected area management effectiveness in four countries. *BioSciences* 53, 833-841.
- Essl F., Latombe G., Lenzner B., Pagad S., Seebens H., Smith K., Wilson J.R.U., Genovesi P., 2020. Biological invasions are one of the most serious threats to global biodiversity and ecosystem integrity. *NeoBiota* 62, 99-121.
- Fontanelli M., Raffaelli M., Martelloni L., Frascioni C., Peruzzi A., 2013. The influence of non-living mulch, mechanical and thermal treatments on weed population and yield of rainfed fresh-market tomato (*Solanum lycopersicum* L.). *Spanish Journal of Agricultural Research* 11(3), 593-602.
- Fried G., 2010. Prioritization of potential invasive alien plants in France. In: *Proceedings of the 2nd International workshop on invasive plants in the Mediterranean type regions of the world*. pp. 120-134. EPPO, Trabzon, Turkey.
- Galasso G., Conti F., Peruzzi L., Ardenghi N.M.G., Banfi E., Celesti-Grapow L., Albano A., Alessandrini A., Bacchetta G., Ballelli S., Bandini Mazzanti M., Barberis G., Bernardo L., Blasi C., Bouvet D., Bovio M., Cecchi L., Del Guacchio E., Domina G., Fascetti S., Gallo L., Gubellini L., Guiggi A., Iamónico D., Iberite M., Jiménez-Mejías P., Lattanzi E., Marchetti D., Martinetto E., Masin R.R., Medagli P., Passalacqua N.G., Peccenini S., Pennesi R., Pierini B., Podda L., Poldini L., Prosser F., Raimondo F.M., Roma-Marzio F., Rosati L., Santangelo A., Scoppola A., Scortegagna S., Selvaggi A., Selvi F., Soldano A., Stinca A., Wagensommer R.P., Wilhelm T., Bartolucci F., 2018. An updated checklist of the vascular flora alien to Italy. *Plant Biosystems* 152(3), 556-592.
- García-Llorente M., Martín-López B., Nunes P.A.L.D., González J.A., Alcorlo P., Montes C., 2011. Analyzing the social factors that influence willingness to pay. *Environmental Management* 48, 418-435.
- Geesing D., Felker P., Bingham R.L., 2000. Influence of mesquite (*Prosopis glandulosa*) on soil nitrogen and carbon development: Implications for global carbon sequestration. *Journal of Arid Environments* 46(2), 157-180.
- Hayes D.G., Anunciado M.B., DeBruyn J.M., Bandopadhyay S., Schaeffer S., English M., Ghimire S., Miles C., Flury M., Sintim H.Y., 2019. Biodegradable plastic mulch films for sustainable specialty crop production. In *Polymers for agri-food applications* (ed. Gutiérrez T.), pp. 183-213. Springer, Cham, Switzerland.
- Hulme P.E., 2003. Biological invasions: Winning the science battles but losing the conservation war? *Oryx* 37, 178-193.
- Hyde M.A., Wursten B.T., Ballings P., Coates Palgrave M., 2020. Flora of Zimbabwe: Home page. Available at <https://www.zimbabweflora.co.zw/index.php> (last accessed 21 January 2020).
- Keeley J.E., 2001. Fire and invasive species in mediterranean-climate ecosystems of California. In *Proceedings of the invasive species workshop: the role of fire in the control and spread of invasive species* (eds. Galley K.E.M., Wilson T.P.), Miscellaneous publication 1181-94. Tall Timbers Research Station, Tallahassee, FL.
- Keeley J.E., 2006. Fire management impacts on invasive plants in the Western United States. *Conservation Biology* 20(2), 375-384.
- Kettenring K.M., Adams C.R., 2011. Lessons learned from invasive plant control experiments: a systematic review and meta-analysis. *Journal of Applied Ecology* 48, 970-979.
- Křivánek M., 2006. Biologické invaze a možnosti jejich předpovědi (Predikční modely pro stanovení invazního potenciálu vyšších rostlin) [Biological invasions and possibilities of their prediction (Prediction models for determination of invasive potential of higher plants)]. *Acta Pruhoniciana* 84, 83-92.
- Kruse R., Bend E., Bierzychudek P., 2004. Native plant regeneration and introduction of non-natives following post-fire rehabilitation with straw mulch and barley seeding. *Forest Ecology and Management* 196, 299-310.
- Lenth R., 2020. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.4.7. <https://CRAN.R-project.org/package=emmeans>.
- MacDonald N.W., Scull B.T., Abella S.R., 2007. Mid-spring burning reduces spotted knapweed and increases native grasses during a Michigan experimental grassland establishment. *Restoration Ecology* 15, 118-128.
- Mack R.N., Lonsdale W.M., 2002. Eradicating invasive plants: hard-won lessons for islands. In *Tuning the tide: the eradication of invasive species* (eds. Veitch C.R., Clout M.N.), 164-172. IUCN, Cambridge, UK.
- Mack R.N., Simberloff D., Lonsdale W.M., Evans H., Clout M., Bazzaz F.A., 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications* 10(3), 689-710.

- Matarczyk J.A., Willis A.J., Vranjic J.A. Ash J.E., 2002. Herbicides, weeds and endangered species: management of bitou bush (*Chrysanthemoides monilifera* ssp. *rotundata*) with glyphosate and impacts on the endangered shrub, *Pimelea spicata*. *Biological Conservation* 108, 133-141.
- McNeely J., 2001. The great reshuffling: human dimensions of invasive alien species, 113-119. IUCN, Gland, Switzerland.
- Milakovic I., Fiedler K., Karrer G., 2014. Management of roadside populations of invasive *Ambrosia artemisiifolia* by mowing. *Weed Research* 54, 256-264.
- Miller K.E., Gorchoff D.L., 2004. The invasive shrub, *Lonicera maackii*, reduces growth and fecundity of perennial forest herbs. *Oecologia* 139, 359-375.
- Mojžiš M., 2002. Energetic requirements of flame weed control. *Research in Agricultural Engineering* 48, 94-97.
- Muñoz M., Torres-Pagán N., Peiró R., Guijarro R., Sánchez-Moreiras A.M., Verdeguer M., 2020. Phytotoxic effects of three natural compounds: pelargonic acid, carvacrol, and cinnamic aldehyde, against problematic weeds in Mediterranean crops. *Agronomy* 10(6), 791.
- Nielsen C., Ravn H.P., Nentwig W., Wade M., 2005. The giant hogweed best practice manual. Guidelines for the management and control of invasive weeds in Europe forest and landscape. Hørsholm, Denmark.
- Oerke E.C., 2006. Crop losses to pests. Centenary review. *The Journal of Agricultural Science* 144, 31-43.
- Peruzzi A., Ginanni M., Fontanelli M., Raffaelli M., Barberi P., 2007. Innovative strategies for on-farm weed management in organic carrot. *Renewable Agriculture and Food Systems* 22(4), 246-259.
- Pimentel D., Lach L., Zuniga R., Morrison D., 2000. Environmental and economic costs of Nonindigenous species in the United States. *Bio-Science* 50, 53-65.
- Pimentel D., Zuniga R., Morrison D., 2005. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecological Economics* 52, 273-288.
- R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Raghu S., Dhileepan K., Treviño M., 2006. Response of an invasive liana to simulated herbivory: implications for its biological control. *Acta Oecologica* 29, 335-345.
- Rask A.M., Larsen S.U., Andreasen C., Kristoffersen P., 2013. Determining treatment frequency for controlling weeds on traffic islands using chemical and non-chemical weed control. *Weed Research* 53, 249-258.
- Rejmánek M., Richardson D.M., 1996. What attributes make some plant species more invasive? *Ecology* 6, 1655-1661.
- Richardson D.M., Pyšek P., Rejmánek M., Barbour M.G., Panetta F.D., West C.J., 2000. Naturalization and invasion of alien plants: concepts and definitions. *Diversity and Distributions* 6, 93-107.
- Roberts P.D., Pullin A.S., 2004. Effectiveness of the control of ragwort (*Senecio*) species: "Are currently recommended herbicides effective for control of ragwort (*Senecio*) species?" CEE review 04-003 (SR5a).
- Roos K., Rödel H.G., Beck E., 2011. Short- and long-term effects of weed control on pastures infested with *Pteridium arachnoideum* and an attempt to regenerate abandoned pastures in South Ecuador. *Weed Research* 51, 165-176.
- Sebastian D.J., Nissen S.J., Sebastian J.R., 2017. Seed bank depletion: the key to long-term downy brome (*Bromus tectorum* L.) management. *Rangeland Ecology & Management* 70, 477-483.
- Shackleton R.T., Foxcroft L.C., Pyšek P., Wood L.E., Richardson D.M., 2020. Assessing biological invasions in protected areas after 30 years: Revisiting nature reserves targeted by the 1980s SCOPE programme. *Biological Conservation* 243, 108424.
- Shaw R.H., 2003. Biological control of invasive weeds in the UK: opportunities and challenges. In *Plant invasions: ecological threats and management solutions* (eds. Child L.E., Brock J.H., Brundu G. et al.), 337-354. Backhuys Publishers, Leiden, The Netherlands.
- Stohlgren T.J., Binkley D., Chong G.W., A. Kalkhan M., Schell L.D., Bull K.A., Otsuki Y., Newman G., Bashkin M., Son Y., 1999. Exotic plant species invade hot spots of native plant diversity. *Ecological Monographs* 69, 25-46.
- Suter M., Siegrist-Maag S., Connolly J., Lüscher A., 2007. Can the occurrence of *Senecio jacobaea* be influenced by management practice? *Weed Research* 47, 262-269.
- Trtikova M., 2009. Effects of competition and mowing on growth and reproduction of the invasive plant *Erigeron annuus* at two contrasting altitudes. *Botanica Helvetica* 119, 1-6.
- Ulloa S.M., Datta A., Bruening C., Gogosc G., Arkebauer T.J., Knezevi S.Z., 2012. Weed control and crop tolerance to propane flaming as influenced by the time of day. *Crop Protection* 31, 1-7.
- Viciani D., Vidali M., Gigante D., Bolpagni R., Villani M., Rosario Acosta A.T., Adorni M., Aleffi M., Allegrezza M.,

Angiolini C., Assini S., Bagella S., Bonari G., Bovio M., Bracco F., Brundu G., Buffa G., Caccianiga M., Carnevali L., Ceschin S., Ciaschetti G., Cogoni A., Di Cecco V., Foggi B., Frattaroli A.R., Genovesi P., Gentili R., Lazzaro L., Lonati M., Lucchese F., Mainetti A., Mariotti M., Minissale P., Paura B., Pellizzari M., Perrino E.V., Pirone G., Poggio L., Poldini L., Poponessi S., Prisco I., Prosser F., Puglisi M., Rosati L., Selvaggi A., Sottovia L., Spampinato G., Stanisci A., Stinca A., Venanzoni R., Lastrucci L. (2020) A first checklist of the alien vegetation in Italy. *Plant Sociology* 57, 29-54.

Villaverde J.J., Sevilla-Morán B., Sandín-España P., López-Goti C., Alonso-Prados J.L., 2014. Biopesticides in the framework of the European Pesticide Regulation (EC) No. 1107/2009. *Pest Management Science* 70, 2-5.

Wearne L.J., Morgan J.W., 2004. Community-level changes in Australian subalpine vegetation following invasion by the non-native shrub *Cytisus scoparius*. *Journal of Vegetation Science* 15, 595-604.

Weidlich E.W.A., Flórido F.G., Sorrini T.B., Brancalion P.H.S., 2020. Controlling invasive plant species in ecological restoration: a global review. *Journal of Applied Ecology* 57, 1806-1817.

Wootton L.S., Halsey S.D., Bevaart K., McGough A., Ondreicka J., Patel P., 2005. When invasive species have benefits as well as costs: managing *Carex kobomugi* (Asiatic sand sedge) in New Jersey's coastal dunes. *Biological Invasions* 7, 1387-3547.

Zappa E., Campodonico P.G., 2005. *Senecio deltoideus* Less. (Asteraceae), specie sud africana sfuggita alla coltivazione nei Giardini Botanici Hanbury. *Informatore Botanico Italiano* 37(1A), 458-459.