

Paleomicrobiology of the human digestive tract: A review

Riccardo Nodari, Michel Drancourt, Rémi Barbieri

▶ To cite this version:

Riccardo Nodari, Michel Drancourt, Rémi Barbieri. Paleomicrobiology of the human digestive tract: A review. Microbial Pathogenesis, 2021, 157, pp.104972. 10.1016/j.micpath.2021.104972. hal-03334114

HAL Id: hal-03334114 https://amu.hal.science/hal-03334114

Submitted on 13 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 Paleomicrobiology of the human digestive tract: a review. 2 3 Riccardo Nodaria, Michel Drancourtb, Rémi Barbierib,c* 4 5 6 ^aDepartment of Biosciences and Pediatric Clinical Research Center "Romeo and 7 Enrica Invernizzi", University of Milan, Milan, 20133, Italy. 8 ^bAix-Marseille Univ., IRD, MEPHI, IHU Méditerranée Infection, Marseille, France. 9 °UMR 7268, Anthropologie Bioculturelle, Droit, Ethique et Santé, Aix Marseille Univ., 10 11 CNRS, EFS, ADES, Marseille, France. 11 12 * Corresponding author: remibarbieri@hotmail.fr 13 IHU - Méditerranée Infection, 19-21 Boulevard Jean Moulin, 13005 Marseille. 14

Abstract

The microbiota is a hot topic of research in medical microbiology, boosted by culturomics and metagenomics, with unanticipated knowledge outputs in physiology and pathology. Knowledge of the microbiota in ancient populations may therefore be of prime interest in understanding factors shaping the coevolution of the microbiota and populations. Studies on ancient human microbiomes can help us understand how the community of microorganisms presents in the oral cavity and the gut was shaped during the evolution of our species and what environmental, social or cultural changes may have changed it. This review cumulates and summarizes the discoveries in the field of the ancient human microbiota, focusing on the remains used as samples and techniques used to handle and analyze them.

Keywords

- 28 Coprolite; dental calculus; microbiome; microbiota; mummified tissue;
- 29 paleomicrobiology; paleogenomics; paleoproteomics.

1. Introduction

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

In the last 20 years, there has been an exponential increase in interest in the microbiome and its effect on human health. We currently know that the microbiome plays a fundamental role in the host via the development, maintenance and priming of the immune system [1–3], defense against pathogens [4,5], metabolism, digestion and the production of vitamins [6]. Furthermore, slight modification of the bacterial equilibrium of the microbiome (dysbiosis) due to changes in diet, physical exercise or antibiotic treatments can lead to major consequences for the host [7–11]. Recent studies have focused their attention on the composition of the microbiome of our ancestors and its evolution throughout history. Particular attention was given to the periods of major revolutions in human lifestyles, such as the Neolithization process (debuted 12,000 years BCE [12]) and the industrialization period (1760-1840 CE). With the Neolithization process, we observed the transition from a nomadic huntergatherer diet characterized by the consumption of only wild and unprocessed food foraged and hunted from the environment [13] to a sedentary one based on agriculture and livestock. Then the industrialization period arose the advent of modern sanitation and Western diet, characterized by food that has been processed and modified and has a low quantity of vegetables and fibers [14–16]. In the literature, there is some confusion about the definition of the terms "microbiota" and "microbiome" and are often used interchangeably. The term "microbiota" refers to all the microbial taxa associated with the host [17], while "microbiome" can refer to "the ecological community of commensal, symbiotic, and pathogenic microorganisms that literally share our body space and have been all but

ignored as determinants of health and disease" [18] or to "the collective genome of 53 our indigenous microbes" [19]. 54 55 Diets of past and present human populations are very variable and mostly depend on resource availability and ecological settings [20]. Comparative studies on the 56 57 microbiota among human populations with different dietary habits have shown how 58 differences in diet are reflected in differences in the microbiota composition [14,16]. 59 For instance, De Filippo and colleagues [21] showed how children from urban 60 Europe, with a modern Western diet, and children from rural Africa, with a diet very 61 high in fibers, present differences in microbiota composition that can be attributed to 62 an evolutionary adaptation to the diet composition, allowing each population to maximize the nutritional value of each diet. Other studies on mouse models have 63 shown how changing diet correlated with changes in the microbial composition of the 64 gut microbiota [22]. Moreover, Groussin et al. [23] demonstrated how microbiota 65 66 composition reconstructed from ancient mammal remains could be used to reconstruct ancient diets of these animals and that information about dietary shift can 67 be recorded by the gut microbiome. For these reasons, ancient microbiota studies 68 69 are not restricted to the microbial diversity and composition of the ancient remains but also to the dietary information that is preserved inside these specimens. The first 70 study to target the ancient DNA of multiple species of bacteria on human remains 71 72 was conducted by Ubaldi et al, analyzing the gut microbiome from the colon of a mummy recovered in Cuzco, Peru, that dated back to the 10th-11th century CE [24]. 73 74 Following by many other PCR-based studies on coprolites and mummified samples [25–27]. In 2008, Tito et al. performed the first study on an ancient microbiota using 75 NGS techniques, and from that point, the number of ancient microbiome analyses 76 77 multiplied [28].

Retrieving information on ancient human microbiomes is not easy and is strictly dependent on the availability of suitable samples for the analysis. Dental calculus and coprolite are the only two human remains that persist long enough to be used for these studies, with the exceptions of frozen and mummified tissues [29]. In this review, we will present the remains used as starting samples for ancient human microbiota studies and the methods used to retrieve this information.

2. Literature search strategy and terminology use

- All the papers used for this review were obtained from PubMed and Google Scholar.
- The keywords used alone or in combination for the search were "paleomicrobiology",
- "ancient", "human", "paleo", "microbiota", "microbiome", "oral", "dental calculus",
- "gut", "intestinal", "feces", "faeces", "paleofeces", "paleofaeces", "coprolite", "latrines",
- 89 "mummy", "mummified", "tissue", "metagenomic", "shotgun", and "proteomics".
- 90 Only studies written in English and on samples of human origin that targeted multiple
- 91 species from the digestive tract microbiota of the host were considered. The results
- of this bibliographical research are shown in Table 1 (last search March 2021).
- 93 As described in the introduction section, the two terms "microbiota" and "microbiome"
- are often used interchangeably, and since a discussion on the terminology was not
- one of the purposes of this review, we used the two terms as synonyms, following
- 96 the definition given by Lederberg [18].

Contamination prevention and authentication

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

Paleomicrobiological studies on ancient molecules must follow standard protocols to avoid modern exogenous contamination and ensure the authenticity of the results [30]. Contaminating molecules can come from the burial site, handling, transport, storage, and laboratory environments; from the materials and instruments used during the analysis; and from the operators at each step of the study [31,32]. Regardless of the starting sample type, all the steps of the analysis of ancient molecules must be performed in a dedicated clean hood and laboratory, ideally exclusively dedicated to work on ancient remains, excluding any positive controls. and including several negative controls that should be handled strictly in parallel with investigated samples [31]. Authentication of ancient DNA can be performed by checking for fragmentation (ancient DNA rarely exceeds 200-300 bp length) [33,34] and degradation (e.g., assessing the PCR success related to the size of the amplicon of a selected DNA sequence) [33]. Multiple bioinformatics tools can contribute to the authentication of ancient molecules. Software like MapDamage 2.0 [35,36] quantify the patterns of damage in NGS datasets generated from ancient DNA and therefore help differentiate endogenous from exogenous/contaminating DNA. Other tools like SourceTracker [33,37] estimate the proportion of DNA sequences that come from the source environments to determine whether the reconstructed microbial communities observed are endogenous to the samples. Moreover, directly comparing the DNA sequences obtained from different tissues with those from soil from the site where samples have been collected through bioinformatic analysis may facilitate the interpretation of the data [34].

3. Ancient Oral Cavity Microbiota.

3.1. Dental calculus

represented in Figure 1.

Dental calculus, or tartar, is a mineralized bacterial biofilm that forms on the teeth as a result of the action of bacteria and saliva in the oral cavity and is very common in adults without active dental hygiene [29]. Dental calculus presents some characteristics that make it the most reliable source for information about the oral microbiota. It is common and relatively abundant in archaeological sites; it has the capability to preserve ancient biomolecules, avoiding environmental contaminants; and it contains a high quantity of DNA originating from the oral microbiome [29,38,39]. In fact, it can be found on teeth in all past and present human populations and can survive for thousands and even millions of years. Accordingly, dental calculus remains were obtained from a 12-8 million-year-old fossil of the ape *Sivapithecus* [40].

3.1.1. Analysis of ancient DNA from dental calculus.

After the recent confirmation of the presence and abundance of DNA in dental calculus [41], multiple studies have successfully extracted and analyzed DNA sequences from this type of sample. Different protocols have been applied for DNA extraction in the past. In general, it is very similar to apply the extraction methods used for other types of samples, such as bones and teeth, with a "lysis step" followed by a "purification step". The lysis step usually relies on decalcification through ethylenediaminetetraacetic acid (EDTA) with or without Tris-HCl and protein digestion using proteinase K with or without sodium-dodecyl-sulfate (SDS), with a

variable incubation time and temperature (usually 24 h at 56 °C) [38,42–46]. Occasionally, in the lysis step, Tris-HCl and SDS are added to the sample to improve decalcification and protein digestion [38,47]. After the lysis step, purification of DNA is performed using the phenol:chloroform:isoamyl alcohol (25:24:1) method [47] preferentially followed by Qiagen MinElute column purification [38,43,45,46] or using commercial kits for DNA purification, such as the EZ1 DNA tissue kit (Qiagen) [44,48] and QIAamp DNA Investigator Kit (Qiagen) [42]. It is also possible to use an in-house silica-based method for the extraction and purification of ancient DNA [20]. Polymerase chain reaction (PCR) amplification and subsequent sequencing can be used to analyze DNA sequences specific for microorganisms from the oral microbial community. PCR primers were specifically designed for five species of bacteria, Actinomyces naeslundii, Fusobacterium nucleatum, Porphyromonas gingivalis, Streptococcus gordonii and Streptococcus mutans, by De La Fuente et al. [47], which were detected in dental calculus from 4,000-year-old to modern samples. Specific primers are useful for the identification of single species of microorganisms, such as pathogens or common microbes of the oral microbiota; to determine the conservation of DNA in the sample; and to study specific genes [49]. The amplification of a conserved bacterial region of the 16S ribosomal RNA (16S rRNA) gene and subsequent high-throughput sequencing yields information about the diversity of bacteria present in the samples, describing the bacterial community of the ancient oral microbiota [20,38,42,50,51]. Using this method, it is possible to identify hundreds of species with a single PCR amplification [49]. Adler et al. [42] used the results of high-throughput sequencing to compare the bacteria present in the oral cavity of Mesolithic, Neolithic, Bronze Age, medieval and modern individuals,

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

reporting a reduction in the bacterial diversity and a shift toward cariogenic bacterial dominance after medieval times, concomitant with the industrial revolution [42]. Following the same principle, regions conserved among any organism of interest can be amplified. PCR sequencing of 16S rRNA and methyl coenzyme M reductase (*mcrA*) genes was used to detect and describe the methanogen community present in the oral human microbiota, allowing the identification of multiple archaea, such as *Candidatus* Methanobrevibacter sp. N13, *Methanobrevibacter oralis* and *Methanoculleus bourgensis* [44]. Unfortunately, this approach has limited applicability when the aDNA is known to be highly degraded and fragmented or when the targeted region contains extensive length polymorphisms that induce a bias toward taxa with the shortest amplicon lengths [43].

In contrast to targeted gene amplification, shotgun sequencing allows the sequencing of random DNA fragments present in the sample independently of their origin (eukarya, bacteria, archaea, and viruses), overcoming targeted sequencing problems [38,43]. This approach allows the analysis of all the sequences inside the sample at the same time without any prior knowledge, providing both taxonomic and functional information [49]. Shotgun metagenomics applied to ancient human dental calculus allowed the characterization of the oral microbiota from multiple archaeological sites [20,38,39], providing information about antibiotic resistance and metabolic functions of ancient bacteria [20,38]. Data obtained with this approach showed a possible correlation between behavior, diet, and health and the oral microbiota composition of two populations of European Neanderthals [20]. This finding suggested a central role of meat consumption in early hominid microbiota composition and that the repertoire of microorganisms preserved in dental calculus could be used to derive the dietary behavior of ancient populations by comparing the

shotgun data obtained to those obtained for other ancient calculus specimens from populations with a known diet. Moreover, the shotgun metagenomic approach also allowed the reconstruction of draft genomes of organisms present in ancient dental calculus, such as the bacterium *Tannerella forsythia* [38] and the archaeon *Methanobrevibacter oralis* [20].

3.1.2. Analysis of ancient protein from dental calculus.

Similar to shotgun metagenomics for nucleotides, shotgun proteomics allows the detection and study of extracted peptides from ancient human dental calculus [52]. Despite the better preservation capability of proteins than DNA molecules, proteomic approaches are relatively new in the field of paleomicrobiology, providing valuable information about the diet, microbiota function and composition, host health and immune response [53,54]. Warriner and colleagues showed how tandem mass spectrometry (MS/MS) used for shotgun proteomics can produce information about the presence of bacterial species-specific proteins, virulence factors, and antigenic proteins, as well as human immune response proteins (antimicrobial gene products) [38,55]. In addition, it has been shown that mass spectrometry for proteomic analysis can be used to describe the oral human metabolome (the collection of all the low-weight metabolites produced by metabolism [46]), providing important information about salivary and microbiome-related proteins [46] and dietary information from ancient populations [56,57].

3.1.3. Microscopical analysis on dental calculus.

Microscopic analysis of dental calculus allowed the detection of aliments and the presence of calcified microorganisms [58,59]. Examination through scanning electron microscopy (SEM) can be performed without treating the sample, allowing

complete preservation [60], while optical microscopy and transmission electron microscopy (TEM) need a pretreatment to be performed that partially or completely disrupts the sample [60]. For SEM, a preliminary step for the decalcification of the sample is performed using hydrochloric acid (HCI) at different concentrations [38,61]. For TEM, ancient samples follow routine procedure, with treatment with osmium tetroxide fixative and epoxy resin embedding [41]. Optical microscopy can be used to distinguish between gram-positive and gram-negative bacteria [38,60,62]. Optical and electron microscopy usually do not allow the identification of species but can provide important pieces of information about the presence, dimension and shape of microorganisms in the samples [59]. Microscopic analysis can also be used for biomolecular investigations, using gold-labeled antibody TEM to detect the presence of DNA [41]; fluorescence microscopy and polyclonal antibodies for the identification of specific bacteria, such as *Streptococcus mutans* [62]; and fluorescent *in situ* hybridization (FISH) to reveal archaea, such as *Candidatus* Methanobrevibacter sp. [44].

3.2. Other sources of ancient oral microbiota.

Despite the vast majority of studies on ancient oral microbiota have been performed on dental calculus, some authors tried to think outside of the box obtaining interesting results. In 2019, Jensen and colleagues analyzed a chewed birch pitch from Denmark dated to 5,700 years ago and revealed the presence of preserved DNA material. Starting from this specimen, they reconstructed a complete ancient human genome and identified microbial DNA sequences that reflected the oral microbiome of the person who chewed the pitch [63]. This work highlighted the

potential utility of chewed birch pitch in the field of paleomicrobiology for the study of ancient microbiota.

Moreover, the analysis of buccal swabs from Inca mummies through shotgun proteomics analysis was used to describe active states of infection on an individual at the time of death by providing information on the immune response and potentially on the pathogens present when combined with DNA amplification [64].

4. Ancient Gut Microbiota.

4.1. Coprolites and palaeofaeces.

The primary sources for information about the ancient gut microbiota are coprolites, which are "any formed fecal mass remains, including mineralized, desiccated, or frozen feces and even the intestinal contents of mummies" [65]. These remains can be preserved for thousands or even millions of years [66], providing information about the gut microbiota components, the presence of pathogens and the diet of past human and animal populations [67]. In fact, traces of aliments consumed by the host, such as bones of small animals, feathers, seafood shells, and plant seeds, are commonly found in coprolites [65,68–71]. Moreover, it is possible to correlate the presence of animal parasites with the diet of an ancient population [70,72,73].

4.1.1. Molecular analysis on coprolites.

DNA analysis can potentially produce additional information about the complete set of species inside the coprolite and the evolution of these organisms. To reduce environmental contamination, extraction of DNA is performed on the core of the coprolites after the removal of a few millimeters (3 mm - 1 cm) of coprolite surface. Multiple protocols and commercial kits are available for DNA extraction from fecal

and soil samples and are also used on ancient samples, e.g., UltraClean Fecal DNA Isolation Kit (MOBIO) [33] and PowerSoil® DNA Isolation Kit (MOBIO) [70,72–74]. Protocol specific for coprolites that does not require commercial kits have also been developed, such as the one from Kuch *et al.* [75]. In 2019, Hagan and colleagues tested the efficacy of 5 different methods for DNA extraction from ancient fecal material [36], indicating that despite every tested method being valuable, methods used for modern microbiota analysis recovered less DNA than methods developed and optimized for ancient bone material [39,76].

Massive sequencing of specific amplified regions of the DNA, such as 16S rDNA,

provided information about the prokaryotic community inside the coprolite [34,70,74]. The same approach can be used for the eukaryotic community, e.g., by using primers that target the 18S ribosomal DNA (18S rDNA) or Internal transcribed spacer (ITS) region for fungal diversity [70,74]. Whole-genome shotgun sequencing does not require previous amplification of the DNA and gives information on the entirety of DNA sequences present in the sample, allowing both taxonomic and functional characterization [28,34,73,77]. Furthermore, pathogen DNA sequences, including those of Clostridium botulinum, Trypanosoma cruzi and human papillomaviruses (HPVs) [34], as well as drug resistance genes [34,77], are detected. Moreover, viral metagenomes from coprolites have been described [72,78].

4.1.2. Microscopic and immunoassay analysis on coprolites.

Rehydration of samples is the preliminary step for subsequent microscopic analysis. Different protocols have been developed for coprolite rehydration [79]. These protocols all rely on using 0.5% trisodium phosphate (Na₃PO₄) aqueous solution sometimes complemented with 5% glycerinated water and formalin (Le Bailly 2010)

or hydrochloric acid (HCI) to dissolve the calcium carbonates [80]. This step can last from 24 h to one week [81,82] and can be combined with a sedimentation step [68,73,77]. Microscopic analysis allows the identification of multicellular parasites, such as helminth eggs [73,77,83]; protozoan cysts [84–86]; and food traces [87,88]. Parasite identification relies on morphometric characterization with light and electron microscopy, which is based on the experience and ability of the operator. Cases have been reported in which pollen grains, which are morphologically similar to pinworm eggs, have been confused for parasitic eggs [89]. The use of confocal laser scanning microscopy (CLSM) has been revealed to be a powerful tool for the identification of parasites in coprolites. This technique allows better morphological identification, detects intrinsic autofluorescent molecules and has the great advantage of being less destructive to the specimen, allowing further analyses to be performed (e.g., molecular analysis) on the same specimen [90]. Moreover, immunofluorescence assays (IFAs) or enzyme-linked immunosorbent assays (ELISAs) have been used for the detection of intestinal protozoa, such as Giardia spp. [84,91–95], Cryptosporidium spp. [84,95,96], and Entamoeba histolytica [93– 95,97,98].

4.1.3. Culture-dependent analysis on coprolites.

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

Finally, a culture-dependent investigation was performed on a 14th-century coprolite found in Belgium in parallel to a metagenomic analysis [77]. A portion from the inner part of the coprolite was inoculated into Schaedler and R2A liquid and solid culture media and maintained for several days at 30 °C under various atmospheres created using gas-generating pouch systems (BD Gas PakTM EZ). Small colonies were observable after 10 days of incubation, and ten different bacterial species

(Paenibacillus macerans, Bacillus jeotgali, Staphylococcus pasteuri, Staphylococcus epidermidis, Staphylococcus cohnii, Micrococcus luteus, Pseudomonas geniculate, Stenotrophomonas maltophilia, Bacillus horti and Clostridium magnum) were identified by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF-MS) and 16S rRNA gene sequencing. Additionally, a portion from the inner part of the coprolite was cultured in BD BACTEC, Lytic/10 Anaerobic and Aerobic bottles enriched with 7 mL of defibrinated sheep blood and incubated at 37 °C. After two days, cultures were diluted and plated onto COS culture plates (bioMérieux) and reincubated at 37 °C under different atmospheres. Eight different bacterial species, Rhodanobacter sp. organism, Paenibacillus sp. organism, Paenibacillus macerans, Paenibacillus thiaminolyticus, Paenibacillus ehimensis, Staphylococcus arlettae, Propionibacterium acnes and Enterobacter cloacae, were isolated and identified from the blood-enriched cultures using MALDI-TOF-MS and 16S rRNA gene sequencing.

4.2. Mummified tissues.

Other than coprolites, natural and artificial mummified tissues are also valuable sources for ancient human microbiota studies. DNA, proteins, and even entire parasites (eggs, larvae, and adults) have been found in mummified tissues. Well-preserved mummies offer the rare possibility to study microbiota both inside and outside of the digestive tract, allowing the identification of specific pathogens of certain organs, i.e., *Helicobacter pylori* [99], and even compare the microbial communities present in different organs from the same individual [25]. Mummified tissues from the digestive tract are used for the identification of the microbial community of the gut, stomach, and oral cavity. Similar to the methods used for

coprolite analysis, mummified tissues can be analyzed for parasite and protozoan detection through microscopy. The rehydration step can be performed following the same methods used for coprolites with trisodium phosphate [79] or with a solution of potassium hydroxide at 4-7% [79,100]. Techniques of 16S rRNA gene high-throughput sequencing and metagenomics performed on mummified digestive tract tissues were used for the characterization of ancient microbiota. The presence of DNA sequences homologous to many organisms of medical interest were also detected, such as *Clostridium botulinum*, Trypanosoma cruzi, HPVs, Pseudomonas spp., and Leishmania donovani [24,101,102]. The genomes of 5,300-year-old Helicobacter pylori [99], Clostridium perfringens, Clostridium sp. Ade.TY, Clostridium algidicarnis, and Pseudomonas veronii [102] were reconstructed and used for phylogenetic analysis to study the evolution of these organisms. Moreover, metagenomic analysis allowed the identification of sequences homologous to bacteriophages [103], metabolic profiles [101], multiple sequences associated with antibiotic resistance and virulence-related genes in old mummified gut tissues [34,102,104]. Data obtained from mummified gut remains could also potentially be used to infer ancient dietary habits [104]. Organs connected to the digestive tract have been studied using microscopy or DNA-related techniques for the presence of pathogens and parasites: studies on gallbladder samples detected Clonorchis sinensis [105], Mycobacterium tuberculosis [106]; studies on liver samples detected hepatitis B virus [107]. Geographical distribution of the specimens analyzed for the gut microbiota are

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

represented in figure 1.

5. Perspectives

We have just begun to understand the relationships between diet, health, pathological conditions, and the composition of our microbiota, but it is clear that the gut microbiota plays a role in the health and disease conditions of humans. This explains the strong interest in medical research for this subject. Combining modern techniques and protocols used in the field of paleomicrobiology for the analysis and the extraction of data from ancient human remains, we can study the past of humankind and its microbiota. An evolutionary framework that leads us to understand how these changes in human lifestyle have influenced changes in the microbiota can be built through the comparison of the microbiota composition among different eras and with data on the microbiota in the modern world. Studies on the microbiota of our ancestors can help us describe the evolutionary path that led humans and their microbiota to where we are now, revealing how changes in diet, innovations in modern medicine, hygiene, and antibiotics have changed the composition and, therefore, how the characteristics of our microbiota have shaped humans as a species.

374 **References**

- N. Kamada, S.-U. Seo, G.Y. Chen, G. Núñez, Role of the gut microbiota in immunity and inflammatory disease, Nat Rev Immunol. 13 (2013) 321–335. https://doi.org/10.1038/nri3430.
- 378 [2] Y. Belkaid, O.J. Harrison, Homeostatic immunity and the microbiota, Immunity. 46 (2017) 562–576.
- 380 [3] S. Sanna, N.R. van Zuydam, A. Mahajan, A. Kurilshikov, A.V. Vila, U. Võsa, Z.
 381 Mujagic, A.A. Masclee, D.M. Jonkers, M. Oosting, Causal relationships among the gut
 382 microbiome, short-chain fatty acids and metabolic diseases, Nature Genetics. 51
 383 (2019) 600–605.
- T. Ichinohe, I.K. Pang, Y. Kumamoto, D.R. Peaper, J.H. Ho, T.S. Murray, A. Iwasaki, Microbiota regulates immune defense against respiratory tract influenza A virus infection, Proceedings of the National Academy of Sciences. 108 (2011) 5354–5359.
- J.M. Pickard, M.Y. Zeng, R. Caruso, G. Núñez, Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease, Immunological Reviews. 279 (2017) 70–89.
- 390 [6] I. Rowland, G. Gibson, A. Heinken, K. Scott, J. Swann, I. Thiele, K. Tuohy, Gut 391 microbiota functions: metabolism of nutrients and other food components, European 392 Journal of Nutrition. 57 (2018) 1–24.
- L. Dethlefsen, S. Huse, M.L. Sogin, D.A. Relman, The pervasive effects of an antibiotic on the human gut microbiota, as revealed by deep 16S rRNA sequencing, PLoS Biol. 6 (2008) e280. https://doi.org/10.1371/journal.pbio.0060280.
- 396 [8] M.T. Alou, J.-C. Lagier, D. Raoult, Diet influence on the gut microbiota and dysbiosis related to nutritional disorders, Human Microbiome Journal. 1 (2016) 3–11.
- 398 [9] M.P. Francino, Antibiotics and the human gut microbiome: dysbioses and accumulation of resistances, Frontiers in Microbiology. 6 (2016) 1543.
- V. Monda, I. Villano, A. Messina, A. Valenzano, T. Esposito, F. Moscatelli, A.
 Viggiano, G. Cibelli, S. Chieffi, M. Monda, Exercise modifies the gut microbiota with positive health effects, Oxidative Medicine and Cellular Longevity. 2017 (2017).
- 403 [11] R.K. Singh, H.-W. Chang, D. Yan, K.M. Lee, D. Ucmak, K. Wong, M. Abrouk, B. Farahnik, M. Nakamura, T.H. Zhu, Influence of diet on the gut microbiome and implications for human health, Journal of Translational Medicine. 15 (2017) 73.
- 406 [12] A. Eddine, R.G. Rocha, N. Mostefai, Y. Karssene, K. De Smet, J.C. Brito, D. Klees, C. Nowak, B. Cocchiararo, S. Lopes, P. van der Leer, R. Godinho, Demographic expansion of an African opportunistic carnivore during the Neolithic revolution, Biology Letters. 16 (2020) 20190560. https://doi.org/10.1098/rsbl.2019.0560.
- 410 [13] J.H. O'Keefe, L. Cordain, Cardiovascular disease resulting from a diet and lifestyle at 411 odds with our Paleolithic genome: how to become a 21st-century hunter-gatherer, 412 Mayo Clin Proc. 79 (2004) 101–108. https://doi.org/10.4065/79.1.101.
- 413 [14] K.M. Maslowski, C.R. Mackay, Diet, gut microbiota and immune responses, Nature Immunology. 12 (2011) 5–9.
- L.T. Stiemsma, L.A. Reynolds, S.E. Turvey, B.B. Finlay, The hygiene hypothesis: current perspectives and future therapies, ImmunoTargets and Therapy. 4 (2015) 143.
- 417 [16] L. Moles, D. Otaegui, The Impact of Diet on Microbiota Evolution and Human Health.
 418 Is Diet an Adequate Tool for Microbiota Modulation?, Nutrients. 12 (2020) 1654.
- 419 [17] L.K. Ursell, J.L. Metcalf, L.W. Parfrey, R. Knight, Defining the human microbiome, 420 Nutrition Reviews. 70 (2012) S38–S44.
- 421 [18] J. Lederberg, A.T. McCray, Ome SweetOmics--A genealogical treasury of words, The Scientist. 15 (2001) 8–8.
- 423 [19] L.V. Hooper, J.I. Gordon, Commensal host-bacterial relationships in the gut, Science. 424 292 (2001) 1115–1118.

- 425 [20] L.S. Weyrich, S. Duchene, J. Soubrier, L. Arriola, B. Llamas, J. Breen, A.G. Morris, K.W. Alt, D. Caramelli, V. Dresely, Neanderthal behaviour, diet, and disease inferred from ancient DNA in dental calculus, Nature. 544 (2017) 357–361.
- 428 [21] C. De Filippo, D. Cavalieri, M. Di Paola, M. Ramazzotti, J.B. Poullet, S. Massart, S.
 429 Collini, G. Pieraccini, P. Lionetti, Impact of diet in shaping gut microbiota revealed by a
 430 comparative study in children from Europe and rural Africa, Proceedings of the
 431 National Academy of Sciences. 107 (2010) 14691–14696.
- 432 [22] P.J. Turnbaugh, M. Hamady, T. Yatsunenko, B.L. Cantarel, A. Duncan, R.E. Ley, M.L. Sogin, W.J. Jones, B.A. Roe, J.P. Affourtit, A core gut microbiome in obese and lean twins, Nature. 457 (2009) 480–484.
- 435 [23] M. Groussin, F. Mazel, J.G. Sanders, C.S. Smillie, S. Lavergne, W. Thuiller, E.J. Alm, Unraveling the processes shaping mammalian gut microbiomes over evolutionary time, Nature Communications. 8 (2017) 1–12.
- 438 [24] M. Ubaldi, S. Luciani, I. Marota, G. Fornaciari, R.J. Cano, F. Rollo, Sequence analysis 439 of bacterial DNA in the colon of an Andean mummy, American Journal of Physical 440 Anthropology: The Official Publication of the American Association of Physical 441 Anthropologists. 107 (1998) 285–295.
- 442 [25] R.J. Cano, F. Tiefenbrunner, M. Ubaldi, C. Del Cueto, S. Luciani, T. Cox, P. Orkand, K.H. Künzel, F. Rollo, Sequence analysis of bacterial DNA in the colon and stomach of the Tyrolean Iceman, American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists. 112 (2000) 297–309.
- 447 [26] S. Luciani, G. Fornaciari, O. Rickards, C.M. Labarga, F. Rollo, Molecular 448 characterization of a pre-Columbian mummy and in situ coprolite, American Journal of 449 Physical Anthropology. 129 (2006) 620–629. https://doi.org/10.1002/ajpa.20314.
- 450 [27] F. Rollo, S. Luciani, I. Marota, C. Olivieri, L. Ermini, Persistence and decay of the 451 intestinal microbiota's DNA in glacier mummies from the Alps, Journal of 452 Archaeological Science. 34 (2007) 1294–1305. 453 https://doi.org/10.1016/j.jas.2006.10.019.
- 454 [28] R.Y. Tito, S. Macmil, G. Wiley, F. Najar, L. Cleeland, C. Qu, P. Wang, F. Romagne, S. Leonard, A.J. Ruiz, K. Reinhard, B.A. Roe, C.M.L. Jr, Phylotyping and Functional Analysis of Two Ancient Human Microbiomes, PLOS ONE. 3 (2008) e3703. https://doi.org/10.1371/journal.pone.0003703.
- 458 [29] C. Warinner, C. Speller, M.J. Collins, C.M. Lewis Jr, Ancient human microbiomes, 459 Journal of Human Evolution. 79 (2015) 125–136.
- 460 [30] A. Cooper, H.N. Poinar, Ancient DNA: do it right or not at all, Science. 289 (2000)
 461 1139. https://doi.org/10.1126/science.289.5482.1139b.
 462 [31] M. Drancourt, D. Raoult, Palaeomicrobiology: current issues and perspectives, Na

- [31] M. Drancourt, D. Raoult, Palaeomicrobiology: current issues and perspectives, Nature Reviews Microbiology. 3 (2005) 23–35. https://doi.org/10.1038/nrmicro1063.
- 464 [32] D.Y. Yang, K. Watt, Contamination controls when preparing archaeological remains 465 for ancient DNA analysis, Journal of Archaeological Science. 32 (2005) 331–336. 466 https://doi.org/10.1016/j.jas.2004.09.008.
- 467 [33] R.Y. Tito, D. Knights, J. Metcalf, A.J. Obregon-Tito, L. Cleeland, F. Najar, B. Roe, K. Reinhard, K. Sobolik, S. Belknap, M. Foster, P. Spicer, R. Knight, C.M.L. Jr, Insights from Characterizing Extinct Human Gut Microbiomes, PLOS ONE. 7 (2012) e51146. https://doi.org/10.1371/journal.pone.0051146.
- 471 [34] T.M. Santiago-Rodriguez, G. Fornaciari, S. Luciani, S.E. Dowd, G.A. Toranzos, I.
 472 Marota, R.J. Cano, Gut microbiome of an 11 th century AD Pre-Columbian Andean
 473 mummy, PloS One. 10 (2015) e0138135.
- 474 [35] H. Jónsson, A. Ginolhac, M. Schubert, P.L.F. Johnson, L. Orlando, mapDamage2.0: 475 fast approximate Bayesian estimates of ancient DNA damage parameters, 476 Bioinformatics. 29 (2013) 1682–1684. https://doi.org/10.1093/bioinformatics/btt193.
- 477 [36] R.W. Hagan, C.A. Hofman, A. Hübner, K. Reinhard, S. Schnorr, C.M. Lewis Jr, K. Sankaranarayanan, C.G. Warinner, Comparison of extraction methods for recovering

- 479 ancient microbial DNA from paleofeces, American Journal of Physical Anthropology. 480 171 (2020) 275–284.
- 481 [37] D. Knights, J. Kuczynski, E.S. Charlson, J. Zaneveld, M.C. Mozer, R.G. Collman, F.D. Bushman, R. Knight, S.T. Kelley, Bayesian community-wide culture-independent microbial source tracking, Nature Methods. 8 (2011) 761–763. https://doi.org/10.1038/nmeth.1650.
- 485 [38] C. Warinner, J.F.M. Rodrigues, R. Vyas, C. Trachsel, N. Shved, J. Grossmann, A.
 486 Radini, Y. Hancock, R.Y. Tito, S. Fiddyment, C. Speller, J. Hendy, S. Charlton, H.U.
 487 Luder, D.C. Salazar-García, E. Eppler, R. Seiler, L.H. Hansen, J.A.S. Castruita, S.
 488 Barkow-Oesterreicher, K.Y. Teoh, C.D. Kelstrup, J.V. Olsen, P. Nanni, T. Kawai, E.
 489 Willerslev, C. von Mering, C.M. Lewis, M.J. Collins, M.T.P. Gilbert, F. Rühli, E.
 490 Cappellini, Pathogens and host immunity in the ancient human oral cavity, Nature
 491 Genetics. 46 (2014) 336–344. https://doi.org/10.1038/ng.2906.

493

494

495

496

497

501

502

503

504

505

506

507

508

515

516

- [39] A.E. Mann, S. Sabin, K. Ziesemer, Å.J. Vågene, H. Schroeder, A.T. Ozga, K. Sankaranarayanan, C.A. Hofman, J.A. Fellows Yates, D.C. Salazar-García, B. Frohlich, M. Aldenderfer, M. Hoogland, C. Read, G.R. Milner, A.C. Stone, C.M. Lewis, J. Krause, C. Hofman, K.I. Bos, C. Warinner, Differential preservation of endogenous human and microbial DNA in dental calculus and dentin, Scientific Reports. 8 (2018) 9822. https://doi.org/10.1038/s41598-018-28091-9.
- 498 [40] I. Hershkovitz, J. Kelly, B. Latimer, B.M. Rothschild, S. Simpson, J. Polak, M.
 499 Rosenberg, Oral bacteria in miocene Sivapithecus, J Hum Evol. 33 (1997) 507–512.
 500 https://doi.org/10.1006/jhev.1997.0149.
 - [41] H.R. Preus, O.J. Marvik, K.A. Selvig, P. Bennike, Ancient bacterial DNA (aDNA) in dental calculus from archaeological human remains, Journal of Archaeological Science. 38 (2011) 1827–1831. https://doi.org/10.1016/j.jas.2011.03.020.
 - [42] C.J. Adler, K. Dobney, L.S. Weyrich, J. Kaidonis, A.W. Walker, W. Haak, C.J.A. Bradshaw, G. Townsend, A. Sołtysiak, K.W. Alt, J. Parkhill, A. Cooper, Sequencing ancient calcified dental plaque shows changes in oral microbiota with dietary shifts of the Neolithic and Industrial revolutions, Nat Genet. 45 (2013) 450–455, 455e1. https://doi.org/10.1038/ng.2536.
- K.A. Ziesemer, A.E. Mann, K. Sankaranarayanan, H. Schroeder, A.T. Ozga, B.W.
 Brandt, E. Zaura, A. Waters-Rist, M. Hoogland, D.C. Salazar-García, M. Aldenderfer,
 C. Speller, J. Hendy, D.A. Weston, S.J. MacDonald, G.H. Thomas, M.J. Collins, C.M.
 Lewis, C. Hofman, C. Warinner, Intrinsic challenges in ancient microbiome
 reconstruction using 16S rRNA gene amplification, Sci Rep. 5 (2015) 16498.
 https://doi.org/10.1038/srep16498.
 - [44] H.T.T. Huynh, V.D. Nkamga, M. Signoli, S. Tzortzis, R. Pinguet, G. Audoly, G. Aboudharam, M. Drancourt, Restricted diversity of dental calculus methanogens over five centuries, France, Sci Rep. 6 (2016) 25775. https://doi.org/10.1038/srep25775.
- 518 [45] A.T. Ozga, M.A. Nieves-Colón, T.P. Honap, K. Sankaranarayanan, C.A. Hofman, G.R. Milner, C.M. Lewis, A.C. Stone, C. Warinner, Successful enrichment and recovery of whole mitochondrial genomes from ancient human dental calculus, Am J Phys Anthropol. 160 (2016) 220–228. https://doi.org/10.1002/ajpa.22960.
- I.M. Velsko, K.A. Overmyer, C. Speller, L. Klaus, M.J. Collins, L. Loe, L.A.F. Frantz, K.
 Sankaranarayanan, C.M. Lewis, J.B.R. Martinez, E. Chaves, J.J. Coon, G. Larson, C.
 Warinner, The dental calculus metabolome in modern and historic samples,
 Metabolomics. 13 (2017) 134. https://doi.org/10.1007/s11306-017-1270-3.
- 526 [47] C.D.L. Fuente, S. Flores, M. Moraga, Dna from Human Ancient Bacteria: A Novel 527 Source of Genetic Evidence from Archaeological Dental Calculus, Archaeometry. 55 528 (2013) 767–778. https://doi.org/10.1111/j.1475-4754.2012.00707.x.
- 529 [48] A. Bringuier, S. Khelaifia, H. Richet, G. Aboudharam, M. Drancourt, Real-Time PCR
 530 Quantification of Methanobrevibacter oralis in Periodontitis, J Clin Microbiol. 51 (2013)
 531 993–994. https://doi.org/10.1128/JCM.02863-12.
- 532 [49] L.S. Weyrich, K. Dobney, A. Cooper, Ancient DNA analysis of dental calculus, J Hum Evol. 79 (2015) 119–124. https://doi.org/10.1016/j.jhevol.2014.06.018.

- 534 [50] T.M. Santiago-Rodriguez, Y. Narganes-Storde, L. Chanlatte-Baik, G.A. Toranzos, R.J. Cano, Insights of the dental calculi microbiome of pre-Columbian inhabitants from Puerto Rico, PeerJ. 5 (2017) e3277.
- [51] T.M. Santiago-Rodriguez, A. Fornaciari, G. Fornaciari, S. Luciani, I. Marota, G.
 Vercellotti, G.A. Toranzos, V. Giuffra, R.J. Cano, Commensal and Pathogenic
 Members of the Dental Calculus Microbiome of Badia Pozzeveri Individuals from the
 11th to 19th Centuries, Genes (Basel). 10 (2019).
 https://doi.org/10.3390/genes10040299.
- 542 [52] J.R. Yates, Mass spectrometry and the age of the proteome, J Mass Spectrom. 33 (1998) 1–19. https://doi.org/10.1002/(SICI)1096-9888(199801)33:1<1::AID-JMS624>3.0.CO;2-9.
- [53] E. Cappellini, M.J. Collins, M.T.P. Gilbert, Biochemistry. Unlocking ancient protein
 palimpsests, Science. 343 (2014) 1320–1322.
 https://doi.org/10.1126/science.1249274.
- 548 [54] R. Barbieri, R. Mekni, A. Levasseur, E. Chabrière, M. Signoli, S. Tzortzis, G. Aboudharam, M. Drancourt, Paleoproteomics of the Dental Pulp: The plague paradigm, PLoS One. 12 (2017) e0180552. https://doi.org/10.1371/journal.pone.0180552.

553

554

555

556

557

558

559

560

561

562 563

564

565

566

570

571

- [55] R.R. Jersie-Christensen, L.T. Lanigan, D. Lyon, M. Mackie, D. Belstrøm, C.D. Kelstrup, A.K. Fotakis, E. Willerslev, N. Lynnerup, L.J. Jensen, E. Cappellini, J.V. Olsen, Quantitative metaproteomics of medieval dental calculus reveals individual oral health status, Nat Commun. 9 (2018) 4744. https://doi.org/10.1038/s41467-018-07148-3.
 - [56] C. Warinner, J. Hendy, C. Speller, E. Cappellini, R. Fischer, C. Trachsel, J. Arneborg, N. Lynnerup, O.E. Craig, D.M. Swallow, A. Fotakis, R.J. Christensen, J.V. Olsen, A. Liebert, N. Montalva, S. Fiddyment, S. Charlton, M. Mackie, A. Canci, A. Bouwman, F. Rühli, M.T.P. Gilbert, M.J. Collins, Direct evidence of milk consumption from ancient human dental calculus, Sci Rep. 4 (2014) 7104. https://doi.org/10.1038/srep07104.
- [57] J. Hendy, C. Warinner, A. Bouwman, M.J. Collins, S. Fiddyment, R. Fischer, R. Hagan, C.A. Hofman, M. Holst, E. Chaves, L. Klaus, G. Larson, M. Mackie, K. McGrath, A.Z. Mundorff, A. Radini, H. Rao, C. Trachsel, I.M. Velsko, C.F. Speller, Proteomic evidence of dietary sources in ancient dental calculus, Proc Biol Sci. 285 (2018). https://doi.org/10.1098/rspb.2018.0977.
- 567 [58] C. Meller, I. Urzua, G. Moncada, C.V. Ohle, Prevalence of oral pathologic findings in an ancient pre-Columbian archeologic site in the Atacama Desert, Oral Diseases. 15 (2009) 287–294. https://doi.org/10.1111/j.1601-0825.2009.01524.x.
 - [59] H.T.T. Huynh, J. Verneau, A. Levasseur, M. Drancourt, G. Aboudharam, Bacteria and archaea paleomicrobiology of the dental calculus: a review, Molecular Oral Microbiology. 31 (2015) 234–242. https://doi.org/10.1111/omi.12118.
- 573 [60] P. Charlier, I. Huynh-Charlier, O. Munoz, M. Billard, L. Brun, G.L. de L. Grandmaison, 574 The microscopic (optical and SEM) examination of dental calculus deposits (DCD). 575 Potential interest in forensic anthropology of a bio-archaeological method, Legal 576 Medicine. 12 (2010) 163. https://doi.org/10.1016/j.legalmed.2010.03.003.
- [61] K. Hardy, S. Buckley, L. Copeland, Pleistocene dental calculus: Recovering
 information on Paleolithic food items, medicines, paleoenvironment and microbes,
 Evolutionary Anthropology: Issues, News, and Reviews. 27 (2018) 234–246.
 https://doi.org/10.1002/evan.21718.
- 581 [62] A. Linossier, M. Gajardo, J. Olavarria, Paleomicrobiological study in dental calculus: Streptococcus mutans, Scanning Microsc. 10 (1996) 1005–1013; discussion 1014.
- T.Z.T. Jensen, J. Niemann, K.H. Iversen, A.K. Fotakis, S. Gopalakrishnan, Å.J.
 Vågene, M.W. Pedersen, M.-H.S. Sinding, M.R. Ellegaard, M.E. Allentoft, L.T.
 Lanigan, A.J. Taurozzi, S.H. Nielsen, M.W. Dee, M.N. Mortensen, M.C. Christensen,
 S.A. Sørensen, M.J. Collins, M.T.P. Gilbert, M. Sikora, S. Rasmussen, H. Schroeder,
 A 5700 year-old human genome and oral microbiome from chewed birch pitch, Nat
 Commun. 10 (2019) 5520. https://doi.org/10.1038/s41467-019-13549-9.

- 589 [64] A. Corthals, A. Koller, D.W. Martin, R. Rieger, E.I. Chen, M. Bernaski, G. Recagno, 590 L.M. Dávalos, Detecting the Immune System Response of a 500 Year-Old Inca 591 Mummy, PLOS ONE. 7 (2012) e41244. https://doi.org/10.1371/journal.pone.0041244.
- 592 [65] K. Reinhard, Coprolite Analysis: The analysis of ancient human feces for dietary data, 593 Karl Reinhard Papers/Publications. (2000). 594 https://digitalcommons.unl.edu/natresreinhard/60.
- 595 [66] P.C. Dentzien-Dias, G. Poinar, A.E.Q. de Figueiredo, A.C.L. Pacheco, B.L.D. Horn, C.L. Schultz, Tapeworm eggs in a 270 million-year-old shark coprolite, PLoS One. 8 (2013) e55007. https://doi.org/10.1371/journal.pone.0055007.
- 598 [67] S. Appelt, M. Drancourt, M. Le Bailly, Human Coprolites as a Source for
 599 Paleomicrobiology, Microbiol Spectr. 4 (2016).
 600 https://doi.org/10.1128/microbiolspec.PoH-0002-2014.
- 601 [68] K.J. Reinhard, V.M. Bryant Jr, Coprolite analysis: a biological perspective on archaeology, Archaeological Method and Theory. (1992) 245–288.
- 603 [69] K.J. Reinhard, J.R. Ambler, C.R. Szuter, Hunter-gatherer use of small animal food 604 resources: coprolite evidence, International Journal of Osteoarchaeology. 17 (2007) 605 416–428.
- [70] R.J. Cano, J. Rivera-Perez, G.A. Toranzos, T.M. Santiago-Rodriguez, Y.M. Narganes Storde, L. Chanlatte-Baik, E. García-Roldán, L. Bunkley-Williams, S.E. Massey,
 Paleomicrobiology: revealing fecal microbiomes of ancient indigenous cultures, PloS
 One. 9 (2014) e106833.
- 610 [71] G.P. Shipley, K. Kindscher, Evidence for the Paleoethnobotany of the Neanderthal: A Review of the Literature, Scientifica. 2016 (2016).
- [72] J.I. Rivera-Perez, R.J. Cano, Y. Narganes-Storde, L. Chanlatte-Baik, G.A. Toranzos,
 Retroviral DNA Sequences as a Means for Determining Ancient Diets, PLoS One. 10
 (2015). https://doi.org/10.1371/journal.pone.0144951.
- R. Wiscovitch-Russo, J. Rivera-Perez, Y.M. Narganes-Storde, E. García-Roldán, L.
 Bunkley-Williams, R. Cano, G.A. Toranzos, Pre-Columbian zoonotic enteric parasites:
 An insight into Puerto Rican indigenous culture diets and life styles, PLoS One. 15
 (2020). https://doi.org/10.1371/journal.pone.0227810.
- [74] T.M. Santiago-Rodriguez, Y.M. Narganes-Storde, L. Chanlatte, E. Crespo-Torres,
 G.A. Toranzos, R. Jimenez-Flores, A. Hamrick, R.J. Cano, Microbial Communities in
 Pre-Columbian Coprolites, PLoS One. 8 (2013).
 https://doi.org/10.1371/journal.pone.0065191.
- 623 [75] M. Kuch, H. Poinar, Extraction of DNA from paleofeces, Methods Mol Biol. 840 (2012) 37–42. https://doi.org/10.1007/978-1-61779-516-9_5.
- 525 [76] J. Dabney, M. Meyer, S. Pääbo, Ancient DNA damage, Cold Spring Harbor Perspectives in Biology. 5 (2013) a012567.
- 527 [77] S. Appelt, F. Armougom, M. Le Bailly, C. Robert, M. Drancourt, Polyphasic analysis of a middle ages coprolite microbiota, Belgium, PloS One. 9 (2014) e88376.
- [78] S. Appelt, L. Fancello, M. Le Bailly, D. Raoult, M. Drancourt, C. Desnues, Viruses in a
 14th-century coprolite, Applied and Environmental Microbiology. 80 (2014) 2648–
 2655.
- 632 [79] M. Camacho, A. Araújo, J. Morrow, J. Buikstra, K. Reinhard, Recovering parasites 633 from mummies and coprolites: an epidemiological approach, Parasites & Vectors. 11 634 (2018) 248.
- 635 [80] P.D. Mitchell, E. Anastasiou, D. Syon, Human intestinal parasites in crusader Acre: 636 evidence for migration with disease in the medieval period, International Journal of 637 Paleopathology. 1 (2011) 132–137.
- 638 [81] M. Seo, D.H. Shin, S.-M. Guk, C.S. Oh, E.-J. Lee, M.H. Shin, M.J. Kim, S.D. Lee, Y.-639 S. Kim, Y.S. Yi, M. Spigelman, J.-Y. Chai, Gymnophalloides seoi Eggs from the Stool 640 of a 17th Century Female Mummy Found in Hadong, Republic of Korea, Para. 94 641 (2008) 467–472. https://doi.org/10.1645/GE-1365.1.

- [82] K.J. Reinhard, M. Camacho, Pinworm Infection at Salmon Ruins and Aztec Ruins:
 Relation to Pueblo III Regional Violence, Korean J Parasitol. 57 (2019) 627–633.
 https://doi.org/10.3347/kjp.2019.57.6.627.
- 645 [83] L.M. Cleeland, M.V. Reichard, R.Y. Tito, K.J. Reinhard, C.M. Lewis Jr, Clarifying 646 prehistoric parasitism from a complementary morphological and molecular approach, 647 Journal of Archaeological Science. 40 (2013) 3060–3066.
- 648 [84] Y.R. Ortega, D. Bonavia, Cryptosporidium, giardia, and cyclospora in ancient 649 Peruvians, Journal of Parasitology. 89 (2003) 635–636.

651

652

653

660 661

662

663

664

665

666

667

668

669

670

671

672

- [85] M.H. Fugassa, M.O. Beltrame, N.H. Sardella, M.T. Civalero, C. Aschero, Paleoparasitological results from coprolites dated at the Pleistocene–Holocene transition as source of paleoecological evidence in Patagonia, Journal of Archaeological Science. 37 (2010) 880–884.
- 654 [86] L. Frías, D. Leles, A. Araújo, Studies on protozoa in ancient remains-A Review, 655 Memórias Do Instituto Oswaldo Cruz. 108 (2013) 1–12.
- [87] K.J. Reinhard, L.F. Ferreira, F. Bouchet, L. Sianto, J.M.F. Dutra, A. Iniguez, D. Leles,
 M. Le Bailly, M. Fugassa, E. Pucu, A. Araújo, Food, parasites, and epidemiological
 transitions: A broad perspective, International Journal of Paleopathology. 3 (2013)
 150–157. https://doi.org/10.1016/j.ijpp.2013.05.003.
 - [88] E. Pucu, J. Russ, K. Reinhard, Diet analysis reveals pre-historic meals among the Loma San Gabriel at La Cueva de Los Muertos Chiquitos, Rio Zape, Mexico (600–800 CE), Archaeological and Anthropological Sciences. 12 (2020) 1–14.
 - [89] M. Camacho, K.J. Reinhard, Confusing a Pollen Grain with a Parasite Egg: an Appraisal of "Paleoparasitological Evidence of Pinworm (Enterobius Vermicularis) Infection in a Female Adolescent Residing in Ancient Tehran," Korean J Parasitol. 57 (2019) 621–625. https://doi.org/10.3347/kjp.2019.57.6.621.
 - [90] J.J. Morrow, C. Elowsky, Application of Autofluorescence for Confocal Microscopy to Aid in Archaeoparasitological Analyses, The Korean Journal of Parasitology. 57 (2019) 581.
 - [91] M.L.C. Gonçalves, A. Araújo, R. Duarte, J.P. da Silva, K. Reinhard, F. Bouchet, L.F. Ferreira, Detection of Giardia duodenalis antigen in coprolites using a commercially available enzyme-linked immunosorbent assay, Trans R Soc Trop Med Hyg. 96 (2002) 640–643. https://doi.org/10.1016/S0035-9203(02)90337-8.
- [92] M. Le Bailly, M.L. Gonçalves, S. Harter-Lailheugue, F. Prodéo, A. Araujo, F. Bouchet,
 New finding of Giardia intestinalis (Eukaryote, Metamonad) in Old World
 archaeological site using immunofluorescence and enzyme-linked immunosorbent
 assays, Memórias Do Instituto Oswaldo Cruz. 103 (2008) 298–300.
 https://doi.org/10.1590/S0074-02762008005000018.
- [93] P.D. Mitchell, E. Stern, Y. Tepper, Dysentery in the crusader kingdom of Jerusalem:
 an ELISA analysis of two medieval latrines in the City of Acre (Israel), Journal of
 Archaeological Science. 35 (2008) 1849–1853.
 https://doi.org/10.1016/j.jas.2007.11.017.
- 683 [94] W.H. Eskew, M.L. Ledger, A. Lloyd, G. Pyles, J. Gosker, P.D. Mitchell, Intestinal 684 Parasites in an Ottoman Period Latrine from Acre (Israel) Dating to the Early 1800s 685 CE, Korean J Parasitol. 57 (2019) 575–580. https://doi.org/10.3347/kjp.2019.57.6.575.
- [95] D. Leles, L. Frías, A. Araújo, B. Brener, A. Sudré, M. Chame, V. Laurentino, Are immunoenzymatic tests for intestinal protozoans reliable when used on archaeological material?, Experimental Parasitology. 205 (2019) 107739.
 https://doi.org/10.1016/j.exppara.2019.107739.
- [96] J.J. Morrow, K.J. Reinhard, Cryptosporidium parvum Among Coprolites from La
 Cueva de los Muertos Chiquitos (600-800 CE), Rio Zape Valley, Durango, Mexico, J
 Parasitol. 102 (2016) 429–435. https://doi.org/10.1645/15-916.
- 693 [97] M.L.C. Goncalves, V.L. da Silva, C.M. de Andrade, K. Reinhard, G.C. da Rocha, M. Le Bailly, F. Bouchet, L.F. Ferreira, A. Araujo, Amoebiasis distribution in the past: first steps using an immunoassay technique, Trans R Soc Trop Med Hyg. 98 (2004) 88–91. https://doi.org/10.1016/S0035-9203(03)00011-7.

697 [98] M. Le Bailly, M.L. Gonçalves, C. Lefèvre, D.C. Roper, J.W. Pye, A. Araujo, F.
698 Bouchet, Parasitism in Kansas in the 1800s: a glimpse to the past through the
699 analysis of grave sediments from Meadowlark cemetery, Memórias Do Instituto
700 Oswaldo Cruz. 101 (2006) 53–56. https://doi.org/10.1590/S0074701 02762006001000009.

- [99] F. Maixner, B. Krause-Kyora, D. Turaev, A. Herbig, M.R. Hoopmann, J.L. Hallows, U. Kusebauch, E.E. Vigl, P. Malfertheiner, F. Megraud, N. O'Sullivan, G. Cipollini, V. Coia, M. Samadelli, L. Engstrand, B. Linz, R.L. Moritz, R. Grimm, J. Krause, A. Nebel, Y. Moodley, T. Rattei, A. Zink, The 5300-year-old Helicobacter pylori genome of the Iceman, Science. 351 (2016) 162–165. https://doi.org/10.1126/science.aad2545.
- [100] M.R. Hidalgo-Argüello, N. Díez Baños, J. Fregeneda Grandes, E. Prada Marcos, PARASITOLOGICAL ANALYSIS OF LEONESE ROYALTY FROM COLLEGIATE-BASILICA OF ST. ISIDORO, LEÓN (SPAIN): HELMINTHS, PROTOZOA, AND MITES, J Parasitol. 89 (2003) 738–743. https://doi.org/10.1645/0022-3395(2003)089[0738:PAOLRF]2.0.CO;2.
- [101] T.M. Santiago-Rodriguez, G. Fornaciari, S. Luciani, S.E. Dowd, G.A. Toranzos, I. Marota, R.J. Cano, Taxonomic and predicted metabolic profiles of the human gut microbiome in pre-Columbian mummies, FEMS Microbiol Ecol. 92 (2016). https://doi.org/10.1093/femsec/fiw182.
- [102] G.A. Lugli, C. Milani, L. Mancabelli, F. Turroni, C. Ferrario, S. Duranti, D. van Sinderen, M. Ventura, Ancient bacteria of the Ötzi's microbiome: a genomic tale from the Copper Age, Microbiome. 5 (2017). https://doi.org/10.1186/s40168-016-0221-y.
- [103] T.M. Santiago-Rodriguez, G. Fornaciari, S. Luciani, S.E. Dowd, G.A. Toranzos, I. Marota, R.J. Cano, Natural mummification of the human gut preserves bacteriophage DNA, FEMS Microbiol Lett. 363 (2016). https://doi.org/10.1093/femsle/fnv219.
 - [104] T.M. Santiago-Rodriguez, G. Fornaciari, S. Luciani, G.A. Toranzos, I. Marota, V. Giuffra, R.J. Cano, Gut Microbiome and Putative Resistome of Inca and Italian Nobility Mummies, Genes. 8 (2017) 310. https://doi.org/10.3390/genes8110310.
 - [105] W.-Q. Liu, J. Liu, J.-H. Zhang, X.-C. Long, J.-H. Lei, Y.-L. Li, Comparison of ancient and modern Clonorchis sinensis based on ITS1 and ITS2 sequences, Acta Tropica. 101 (2007) 91–94. https://doi.org/10.1016/j.actatropica.2006.08.010.
 - [106] H.D. Donoghue, O.Y.-C. Lee, D.E. Minnikin, G.S. Besra, J.H. Taylor, M. Spigelman, Tuberculosis in Dr Granville's mummy: a molecular re-examination of the earliest known Egyptian mummy to be scientifically examined and given a medical diagnosis, Proceedings of the Royal Society B: Biological Sciences. 277 (2010) 51–56. https://doi.org/10.1098/rspb.2009.1484.
 - [107] G.K. Bar-Gal, M.J. Kim, A. Klein, D.H. Shin, C.S. Oh, J.W. Kim, T.-H. Kim, S.B. Kim, P.R. Grant, O. Pappo, M. Spigelman, D. Shouval, Tracing hepatitis B virus to the 16th century in a Korean mummy, Hepatology. 56 (2012) 1671–1680. https://doi.org/10.1002/hep.25852.
- [108] F. Maixner, A. Thomma, G. Cipollini, S. Widder, T. Rattei, A. Zink, Metagenomic Analysis Reveals Presence of Treponema denticola in a Tissue Biopsy of the Iceman, PLoS One. 9 (2014). https://doi.org/10.1371/journal.pone.0099994.
- [109] M. Mackie, J. Hendy, A.D. Lowe, A. Sperduti, M. Holst, M.J. Collins, C.F. Speller, Preservation of the metaproteome: variability of protein preservation in ancient dental calculus, STAR: Science & Technology of Archaeological Research. 3 (2017) 58–70. https://doi.org/10.1080/20548923.2017.1361629.
- [110] K.A. Ziesemer, J. Ramos-Madrigal, A.E. Mann, B.W. Brandt, K. Sankaranarayanan,
 A.T. Ozga, M. Hoogland, C.A. Hofman, D.C. Salazar-García, B. Frohlich, G.R. Milner,
 A.C. Stone, M. Aldenderfer, C.M. Lewis, C.L. Hofman, C. Warinner, H. Schroeder,
 The efficacy of whole human genome capture on ancient dental calculus and dentin,
 American Journal of Physical Anthropology. 168 (2019) 496–509.
 https://doi.org/10.1002/ajpa.23763.
- 750 [111] A. Tett, K.D. Huang, F. Asnicar, H. Fehlner-Peach, E. Pasolli, N. Karcher, F. Armanini, P. Manghi, K. Bonham, M. Zolfo, F. De Filippis, C. Magnabosco, R. Bonneau, J.

Lusingu, J. Amuasi, K. Reinhard, T. Rattei, F. Boulund, L. Engstrand, A. Zink, M.C. Collado, D.R. Littman, D. Eibach, D. Ercolini, O. Rota-Stabelli, C. Huttenhower, F. Maixner, N. Segata, The Prevotella copri Complex Comprises Four Distinct Clades Underrepresented in Westernized Populations, Cell Host & Microbe. 26 (2019) 666-679.e7. https://doi.org/10.1016/j.chom.2019.08.018.

- [112] I.M. Velsko, J.A. Fellows Yates, F. Aron, R.W. Hagan, L.A.F. Frantz, L. Loe, J.B.R. Martinez, E. Chaves, C. Gosden, G. Larson, C. Warinner, Microbial differences between dental plaque and historic dental calculus are related to oral biofilm maturation stage, Microbiome. 7 (2019) 102. https://doi.org/10.1186/s40168-019-0717-3.
- [113] M. Borry, B. Cordova, A. Perri, M. Wibowo, T. Prasad Honap, J. Ko, J. Yu, K. Britton, L. Girdland-Flink, R.C. Power, I. Stuijts, D.C. Salazar-García, C. Hofman, R. Hagan, T. Samdapawindé Kagoné, N. Meda, H. Carabin, D. Jacobson, K. Reinhard, C. Lewis, A. Kostic, C. Jeong, A. Herbig, A. Hübner, C. Warinner, CoproID predicts the source of coprolites and paleofeces using microbiome composition and host DNA content, PeerJ. 8 (2020). https://doi.org/10.7717/peerj.9001.
- [114] D. Chessa, M. Murgia, E. Sias, M. Deligios, V. Mazzarello, M. Fiamma, D. Rovina, G. Carenti, G. Ganau, E. Pintore, M. Fiori, G.L. Kay, A. Ponzeletti, P. Cappuccinelli, D.J. Kelvin, J. Wain, S. Rubino, Metagenomics and microscope revealed T. trichiura and other intestinal parasites in a cesspit of an Italian nineteenth century aristocratic palace, Sci Rep. 10 (2020). https://doi.org/10.1038/s41598-020-69497-8.
- [115] J. Neukamm, S. Pfrengle, M. Molak, A. Seitz, M. Francken, P. Eppenberger, C. Avanzi, E. Reiter, C. Urban, B. Welte, P.W. Stockhammer, B. Teßmann, A. Herbig, K. Harvati, K. Nieselt, J. Krause, V.J. Schuenemann, 2000-year-old pathogen genomes reconstructed from metagenomic analysis of Egyptian mummified individuals, BMC Biology. 18 (2020) 108. https://doi.org/10.1186/s12915-020-00839-8.
- [116] A. Philips, I. Stolarek, L. Handschuh, K. Nowis, A. Juras, D. Trzciński, W. Nowaczewska, A. Wrzesińska, J. Potempa, M. Figlerowicz, Analysis of oral microbiome from fossil human remains revealed the significant differences in virulence factors of modern and ancient Tannerella forsythia, BMC Genomics. 21 (2020) 402. https://doi.org/10.1186/s12864-020-06810-9.
- [117] R.F. Rifkin, S. Vikram, J.-B. Ramond, A. Rey-Iglesia, T.B. Brand, G. Porraz, A. Val, G. Hall, S. Woodborne, M. Le Bailly, M. Potgieter, S.J. Underdown, J.E. Koopman, D.A. Cowan, Y. Van de Peer, E. Willerslev, A.J. Hansen, Multi-proxy analyses of a mid-15th century Middle Iron Age Bantu-speaker palaeo-faecal specimen elucidates the configuration of the 'ancestral' sub-Saharan African intestinal microbiome, Microbiome. 8 (2020) 62. https://doi.org/10.1186/s40168-020-00832-x.
- [118] D.K. Jacobson, T.P. Honap, C. Monroe, J. Lund, B.A. Houk, A.C. Novotny, C. Robin, E. Marini, C.M. Lewis, Functional diversity of microbial ecologies estimated from ancient human coprolites and dental calculus, Philosophical Transactions of the Royal Society B: Biological Sciences. 375 (2020) 20190586. https://doi.org/10.1098/rstb.2019.0586.
- [119] S. Sabin, H.-Y. Yeh, A. Pluskowski, C. Clamer, P.D. Mitchell, K.I. Bos, Estimating molecular preservation of the intestinal microbiome via metagenomic analyses of latrine sediments from two medieval cities, Philosophical Transactions of the Royal Society B: Biological Sciences. 375 (2020). https://doi.org/10.1098/rstb.2019.0576.
- [120] R. Eisenhofer, H. Kanzawa-Kiriyama, K. Shinoda, L.S. Weyrich, Investigating the demographic history of Japan using ancient oral microbiota, Philosophical Transactions of the Royal Society B: Biological Sciences. 375 (2020) 20190578. https://doi.org/10.1098/rstb.2019.0578.
- [121] A. Kazarina, E. Petersone-Gordina, J. Kimsis, J. Kuzmicka, P. Zayakin, Ž. Griškjans, G. Gerhards, R. Ranka, The Postmedieval Latvian Oral Microbiome in the Context of Modern Dental Calculus and Modern Dental Plaque Microbial Profiles, Genes. 12 (2021). https://doi.org/10.3390/genes12020309.

806 807 808	[122]	QGIS.org, QGIS Geographic Information System. QGIS Association. http://www.qgis.org., (2020). http://www.qgis.org.
809		

810	Funding
811	Not applicable.
812	
813	Author's contributions
814	RN: Investigation, Writing - Original Draft. MD: Writing - Review & Editing. RB:
815	Writing - Original Draft, Supervision, Review & Editing.
816	
817	Declaration of competing interest
818	The authors have no competing interests to declare.

Table 1. Scientific studies performed on ancient human digestive tract microbiota catalogued per archaeological site, dating, sample type, and method used for the analysis. Only studies written in English, published before March 2021, and on samples of human origin that targeted multiple species from the digestive tract microbiota of the host were considered.

Archaeological site	Present-day country	Dating	Sample type	Method	References
Cuzco	Peru	10th-11th century CE	Mummified colon	PCR-based techniques	[24]
Similaun mountain	Italy	3400-3100 BCE	Mummified stomach and colon	PCR-based techniques	[25]
Cuzco	Peru	980-1170 CE	coprolites	PCR-based techniques	[26]
Forni glacier (Alta Val di Pejo, northern Italy)	Italy	1918 CE	Mummified colon	PCR-based techniques	[27]
Similaun mountain	Italy	3400-3100 BCE	Mummified colon	PCR-based techniques	[27]
La Cueva de los Chiquitos Muertos", Rio Zape, Durango Durango	Mexico	600 - 700 CE	coprolites	Shotgun metagenomic	[28]
Caserones	Chile	400 CE	coprolites	16S rRNA Gene High-Throughput Sequencing	[33]
Hind's Cave	Texas	6000 BCE	coprolites	16S rRNA Gene High-Throughput Sequencing	[33]
La Cueva de los Chiquitos Muertos", Rio Zape, Durango Durango	Mexico	600 - 700 CE	coprolites	16S rRNA Gene High-Throughput Sequencing	[33]
Benzingerode-Heimburg	Germany	2150-1600 BCE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Dudka	Poland	5550-3450 BCE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Halberstadt-Sonntagsfeld	Germany	5400-4725 BCE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Northamptonshire	England	900-1150 CE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Quedlinburg XII	Germany	2450-2000 BCE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
York	England	1250-1350 CE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
York	England	1000-1600 CE	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Yorkshire	Engalnd	Bronze age	dental calculus	16S rRNA Gene High-Throughput Sequencing	[42]
Sorcé, island of Vieques	Puerto Rico	180 - 600 CE	coprolites	16s rRNA gene and ITS region PCR- based techniques	[74]
Tecla, Guayanilla	Puerto Rico	180 - 600 CE	coprolites	16s rRNA gene and ITS region PCR-	[74]
Namur	Belgium	14th-century	coprolites	based techniques shotgun metagenomic	[78]
Namur	Belgium	14th-century	coprolites	culture-base method	[77]
Namur	Belgium	14th-century	coprolites	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[77]
Sorcé, island of Vieques	Puerto Rico	5 to 1170 CE	coprolites	16S rRNA Gene High-Throughput Sequencing	[70]
Tecla, Guayanilla	Puerto Rico	5 to 1170 CE	coprolites	18S rRNA Gene High-Throughput Sequencing	[70]
Similaun mountain	Italy	3400-3100 BCE	gingival tissue biopsy	PCR-based techniques	[108]
Similaun mountain	Italy	3400-3100 BCE	Mouth swab	PCR-based techniques	[108]
St. Petri church and convent complex in Dalheim	Germany	950-1200 CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[38]
St. Petri church and convent complex in Dalheim	Germany	950-1200 CE	dental calculus	Shotgun metagenomic	[38]
Cuzco	Peru	11th century CE	paleofaeces	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[34]
Cuzco	Peru	11th century CE	mummified colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[34]
Anse à la Gourde	Guadalupe	975–1395 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]

				4CC aDNIA Consultable Throughout	
Camino del Molino, Caravaca de la Cruz, Murcia	Spain	2340-2920 BCE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Lavoutte	St. Lucia	990–1255 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Middenbeemster	Netherlands	1611–1866 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Rupert's Valley	St. Helena	1840–1872 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Samdzong	Nepal	400–650 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Tickhill, Yorkshire	UK	1450–1600 CE	Dental calculus	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[43]
Avosnes	France	1300 CE	dental calculus	PCR-based techniques	[44]
Douai	France	1710-1712 CE	dental calculus	PCR-based techniques	[44]
Forbach	France	1813 CE	dental calculus	PCR-based techniques	[44]
Les Fedons	France	1590 CE	dental calculus	PCR-based techniques	[44]
Martigues	France	1720-1721 CE	dental calculus	PCR-based techniques	[44]
Saint-Mitre-les-Remparts	France	1500-1700 CE	dental calculus	PCR-based techniques	[44]
Cuzco	Peru	980-1170 CE	Mummified descending colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[103]
Cuzco	Peru	14-15th century CE	Mummified abdominal viscera	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[103]
Cuzco	Peru	1410–1530 CE	Mummified transverse colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[103]
Cuzco	Peru	980-1170 CE	Mummified descending colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[101]
Cuzco	Peru	14-15th century CE	Mummified abdominal viscera	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[101]
Cuzco	Peru	1410–1530 CE	Mummified transverse colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[101]
Similaun mountain	Italy	3400-3100 BCE	Mummified distal gut	shotgun metagenomic	[102]
Leicester, UK (NGR SK 586038), Oxford Street	UK	2nd–5th century CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[109]
Rome, Isola Sacra	Italy	1st–3rd century CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[109]
York, UK (NGR: SE 59324510 & SE 59285095), Driffield Terrace	UK	2nd–4th century CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[109]
Sorcé, island of Vieques	Puerto Rico	10 - 385 CE	Dental calculus	16S rRNA gene high-throughput	[50]
Cuzco	Peru	980-1170 CE	Mummified descending colon	sequencing 16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[104]
Cuzco	Peru	14-15th century CE	Mummified abdominal viscera	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[104]
Cuzco	Peru	1410–1530 CE	Mummified transverse colon	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[104]
Naples, Basilica of St. Domenico Maggiore	Italy	15-16th century CE	Mummified colon and abdominal viscera	16S rRNA Gene High-Throughput Sequencing and Shotgun metagenomic	[104]
Radcliffe Infirmary Burial Ground collection	England	1770-1855 CE	Dental calculus	mass spectrometry (MS)-based metabolites characterization	[46]
Tjærby, Jutland, Denmark	Denmark	1100-1450 CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[55]
Anse à la Gourde	Guadalupe	975-1395 CE	dental calculus	shotgun metagenomic	[39]
Arbulag Soum, Khövsgöl,	Mongolia	930-1650 BCE	dental calculus	shotgun metagenomic	[39]
Camino del Molino, Caravaca de la Cruz, Murcia	Spain	2340-2920 BCE	dental calculus	shotgun metagenomic	[39]
Kilteasheen	Ireland	600-1300 CE	dental calculus	shotgun metagenomic	[39]
Middenbeemster	Netherlands	1611-1866 CE	dental calculus	shotgun metagenomic	[39]
Norris Farms	Illinois (USA)	1300 CE	dental calculus	shotgun metagenomic	[39]
Samdzong	Nepal	400-650 CE	dental calculus	shotgun metagenomic	[39]
	ера.	.00 050 02			[00]

Anse à la Gourde	Guadalupe	975-1395 CE	dental calculus	shotgun metagenomic	[110]
Arbulag Soum, Khövsgöl,	Mongolia	930-1650 BCE	dental calculus	shotgun metagenomic	[110]
Camino del Molino, Caravaca de la Cruz, Murcia	Spain	2340-2920 BCE	dental calculus	shotgun metagenomic	[110]
Kilteasheen	Ireland	600-1300 CE	dental calculus	shotgun metagenomic	[110]
Middenbeemster	Netherlands	1611-1866 CE	dental calculus	shotgun metagenomic	[110]
Norris Farms	Illinois (USA)	1300 CE	dental calculus	shotgun metagenomic	[110]
Samdzong	Nepal	400-650 CE	dental calculus	shotgun metagenomic	[110]
site of Syltholm on the island of Lolland	Denmark	3700 BCE	chewed birch pitch	shotgun metagenomic	[63]
San Pietro di Pozzeveri	Italy	11th to 19th century CE	Dental calculus	16S rRNA gene High-Throughput	[51]
La Cueva de los Chiquitos Muertos'', Rio Zape, Durango Durango	Mexico	700 CE	coprolites	Sequencing Shotgun metagenomic	[111]
Similaun mountain	Italy	3400-3100 BCE	intestinal mummified tissue	Shotgun metagenomic	[111]
Radcliffe Infirmary Burial Ground collection	England	1770–1855 CE	dental calculus	mass spectrometry (MS)-based proteomics profiling	[112]
Radcliffe Infirmary Burial Ground	England	1770–1855 CE	dental calculus	shotgun metagenomic	[112]
collection La Cueva de los Chiquitos Muertos", Rio Zape, Durango	Mexico	700 CE	coprolites	Shotgun metagenomic	[113]
Durango Surrey	UK	Post medieval	coprolites	Shotgun metagenomic	[113]
palazzo ducale Sassari	Italy	1800-1850 CE	cesspit sediment	16S and 18S rRNA Gene High- Throughput Sequencing and Shotgun metagenomic	[114]
La Cueva de los Chiquitos Muertos", Rio Zape, Durango Durango	Mexico	700 CE	coprolites	Shotgun metagenomic	[36]
Abusir	Egypt	340-395 CE	Dental calculus	Shotgun metagenomic	[115]
Abusir	Egypt	138-200 CE	Dental calculus	Shotgun metagenomic	[115]
Abusir	Egypt	2196-2045 BCE	Dental calculus	Shotgun metagenomic	[115]
Abusir	Egypt	109-40 BCE	Dental calculus	Shotgun metagenomic	[115]
Abusir	Egypt	369-211 BCE	Dental calculus	Shotgun metagenomic	[115]
Brzeg	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Dziekanowice	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Gniezno	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Gołuń	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Końskie	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Kowalewko	Poland	100-400 CE	teeth	shotgun metagenomic	[116]
Ląd	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Łęgowo	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Markowice	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Masłomęcz	Poland	100-400 CE	teeth	shotgun metagenomic	[116]
Niemcza	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Obłaczkowo	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Opole	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Ostrów Lednicki	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Płońsk	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Poznań–Śródka	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Sowinki	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Warszawa	Poland	900-1400 CE	teeth	shotgun metagenomic	[116]
Bushman Rock Shelter (BRS) in Limpopo Province	South Africa	1460 CE	paleo-faecal sample	Shotgun metagenomic	[117]
western Belize: Chan Chich	Belize	713-885 CE	Dental calculus	Shotgun metagenomic	[118]
western Belize: Chan	Belize	170 BCE-50 CE	Dental calculus	Shotgun metagenomic	[118]
	-				,

Lu Maccioni (Alghero)	Italy	1126-825 BCE	Dental calculus	Shotgun metagenomic	[118]
Capo Pecora (Arbus)	Italy	1384-936 BCE	Dental calculus	Shotgun metagenomic	[118]
Perdalba (Sardara)	Italy	1400-850 BCE	Dental calculus	Shotgun metagenomic	[118]
Riga	Latvia	1356 CE	Latrine sediment	Shotgun metagenomic	[119]
Jerusalem	Israel/State of Palestine	1400-1500 CE	Latrine sediment	Shotgun metagenomic	[119]
Ebishima shell mound in Iwate prefecture	Japan	1000-400 BCE	Dental calculus	Shotgun metagenomic	[120]
Ikawazu, Aichi prefecture,	Japan	1000-400 BCE	Dental calculus	Shotgun metagenomic	[120]
Miyano, Iwate prefecture	Japan	1000-400 BCE	Dental calculus	Shotgun metagenomic	[120]
Ikenohata-Shichikencho site, Taito-ku, Tokyo	Japan	1000-400 BCE	Dental calculus	Shotgun metagenomic	[120]
Riga	Latvia	1500-1700 CE	Dental calculus	Shotgun metagenomic	[121]
Cesis	Latvia	1600-1700 CE	Dental calculus	Shotgun metagenomic	[121]
Kuldiga	Latvia	1500-1700 CE	Dental calculus	Shotgun metagenomic	[121]
Jelgava	Latvia	1500-1700 CE	Dental calculus	Shotgun metagenomic	[121]

Figure 1. Geographical localization of ancient human remains analyzed for microbiota studies in the World. Colour indicate the dating of the analyzed samples from 6000 BCE to 1918 CE. Oral microbiota and gut microbiota studies are indicated with two symbols representing a tooth and the gut respectively. The corresponding references are listed in Table 1. Map generated with QGIS software [122]. In the upper part of the figure, 6 pictures representative of the original samples used to study the digestive tract microbiota [39,63,70,109,113,116].

