



**HAL**  
open science

## Disinfection of gloved hands during routine care

A. Vogel, P. Brouqui, S. Boudjema

► **To cite this version:**

A. Vogel, P. Brouqui, S. Boudjema. Disinfection of gloved hands during routine care. *New Microbes and New Infections*, 2021, 41, pp.100855. <10.1016/j.nmni.2021.100855>. <hal-03334143>

**HAL Id: hal-03334143**

**<https://amu.hal.science/hal-03334143v1>**

Submitted on 6 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC-ND 4.0 - Attribution - Non-commercial use - No Derivative Works - International License

# Disinfection of gloved hands during routine care

A. Vogel, P. Brouqui and S. Boudjema

Aix-Marseille Université, IRD, MEPHI, IHU-Méditerranée Infection, Marseille, France

## Abstract

Healthcare-associated infections are a major issue in public health. After several decades of hand hygiene programmes, it is time to admit that we have failed to achieve our goal. One of the reasons is the overuse of gloves, which is in part justified by the nature of the nursing care. Several experimental studies supported the effectiveness of disinfecting gloves, but evidence for routine feasibility and effectiveness is lacking. The aim of this study was to evaluate the effectiveness of alcohol-based disinfection of gloves during nursing care. Swabs were taken from the most contaminated part of both hands at different times: T0 (before wearing gloves), T10 (after 10 minutes of nursing care) from both gloves, T10A (just after the gloves were disinfected for 30 seconds with bedside disposable hydroalcoholic solution) and when possible at T20 and T20A. After 10 minutes of nursing care, gloves were contaminated in 72.5% of cases. After alcohol-based disinfection, gloves became free of microorganism in 79.3% of cases. Alcohol-based disinfection of gloves during routine care is effective and appears to be a reasonable alternative to current recommendations.

© 2021 Published by Elsevier Ltd.

**Keywords:** Disinfection, gloves, hand hygiene, nosocomial infection, nursing research, pathogen

**Original Submission:** 13 October 2020; **Revised Submission:** 4 February 2021; **Accepted:** 22 February 2021

**Article published online:** 3 March 2021

**Corresponding author:** S. Boudjema, IHU Méditerranée Infection  
19-21 Boulevard Jean Moulin, 13005 Marseille (France)/ Aix Marseille  
University, Chemin des Bourrelly 13015 Marseille, France.  
**E-mail:** [sophia.boudjema@univ-amu.fr](mailto:sophia.boudjema@univ-amu.fr)


## Introduction

Hand hygiene, particularly the use of alcohol-based hand rubs, is currently recognized as the main barrier against cross-transmission of infections. However, despite worldwide investment in hand hygiene promotion, hospital-acquired infections have not been significantly lowered. One of the reasons of the still-insufficient practice of alcohol-based hand rub is the overuse of gloves, which is derived from the healthcare behavioral adaptation of institutional protocols to routine care. A video analysis during routine care pointed out the complex behaviors of healthcare workers (HCWs) regarding hand hygiene and wearing gloves. HCWs consider hand disinfection and wearing gloves to be two different practices [1]. The World Health Organization recommends that

gloves be changed between caring for patients and between providing different types of care [2]. However, these recommendations are not easily applicable in routine care; indeed, Chau *et al.* [3] reported that gloves were not changed in 75% of required situations.

Gloves are required when it can be reasonably anticipated that contact with blood or other potentially infectious materials could occur. These situations cannot be anticipated, which results in disposable gloves being unjustifiably worn and hand disinfection being compromised [1]. Contaminated gloves have been shown to support cross-transmission of methicillin-resistant *Staphylococcus aureus* [4]. Disinfecting gloves has been reported to significantly decrease catheter contamination and catheter-associated bloodstream infections [5]. Similarly, disinfecting gloves has reduced the incidence of methicillin-resistant *S. aureus* [6]. As a result, in their excellent review, Kampf and Lemmen [7] emphasized promoting disinfecting gloves with an alcohol-based solution, which might result in a significant reduction of cross-transmission. These authors concluded that disinfecting gloves is as efficient as hand disinfection for alcohol-sensitive microorganisms and suggested that this practice be promoted.





**FIG. 2.** Microorganisms on nurses' hands and gloves. Samples were collected 40 times: time T0 (before wearing gloves), T10 (after 10 minutes of nursing care) and T10A (just after gloves were disinfected for 30 seconds with bedside disposable hydroalcoholic solution).

pathogenic agent such as *Acinetobacter lwoffii*, *Pseudomonas putida* or *Proteus mirabilis*, meaning that HCWs' hands were contaminated by pathogenic bacteria before providing care to their patients. In four situations, the same bacteria that were recovered on the nurses' hands were also recovered at T10 on the gloves, suggesting possible cross-contamination of the gloves by the caregivers' hands (two *Staphylococcus pasteurii*; one each *Staphylococcus epidermidis* and *Micrococcus luteus*).

At T10, among the 18 nursing care visits for which both hands were sterile at T0, 17 (94%) of 18 had at least one contaminated glove, and five (29.4%) of 17 were contaminated by pathogenic *Enterococcus faecium* ( $n = 3$ ), *P. mirabilis* ( $n = 1$ ) and *A. lwoffii* ( $n = 1$ ), suggesting that during the first 10 minutes of nursing care, a third of the HCWs contaminated at least one of their gloves with pathogenic bacteria, mostly enterobacteria, from the digestive tract. Among 12 nursing care visits for which a hand was contaminated at T0, all 12 were contaminated at T10.

Finally, in the first 10 minutes of nursing care, 29 (72.5%) of 40 HCWs had at least one glove contaminated. Disinfection of the gloves with a hydroalcoholic solution for 30 seconds sterilized both gloves in 79.3% (23/29) of cases. No difference was found between nurses and assistant nurses in the ratio of glove contamination or in the efficacy of alcohol-based rubs.

## Discussion

In this study, 70% (28/40) of HCWs had both hands free of microorganism before coming into contact with patients, which

likely reflects the specificity of this ward in which hand hygiene is regularly promoted [9]. Our average hand hygiene compliance rate, as measured by our automated surveillance system (last 6 months), is 35%; one of the explanations for this result is the overuse and misuse of gloves, as revealed by a video survey. Gloves should be used only in case of contact with dirty matter. This situation during nursing should be anticipated; nurses who prefer to wear gloves in any circumstances should be challenged [1].

It is unrealistic to believe that campaigns promoting better use of gloves will be more effective than promoting hand hygiene. Some authors, among them nurse practitioners, suggested disinfecting gloves as an alternative, and several experimental studies have supported the effectiveness of disinfecting gloves [10,11]. We showed that glove disinfection with hydroalcoholic solution is efficient in 79.3% of cases. Although the disinfection rate in experimental studies is higher than that reported here, the experimental context, such as the bacterial loads and disinfection protocols used in these studies, is not transferrable to clinical practice and does not take into consideration the quality of hand-rub practice during care [7,12–15]. Our data only reflect practice in our ward; larger observational studies are needed to confirm the feasibility of such an alternative practice.

Compliance with hand hygiene was reported to be impaired by wearing gloves [16]. It has been recently reported that disinfecting gloved hands significantly improved compliance with hand hygiene, from 31% to 65% [17]. In addition to the growing body of literature on the effectiveness of glove disinfection with

hydroalcoholic solution, our data convinced us, as well as our colleagues [7,17], that regular disinfection of gloves during care visits might be an alternative to avoid glove misuse and will be more easily accepted by HCWs.

## Conclusions

In routine care, disinfection with hydroalcoholic solution is a feasible and efficient way to decontaminate gloves, which suggests that cross-transmission to the environment and patients can be lowered by using this method.

## Conflict of interest

None declared.

## Acknowledgements

The authors thank the HCWs of the medical unit for their investment and participation in this study. English-language editing services were provided by [cookietrad@gmail.com](mailto:cookietrad@gmail.com). Funded in part by ANR-15-CE36-0004-01 entitled MediHand-Trace (MHT) and Agence Nationale de la Recherche (ANR), IHU Mediterranée Infection 10-IAHU-03.

## References

- [1] Boudjema S, Tarantini C, Peretti-Watel P, Brouqui P. Merging video coaching and an anthropologic approach to understand health care provider behavior toward hand hygiene protocols. *Am J Infect Contr* 2017;45:487–91.
- [2] World Health Organization (WHO). WHO guidelines on hand hygiene in health care. Available at: 15 January 2009. <https://www.who.int/publications/i/item/9789241597906>.
- [3] Chau JP, Thompson DR, Twinn S, Lee DT, Pang SW. An evaluation of hospital hand hygiene practice and glove use in Hong Kong. *J Clin Nurs* 2011;20:1319–28.
- [4] McBryde ES, Bradley LC, Whitby M, McElwain DL. An investigation of contact transmission of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 2004;58:104–8.
- [5] Kocent H, Corke C, Alajeel A, Graves S. Washing of gloved hands in antiseptic solution prior to central venous line insertion reduces contamination. *Anaesth Intensive Care* 2002;30:338–40.
- [6] Matsumoto K, Shigemi A, Yaji K, Shimodozono Y, Takeda Y, Ikawa K, et al. Reduction in the incidence of MRSA with use of alcohol-based hand rub solutions and gloves. *J Infect Chemother* 2012;18:269–71.
- [7] Kampf G, Lemmen S. Disinfection of gloved hands for multiple activities with indicated glove use on the same patient. *J Hosp Infect* 2017;97:3–10.
- [8] Seng P, Drancourt M, Gouriet F, La SB, Fournier PE, Rolain JM, et al. Ongoing revolution in bacteriology: routine identification of bacteria by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Clin Infect Dis* 2009;49:543–51.
- [9] Dufour JC, Reynier P, Boudjema S, Soto AA, Giorgi R, Brouqui P. Evaluation of hand hygiene compliance and associated factors with a radio-frequency-identification–based real-time continuous automated monitoring system. *J Hosp Infect* 2017;95:344–51.
- [10] Assadian O, Humphreys PN, Ousey KJ. Disinfection of artificially contaminated gloved hands reduces transmission of *Staphylococcus epidermidis* to catheter valves. *J Hosp Infect* 2018;100:e57–9.
- [11] Newsom SW, Rowland C. Application of the hygienic hand-disinfection test to the gloved hand. *J Hosp Infect* 1989;14:245–7.
- [12] Pitten FA, Muller P, Heeg P, Kramer A. [The efficacy of repeated disinfection of disposable gloves during usage]. *Zentralbl Hyg Umweltmed* 1999;201:555–62.
- [13] Janota J, Šebková S, Višňovská M, Kudláčková J, Hamplová D, Zach J. Hand hygiene with alcohol hand rub and gloves reduces the incidence of late onset sepsis in preterm neonates. *Acta Paediatr* 2014;103:1053–6.
- [14] Ng PC, Wong HL, Lyon DJ, So KW, Liu F, Lam RKY, et al. Combined use of alcohol hand rub and gloves reduces the incidence of late onset infection in very low birthweight infants. *Arch Dis Child Fetal Neonatal* 2004;89:F336–40.
- [15] Scheithauer S, Häfner H, Seef R, Seef S, Hilgers RD, Lemmen S. Disinfection of gloves: feasible, but pay attention to the disinfectant/glove combination. *J Hosp Infect* 2016;94:268–2672.1.
- [16] Florea O, Dufour J-C, Magnin C, Brouqui P, Boudjema S. Improving health care workers compliance with traceability by recording the nursing process at the point of care using a personal digital assistant with a barcode. *J Nurs Care* 2020;9:500.
- [17] Fehling P, Hasenkamp J, Unkel S, Thalmann I, Hornig S, Trümper L, et al. Effect of gloved hand disinfection on hand hygiene before infection-prone procedures on a stem cell ward. *J Hosp Infect* 2019;103:321–7.