



The *Azospirillum brasilense* Type VI secretion system promotes cell aggregation, biocontrol protection against phytopathogens and attachment to the microalgae *Chlorella sorokiniana*

Fabricio Cassan, Anahí Coniglio, Edgar Amavizca, Guillermo Maroniche, E. Cascales, Yoav Bashan, Luz De-bashan

► **To cite this version:**

Fabricio Cassan, Anahí Coniglio, Edgar Amavizca, Guillermo Maroniche, E. Cascales, et al.. The *Azospirillum brasilense* Type VI secretion system promotes cell aggregation, biocontrol protection against phytopathogens and attachment to the microalgae *Chlorella sorokiniana*. *Environmental Microbiology*, 2021, 10.1111/1462-2920.15749 . hal-03336446

HAL Id: hal-03336446

<https://amu.hal.science/hal-03336446>

Submitted on 5 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Submitted to Environmental Microbiology

SI: Multifaceted Plant-Microbe Interactions

The *Azospirillum brasilense* Type VI secretion system promotes cell aggregation, biocontrol protection against phytopathogens and attachment to the microalgae *Chlorella sorokiniana*

Fabricio D. Cassan¹, Anahí Coniglio^{1,7}, Edgar Amavizca^{3,7}, Guillermo Maroniche⁵, Eric Cascales⁶, Yoav Bashan^{2,3,4†}, Luz E. de-Bashan^{2,3,4,*}

¹Laboratorio de Fisiología Vegetal y de la interacción Planta-Microorganismo. Instituto de Investigaciones Agrobiotecnológicas (INIAB). Universidad Nacional de Río Cuarto. Ruta 36, Km 601, Río Cuarto, Córdoba Argentina

²The Bashan Institute of Science, 1730 Post Oak Court, Auburn AL 36830, USA

³Environmental Microbiology Group, Northwestern Center for Biological Research (CIBNOR), Calle. IPN 195, La Paz, B.C.S. 23096, Mexico

⁴Dept. of Entomology and Plant Pathology, 301 Funchess Hall, Auburn University, Auburn, AL 36849, USA

⁵Facultad de Ciencias Agrarias, Universidad Nacional de Mar del Plata, Buenos Aires, Argentina

⁶Laboratoire d'Ingénierie des Systèmes Macromoléculaires, Institut de Microbiologie, Bioénergies et Biotechnologie, Aix-Marseille Université – CNRS UMR7255, 31 Chemin Joseph Aiguier, CS7071, 13402 Marseille Cedex 09, France

Anahí Coniglio, and Edgar Amavizca contributed equally to this study.

Running title: T6SS in microalgae-bacteria association

*Author for correspondence: email: luz@bashanfoundation.org; legonzal04@cibnor.mx; Tel: +1-256-307-1963 (USA); +52-612-123-8484, ext. 3419, 3420 (Mexico)

Number of Figures: 7

Number of Tables: 1

Supplementary material: 2 Figure, 1 Table

1
2
3
4
5

† Dedication: This study is dedicated to the memory of Dr. Yoav Bashan, leading figure in the field of Plant Growth-Promoting Bacteria (PGPB) for environmental purposes, and founder of the Bashan Institute of Science, USA. Prof. Bashan passed away during the edition of the manuscript.

Summary

The plant-growth promoting bacterium *Azospirillum brasilense* is able to associate with the microalgae *Chlorella sorokiniana*. Attachment of *A. brasilense* increases the metabolic performances of the microalgae. Recent genome analyses have revealed that the *A. brasilense* Az39 genome contains two complete sets of genes encoding Type VI secretion systems (T6SS), including the T6SS1 that is induced by the indole-3-acetic acid (IAA) phytohormone. The T6SS is a multiprotein machine, widespread in gram-negative bacteria, that delivers protein effectors in both prokaryotic and eukaryotic cells. Here we show that the *A. brasilense* T6SS is required for *Chlorella-Azospirillum* synthetic mutualism. Our data demonstrate that the T6SS is an important determinant to promote production of carbohydrates and photosynthetic pigments by the microalgae. We further show that this is likely due to the role of the T6SS during the attachment stage and for the production of IAA phytohormones. Finally, we demonstrate that the *A. brasilense* T6SS provides antagonistic activities against a number of plant pathogens such as *Agrobacterium*, *Pectobacterium*, *Dickeya* and *Ralstonia* species *in vitro*, suggesting that, in addition to promoting growth, *A. brasilense* might confer T6SS-dependent bio-control protection to microalgae and plants against bacterial pathogens.

Significance

Azospirillum brasilense, a plant growth promoting bacterium, increases the metabolic performances of plants and microalgae. Here we show that the *A. brasilense* type VI secretion system (T6SS) is an important determinant of *Azospirillum* attachment to the microalgae *Chlorella sorokiniana*, and hence that a mutant defective in T6SS decreases metabolite production in the microalga. In addition, we show that the T6SS confers antibacterial activity against competitor

1 bacterial cells, including phytopathogens *in vitro*, suggesting that the T6SS is also an important
2 determinant for bioprotection. This study therefore demonstrates that a secretion system, involved
3 in the secretion of bacterial effectors and toxins, participates in processes that are beneficial to
4 plant and microalgae cells.

5 **Keywords:** Attachment; aggregation; Plant Growth Promoting Bacteria (PGPB); *Azospirillum*;
6 bacteria-microalgae interaction; *Chlorella*; Type VI secretion system (T6SS)

7

1 **Introduction**

2 *Azospirillum* spp. is one of the best-characterized genera of plant growth-promoting bacteria
3 (PGPB) and can colonize more than 130 plant species in 37 families (Pereg *et al.*, 2016). The
4 association of *Azospirillum* spp. with these plants significantly improves growth, development,
5 and in many cases, yield under field conditions (Cassan *et al.*, 2020). There is no single mechanism
6 involved in promoting plant growth with *Azospirillum*, but rather a combination of mechanisms in
7 different cases of inoculation. These mechanisms work together or in tandem, and the phenomenon
8 is commonly known as “multiple mechanism theory” (Bashan and de-Bashan, 2010). The
9 production of indole-3-acetic acid (IAA) by *Azospirillum* spp. is one of the main mechanisms
10 proposed for the effect of this bacterial species on plant growth (Bashan and de-Bashan, 2010).
11 The *A. brasilense* Az39 used in this study was isolated in 1982 from surface-sterilized wheat
12 seedlings in Marcos Juarez, Argentina. This strain was selected for inoculant formulation, based
13 on its ability to increase crop yields of maize and wheat under agronomic conditions (Díaz-Zorita
14 and Fernández-Canigia, 2009). Currently, *A. brasilense* Az39 is one of the most used strains for
15 crops such as maize and sorghum in Argentina and South America (Cassan *et al.*, 2020).

16 For decades, *Azospirillum* spp. has been known to associate to roots of higher plants
17 (Bashan and de-Bashan, 2010), such as cereals, tomato, pepper, cotton, and soybean, and with the
18 microalgae *Chlorella sorokiniana* (Levanony *et al.*, 1989; Bashan *et al.*, 1991; Assmus *et al.*,
19 1995). The first step of the association corresponds to attachment to the root, a critical step because
20 all azospirilla are highly motile bacteria *in vitro* (Bashan and Holguin, 1994; Alexandre *et al.*,
21 2000), in soil (Bashan and Levanony, 1987), and in water (Puente *et al.*, 1999). In addition to
22 motility, root colonization also requires the production of surface polysaccharides (Jofre *et al.*,
23 2004). After attachment, azospirilla colonize the roots. Host colonization usually requires the

1 delivery of effector proteins by dedicated secretion systems. Up to eleven secretion systems (Types
2 I–XI) have been described so far in gram-negative bacteria (Costa *et al.*, 2015). Genome analyses
3 show that *A. brasilense* strains are lacking Type III and Type IV secretion systems.
4 Transcriptomics and bioinformatics analyses show that *A. brasilense* strains Sp245 and Az39
5 contain gene clusters encoding putative Type VI secretion systems (T6SSs) (Van Puyvelde *et al.*,
6 2011; Rivera *et al.*, 2014; Fig. 1; Table S1). The T6SS is a widespread, multi-protein apparatus,
7 found in gram-negative bacteria (Bingle *et al.*, 2008; Cascales, 2008). The T6SS is not only
8 distributed in human, animal, and plant pathogenic bacteria where it is involved in virulence and
9 pathogenesis, but also found in commensal and symbiotic PGPB (Bingle *et al.*, 2008; Russell *et*
10 *al.*, 2014). In these bacteria, the T6SS participates to intermicrobial rivalry in bacterial and fungal
11 competition, biofilm formation, and establishment of symbiosis (Jiménez-Guerrero *et al.*, 2013;
12 Durand *et al.*, 2014; Russell *et al.*, 2014; Ryu, 2015; Bernal *et al.*, 2017; Trunk *et al.*, 2019;
13 Gallegos-Monterossa and Coulthurst, 2021). At the molecular level, the T6SS acts like a cellular
14 “biological syringe”: it assembles a bacteriophage tail-like structure anchored to the cell envelope
15 by a membrane complex (Zoued *et al.*, 2014; Coulthurst, 2019; Cherrak *et al.*, 2019). The tail-like
16 structure consists of an inner tube made of hexamers of the Hcp protein capped by a puncturing
17 VgrG-PAAR complex and wrapped in the TssBC contractile sheath (Zoued *et al.*, 2014; Cherrak
18 *et al.*, 2019; Basler, 2015). Effector toxins are loaded onto the VgrG or PAAR proteins or within
19 the Hcp tube (Hernandez *et al.*, 2020; Jurenas and Journet, 2021). Contraction of the sheath propels
20 the inner tube towards the target cell (Basler, 2015; Brackmann *et al.*, 2017). Penetration of the
21 tube into the target cell allows delivery of the effectors (Hernandez *et al.*, 2020; Jurenas and
22 Journet, 2021). While the mechanisms of action and the role of T6SS in animal and plant pathogens
23 have been well described in recent years, it has been less studied in plant symbiotic and associative

bacteria (Jiménez-Guerrero *et al.*, 2013; Ryu, 2015; Wu *et al.*, 2018; Wu *et al.*, 2019; Bernal *et al.*, 2018; Borrero de Acuña and Bernal, 2021). In *Rhizobium leguminosarum*, the T6SS affects the attachment, nodulation, and symbiosis processes (Roest *et al.*, 1997; Bladergroen *et al.*, 2003). In *Pseudomonas fluorescens*, the T6SS is involved in bacterial competition, motility, biofilm formation, and is critical to persist in the rhizosphere microbiome (Decoin *et al.*, 2014; 2015; Gallique *et al.*, 2017; Durán *et al.*, 2021). *P. putida* produces three T6SSs, including K1-T6SS that confers antibacterial activity against numerous bacterial phytopathogens, and hence can be used as a biocontrol agent against plant pathogens (Bernal *et al.*, 2017). Finally, *Agrobacterium tumefaciens* was shown to deploy a T6SS to destroy competitors and to increase colonization *in planta* (Ma *et al.*, 2014; Wu *et al.*, 2018; Wu *et al.*, 2019). *A. brasilense* Az39 contains two gene clusters encoding T6SSs. A transcriptomic study of *A. brasilense* Sp245 showed that the T6SS1 gene cluster is induced by indole-3-acetic acid (IAA) (Van Puyvelde *et al.*, 2011). The activation of T6SS1 by a phytohormone in *A. brasilense* prompted us to investigate the role of this T6SS in interbacterial competition and its interaction with microalgal cells.

The freshwater unicellular microalgae *Chlorella* spp. can associate with several bacterial species (Imase *et al.*, 2008), including *Azospirillum* spp. (Hernandez *et al.*, 2009; de-Bashan *et al.*, 2015). The *Chlorella*-*Azospirillum* association is an established model system of synthetic mutualism to study cellular mechanisms during plant-bacteria interaction (de-Bashan and Bashan, 2008; de-Bashan *et al.*, 2016). This association significantly promotes the production of carbohydrates, fatty acids, lipids, and pigments of the microalgae, both autotrophically and heterotrophically (de-Bashan *et al.*, 2002, 2015; Choix *et al.*, 2012a, b; Leyva *et al.*, 2014).

Previously, we tested the attachment and mutual aggregation between the microalga *Chlorella sorokiniana* and *Azospirillum brasilense* immobilized in alginate beads to facilitate the

initial stage of formation of this synthetic association (de-Bashan and Bashan, 2008; de-Bashan *et al.*, 2015; de-Bashan *et al.*, 2016). Attachment of *Azospirillum brasilense* to *Chlorella* spp. cells involve fibrillar connections (Lebsky *et al.*, 2001; de-Bashan *et al.*, 2011), similar to what was observed during the attachment of *Azospirillum* spp. to cells of higher plants (Bashan *et al.*, 1986; Bashan *et al.*, 1991). However, we currently lack molecular details on how *Azospirillum* spp. efficiently colonizes microalgae and plants.

In this study, we show that the *A. brasilense* T6SS1 has a positive effect on growth of the microalgae *Chlorella*. We further demonstrate that a *A. brasilense* strain defective for T6SS1 presents a reduced enhancement of the production of microalgal metabolites such as lipids, carbohydrates, and photosynthetic pigments compared to wild-type *A. brasilense*. Finally, we demonstrate that the *Azospirillum* T6SS provides anti-bacterial activity against *E. coli* and several plant pathogens, including *Agrobacterium tumefaciens*, *Pectobacterium carotovorum* and *Dickeya dadantii*. These data support the idea that the T6SS is required for the establishment of an efficient synthetic mutualism between *Azospirillum* and *Chlorella* and confers protection of the microalgae - and potentially of plants - against pathogens.

Results

Construction of the A. brasilense Az39Δhcp-E mutant

The Hcp protein is an essential component of the T6SS and its encoding gene is present as a single copy in the T6SS1 gene cluster of *A. brasilense* Az39. It is located upstream of the *tssE* gene, which encodes an important component of the T6SS assembly baseplate (Brunet *et al.*, 2015) (Fig. 1). Both genes are in tandem in chromid 1 in *A. brasilense* Az39 (GenBank, CP007794.1, AbAz39_p1, WP_051658325.1) and share high identity to those encoded within the genomes of

1 *A. brasilense* FP2 (99%), Sp245 (98%), Sp7 (99%), and *Azospirillum* sp. B510 (64%). We used
2 targeted mutagenesis to construct an isogenic mutant of Az39. For this, a genomic fragment
3 overlapping with the *hcp* and *tssE* genes was replaced with a gentamycin-resistant cassette by
4 homologous recombination (Fig. S1). The genomic structure of the wild-type strain and its
5 isogenic $\Delta hcp-tssE$ ($\Delta hcp-E$) mutant was confirmed by PCR using several combinations of primers
6 (Fig. S1).

7
8 *Phenotypic characterization of the A. brasilense Az39 $\Delta hcp-E$ strain for cell growth, IAA*
9 *production, motility, cell aggregation and biofilm formation*

10 To characterize the Az39 $\Delta hcp-E$ mutant strain, we measured its growth rate in rich and minimum
11 media, its production of indole-3 acetic acid, and its capacity to swim, swarm and to form biofilm
12 compared to the wild-type strain.

13 *Growth rate* - The growth rate and the number of colony-forming units (CFU) of the *A. brasilense*
14 wild-type strain and its *hcp-tssE* mutant were compared in two different media. In both rich (LB)
15 and minimal (MMAB) media, the two strains exhibited similar growth patterns for 80 hours (Fig.
16 S2). Wild-type *A. brasilense* Az39 had the highest number of living cells (4.63×10^9 CFU·mL⁻¹ in
17 LB medium, and 1.85×10^9 CFU·mL⁻¹ in MMAB) during the exponential growth phase (24 h after
18 inoculation) and fewer cells (0.75×10^9 CFU·mL⁻¹ in LB; 0.75×10^9 CFU·mL⁻¹ in MMAB) during
19 the stationary growth phase (>36 h after inoculation) (Fig S2). The *A. brasilense* Az39 $\Delta hcp-E$
20 mutant strain exhibited a similar pattern, with a higher number of cells during the exponential
21 growth phase (2×10^9 CFU·mL⁻¹ in LB; 1.6×10^9 CFU·mL⁻¹ in MMAB) and fewer living cells
22 during the stationary growth phase (1.5×10^9 CFU·mL⁻¹ in LB; 1.06×10^9 CFU·mL⁻¹) (Fig. S2).

1 *Indole-3-acetic acid (IAA) production* - In minimum medium, the mutant strain produced
2 significantly less IAA during exponential growth compared to wild-type Az39, with a maximum
3 difference at 24 h of culture (Az39 Δ hcp-E: $56.77 \pm 2.59 \mu\text{g}\cdot\text{mL}^{-1}$; Az39: $79.98 \pm 2.14 \mu\text{g}\cdot\text{mL}^{-1}$),
4 but the accumulation of the phytohormone in the culture medium was restored to wild-type levels
5 during the stationary growth phase (Az39 Δ hcp-E: $79.82 \pm 1.60 \mu\text{g}\cdot\text{mL}^{-1}$, Az39: 79.45 ± 4.94
6 $\mu\text{g}\cdot\text{mL}^{-1}$), showing a saturation kinetic (Fig. 2).

7 *Swimming, swarming, aggregation and biofilm formation* - Previous reports have shown that
8 mutation within T6SS gene clusters confer defects in swimming, swarming or biofilm formation
9 (Aschtgen *et al.*, 2008; de Pace *et al.*, 2010; Decoin *et al.*, 2014; Bouteiller *et al.*, 2020). The
10 Az39 Δ hcp-E mutant strain exhibited better swimming capacity compared to its parental strain in
11 swim and MMAB culture media (Fig. 3a). By contrast, both strains swarmed at similar rates (Fig.
12 3b). Cell aggregation was reduced for the Az39 Δ hcp-E mutant in MMAB and LB culture media
13 with an average reduction of 30% in comparison to the wild-type strain (Table 1). Finally, we
14 observed that the Az39 Δ hcp-E mutant strain produced ~50 % less biofilm after 48 and 72 h of
15 incubation, indicating a reduction in attachment capacity between cells. After 96 h of incubation,
16 the mutant showed a reduction of 23.15 % of biofilm production compared to the wild-type strain,
17 but this difference was not significant (Fig. 3c). Altogether, these data showed that the *A.*
18 *brasilense* T6SS1 decreases the ability to swim and confers an increased ability to adhere to abiotic
19 surfaces, at least in the early stages of the attachment phase.

20
21 *The A. brasilense T6SS provides anti-bacterial activity against competitors, including*
22 *phytopathogens.*

A number of reports have demonstrated that the T6SS is involved in regulating bacterial communities by targeting effector toxins into competing bacteria sharing the same ecological niche (Hood *et al.*, 2010; Wexler *et al.*, 2016; Sana *et al.*, 2016; Anderson *et al.*, 2017; Chassaing and Cascales, 2018; Allsopp *et al.*, 2020; Wood *et al.*, 2020; Duran *et al.*, 2021). We therefore asked whether the *A. brasilense* T6SS might confer protection to the microalgae or plants by delivering anti-bacterial toxins into phytopathogens. We performed *in vitro* competition experiments in minimal medium against *Escherichia coli* K-12, the soil bacterium *Ralstonia eutropha* and selected plant pathogens such as *Agrobacterium tumefaciens*, *Pectobacterium carotovorum*, *Dickeya dadantii* and *Ralstonia solanacearum*. After 8 h of co-incubation with wild-type *A. brasilense* Az39 and its isogenic $\Delta hcp-E$ mutant, surviving prey cells were counted on selective medium. Fig. 4 shows that the wild-type *A. brasilense* strain has T6SS-dependent anti-bacterial activity against all the strains tested as only 7-15% of cells survive against the wild-type *A. brasilense* compared to the $\Delta hcp-E$ mutant. The T6SS activity was significantly increased in presence of indole-3-acetic acid, previously shown to induce the T6SS gene cluster (Van Puyvelde *et al.*, 2011). In presence of IAA, only 0.1-1% of the prey cells survived co-incubation with wild-type *A. brasilense* cells whereas their recovery was not affected when co-incubated with $\Delta hcp-E$ mutant cells (Fig. 4). We conclude that the *A. brasilense* T6SS has anti-bacterial activity against several strains and therefore might confer protection to plant and microalgae against bacterial pathogens.

Contribution of the T6SS for C. sorokiniana/A. brasilense association

Microalgae growth - To test the role of the *A. brasilense* T6SS for mutualistic growth of *Chlorella*, we followed the growth of the microalgae alone or in presence of the *A. brasilense* Az39 wild-type

1 strain or its isogenic $\Delta hcp-E$ mutant (Fig. 5a). When cultured alone, the *C. sorokiniana* population
2 increased for the first two days and then slowly decreased. After six days, the population reached
3 its lowest numbers (Fig. 5a, open circles). In the presence of the *A. brasilense* wild-type or mutant
4 strains, the *Chlorella* population increased for 4 days and then declined (Fig. 5a, closed squares
5 and triangles, respectively). The growth rate of the microalgae after 4 days with either *A. brasilense*
6 strain was ~3.5 higher than microalgae cultured alone (Fig. 5a). For 6 days, the interaction with
7 both types of bacteria supported larger populations of microalgae, compared to the microalgae that
8 were cultured alone.

9
10 *Cell aggregation - Chlorella-Azospirillum* association and aggregation were probed by
11 fluorescence in situ hybridization (FISH) (Fig. 5b). Aggregates of the two bacterial strains grown
12 with *C. sorokiniana* showed different dynamics throughout the experiment. The initial distribution
13 of cells represents the outset of the microorganism's interaction (Fig. 5b panels a, e, i, m, q). In all
14 cases, the aggregates are increasing in size from day 1 (Fig. 5b panels a-t; Fig. 5c). The difference
15 in aggregates depending on the bacterial strain is evident at day 4. When *C. sorokiniana* was co-
16 cultured with *A. brasilense* Az39 large aggregates composed of bacteria and microalgae were
17 found (Fig. 5b panel d; Fig. 5c). Co-cultures of *C. sorokiniana* with *A. brasilense* Az39 $\Delta hcp-E$
18 showed aggregates mainly composed of microalgae (Fig. 5b panel h) and significantly smaller
19 than the aggregates formed with the wild-type Az39 strain (Fig. 5c). *C. sorokiniana* alone showed
20 smaller aggregates (5b panel t; Fig. 5c). *A. brasilense* Az39 growing alone displayed aggregates
21 increasing in size (Fig. 5b panels i-l; Fig. 5c); however, *A. brasilense* Az39 $\Delta hcp-E$ showed less
22 aggregation than wild-type, maintaining a similar size of aggregates at days 2 and 4 (Fig. 5b panels
23 m-p; Fig. 5c), in agreement with the defect in cell aggregation observed previously (Table 1).

1
2 *Accumulation of carbohydrates, lipids and photosynthetic pigments* - To better understand the
3 effect of the *A. brasilense* T6SS during mutualistic growth with *C. sorokiniana*, we measured the
4 accumulation of carbohydrates, lipids and pigments in the microalgae alone or in presence of the
5 *A. brasilense* Az39 wild-type strain or its isogenic $\Delta hcp-E$ mutant (Fig. 6 and 7). Fig. 6a shows
6 that co-culture of *Chlorella* with the wild-type *A. brasilense* Az39 strain significantly increased
7 the accumulation of carbohydrates in the microalgae-bacteria association only at days 1 and 4 of
8 co-culturing, while co-culture with *A. brasilense* Az39 $\Delta hcp-E$ cells had no effect on carbohydrates
9 accumulation (Fig. 6a). Regarding lipids production, co-culture of *C. sorokiniana* with either *A.*
10 *brasilense* strain significantly improved the production of lipids in the microalgae, although the
11 wild-type induced higher production of lipids at days 1, 2, and 6, while it was higher with the
12 mutant at day 4 (Fig. 6b). The effect of the *A. brasilense* T6SS on microalgae photosynthesis was
13 indirectly measured by quantifying major and auxiliary photosynthetic pigments: chlorophyll *a*
14 and *b*, violaxanthin, lutein and β -carotene (Fig. 7c-f). The accumulation of these photosynthetic
15 pigments followed a similar pattern. After 1 day of co-culture, the presence of *A. brasilense* strains
16 did not improve the synthesis of pigments. However, the presence of the Az39 wild-type strain
17 increased the production of pigments by *Chlorella* on days 2, 4 and 6. By contrast, co-culture of
18 *Chlorella* with the *A. brasilense* $\Delta hcp-E$ strain did not improve pigment production during the four
19 first days.

22 **Discussion**

1 In this study, we provide the first characterization of a Type VI secretion system from the
2 plant-growth promoting bacterium *Azospirillum brasilense* (Fig. 7). We show that the *A. brasilense*
3 T6SS1 is involved in attachment to the microalgae *Chlorella sorokiniana*, participates to the
4 formation of an efficient algae/bacterium consortium, and in the production of the IAA
5 phytohormone. Through its role on attachment and IAA levels, which exert major changes on
6 growth and the metabolism of the microalgae the T6SS indirectly allows increased lipid
7 metabolism in the microalgae. We also reveal that the *A. brasilense* T6SS confers protection
8 against bacterial phytopathogens. This study therefore demonstrates that the T6SS is an important
9 beneficial determinant of *Azospirillum/Chlorella* interaction by mediating attachment,
10 improvement of the metabolic performances and bio-control (Fig. 7).

11 We have shown that the T6SS participates to biofilm formation on abiotic surfaces and helps
12 attachment of *Azospirillum* to *Chlorella* cells and formation of *Chlorella* clusters. In addition, we
13 observed a role of the T6SS in IAA production. Attachment and IAA production are key in
14 *Azospirillum*–plant cell interactions (Bashan and de-Bashan, 2010). Attachment of *Azospirillum*
15 to cells is a major, active and complex phenomenon (Bashan, Levanony and Klein 1986; Gafni *et*
16 *al.* 1986). The initial interaction of *Azospirillum* spp. with plant cells is mainly mediated by
17 attachment of the bacterium, which employs a variety of piliated appendages at the cell surface to
18 promote stable interactions between the eukaryotic and prokaryotic cells (Pereg *et al.*, 2016);
19 Attachment between bacteria and microalgae was previously observed (de-Bashan *et al.*, 2011,
20 2015, 2016), but the question remains on whether this attachment is significant for the
21 development of this association by creating additional massive microalgae-bacteria aggregates.
22 Active attachment to plants and subsequent colonization were observed with the roots of cereals,
23 tomato, pepper, cotton and soybean (Bashan *et al.* 1989; Bashan, Singh and Levanony 1989;

1 Bashan, Levanony and Whitmoyer 1991; Levanony *et al.* 1989; Assmus *et al.* 1995; Pereg Gerk,
2 Gilchrist and Kennedy 2000). The initial step of plant root colonization by *Azospirillum* spp.
3 necessitates flagella and exopolysaccharides to create very stable interaction between the
4 eukaryotic and prokaryotic cells (Pereg, de-Bashan and Bashan 2016). These structures are also
5 needed for adhesion to non-living substrates (Bashan and Levanony 1988; Bashan and Holguin
6 1993). The polar flagellum of *A. brasilense*, which is primarily used for swimming, is involved in
7 the initial attachment process of the bacteria to wheat roots (Croes *et al.* 1991). Then the attachment
8 process involves two distinct steps. The first stage is fast and corresponds to a weak adhesion and
9 is mediated by host cell surface hydrophobicity, charges and lectins (Castellanos, Ascencio and
10 Bashan 1998, 2000). The second stage takes longer time but the attachment is irreversible. This
11 stage is mediated by extracellular surface polysaccharides and mucigel-like substances (Gafni *et*
12 *al.* 1986; Bashan and Levanony 1988; Bashan, Levanony and Whitmoyer 1991; Levanony and
13 Bashan 1991; Michiels *et al.*, 1989; Puente *et al.* 1999; Pereg Gerk, Gilchrist and Kennedy 2000;
14 Lerner *et al.*, 2009). Other determinants of *Azospirillum* attachment are the *flcA* regulator (Pereg
15 Gerk *et al.* 1998) that is also involved in stress response and carbohydrate and nitrogen metabolism
16 (Hou *et al.* 2014), chemotaxis pathways (Bible, Russell and Alexandre 2012; Pereg, de-Bashan
17 and Bashan 2016), Cpa pili (Wisniewski-Dye *et al.* 2011) as well as genes related to the surface
18 properties of *Azospirillum*, such as *noeJ* (mannose-6-phosphate isomerase) and *noeL* (GDP-
19 mannose 4,6-dehydratase) and pathways involved in lipopolysaccharide core processing (dTDP-
20 rhamnose biosynthesis pathway; Jofre *et al.*, 2004) or energy taxis (Greer-Phillips, Stephens and
21 Alexandre 2004). Although the T6SS might be a new molecular determinant of the adhesion of
22 *Azospirillum* to microalgae cells, it remains to define whether the effect of the T6SS mutation on
23 adhesion is due to defect in flagellar synthesis or expression of attachment genes. While cross-talk

1 between the T6SS and flagellar genes have been demonstrated in the *P. fluorescens* PGPB
2 (Bouteiller et al., 2020), the first hypothesis is unlikely as we have shown here that deletion of the
3 *hcp-tssE* genes does not affect swarming rates and increases swimming rates. The second
4 hypothesis merits investigation, as previous studies have shown that the inactivation of the T6SS
5 may affect expression of fimbrial genes in pathogenic *E. coli* strains (de Pace et al., 2010).

6 Previous studies have shown that *A. brasilense* enhance the growth and induce an increased
7 accumulation of carbohydrates, starch (Choix et al., 2012a, b), fatty acids, total lipids (Leyva et
8 al., 2014; Peng et al., 2020) and pigments (de-Bashan et al. 2002; Peng et al., 2020) in *Chlorella*
9 spp. The effect has been mostly attributed to the IAA production by the bacteria (de-Bashan and
10 Bashan, 2008; de-Bashan et al., 2011, 2015; Peng et al., 2020). Since the concentration of IAA
11 produced by the wild-type and T6SS *A. brasilense* strains is similar at the times of the experiments
12 with the microalgae (after 30 hours), the reduction of lipids and photosynthetic pigments induced
13 by the T6SS mutant can be attributed to its lower attachment capacity. It was previously shown
14 that attachment is not essential for C and N compounds transfer between individual cells of *C.*
15 *sorokiniana* and *A. brasilense* (de-Bashan et al., 2016); however, the metabolites produced by
16 *Azospirillum*, might be transient and probably a close cell proximity is required to achieve a more
17 effective mass transfer (Peng et al., 2020).

18 We have also shown that mutation in the *A. brasilense* T6SS affects IAA production. IAA, as a
19 phytohormone, has no important role in bacterial metabolism (Spaepen and Vanderleyden, 2011),
20 but is widespread among PGPB (Duca et al., 2014) and increases plant growth. The production of
21 IAA during the interaction between *Azospirillum* spp. and *Chlorella* spp. and its effect on growth
22 was reported several times (de-Bashan et al., 2008; Palacios et al., 2016a, b, Peng et al., 2020).
23 The role of the T6SS in *Azospirillum*/plant interactions suggests that the expression of this cluster

1 and/or the activity of the machine is responsive to environmental or plant cues. It would be
2 therefore interesting to determine what are the environmental signals in addition to IAA (Van
3 Puyvelde *et al.*, 2011), and to identify the regulators that control the expression of this T6SS gene
4 cluster.

5 The role of the *A. brasilense* T6SS in plant/bacterium interactions is supported by the observation
6 that T6SS genes are up-regulated in presence of the IAA phytohormone (Van Puyvelde *et al.*,
7 2011). However, contrarily to T2SS or T3SS present in plant-associated bacteria, the *A. brasilense*
8 T6SS is not deployed for pathogenic purposes. Interestingly, T6SS gene clusters are conserved in
9 proteobacterial and bacteroidetes genomes, including non-pathogenic bacteria such as gut
10 symbiots (Wexler *et al.*, 2016), soil and marine bacteria. The presence of a T6SS in these strains
11 suggests that the T6SS could be used for promoting commensal or mutualistic relationships
12 between bacteria and eukaryotes (Jani and Cotter 2010; Konovalova, Petters and Sogaard-
13 Andersen 2010; Jiménez-Guerrero *et al.* 2013). Indeed, the *Rhizobium leguminosarum* bv. *trifolii*
14 T6SS has a role in the nodulation process by blocking the colonization/infection process of non-
15 host peas (Roest *et al.* 1997; Bladergroen *et al.* 2003). Whereas it has been proposed that RbsB is
16 a T6SS effector delivered by *R. leguminosarum* into pea cells (Bladergroen *et al.* 2003), the
17 effector(s) delivered by the *A. brasilense* T6SS into plant/algae cells remain to be identified.

18 Finally, our data show that the T6SS provides antagonistic activities to *A. brasilense* against a
19 number of bacterial strains including phytopathogens such as *Agrobacterium*, *Pectobacterium*,
20 *Dickeya* and *Ralstonia in vitro*. This demonstrates that, in addition to promoting growth, *A.*
21 *brasilense* likely confers protection against harmful pathogens in a T6SS-dependent manner.
22 Effectors with DNase, phospholipase and peptidoglycan hydrolase activities have been shown to
23 be delivered into bacterial cells (Russel *et al.*, 2014; Jurenas and Journet, 2021). The *A. brasilense*

1 T6SS effector repertoire needs to be determined. It has been shown that the effector genes are
2 usually located within the T6SS gene cluster or within *hcp/vgrG* islands scattered in the genome
3 (Durand *et al.*, 2014). Indeed, the *vgrG* gene encoded within the cluster is fused to an additional
4 domain resembling *Bacillus subtilis* YwqJ, a potent toxin with putative deaminase activity, and is
5 followed by a gene resembling its cognate YwqK antitoxin (Brantl and Müller, 2019; Kobayashi,
6 2021) (Fig. 1).

7 In summary, the *Chlorella* spp.–*Azospirillum* spp. association has been used as a model system to
8 study aggregation, metabolic, and molecular effect occurring during plant-bacteria interaction (de-
9 Bashan and Bashan, 2008). The present work extends the potential to study a molecular
10 mechanism occurring between a bacterium and a microalga, specifically by showing that a T6SS
11 mutant is also deficient in attachment capabilities and produces less microalgal aggregation. This
12 T6SS mutant, while promoting the growth of microalgae similar to its parental wild-type *A.*
13 *brasilense* strain also impacts lipid, and pigment metabolism. This study has thus far broader
14 impacts than observing interactions between microalgae and bacteria *in vitro*, by demonstrating
15 that a widespread secretion system, known almost exclusively from studies of pathogenesis and
16 antibacterial competition, is participating in processes that are beneficial to the plant/microalga
17 cells.

19 **Experimental procedures**

20 *Microorganisms and growth conditions*

21 The unicellular microalga *Chlorella sorokiniana* Shih. et Krauss (UTEX 2714, University of
22 Texas, Austin, TX; formerly *C. vulgaris* UTEX 2714, Bashan *et al.* (2016)) was used in this study.
23 *C. sorokiniana* was axenically grown at 27 ± 2 °C, with stirring at 120 rpm under a light intensity

of 60 $\mu\text{mol photon}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ in sterile C30 minimal medium (composition (in $\text{g}\cdot\text{L}^{-1}$): KNO_3 (25), $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$ (10), KH_2PO_4 (4), K_2HPO_4 (1), $\text{FeSO}_4\cdot 7\text{H}_2\text{O}$ (1), and (in $\mu\text{g}\cdot\text{L}^{-1}$): H_3BO_3 (2.86), $\text{MnCl}_2\cdot 4\text{H}_2\text{O}$ (1.81), $\text{ZnSO}_4\cdot 7\text{H}_2\text{O}$ (0.11), $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$ (0.09), and NaMoO_4 (0.021) at pH 5.25), and transferred to new medium every 7 days. Periodically, the cultures were checked for purity and absence of bacterial contamination on nutrient agar (#N9405, Sigma-Aldrich, St Luis, MO). The wild-type *Azospirillum brasilense* Az39 strain (Instituto de Microbiología y Zoología Agrícola del INTA-IMyZA, Castelar, Buenos Aires, Argentina, strain Az39 (WDCM31), Rivera *et al.* (2014)) and its isogenic *Az39* mutant (this study, see construction below) were grown in BTB medium containing gluconic acid as a carbon source (Bashan, Trejo and de-Bashan 2011) for 18 h, at 35 ± 2 °C and stirred at 120 rpm, or in MMAB minimal medium (composition (in $\text{g}\cdot\text{L}^{-1}$): K_2HPO_4 (3.0), NaH_2PO_4 (1.0), NH_4Cl (1.0), $\text{MgSO}_4\cdot 7\text{H}_2\text{O}$ (0.3), KCl (0.15), $\text{CaCl}_2\cdot 2\text{H}_2\text{O}$ (0.01), $\text{FeSO}_4\cdot 7\text{H}_2\text{O}$ (0.0025), sodium malate (5.0), biotin (0.005) and microelements) at 28°C.

Construction of the A. brasilense Az39 mutant

Genomic analysis of the Type VI secretion system in A. brasilense Az39 - Using the genome sequence of *A. brasilense* Az39 (Rivera *et al.* 2014), two complete sets of genes encoding putative T6SSs were detected and analysed by RAST (Aziz *et al.* 2008), KEGG (Kanehisa *et al.* 2012), SecReT6 (Li *et al.*, 2015) and BastionHub (Wang *et al.*, 2021). The T6SS1 gene cluster, which was studied here, is shown in Fig. 1 and T6SS proteins are listed in Table S1. The gene cluster encodes all the core-component genes, *tssA-tssM*, required to assemble a fully functional T6SS (Cascales, 2008).

Az39 mutant construction - The *hcp* and *tssE* genes were simultaneously deleted by homologous recombination. Fragments upstream and downstream of the *hcp-tssE* region were

1 amplified by PCR from genomic DNA using primer pairs Az39-5'HcpFW/Az39-B-5'HcpRV and
 2 Az39-B-3'HcpFW/Az39-3'HcpRV, respectively (Table S2) cloned into the pGemT-Easy vector
 3 (Promega, Madison, WI) and sequenced. The upstream fragment was inserted into the downstream
 4 fragment-containing plasmid, using *Bam*HI, and *Pst*I restriction enzymes to yield the pGemT-Hcp-
 5 UD vector. Then, a gentamicin-resistance cassette (Gm^R), excised from pME3280a (Zuber *et al.*,
 6 2003) with *Xho*I, was inserted in reverse orientation into the *Bam*HI site of pGemT-Hcp-UD after
 7 blunting ends with T4 DNA polymerase (New England Biolabs, Ipswich, MA). The resulting
 8 upstream-Gm^R–downstream construction was finally sub-cloned into pK18mobSacB (Schäfer *et*
 9 *al.*, 1994), using *Pst*I and *Sph*I to obtain pK18-Hcp-UGD. This suicide plasmid, carrying the
 10 mutagenesis cassette, was mobilized into *A. brasilense* Az39 by biparental mating with
 11 *Escherichia coli* S17-1 as the donor (Simon *et al.*, 1983). Double recombinant clones, resistant to
 12 gentamicin and sensitive to kanamycin, were recovered without the need for *sacB* counter
 13 selection. To confirm the correct replacement of the targeted genome region, PCR was used with
 14 several primer combinations, and the amplified fragments were visualized by TAE-agarose
 15 electrophoresis. The phenotypes of the mutant were analyzed in three independent clones (C1, C2,
 16 and C3) to exclude the presence of additional mutations (Fig. S1). Fig. S1 describes the genomic
 17 arrangement of the wild-type, the two possible simple recombinants (R1 or R2), and the double
 18 recombinant (R1+R2) of *A. brasilense* Az39. The site of annealing and the product length of the
 19 different diagnostic primers (P1, P2, P3, P4, P5, P6, GmR, and GmF) on the four possible
 20 arrangements are indicated. The inserted table indicates the PCR products expected for each strain,
 21 and the agarose gel shows the actual results of the PCR. A PCR was performed using P3 and P6
 22 primers that uniquely anneal on the genomic region that flanks the modified region.

1 *Strain characterization*

2 *Bacterial Growth* - Growth curves of *A. brasilense* Az39 and mutant strain Az39 Δ hcp-E were
3 determined by measuring the turbidity of cell suspensions (OD₅₉₅) and cell number (CFU.mL⁻¹)
4 over time. Sample of 0.5 mL of an overnight culture (OD₅₉₅ = 1.0, ~10⁹ CFU.mL⁻¹) were
5 inoculated to 200-mL Erlenmeyer flasks containing 50 mL of Luria-Bertani medium (LB) or
6 MMAB minimal medium (Vanstockem *et al.*, 1987) supplemented with 100 µg.mL⁻¹ of L-
7 tryptophan. Cultures were incubated at 37 °C and constant stirring at 180 rpm for 80 h. Turbidity
8 of cell suspensions was spectrophotometrically measured at 595 nm over time (ZelTec ZL5000P,
9 Zel Technologies, Hampton, VA) at 5 h intervals. At the same time, cell counts were performed
10 on LB agar plates supplemented with Congo red (1.5%, v:v), after incubation for 72 h at 37 °C.

11
12 *Indole-3-acetic acid production* - Pure cultures of *A. brasilense* Az39 and its Δ hcp-E derivative
13 were cultivated in LB and MMAB media, supplemented with L-tryptophan, at 37 °C, under
14 constant stirring at 180 rpm for 48 h corresponding to stationary stage of growth of this species.
15 Cultures were centrifuged at 15,650 × g for 10 min and cells were discarded. Quantification of
16 IAA was performed by spectrophotometry (Glickmann and Dessaux, (1995) and confirmed by
17 HPLC (Rivera *et al.*, 2018). Briefly, aliquots of 1000 µL of bacterial culture were centrifuged at
18 11,300 × g for 10 min. Subsequently, samples were filtered (0.2 µm), and 500 µL of supernatant
19 were mixed with 500 µL of Salkowski's reagent (7.9 M H₂SO₄ and 12.5 g.L⁻¹ FeCl₃) and gently
20 shaken in inverted position at least 10 times. Samples were incubated in the dark for 30 min and
21 the absorbance at 530 nm was measured. An aliquot of filtered supernatants was injected with a
22 final volume of 20 µL in an HPLC Waters 600-MS device (Waters Inc., USA) equipped with an
23 U6K injector and C18 reverse phase column (Purospher STAR RP C-18 3 mm, Lichrocart 55-4)

1 heated at 30°C, coupled to a system with UV-VIS Waters 486 detector (Waters Inc., USA) set at
2 265 nm. The elution was performed with a mixture of ethanol: acetic acid: water (Et-OH/H-
3 Ac/H₂O) (12: 1: 87) as mobile phase at a flow rate of 1 mL·min⁻¹ at 30°C. The retention time for
4 IAA was 10.1-10.3 minutes and the quantification was performed by integration of the peak area
5 corresponding to the retention time (RT) using an integration software (Waters Inc. USA). The
6 IAA concentration was expressed in µg·mL⁻¹

7
8 *Motility* - Motility was assayed under different concentrations of agar-agar: 1 µL of bacterial
9 culture in late exponential phase was placed in the center of Petri dishes containing minimal swim
10 motility medium (Atkinson *et al.*, 2006) and MMAB medium as previously described
11 (Vanstockem *et al.*, 1987). Minimal swim motility agar plates contained (g·L⁻¹) tryptone (10.0)
12 and NaCl (5.0). Both media were prepared with 0.3% (w/v) agar to observe swimming and 0.7%
13 (w/v) agar for determining swarming. The plates were inverted and incubated at 28 ± 2 °C for 72
14 h to avoid drying the agar at higher temperatures. The diameter of the displacement halo was then
15 measured (Atkinson *et al.*, 1999; Hall and Krieg, 1983).

16
17 *Cell aggregation and biofilm production* - Cell aggregation was measured according to Madi and
18 Henis (1989) with modifications (Burdman *et al.*, 1998). Briefly, 5 mL of stationary cultures of
19 both strains obtained in MMAB and LB culture medium at 37°C and constant stirring at 180 rpm
20 for 48 hours were transferred to 10 mL capacity conical tubes and allowed to stand for 20 min.
21 Then, turbidity was measured at 540 nm using a Zeltec ZL5000P spectrophotometer (OD₁). The
22 culture was homogenized for 1 min and the turbidity was measured again (OD₂). The aggregation
23 percentage was calculated according to the following equation %AP (OD₂ - OD₁) × 100/OD₂.

Biofilm formation was measured according to O'Toole & Kolter (1998). Briefly, 13 μ L of late exponential cultures were inoculated in hemolysis tubes containing 1,300 μ L of sterile LB medium and incubated at $37 \pm 1^\circ\text{C}$ for 48, 72, and 96 h, without stirring. At each sampling time, the culture medium was carefully removed from the tube by micropipetting, leaving only the biofilm. The biofilm was rinsed three times with 1,300 μ L of sterile 0.85 % saline solution. Once the culture medium was removed, the biofilm was stained with 0.1% crystal violet (C3886, Sigma-Aldrich) solution (1,300 μ L for 15 min at room temperature, $24 \pm 2^\circ\text{C}$). The stained biofilm was rinsed three times with sterile distilled water. The biofilm was re-suspended in 1,300 μ L of 96% ethanol containing three glass beads (3 mm diam) in each tube and stirred vigorously by a vortex for ~1 min. Finally, absorbance was measured at 560 nm by spectrophotometry (ZelTec ZL5000P, Zel Technologies).

Anti-bacterial competition assay.

Spontaneous *Agrobacterium tumefaciens*, *Pectobacterium carotovorum*, *Dickeya dadantii*, *Ralstonia eutropha* and *Ralstonia solanacearum* strains resistant to nalidixic acid were obtained after 5 consecutive liquid growth cultures in presence of increasing concentrations of nalidixic acid (2, 5, 10, 20 and 30 μ M) and plating on agar plates supplemented with 40 μ M nalidixic acid. *Azospirillum brasilense* and nalidixic acid-resistant recipient cells were grown in MMAB supplemented - or not - with 400 μ M indole-3-acetic acid (IAA) to an $\text{OD}_{595} \sim 1$ and concentrated in MMAB to a final OD_{595} of 10. *A. brasilense* attacker cells were mixed with nalidixic resistant recipient cells at a 4:1 ratio and 15- μ L drops of the mixture were spotted on dried minimum MMAB or MMAB/IAA agar plates. After incubation for 8 h at 28°C , the spots were scratched off, cells

were re-suspended in LB to an OD₅₉₅ of 0.5, and the surviving recipient cells were counted after plating serial dilutions on LB agar plates supplemented with 40 µM nalidixic acid.

Immobilization of microorganisms in alginate beads

Before immobilization in alginate beads, 10 mL of axenic microalgae culture was inoculated into 90 mL sterile C30 medium and incubated at 27 ± 2 °C, with stirring at 120 rpm under 60 µmol photons·m⁻²·s⁻¹ light intensity for six days. The two strains of *A. brasilense* were grown in BTB culture medium containing gluconic acid as a carbon source (Bashan *et al.*, 2011) for 18 h, at 35 ± 2 °C and stirred at 120 rpm. Immobilization procedure was done as described by de-Bashan *et al.* (2015). Briefly, after growth period, each culture was harvested by centrifugation (Hemle Z 200A, Wehingen, Germany) at $2,000 \times g$, washed twice, resuspended in sterile saline solution (0.85% NaCl) and cell density was adjusted to 5×10^6 cells·mL⁻¹ for microalgae and $\sim 10^9$ CFU·mL⁻¹ (OD_{540 nm} = 1) for bacteria. Then, 40 mL of axenic cultures of *C. sorokiniana* or *A. brasilense* Az39 or *A. brasilense* Az39Δhcp-E were mixed with 160 mL of sterile 1.5% alginate solution (~1200 cP, #05218295, MP Biomedicals, Santa Ana, CA) and stirred for 15 min until the alginate slurry was completely dissolved. Beads (highly homogenous; 3 mm diam) were produced in 2% CaCl₂, using automatic bead-forming equipment under constant pressure to ensure uniformity of each bead (de-Bashan and Bashan, 2010). The beads were cured for 30 min in the same 2% CaCl₂ solution to improve hardening. They were washed in sterile 0.85% saline solution. This procedure routinely produces $\sim 1 \times 10^6$ cells·bead⁻¹ for each organism with low variability (de-Bashan *et al.*, 2004). For microalgae and bacteria co-immobilized in the same bead, 20 mL of each culture was mixed in 160 mL alginate to form the beads, as previously described. Because immobilization usually reduces the number of *A. brasilense* in beads, the beads were incubated again for 24 h in

1 10% (v/v) diluted nutrient broth (N7519, Sigma-Aldrich) to reach an initial population of 1×10^6
2 cells·bead⁻¹.

3 4 *Experimental conditions to assess the performance of microalgae.*

5 After the second incubation, beads were washed twice with sterile saline solution. In preparation
6 of each experiment, 15 g of beads from each treatment (microorganisms immobilized alone or co-
7 immobilized) were added to 500 mL Erlenmeyer flasks containing 200 mL of synthetic growth
8 medium (SGM) with the following ingredients (in mg·L⁻¹): NaCl (7), CaCl₂ (4), MgSO₄·7H₂O
9 (2), K₂HPO₄ (217), KH₂PO₄ (8.5), Na₂HPO₄ (33.4), NH₄Cl (191) and incubated under 90 μmol
10 photon·m⁻² s⁻¹ light intensity and stirred at 122 rpm at 29 ± 2 °C for 6 d.

11 12 *Quantifying microalgae cells*

13 In each experiment for each sampling time, and each experimental replicate, three beads per flask
14 were solubilized in 1 mL of 4% NaHCO₃ solution for 20 min at ambient temperature (25 ± 4°C).
15 *C. sorokiniana* cells were counted under a light microscope, using a Neubauer hemocytometer
16 (Bright line counting chamber, Hausser Scientific, Horsham, PA) connected to an image analyser
17 (Image ProPlus 6.3, Media Cybernetics, Silver Spring, MD). Growth rate of *C. sorokiniana* (μ)
18 was defined as: $\mu = (\ln N_{t_1} - \ln N_{t_0}) / (t_1 - t_0)$, where N_{t_1} is the number of cells at sampling time,
19 and N_{t_0} is the number of cells at the beginning of the experiment, t_1 is sampling time and t_0 the
20 beginning of the experiment (Oh-Hama and Miyachi, 1992).

21 22 *Analytical methods*

23 *Total carbohydrates*

For each replicate of each treatment ($n = 15$), carbohydrates extraction and analysis were done following Choix *et al.* (2012a). Briefly, 1 g of alginate beads were sampled after 1, 2, 4, and 6 d, washed in distilled water, dried at 80 °C for 12 h, and ground with a mortar and pestle, which yielded 10 mg samples. These 10 mg ground samples were re-suspended in 5 mL of H₂SO₄ 1 M and sonicated for 4 min at 22.5 kHz with an ultrasonic cell disruptor (Misonix, Farmingdale, NY). Carbohydrates were extracted by acid hydrolysis of the sample at 100°C for 60 min. The quantification was done by the phenol-sulfuric method adapted to microplate (Dubois *et al.*, 1956; Masuko *et al.*, 2005).

Total lipid analysis

Lipid extraction followed the method of Leyva *et al.* (2014), which was developed for microalgae and based on Bligh and Dyer (1959). Quantification of total lipids was done as previously described (Pande *et al.*, 1963).

Pigment analysis

Extraction of pigments (chlorophyll *a* and *b*, violaxanthin, and lutein) was performed with HPLC grade methanol, after dissolving the beads. Identification of pigments used the HPLC method (Vidussi *et al.*, 1996). The HPLC system (Agilent 1100; Agilent Technologies, Santa Clara, CA) was equipped with a reverse phase column (Hypersil BDS C8, 5 µm particle size, 100×45 mm; Thermo Scientific, Waltham, MA) and was run isocratically. The mobile phase included two solutions. Solution A was a mixture of 70:30 methanol:1N ammonium acetate. Solution B was 100% HPLC grade methanol. Injection volume was 20 µL; flow rate was 0.5 mL·min⁻¹. The detector was a diode array with a wavelength of 190–900 nm, with the capacity to detect at five

fixed wavelengths. Identification criteria for pigments were retention time of the standard pigments (International Agency for ^{14}C determinations, Denmark, DHI-Denmark, PPS-CHLA) and absorbance at 350–750 nm from the diode detector. Results are expressed as $\mu\text{g}\cdot\text{g}^{-1}$.

Fluorescent in situ hybridization (FISH)

The FISH procedure described by Palacios *et al.* (2019), was used to analyze the patterns of cell aggregation during the interaction of *C. sorokiniana* and *A. brasilense* strains. Approximately 100 images were acquired with a digital camera (Evolution FV Cooled Color; Media Cybernetics). Five random images per treatment were used for quantification and measurement of clusters with an Image ProPlus 4.5 software (Media Cybernetics).

Experimental design and statistical analyses

Two types of experiments were conducted. (a) *In vitro* characterization of the two bacteria, strains were performed in batch culture in four replicates. Each experiment was independently repeated three times. (b) The setup of the experiments of microalgae-bacteria was by batch culturing in Erlenmeyer flasks in five replicates, where one flask served as a replicate. Each of these experiments was independently repeated two times. Each experimental setup contained the following treatments in beads containing: (1) *A. brasilense* Az39, (2) *A. brasilense* Az39 Δ hcp-E, (3) *C. sorokiniana*, (4) co-immobilization of *C. sorokiniana* with *A. brasilense* Az39, (5) co-immobilization of *C. sorokiniana* with the mutant strain Az39 Δ hcp-E ($n = 20$). The results from each treatment from two repetitions were combined. For characterization of the mutant strain, each pair of values at every sampling time were analyzed by Student's *t*-test. For analysis of the effect of *A. brasilense* strains, the data were first analyzed by one-way ANOVA and then by HSD

Tukey's test. Significance was set at $P < 0.05$, using Statistica 6.0 (Tibco Software, Palo Alto, CA), Prism 5.0 (GraphPad Software, La Jolla, CA), and Sigmaplot 13.0 (Systat Software, Chicago, IL).

Disclosure Statement

No potential conflict of interest was reported by the authors.

Author's contribution

Fabricio D. Cassan. Envisioned the project with Luz de-Bashan. Managed the construction of the mutant and its testing in Argentina. Helped with the writing of the manuscript.

Anahí Coniglio. Measured bacterial growth, IAA concentration, biofilm formation and performed swimming and swarming experiments.

Edgar Amavizca. Performed all experiments and FISH analysis related to microalgae-bacteria interaction.

Guillermo Maroniche. Constructed the mutant.

Eric Cascales. Edited the final version.

Yoav Bashan. General supervision of the microalgae-bacteria studies in Mexico. Wrote the initial draft of the manuscript.

Luz E. de-Bashan. Envisioned the project with Fabricio Cassan. Supervised the Mexican part of the project on microalgae-bacteria interaction. Edited the final version.

All authors read and approved the final version of the manuscript.

Acknowledgments

At UNRC (Argentina), we thank Alejandro Peticari of INTA-IMYZA for providing *Azospirillum brasilense* Az39. At CIBNOR (Mexico), we thank Francisco Hernandez for pigment analysis and Ariel Cruz for scanning electron microscopy. At Universidad Autonoma de Chihuahua (Mexico), we thank Oskar Palacios for feedback and suggestion to the manuscript.

Funding

This study was supported by the CONICET (PPI-112-201001-00454-KA1) and FONCyT (PICT-2012-1051) of Argentina; by Consejo Nacional de Ciencia y Tecnología of Mexico (CONACYT-Basic Science-2015, contract 251102 and CONACYT Basic Science-2017, contract 284562), and by the Bashan Institute of Science of the USA. Work in EC laboratory is supported by the CNRS, the Aix-Marseille University, and the Fondation Bettencourt-Schueller.

References

- Alexandre, G., Greer, S.E., and Zhulin, I.B. (2000) Energy taxis is the dominant behavior in *Azospirillum brasilense*. *J. Bacteriol.* **182**: 6042–6048.
- Allsopp, L.P., Bernal, P., Nolan, L.M., and Filloux, A. (2020) Causalities of war: The connection between type VI secretion system and microbiota. *Cell Microbiol.* **22**: e13153.
- Anderson, M.C., Vonaesch, P., Saffarian, A., Marteyn, B.S., and Sansonetti, P.J. (2017) *Shigella sonnei* encodes a functional T6SS used for interbacterial competition and niche occupancy. *Cell Host Microbe.* **21**: 769-776.e3.
- Aschtgen, M.S., Bernard, C.S., De Bentzmann, S., Lloubès, R., and Cascales, E. (2008) SciN is an outer membrane lipoprotein required for type VI secretion in enteroaggregative *Escherichia coli*. *J Bacteriol.* **190**: 7523-31.

1 Assmus, B., Hutzler, P., Kirchhof, G., Amann, R., Lawrence, J.R., and Hartmann, A. (1995) *In*
2 *Situ* localization of *Azospirillum brasilense* in the rhizosphere of wheat with fluorescently
3 labeled, rRNA-targeted oligonucleotide probes and scanning confocal laser microscopy.
4 *Appl. Environ Microbiol* **61**: 1013–1019.

5 Atkinson, S., Chang, C.-Y., Sockett, R.E., Cámara, M., and Williams, P. (2006) Quorum sensing
6 in *Yersinia enterocolitica* controls swimming and swarming motility. *J Bacteriol* **188**:
7 1451–1461.

8 Atkinson, S., Throup, J.P., Stewart, G.S.A.B., and Williams, P. (1999) A hierarchical quorum
9 sensing system in *Yersinia pseudotuberculosis* is involved in the regulation of motility and
10 clumping. *Mol Microbiol* **33**: 1267–1277.

11 Aziz, R.K. *et al.* (with 25 co-authors). (2008) The RAST server: rapid annotations using
12 subsystems technology. *BMC Genomics* **9**: 75.

13 Bashan, Y., and de-Bashan, L.E. (2010) How the plant growth-promoting bacterium *Azospirillum*
14 promotes plant growth – a critical assessment. *Adv Agron* **108**: 77–136.

15 Bashan, Y., and Holguin, G. (1994) Root-to-root travel of the beneficial bacterium *Azospirillum*
16 *brasilense*. *Appl Environ Microbiol* **60**: 2120–2131.

17 Bashan, Y., and Levanony, H. (1987) Horizontal and vertical movement of *Azospirillum brasilense*
18 Cd in the soil and along the rhizosphere of wheat and weeds in controlled and field
19 environments. *J Gen Microbiol* **133**: 3473–3480.

20 Bashan, Y., and Levanony, H. (1988) Active attachment of *Azospirillum brasilense* Cd to quartz
21 sand and to a light-textured soil by protein bridging. *J Gen Microbiol* **134**: 2269–2279.

22 Bashan, Y., Levanony, H., and Klein, E. (1986) Evidence for a weak active external adsorption of
23 *Azospirillum brasilense* Cd to wheat roots. *J Gen Microbiol* **132**: 3069–3073.

- 1 Bashan, Y., Levanony, H., and Whitmoyer, R.E. (1991) Root surface colonization of non-cereal
2 crop plants by pleomorphic *Azospirillum brasilense* Cd. *J Gen Microbiol* **137**: 187–196.
- 3 Bashan, Y., Trejo, A., and de-Bashan, L.E. (2011) Development of two culture media for mass
4 cultivation of *Azospirillum* spp. and for production of inoculants to enhance plant growth.
5 *Biol Fertil Soils* **47**: 963–969.
- 6 Bashan, Y., Bustillos, J.J., Leyva, L.A., Hernandez, J.-P., and Bacilio, M. (2015) Increase in
7 auxiliary photoprotective photosynthetic pigments in wheat seedlings induced by
8 *Azospirillum brasilense*. *Biol Fertil Soils* **42**: 279–285.
- 9 Bashan, Y., Lopez, B.R., Huss, V.A.R., Amavizca, E., and de-Bashan, L.E. (2016) *Chlorella*
10 *sorokiniana* (formerly *C. vulgaris*) UTEX 2714, a non-thermotolerant microalga useful for
11 biotechnological applications and as a reference strain. *J Appl Phycol* **28**:113–121.
- 12 Basler, M. (2015) Type VI secretion system: secretion by a contractile nanomachine. *Philos Trans*
13 *R Soc Lond B Biol Sci* **370**: 20150021.
- 14 Bernal, P., Allsopp, L.P., Filloux, A., and Llamas, M.A. (2017) The *Pseudomonas putida* T6SS is
15 a plant warden against phytopathogens. *ISME J* **11**: 972-987.
- 16 Bernal, P., Llamas, M.A., and Filloux, A. (2018) Type VI secretion systems in plant-associated
17 bacteria. *Environ Microbiol* **20**: 1-15.
- 18 Bible, A., Russell, M.H., and Alexandre, G. (2012) The *Azospirillum brasilense* Che1 chemotaxis
19 pathway controls swimming velocity, which affects transient cell-to-cell clumping. *J*
20 *Bacteriol* **194**: 3343-55.
- 21 Bingle, L.E, Bailey, C.M., and Pallen, M.J. (2008) Type VI secretion: a beginner’s guide. *Curr Op*
22 *Microbiol* **11**: 1–6.

- 1 Bladergroen, M.R., Badelt, K., and Spaink, H.P. (2003) Infection-blocking genes of a symbiotic
2 *Rhizobium leguminosarum* strain that are involved in temperature-dependent protein
3 secretion. *Mol Plant Microbe Inter***16**: 53–64.
- 4 Bligh, G.E., and Dyer, J.W. (1959) A rapid method for total lipid extraction and purification. *Can*
5 *J Biochem Physiol* **37**: 911–917.
- 6 Borrero de Acuña, J.M., and Bernal, P. (2021) Plant holobiont interactions mediated by the type
7 VI secretion system and the membrane vesicles: promising tools for a greener agriculture.
8 *Environ Microbiol* **23**: 1830-1836.
- 9 Bouteiller, M., Gallique, M., Bourigault, Y., Kosta, A., Hardouin, J., Massier, S., Konto-Ghiorgi,
10 Y., Barbey, C., Latour, X., Chane, A., Feuilloley, M., and Merieau, A. (2020) Crosstalk
11 between the type VI secretion system and the expression of class IV flagellar genes in the
12 *Pseudomonas fluorescens* MFE01 strain. *Microorganisms* **8**: 622.
- 13 Brackmann, M., Nazarov, S., Wang, J., and Basler, M. (2017) Using force to punch holes:
14 mechanics of contractile nanomachines. *Trends Cell Biol* **27**: 623-632.
- 15 Brantl, S., and Müller, P. (2019). Toxin-Antitoxin Systems in *Bacillus subtilis*. *Toxins* (Basel). **11**:
16 262.
- 17 Brunet, Y.R., Zoued, A., Boyer, F., Douzi, B., and Cascales, E. (2015) The Type VI secretion
18 TssEFGK-VgrG phage-like baseplate is recruited to the TssJLM membrane complex via
19 multiple contacts and serves as assembly platform for tail tube/sheath polymerization. *PLoS*
20 *Genetics* **11**: e1005545.
- 21 Burdman, S., Jurkevitch, E., Schwartsburd, B., Hampel, M. and Okon, Y. (1998). Aggregation in
22 *Azospirillum brasilense*: effects of chemical and physical factors and involvement of
23 extracellular components. *Microbiology*, **144**: 1989-1999.

- 1 Cascales, E. (2008) The type VI secretion toolkit. *EMBO Reports* **9**: 735–741.
- 2 Cassan, F., Coniglio, Anahí., López, G. Molina, R., Nievas, S., Le Noir de Carlan, C., Donadio,
3 F., Torres, D., Rosas, S., Olivera Pedrosa, F., Maltempi de Souza, E., Díaz Zorita. M., de
4 Bashan, L.E., and Mora, V. (2020). Everything you must know about *Azospirillum* and its
5 impact on science, research, agriculture and beyond. *Biol Fertil Soils* **56**: 461-479
- 6 Castellanos, T., Ascencio, F., and Bashan, Y. (1998) Cell-surface lectins of *Azospirillum* spp. *Curr*
7 *Microbiol* **36**: 241-4.
- 8 Castellanos, T., Ascencio, F., and Bashan Y. (2000) Starvation-induced changes in the cell surface
9 of *Azospirillum lipoferum*. *FEMS Microbiol Ecol* **33**: 1-9.
- 10 Chassaing, B., and Cascales, E. (2018) Antibacterial weapons: targeted destruction in the
11 microbiota. *Trends Microbiol* **26**: 329-338.
- 12 Cherrak, Y., Flaugnatti, N., Durand, E., Journet, L., and Cascales, E. (2019) Structure and activity
13 of the type VI secretion system. *Microbiol Spectr* **7**: PSIB-0031-2019.
- 14 Choix, F.J., de-Bashan, L.E., and Bashan, Y. (2012a) Enhanced accumulation of starch and total
15 carbohydrates in alginate-immobilized *Chlorella* spp. induced by *Azospirillum brasilense*.
16 I. Autotrophic conditions. *Enzyme Microbl Technol* **51**: 294–299.
- 17 Choix, F.J., de-Bashan, L.E., and Bashan Y. (2012) Enhanced accumulation of starch and total
18 carbohydrates in alginate-immobilized *Chlorella* spp. induced by *Azospirillum brasilense*.
19 II. Heterotrophic conditions. *Enzyme Microb Technol* **51**: 300–309.
- 20 Costa, T.R., Felisberto-Rodrigues, C., Meir, A., Prevost, M.S., Redzej, A., Trokter, M., and
21 Waksman, G. (2015) Secretion systems in Gram-negative bacteria: structural and
22 mechanistic insights. *Nature Rev Microbiol* **13**: 343–359.

- 1 Coulthurst, S. (2019) The Type VI secretion system: a versatile bacterial weapon. *Microbiology*
2 **165**: 503-515.
- 3 Croes, C., Van Bastelaere, E., DeClercq, E., Eyers, M., Vanderleyden, J., and Michiels, K. (1991)
4 Identification and mapping of loci involved in motility, adsorption to wheat roots, colony
5 morphology, and growth in minimal medium on the *Azospirillum brasilense* Sp7 90-MDa
6 plasmid. *Plasmid* **26**: 83-93.
- 7 de-Bashan, L.E., and Bashan, Y. (2008) Joint immobilization of plant growth-promoting bacteria
8 and green microalgae in alginate beads as an experimental model for studying plant-
9 bacterium interactions. *Appl Environ Microbiol* **74**: 6797–6802.
- 10 de-Bashan, L.E., and Bashan, Y. (2010) Immobilized microalgae for removing pollutants: Review
11 of practical aspects. *Bioresour Technol* **101**: 1611–1627.
- 12 de-Bashan, L.E., Antoun, H., and Bashan, Y. (2008) Involvement of indole-3-acetic-acid produced
13 by the growth-promoting bacterium *Azospirillum* spp. in promoting growth of *Chlorella*
14 *vulgaris*. *J Phycol* **44**: 938–947.
- 15 de-Bashan, L.E., Hernandez, J.-P., and Bashan, Y. (2015) Interaction of *Azospirillum* spp. with
16 microalgae; a basic eukaryotic–prokaryotic model and its biotechnological applications. In
17 *Handbook for Azospirillum. Technical issues and protocols* (Cassán, F.D., Okon, Y. &
18 Creus, C.M., editors) 367–388. Springer International Publishing, Switzerland.
- 19 de-Bashan, L.E., Bashan, Y., Moreno, M., Lebsky, V.K., and Bustillos, J.J. (2002) Increased
20 pigment and lipid content, lipid variety, and cell and population size of the microalgae
21 *Chlorella* spp. when co-immobilized in alginate beads with the microalgae-growth-
22 promoting bacterium *Azospirillum brasilenses*. *Can J Microbiol* **8**: 514–552.

- de-Bashan, L.E., Hernandez, J.-P., Morey, T., and Bashan, Y. (2004) Microalgae growth-promoting bacteria as “helpers” for microalgae: a novel approach for removing ammonium and phosphorus from municipal wastewater. *Water Res* **38**: 466–474.
- de-Bashan, L.E., Mayali, X., Bebout, B.M., Weber, P.K., Detweiler, A., Hernandez, J.-P., Prufert-Bebout, L., and Bashan, Y. (2016) Establishment of stable synthetic mutualism without co-evolution between microalgae and bacteria demonstrated by mutual transfer of metabolites (NanoSIMS isotopic imaging) and persistent physical association (Fluorescent in situ hybridization). *Algal Res* **15**: 179–186.
- de-Bashan, L.E., Schmid, M., Rothballer, M., Hartmann, A., and Bashan, Y. (2011) Cell-cell interaction in the eukaryote-prokaryote model using the microalgae *Chlorella vulgaris* and the bacterium *Azospirillum brasilense* immobilized in polymer beads. *J Phycol* **47**: 1350–1359.
- Decoin, V., Barbey, C., Bergeau, D., Latour, X., Feuilloley, M.G., Orange, N., and Merieau, A. (2014) A type VI secretion system is involved in *Pseudomonas fluorescens* bacterial competition. *PLOS One* **9**: 89411.
- Decoin, V., Gallique, M., Barbey, C., Le Mauff, F., Poc, C.D., Feuilloley, M.G., Orange, N., and Merieau, A. (2015) A *Pseudomonas fluorescens* type 6 secretion system is related to mucoidy, motility and bacterial competition. *BMC Microbiol.* **15**: 72.
- de Pace, F., Nakazato, G., Pacheco, A., de Paiva, J.B., Sperandio, V., and da Silveira, W.D. (2010) The type VI secretion system plays a role in type 1 fimbria expression and pathogenesis of an avian pathogenic *Escherichia coli* strain. *Infect Immun* **78**: 4990-8.
- Díaz-Zorita, M., and Fernández-Canigia, M. (2009) Field performance of a liquid formulation of *Azospirillum brasilense* on dryland wheat productivity. *Eur J Soil Biol* **45**: 3–11.

- 1 Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A. and Smith, F. (1956). Colorimetric method
2 for determination of sugars and related substances. *Anal Chem* **28**: 350–356.
- 3 Duca, D., Lorv, J., Patten, C.L., Rose, D., and Glick, B.R. (2014) Indole-3-acetic acid in plant–
4 microbe interactions. *Antonie van Leeuwenhoek* **106**: 85–125.
- 5 Durán, D., Bernal, P., Vazquez-Arias, D., Blanco-Romero, E., Garrido-Sanz, D., Redondo-Nieto,
6 M., Rivilla, R., and Martín, M. (2021) *Pseudomonas fluorescens* F113 type VI secretion
7 systems mediate bacterial killing and adaption to the rhizosphere microbiome. *Sci Rep* **11**:
8 5772.
- 9 Durand, E., Cambillau, C., Cascales, E., and Journet, L. (2014) VgrG, Tae, Tle, and beyond: the
10 versatile arsenal of Type VI secretion effectors. *Trends Microbiol* **22**: 498–507.
- 11 Gafni, R., Okon, Y., Kapulnik, Y., and Fischer, M. (1986) Adsorption of *Azospirillum brasilense*
12 to corn roots. *Soil Biol Biochem* **18**: 69–75
- 13 Gallegos-Monterrosa, R., and Coulthurst, S.J. (2021) The ecological impact of a bacterial weapon:
14 microbial interactions and the Type VI secretion system. *FEMS Microbiol Rev* doi:
15 10.1093/femsre/fuab033.
- 16 Gallique, M., Decoin, V., Barbey, C., Rosay, T., Feuilloley, M.G., Orange, N., and Merieau, A.
17 (2017) Contribution of the *Pseudomonas fluorescens* MFE01 type VI secretion system to
18 biofilm formation. *PLoS One* **12**: e0170770.
- 19 Glickmann, E., and Dessaux, Y. (1995) A Critical Examination of the specificity of the Salkowski
20 Reagent for indolic compounds produced by phytopathogenic bacteria. *Appl Environ*
21 *Microbiol* **61**: 793–796.
- 22 Greer-Phillips, S.E., Stephens, B.B., and Alexandre, G. (2004) An energy taxis transducer
23 promotes root colonization by *Azospirillum brasilense*. *J Bacteriol* **186**: 6595–6604.

1 Hall, P.G., and Krieg, N.R. (1983) Swarming of *Azospirillum brasilense* on solid media. *Can J*
2 *Microbiol* **29**: 1592–1594.

3 Hernandez, J.-P., de-Bashan, L.E., Rodriguez, D.J., Rodriguez, Y., and Bashan Y. (2009) Growth
4 promotion of the freshwater microalga *Chlorella vulgaris* by the nitrogen-fixing, plant
5 growth-promoting bacterium *Bacillus pumilus* from arid zone soils. *Eur J Soil Biol* **45**: 88–
6 93.

7 Hernandez, R.E., Gallegos-Monterrosa, R., and Coulthurst, S.J. (2020) Type VI secretion system
8 effector proteins: Effective weapons for bacterial competitiveness. *Cell Microbiol* **22**:
9 e13241.

10 Hood, R.D., Singh, P., Hsu, F., Güvener, T., Carl, M.A., Trinidad, R.R., Silverman, J.M., Ohlson,
11 B.B., Hicks, K.G., Plemel, R.L., Li, M., Schwarz, S., Wang, W.Y., Merz, A.J., Goodlett,
12 D.R., and Mougous, J.D. (2010) A type VI secretion system of *Pseudomonas aeruginosa*
13 targets a toxin to bacteria. *Cell Host Microbe* **7**: 25-37.

14 Hou, X., McMillan, M., Coumans, J.V.F., Poljak, A., Raftery, M.J., and Pereg, L. (2014) Cellular
15 responses during morphological transformation in *Azospirillum brasilense* and its *flcA*
16 knockout mutant. *PLOS One* **9**: e114435.

17 Imase, M., Watanabe, K., Aoyagi, H., and Tanaka, H. (2008) Construction of an artificial
18 symbiotic community using a *Chlorella*-symbiont association as a model. *FEMS Microbiol*
19 *Ecol* **63**: 273–282.

20 Jani, A.J., and Cotter, P.A. (2010) Type VI secretion: not just for pathogenesis anymore. *Cell Host*
21 *Microbe* **8**: 2-6.

22 Jiménez-Guerrero, I., Cubo, M.T., Pérez-Montaña, F., López-Baena, F.J., Guash-Vidal, B., Ollero,
23 F.J., Bellogín, R., and Espuny, M.R. (2013) Bacterial protein secretion systems.

Implications in beneficial associations with plants. In *Beneficial Plant-microbial Interactions, Ecology and Applications* (González-López, J., ed) 183–213. CRC Press, Boca Raton, FL.

Jofre, E., Lagares, A., and Mori, G. (2004) Disruption of dTDP-rhamnose biosynthesis modifies lipopolysaccharide core, exopolysaccharide production, and root colonization in *Azospirillum brasilense*. *FEMS Microbiol Lett* **231**: 267–275.

Jurénas, D., and Journet, L. (2021) Activity, delivery, and diversity of Type VI secretion effectors. *Mol Microbiol* **115**: 383-394.

Kanehisa, M., Goto, M., Sato, Y., Furumichi, M., and Tanabe, M. (2012) KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acid Res* 40: D109–D114.

Kobayashi, K. (2021) Diverse LXG toxin and antitoxin systems specifically mediate intraspecies competition in *Bacillus subtilis* biofilms. *PLoS Genet* **17**: e1009682.

Konovalova, A., Petters, T., and Søgaard-Andersen, L. (2010) Extracellular biology of *Myxococcus xanthus*. *FEMS Microbiol Rev* **34**: 89-106.

Lebsky, V.K., Gonzalez-Bashan, L.E., and Bashan, Y. (2001) Ultrastructure of interaction in alginate beads between the microalga *Chlorella vulgaris* with its natural associative bacterium *Phyllobacterium myrsinacearum* and with the plant growth-promoting bacterium *Azospirillum brasilense*. *Can J Microbiol* **47**: 1–8.

Lerner, A., Castro-Sowinski, S., Valverde, A., Lerner, H., Dror, R., Okon, Y., and Burdman, S. (2009) The *Azospirillum brasilense* Sp7 *noeJ* and *noeL* genes are involved in extracellular polysaccharide biosynthesis. *Microbiology* **155**: 4058–4068.

- 1 Levanony, H., Bashan, Y., Romano, B., and Klein, E. (1989) Ultrastructural localization and
2 identification of *Azospirillum brasilense* Cd on and within wheat root by immuno-gold
3 labeling. *Plant Soil* **117**: 207–218.
- 4 Leyva, L.A., Bashan, Y., Mendoza, A., and de-Bashan, L.E. (2014) Accumulation of fatty acids
5 in *Chlorella vulgaris* under heterotrophic conditions in relation to activity of acetyl-CoA
6 carboxylase, temperature, and co-immobilization with *Azospirillum brasilense*.
7 *Naturwissenschaften* **101**: 819–830.
- 8 Li, J., Yao, Y., Xu, H.H., Hao, L., Deng, Z., Rajakumar, K., and Ou, H.Y. (2015) SecReT6: a web-
9 based resource for type VI secretion systems found in bacteria. *Environ Microbiol* **17**:
10 2196-202.
- 11 Ma, L.S., Hachani, A., Lin, J.S., Filloux, A., and Lai, E.M. (2014) *Agrobacterium tumefaciens*
12 deploys a superfamily of type VI secretion DNase effectors as weapons for interbacterial
13 competition in planta. *Cell Host Microbe* **16**: 94-104.
- 14 Madi, L. and Henis, Y. (1989). Aggregation in *Azospirillum brasilense* Cd: conditions and factors
15 involved in cell-to-cell adhesion. *Plant Soil* **115**: 89-98.
- 16 Masuko, T., Minami, A., Iwasaki, N., Majima, T., Nishimura, S.I. and Lee, Y.C. (2005).
17 Carbohydrate analysis by a phenol–sulfuric acid method in microplate format. *Anal*
18 *Biochem* **339**: 69–72.
- 19 Michiels, K., De Troch, P., Onyeocha, I., Van Gool, A., Elmerich, C., and Vanderleyden, J. (1989)
20 Plasmid localization and mapping of two *Azospirillum brasilense* loci that affect
21 exopolysaccharide synthesis. *Plasmid* **21**: 142-6.
- 22 Oh-Hama, T., and Miyachi, S. (1992) *Chlorella*. In Micro-algae Biotechnology (Borowitzka, M.A.
23 & Borowitzka, L.J., eds) 3–26. Cambridge University Press, Cambridge UK.

- O'Toole, G., and Kolter, R. (1998) Initiation of biofilm formation in *Pseudomonas fluorescens* WCS 365 proceeds via multiple, convergent signaling pathways: a genetic analysis. *Mol Microbiol* **28**: 449–461.
- Palacios, O.A., Choix, F.J., Bashan, Y., and de-Bashan, L.E. (2016a) Influence of tryptophan and indole-3-acetic acid on starch accumulation in the synthetic mutualistic *Chlorella sorokiniana*-*Azospirillum brasilense* system under heterotrophic conditions. *Res Microbiol* **167**: 367–379.
- Palacios, O.A., Gomez-Anduro, G., Bashan, Y., and de-Bashan, L.E. (2016b) Tryptophan, thiamine, and indole-3-acetic acid exchange between *Chlorella sorokiniana* and the plant growth-promoting bacterium *Azospirillum brasilense*. *FEMS Microbiol Ecol* **92**: fiw077
- Palacios, O.A., Lopez, B.R., Bashan, Y., and de-Bashan, L.E. (2019) Early changes in nutritional conditions affect formation of synthetic mutualism between *Chlorella sorokiniana* and the bacterium *Azospirillum brasilense*. *Microb Ecol* **77**: 980-992.
- Pande, S.V., Parvin, R.K., and Venkitasubramanian, T.A. (1963) Microdetermination of lipids and serum total fatty acids. *Anal Biochem* **6**: 415–423.
- Peng, H., de-Bashan, L.E., Bashan, Y., and Higgins, B.T. (2020) Indole-3-acetic acid from *Azospirillum brasilense* promotes growth in green algae at the expense of energy storage products. *Algal Res* **47**:101845
- Pereg, L., de-Bashan, L.E., and Bashan, Y. (2016) Assessment of affinity and specificity of *Azospirillum* for plants. *Plant Soil* **399**: 389–414.
- Pereg Gerk, L., Paquelin, A., Gounon, P., Kennedy, I.R., and Elmerich, C. (1998) A transcriptional regulator of the LuxR-UhpA family, FlcA, controls flocculation and wheat root surface colonisation by *Azospirillum brasilense* Sp7. *Mol Plant Microbe Inter* **11**: 177–187.

- 1 Puente, M.E., Holguin, G., Glick, B.R., and Bashan Y. (1999) Root surface colonization of black
2 mangrove seedlings by *Azospirillum halofraeference* and *Azospirillum brasilense* in
3 seawater. *FEMS Microbiol Ecol* **29**: 283–292.
- 4 Rivera, D. *et al.* (17 co-authors) (2014) Complete genome sequence of the model rhizosphere strain
5 *Azospirillum brasilense* Az39, successfully applied in agriculture. *Genome Ann* **2**: e00683-
6 14.
- 7 Rivera, D., Mora, V., Lopez, G., Rosas, S., Spaepen, S., Vanderleyden, J. and Cassan, F. (2018).
8 New insights into indole-3-acetic acid metabolism in *Azospirillum brasilense*. *J Appl*
9 *Microbiol* **125**: 1774-1785.
- 10 Roest, H.P., Mulders, I.H.M., Spaink, H.P., Wijffelman, C.A., and Lugtenberg, B.J.J. (1997) A
11 *Rhizobium leguminosarum* biovar *trifolii* locus not localized on the sym plasmid hinders
12 effective nodulation on plants of the pea cross-inoculation group. *Mol Plant Microbe Inter*
13 **7**: 938–941.
- 14 Russell, A.B., Peterson, S.B., and Mougous, J.D. (2014) Type VI secretion system effectors:
15 poisons with a purpose. *Nature Rev Microbiol* **12**: 137–148.
- 16 Ryu, C.-M. (2015) Against friend and foe: Type 6 effectors in plant-associated bacteria. *J*
17 *Microbiol* **53**: 201–208.
- 18 Sana, T.G., Flaugnatti, N., Lugo, K.A., Lam, L.H., Jacobson, A., Baylot, V., Durand, E., Journet,
19 L., Cascales, E., and Monack, D.M. (2016) *Salmonella Typhimurium* utilizes a T6SS-
20 mediated antibacterial weapon to establish in the host gut. *Proc Natl Acad Sci U S A*. **113**:
21 E5044-51.
- 22 Schäfer, A., Tauch, A., Jäger, W., Kalinowski, J., Thierbach, G., and Pühler, A. (1994) Small
23 mobilizable multi-purpose cloning vectors derived from the *Escherichia coli* plasmids

pK18 and pK19: selection of defined deletions in the chromosome of *Corynebacterium glutamicum*. *Gene* **145**: 69–73.

Simon, R., Priefer, U., and Puhler, A. (1983) A broad host range mobilization system for *in vivo* genetic-engineering - transposon mutagenesis in gram-negative bacteria. *Bio/Technology* **1**: 784–91.

Spaepen, S., and Vanderleyden, J. (2011) Auxin and plant-microbe interactions. *Cold Spring Harbor Perspectives in Biology*, doi: 10.1101/schperspect.a001438.

Trunk, K., Coulthurst, S.J., and Quinn, J. (2019) A new front in microbial warfare - delivery of antifungal effectors by the type VI secretion system. *J Fungi* (Basel). **5**: 50.

Van Puyvelde, S., Cloots, L., Engelen, K., Das, F., Marchal, K., Vanderleyden, J., and Spaepen, S. (2011) Transcriptome analysis of the rhizosphere bacterium *Azospirillum brasilense* reveals an extensive auxin response. *Microb Ecol* **61**: 723–728.

Vanstockem, M., Michiels, K., Vanderleyden, J., and Van Gool, A. (1987) Transposon mutagenesis of *Azospirillum brasilense* and *Azospirillum lipoferum*: physiological analysis of Tn5 and Tn5-mob insertional mutants. *Appl Environ Microbiol* **53**: 1387–1405.

Vidussi, F., Claustre, H., Bustillos-Guzman, J., Cailliau, C., and Marty, J.C. (1996) Determination of chlorophylls and carotenoids of marine phytoplankton: separation of chlorophyll a from divinyl-chlorophyll a and zeaxanthin from lutein. *J Plankton Res* **18**: 2377–2382.

Wang, J., Li, J., Hou, Y., Dai, W., Xie, R., Marquez-Lago, T.T., Leier, A., Zhou, T., Torres, V., Hay, I., Stubenrauch, C., Zhang, Y., Song, J., and Lithgow, T. (2021) BastionHub: a universal platform for integrating and analyzing substrates secreted by Gram-negative bacteria. *Nucleic Acids Res* **49**: D651-D659.

- Wexler, A.G., Bao, Y., Whitney, J.C., Bobay, L.M., Xavier, J.B., Schofield, W.B., Barry, N.A., Russell, A.B., Tran, B.Q., Goo, Y.A., Goodlett, D.R., Ochman, H., Mougous, J.D., and Goodman, A.L. (2016) Human symbionts inject and neutralize antibacterial toxins to persist in the gut. *Proc Natl Acad Sci USA* **113**: 3639-44.
- Wisniewski-Dyé, F. *et al.* (with 25 co-authors) (2011) *Azospirillum* genomes reveal transition of bacteria from aquatic to terrestrial environments. *PLOS Genetics* **7**: e1002430.
- Wood, T.E., Aksoy, E., and Hachani, A. (2020) From welfare to warfare: the arbitration of host-microbiota interplay by the type VI secretion system. *Front Cell Infect Microbiol* **10**: 587948.
- Wu, C.F., Santos, M.N.M., Cho, S.T., Chang, H.H., Tsai, Y.M., Smith, D.A., Kuo, C.H., Chang, J.H., and Lai, E.M. (2019) Plant-pathogenic *Agrobacterium tumefaciens* strains have diverse type VI effector-immunity pairs and vary in in-planta competitiveness. *Mol Plant Microbe Interact* **32**: 961-971.
- Wu, C.F., Smith, D.A., Lai, E.M., and Chang, J.H. (2018) The *Agrobacterium* type VI secretion system: a contractile nanomachine for interbacterial competition. *Curr Top Microbiol Immunol* **418**: 215-231.
- Zuber, S., Carruthers, F., Keel, C., Mattart, A., Blumer, C., Pessi, G., Gigot-Bonnefoy, C., Schnider-Keel, U., Heeb, S., Reimmann, C., and Haas, D. (2003) GacS sensor domains pertinent to the regulation of exoproduct formation and to the biocontrol potential of *Pseudomonas fluorescens* CHA0. *Mol Plant Microbe Inter* **16**: 634–644.
- Zoued, A., Brunet, Y.R., Durand, E., Aschtgen, M.S., Logger, L., Douzi, B., Journet, L., Cambillau, C., and Cascales E. (2014) Architecture and assembly of the Type VI secretion system. *Bioch Biophys Acta* **1843**:1664–1673.

1

2

Table 1. Aggregation percentage of *A. brasilense* Az39 wild-type and its isogenic Az39 Δ hcp-*E* mutant.

Cell aggregation was measured after growth for 48 h in LB or MMAB culture medium. The reduction of the aggregation percentage in the Az39 Δ hcp-*E* relative to its parental Az39 strain is indicated below.

* Different letters differ significantly at P<0.05 (Tukey's post-hoc test).

Strain	Aggregation percentage	
	LB	MMAB
<i>A. brasilense</i> Az39	13.2 \pm 2.6 c	26.5 \pm 1.4 a
<i>A. brasilense</i> Az39 Δ hcp- <i>E</i>	8.5 \pm 1.1 d	18.2 \pm 0.9 b
Reduction of aggregation activity (Az39/Az39 Δ hcp- <i>E</i>)	35.6%	31.3%

Figure legends

Fig. 1. Schematic representation of the *Azospirillum brasilense* Az39 T6SS1 gene cluster.

T6SS core components are shown in green, while accessory genes are shown in white. Potential effector and immunity genes are shown in blue. The position of the *hcp* and *tssE* gene deletion used in this work is indicated

Fig. 2. Bacterial production of auxins. Expressed as concentration of IAA-like molecules ($\mu\text{g}\cdot\text{mL}^{-1}$) in MMAB cultures of *A. brasilense* Az39 (●) and Az39 Δ *hcp*-E mutant (■). Pair of values at each sampling time sharing the same letter do not differ significantly from each other by Student's *t*-test at $P < 0.05$.

Fig. 3. Swimming and swarming motility, biofilm formation and cell aggregation. Swimming (a) and swarming (b) rates, of *A. brasilense* Az39 and its isogenic Az39 Δ *hcp*-E mutant after 72 h incubation at $28 \pm 2^\circ\text{C}$ in MMAB or Swim media. The bars represent the mean and SD of growth in diameter (cm) of the swimming and swarming displacement halos, respectively. The percentage of the difference between the mutant and its parental strain is shown on top of the bars. (c) Biofilm production of Az39 and Az39 Δ *hcp*-E after 96-h incubation at $28 \pm 2^\circ\text{C}$ in LB medium. The bars represent the mean and standard deviation of absorbance at OD_{560nm}. The percentage of the difference between the mutant and its parental strain is shown on top of the bars. (d) Cell aggregation was measured after growth for 48 h in LB or MMAB culture medium. The reduction of the aggregation percentage in the mutant relative to its parental strain is indicated on top of the

bars. Each pair of columns denoted by a different letter differs significantly from each other by Student's *t*-test at $P < 0.05$.

Fig. 4. Antibacterial competition assay. The indicated nalidixic acid-resistant strains were mixed with wild-type or $\Delta hcp-E$ *A. brasilense* attacker cells in presence or not of the T6SS gene cluster inducer IAA. The number of surviving recipient cells, counted on nalidix acid plates, is indicated (expressed as \log^{10} of CFU).


Fig. 5. Effect of co-immobilization of *A. brasilense* strains on *C. sorokiniana* growth and cell aggregation. (a) The *C. sorokiniana* cell concentration (expressed as 10^5 cells.mL⁻¹) from cultures of the microalgae alone (○) or grown in presence of *A. brasilense* wild-type (■) or $\Delta hcp-E$ mutant (▲) cells is plotted against time (in days). Cell concentrations along curves having different capital letters differ significantly, using one-way ANOVA and Tukey's post hoc analysis at $P < 0.05$. Cell concentrations at each sampling time that are denoted by different lower-case letters differ significantly, using one-way ANOVA and Tukey's post hoc analysis at $P < 0.05$. Whisker lines represent standard errors (SE). Absence of a whisker line indicates negligible SE. μ = growth rate. (b) Cell aggregation was observed by fluorescence in situ hybridization (FISH) at time 0, 1, 2 and 4 days. (a-d) *A. brasilense* Az39 and *C. sorokiniana* 2714. (e-h) *A. brasilense* Az39 $\Delta hcp-E$ and *C. sorokiniana* 2714. (i-l) *A. brasilense* Az39 alone. (m-p) *A. brasilense* Az39 $\Delta hcp-E$ alone. (q-t) *C. sorokiniana* alone. All micrographs were taken by epifluorescence microscopy, *C. sorokiniana* was not labeled and appeared in red to orange color, while bacterial strains were labeled with specific probes targeting eubacteria (EUB338Mix FITC) and specific probe for *A. brasilense* (Abrax 1420 Cy3) appeared in green to yellow. (c) Size of aggregates created during the





interaction, measured from the results obtained by fluorescence *in situ* hybridization (FISH). Treatments analyzed at each hour denoted by different lowercase letter differ significantly at $P < 0.05$ in one-way analysis of variance, according to Tukey's post hoc analysis. Different capital letter at each day of incubation differs significantly at $P < 0.05$ in Tukey's post hoc analysis.


Fig. 6. Effect of *A. brasilense* co-immobilization on *C. sorokiniana* carbohydrates, lipids, and photosynthetic pigments production. Effect of co-immobilization of wild-type *A. brasilense* Az39 and its isogenic Az39 Δ *hcp-E* mutant on accumulation of (a) carbohydrates, (b) total lipids, (c) chlorophyll *a*, (d) chlorophyll *b*, (e) violaxanthin, and (f) lutein in *Chlorella sorokiniana*. Significant differences (one-way ANOVA and Fisher's post hoc analysis at $P < 0.05$) are indicated by different capital letters on top of each columns. Whisker lines represent SE.



Fig. 7. Schematic representation of the roles of *A. brasilense* T6SS1. The *A. brasilense* T6SS1 is involved in *Azospirillum*/microalgae, *Azospirillum*/*Azospirillum* and *Azospirillum*/bacteria interactions. *Azospirillum*/microalgae interaction: the *A. brasilense* (orange cells) T6SS is required for efficient attachment to the microalgae *C. sorokiniana* (green cell) and participates to the increase of the microalgae population and carbohydrates and photosynthetic pigments production. *Azospirillum*/*Azospirillum* interaction: the T6SS is required for biofilm formation and indole-3-acetic acid (IAA, purple triangles) production. IAA positively regulates the expression of the *A. brasilense* T6SS1 gene cluster. *Azospirillum*/bacteria interactions: *A. brasilense* uses its T6SS to eliminate bacteria sharing the same niche (blue cell). By targeting plant pathogens, it confers protection to plants and microalgae.

















c


Treatments	Aggregate size (μm ²) *		
	1 day	2 day	4 day
<i>C. sorokiniana</i>	8.96 ± 1.28 aA	44.80 ± 3.91 bA	94.27 ± 7.37 cA
<i>C. sorokiniana</i> + <i>A. brasilense</i> Az39	16.99 ± 1.70 aC	42.31 ± 8.49 bA	170.44 ± 24.01 cB
<i>C. sorokiniana</i> + <i>A. brasilense</i> Az39Δhcp-E	11.22 ± 0.68 aA	39.50 ± 2.28 bA	121.89 ± 2.89 cC
<i>A. brasilense</i> Az39	4.13 ± 0.67 aB	26.40 ± 1.75 bB	36.80 ± 2.29 cD
<i>A. brasilense</i> Az39Δhcp-E	3.41 ± 0.53 aB	16.00 ± 1.22 bC	19.50 ± 2.42 bE



■ *C. sorokiniana* + *A. brasilense* Az39Δhcp-E
 ■ *C. sorokiniana* + *A. brasilense* AZ39
 ▨ *C. sorokiniana*

Incubation time (D)

Azospirillum brasilense



Chlorella sorokiniana



T6SS



IAA



T6SS effector