



Mass spectrometry-based top-down and bottom-up approaches for proteomic analysis of the Moroccan *Buthus occitanus* scorpion venom

Khadija Daoudi, Christian Malosse, Ayoub Lafnune, Bouchra Darkaoui, Salma Chakir, Jean-Marc Sabatier, Julia Chamot-Rooke, Rachida Cadi, Naoual Oukkache

► To cite this version:

Khadija Daoudi, Christian Malosse, Ayoub Lafnune, Bouchra Darkaoui, Salma Chakir, et al.. Mass spectrometry-based top-down and bottom-up approaches for proteomic analysis of the Moroccan *Buthus occitanus* scorpion venom. FEBS Open Bio, 2021, 11 (7), pp.1867-1892. 10.1002/2211-5463.13143 . hal-03358052

HAL Id: hal-03358052

<https://amu.hal.science/hal-03358052>

Submitted on 23 Nov 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Mass spectrometry-based top-down and bottom-up approaches for proteomic analysis of the Moroccan *Buthus occitanus* scorpion venom

Khadija Daoudi^{1,2} , Christian Malosse³, Ayoub Lafnoute^{1,2}, Bouchra Darkaoui^{1,2}, Salma Chakir¹, Jean-Marc Sabatier⁴, Julia Chamot-Rooke³, Rachida Cadi² and Naoual Oukkache¹

¹ Laboratory of Venoms and Toxins, Pasteur Institute of Morocco, Casablanca, Morocco

² Laboratory of Molecular Genetics, Physiopathology and Biotechnology, Faculty of Sciences Ain Chock, Hassan II University of Casablanca, Morocco

³ Mass spectrometry for Biology Unit, Institut Pasteur, CNRS USR 2000, Paris, France

⁴ Laboratory INSERM UMR 1097, University of Aix Marseille, France

Keywords

bottom-up; *Buthus occitanus* scorpion; top-down; toxins; venom; venomous

Correspondence

K. Daoudi, Laboratory of Venoms and Toxins, Pasteur Institute of Morocco, 1 Place Louis Pasteur, Casablanca 20250, Morocco.

Laboratory of Molecular Genetics, Physiopathology and Biotechnology, Faculty of Sciences Ain Chock, Hassan II University of Casablanca, B.P 5366 Maarif, Casablanca, Morocco.

E-mail: khadija.daoudi1-etu@etu.univh2c.ma

(Received 29 December 2020, revised 18 February 2021, accepted 11 March 2021)

doi:10.1002/2211-5463.13143

Edited by Alberto Alape-Girón

Buthus occitanus (*B. occitanus*) is one of the most dangerous scorpions in the world. Despite the involvement of *B. occitanus* scorpion in severe cases of envenomation in Morocco, no study has focused yet on the proteomic composition of the Moroccan *B. occitanus* scorpion venom. Mass spectrometry-based proteomic techniques are commonly used in the study of scorpion venoms. The implementation of top-down and bottom-up approaches for proteomic analyses facilitates screening by allowing a global view of the structural aspects of such complex matrices. Here, we provide a partial overview of the venom of *B. occitanus* scorpion, in order to explore the diversity of its toxins and hereafter understand their effects. To this end, a combination of top-down and bottom-up approaches was applied using nano-high liquid chromatography coupled to nano-electrospray tandem mass spectrometry (nano-LC-ESI MS/MS). The LC-MS results showed that *B. occitanus* venom contains around 200 molecular masses ranging from 1868 to 16 720 Da, the most representative of which are those between 5000 and 8000 Da. Interestingly, combined top-down and bottom-up LC-MS/MS results allowed the identification of several toxins, which were mainly those acting on ion channels, including those targeting sodium (NaScTx), potassium (KScTx), chloride (ClScTx), and calcium channels (CaScTx), as well as antimicrobial peptides (AMPs), amphipathic peptides, myotropic neuropeptides, and hypothetical secreted proteins. This study reveals the molecular diversity of *B. occitanus* scorpion venom and identifies components that may have useful pharmacological activities.

Abbreviations

ACN, acetonitrile; AMP, antimicrobial peptides; *B. occitanus*, *Buthus occitanus*; CaScTx, neurotoxins affecting calcium channels; ClScTx, neurotoxins affecting chloride channels; Da, Dalton; EThcD, Electron-Transfer/Higher-Energy Collision Dissociation; FA, formic acid; HCD, higher-energy C-trap dissociation; IAA, iodoacetamide; kDa, kilodalton; KScTx, neurotoxins affecting potassium channels; LC-MS/MS, liquid chromatography coupled to tandem mass spectrometry; LC-MS, liquid chromatography coupled to mass spectrometry; MS, mass spectrometry; MW, molecular weight; nano-LC-ESI MS/MS, nano-liquid chromatography coupled to electrospray tandem mass spectrometry; NaScTx, neurotoxins affecting sodium channels; Q, quadrupole; TIC, total ion chromatogram.

Each year, scorpion stings record new cases of envenomation over the world with an incidence of more than 1.5 million and over 2600 deaths, mainly in tropical and subtropical countries of South America, Asia, and North Africa [1]. Most of these envenomation cases were caused by scorpions belonging to the Buthidae family, which contains dangerous species known by their lethal venoms [2]. The venom of these family members contains a heterogeneous cocktail of compounds, including inorganic substances, enzymes, mucopolysaccharides, allergenic compounds, and peptides with high toxicity toward ionic channels of excitable cells [3–6]. In Morocco, 26 819 cases of scorpion stings were reported in 2019 by the Poison Control and Pharmacovigilance Center of Morocco, with an incidence of 75.3 cases per 100 000 inhabitants [7]. These statistics are due to the diversified scorpion fauna represented by over 50 species, mainly widespread in the middle and southwestern provinces of the kingdom [8]. Among these species, the yellow scorpion *Buthus occitanus* (*B.occitanus*) seems to be one of the most dangerous scorpions, on account of its toxic venom causing the majority of envenomation cases [9]. Although several studies had been carried out on this venom [10–13], no study has yet focused on the proteomic composition of the Moroccan *B. occitanus* scorpion venom despite its medical importance. Moreover, there are various strategies to screen scorpion venoms, from using conventional strategies for targeting one single toxin, to applying the most throughput equipment of screening for a detailed view of all toxic components. Nowadays, mass spectrometry-based proteomic approaches are still one of the most fundamental tools to decrypt the complexity of such matrices, owing to the revolutionary advances in instrumentation and software, in addition to improvement in omics strategies (peptidomic, proteomic, transcriptomic, and genomic) [14–19]. Among the approaches that have improved significantly the proteomics workflow, there are the top-down process, which designates a rapid analytical workflow of intact proteins, and the bottom-up approach, which requires prior proteolytic digestion of proteins before mass spectrometry analysis. These approaches lead to acquiring mass fingerprints, primary structural information, and post-translational modifications [20–23]. The application of these approaches, singly or complementary, in several proteomic studies has increased the number of characterized venoms and identified toxins [24–29]. In this context, this work aimed to ensure an overview of the peptidome of *B. occitanus* scorpion (< 30 kDa), so exploring its toxins arsenal, using a combination of the top-down and bottom-up approaches applied on nano-

high liquid chromatography coupled to a nano-electrospray tandem mass spectrometry (nano-LC-ESI MS/MS).

Materials and methods

Venom preparation

Venom milking

Specimens of *B. occitanus* were collected from the region of Oualidia (32°44'N 9°01'W), in eastern Morocco. The crude venom was milked by electrical stimulation, pooled, centrifuged at 10 000 *g* for 20 min, freeze-dried, and stored at –20 °C until use [30].

Venom Reduction/Alkylation

At first, 2 mg of *B. occitanus* crude venom was subjected to a 30 kDa ‘cutoff’ filter (Amicon® Ultra Centrifugal Filters, Merck Millipore, Tulagreen, Ireland), then centrifuged at 16 900 *g* for 15 min.

Disulfide-bridged half-cysteine residues of this venom filtrate were reduced by 10 mM of DTT in ammonium bicarbonate buffer (50 mM, pH 8.3), for 45 min at a temperature of 56 °C. Cysteine residues were carboxamidomethylated by incubation with 50 mM iodoacetamide [IAA in ammonium bicarbonate (50 mM, pH 8.3)] for 1 h in the dark. Then, these proteins/peptides were desalted by ZipTip C4 (Millipore Corporation - Billerica, USA) and concentrated on a Savant SpeedVac (Thermo Scientific, San Jose, CA, USA).

Mass spectrometry-based proteomic approaches

Top-down proteomics

Intact and reduced/alkylated *B. occitanus* venom filtrates were carried out on an Orbitrap Fusion™ Lumos™ mass spectrometer (Thermo Scientific™ Waltham, MA, USA), equipped with a Dionex HPLC (Fig. 1).

For the online peptide fractionation, 2 µg of samples was loaded to a C4 µ-precolumn cartridge (300 µm i.d. × 5 mm, C4 PepMap 300 particles with 5 µm size and 300 Å pores); the column was equilibrated with solution A [0.1% (v/v) formic acid (FA)]. The separation was maintained over 120 min at 250 nL·min^{–1}, using a linear gradient from 5% to 60% of solution B [acetonitrile (ACN) and 0.1% (v/v) FA].

Proteins/peptides were eluted directly from the column into the mass spectrometer and operated in positive mode with a spray voltage of 1.6 kV. MS spectra were acquired at a resolution setting of 120 000.

MS/MS analysis was performed on data-dependent acquisition, the top 10 abundant precursor ions were




Fig. 1. Experimental workflow performed in this study. At first, *B. occitanus* venom was milked by electrical stimulation and applied to a 30 kDa filter. For the top-down venomics, the flow-through containing toxins < 30 kDa was analyzed by the Thermo Scientific™ Orbitrap Fusion Lumos Tribrid Mass Spectrometer. For the bottom-up approach, two digest methods were achieved: 1) in-solution digestion, the flow-through containing toxin < 30 kDa was directly reduced with DTT, alkylated with IAA, and digested with trypsin; and 2) in-gel digestion, the unstained gel was excised to small cubes, reduced, alkylated, and digested. The digest peptides were then desalted with ZipTip and applied to the Orbitrap Q-Exactive mass spectrometer.

selected for an EThcD fragmentations (Electron-Transfer/Higher-Energy Collision Dissociation) with a dynamic exclusion time of 90 s. MS/MS spectra were acquired at a resolution setting of 120 000, and the mass range was set from 150 to 2000 m/z .

Bottom-up proteomics

In-solution digestion

Reduced/alkylated venom filtrate was digested overnight at a temperature of 37 °C with 0.1 µg of trypsin (Promega, Madison, WI, USA). Tryptic digests were analyzed on a Q-Exactive Plus instrument (Thermo Fisher Scientific, Bremen, Germany) coupled to an EASY-nLC 1200 chromatography system (Thermo Fisher Scientific). Two micrograms was loaded on an in-house packed 50-cm nano-HPLC column (75 µm inner diameter) filled with C18 resin (1.9 µm particles, 100 Å pore size, Reprosil-Pur Basic C18-HD resin; Maisch GmbH, Ammerbuch-Entringen, Germany) and equilibrated in 97% solvent A and 3% solvent B (ACN, 0.1% (v/v) FA).

Peptides were eluted at 250 nL·min⁻¹, using 3–22% gradient of solvent B for 112 min, then 22–38% gradient of

solvent B for 35 min, and finally 38–60% gradient of solvent B for 15 min. The instrument method for the Q-Exactive Plus was set up in the data-dependent acquisition mode. MS and MS-MS spectra were acquired at a resolution of 60 000, 10 of the most abundant precursor ions were selected for HCD fragmentation with collision energy adjusted to 27. Mono-charged precursors and those with a charge state of > 7 were excluded.

In-gel digestion

At first, 2 mg of venom filtrate was unfolded for 5 min at 95 °C in sample buffer (LDS sample buffer) and then subjected to a SDS/PAGE using a 4–20% of polyacrylamide gel (SDS Precast Gel RunBlue, 4–20%, 12 well; Expedeon, CA, USA). The electrophoresis was performed, on a Bio-Rad system, at a constant voltage of 140 V, and the separated proteins were stained with Coomassie Brilliant Blue R (InstantBlue; Expedeon, CA, USA).

Stained bands corresponding to proteins/peptides with masses < 30 kDa (Fig. S1) were manually excised into equal small cubes of 1 mm³, then washed with Milli-Q water, ammonium bicarbonate 50 mM, and ACN 50%.

Subsequently, the slices were submitted to an in-gel reduction with DTT (10 mM) in ammonium bicarbonate buffer (50 mM, pH 8.3) for 45 min at a temperature of 56 °C. Reduced slices were alkylated with IAA (50 mM) in ammonium bicarbonate (50 mM, pH 8.3) buffer for 20 min in the dark, followed by an overnight digestion with 0.1 µg of trypsin (Promega) at a temperature of 37 °C [31]. The enzymatic reaction was stopped by adding 5 µL of FA 5%, and desalted by loading the peptides onto ZipTip C18. After drying, digested peptides were dissolved in 100 µL of 0.1% (v/v) FA and applied on a liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS) system, composed of a nano-flow HPLC pump and an Orbitrap Q-Exactive mass spectrometer (Thermo Scientific) with a nano-electrospray ion source, as described in the section above.

Data analysis

The top-down liquid chromatography coupled to mass spectrometry (LC-MS) data analysis of native *B. occitanus* venom filtrate was deconvoluted using the Xtract algorithm within Thermo Scientific XCALIBUR 2.2 software (Thermo Fisher Scientific).

For protein identification, data from both of the venom nano-LC-MS/MS approaches were processed using the PROTEOME DISCOVER 2.2 software (Thermo Fisher Scientific), against the UniProtKB database, downloaded in 2016 10 11, taxon identifier: 6855 and 4309 entries.

Parameters of processing were as follows: a mass tolerance of MS set at 50 p.p.m. and 0.3 Da for MS/MS. One unique peptide was required for protein identification, minimum peptide length was required at five amino acids, and the false discovery rate cutoff was 1%. Trypsin was chosen

as the specific enzyme, with a maximum number of two missed cleavages for the bottom-up analysis. Variable modifications included oxidation of methionine and carbamidomethylation, while no fixed modification was set.

Results

Mass spectrometry-based proteomic approaches

The whole proteomic approaches are based only on the UniProtKB database-dependent analysis without any manually *de novo* sequence annotation; therefore, the majority of reported peptide annotations are still an approximation. Also, it is important to stress that the relative abundances and the percentages of the described peptides are purely based on total number counts and not concentrations as long as no quantitative analysis was performed.

Top-down proteomics

The total ion chromatogram (TIC) generated from the top-down LC-MS analysis of native *B. occitanus* venom filtrate (Fig. 2) gave a partial picture of the venom complexity, with around 60 peaks, most of them detected with high relative abundance.

The mass fingerprint of *B. occitanus* venom was generated from a manual deconvolution of spectra gained by top-down LC-MS approach, thus detecting a total of 197 monoisotopic masses ranging from 1868 to 16 720 Da (Table 1). We get one mass less than 2000 Da, 28 molecular masses ranging between 2000 and 5000 Da, 147 mass values from 5000 to 8000 Da, and 21 masses for those over 8000 Da.





Fig. 2. TIC of native *B. occitanus* venom filtrate, generated from top-down mass spectrometry analysis (MS1). The x-axis represents the relative abundance (%), and the y-axis, the retention time (min). Spectra were deconvoluted, and generated monoisotopic masses were distributed according to their MW.

Table 1. List of the 197 monoisotopic masses detected by the top-down LC-MS analysis.

Retention time (min)	MW (Da)
0–10	N.D
10–20	1868.0157
20–30	2208.2634; 2506.4634
30–40	2813.4212; 2851.4287; 2966.3848; 3124.4545; 3219.5691; 3233.4756; 3461.4966; 3486.7774; 3538.283; 3550.4334; 3670.8935; 3718.7023; 3823.4412; 3937.8078; 4093.8732; 4321.8654; 4366.9752; 4366.986; 5731.6152; 5919.5155.
40–50	3522.2898; 3614.8741; 3807.4466; 3937.7725; 4333.933; 4366.9856; 4568.7172; 4572.9253; 5185.3781; 6148.8879; 6423.7104; 6439.6786; 6527.7246; 6539.6502; 6541.7326; 6595.7719; 6606.8166; 6610.768; 6611.7946; 6635.0442; 6744.712; 6829.8098; 6831.8926; 6832.876; 6860.9183; 6861.9012; 6872.9404; 6876.9037; 6877.9284; 6893.9821; 6940.948; 6952.1809; 6974.2357; 6979.0052; 6995.0399; 6997.024; 7014.2508; 7016.0204; 7022.0148; 7024.0653; 7107.2902; 7152.0763; 7162.3796; 7177.1647; 7218.3026; 7220.0387; 7220.2052; 7243.2414; 7297.2395; 7393.2604.
50–60	6488.9021; 6609.8127; 6611.7977; 6629.8447; 6677.8651; 6749.8876; 6765.9533; 6779.2433; 6807.922; 6823.1194; 6836.974; 6837.8837; 6862.9698; 6879.9966; 6907.3347; 6919.9628; 6972.7789; 7007.0404; 7011.1444; 7012.1231; 7020.055; 7028.0976; 7035.2491; 7024.1049; 7051.0799; 7061.1245; 7062.1114; 7069.1111; 7079.1299; 7082.3444; 7115.0302; 7115.2113; 7122.274; 7130.9674; 7143.0368; 7250.1077; 7262.1172; 7266.1721; 7268.152; 7283.1496; 7307.2070; 7328.1353; 7394.3224; 7394.5252; 7400.289; 7416.5358; 7435.2763; 7449.3831; 7468.4297; 7491.1348; 7506.1972; 7534.4067; 7607.5077; 7681.4621; 7777.5363; 7840.6401; 7894.5677; 7912.5297; 7924.5736; 7943.5256; 8174.6428; 8344.5958; 9875.9204; 6896.9694; 6880.9937; 7016.998; 7056.1905; 7074.1478; 7104.0354; 7122.2913; 7115.9848; 7175.0715; 7309.2612; 7414.4224; 7600.5; 7654.5083; 7798.6334; 7817.6424; 7832.6366; 7833.6635; 8140.6441; 8159.4822; 8345.5484; 9959.0054; 11068.3376; 16720.7335.
60–70	6896.9694; 6880.9937; 7016.998; 7056.1905; 7074.1478; 7104.0354; 7122.2913; 7115.9848; 7175.0715; 7309.2612; 7414.4224; 7600.5; 7654.5083; 7798.6334; 7817.6424; 7832.6366; 7833.6635; 8140.6441; 8159.4822; 8345.5484; 9959.0054; 11068.3376; 16720.7335.
70–80	6809.9307; 6857.9428; 6859.9368; 6865.9432; 6875.9565; 6880.9796; 6982.0159; 6913.9378; 7009.0523; 7104.9914; 7172.1987; 7200.1528; 7214.1558; 7316.2804; 7377.2599; 7300.0933; 7394.5084.
80–90	7377.2678; 7301.1747; 9140.1069; 11377.1636; 12971.6074; 13004.7435.
90–100	7390.4025; 7466.4483; 7482.4543; 7500.4753; 7704.4655; 7791.5128; 7792.5813; 8672.6993; 8882.0067; 8978.0645; 14577.4253.
100–110	9302.1043; 12990.2825; 12985.6009.
110–120	N.D

N.D: not determined.

Fig. 3. Molecular mass distribution of the monoisotopic masses from MS1 spectra deconvolution. 197 components were detected, with their MW ranging from 1868 to 16 720 Da. These peptides distributed from 1000 to 17 000 Da with 1000 Da mass range windows. The x-axis represents the MW in Da, and the y-axis represents the percentage (%) based on total number counts.

The most representative molecular masses were those from 5000 to 8000 Da, followed by those between 2000 and 5000 Da, which represents respectively 74% and 10% of the total number of measured molecular masses (Fig. 3).

The analysis of reduced/alkylated *B. occitanus* venom filtrate by tandem mass spectrometry allowed the identification of 68 peptides with a molecular weight (MW) from 1959.13 to 7943.53 Da. The detected experimental sequences are shown in Table 2;

Table 2. List of the identified peptides by top-down analysis of the reduced/alkylated *B. occitanus* venom filtrate. Data sets generated from the mass spectrometer were analyzed by the PROTEOME DISCOVER 2.2 software, against UniProtKB/Swiss-Prot database. The amino acids sequences colored in black were those detected by the analysis. Peptide entries in bold were identified by both top-down and bottom-up approaches.

Category	Accession	Description	Identified Sequence
NaScTx	P59356	Alpha-like toxin Lqh6	MVRDGYIAQPENCVYHCIPDCDCLCKDNGGTGGHCGFKLGHGIACWCNALPDNVGIIV DGVKCHK
	P13488	Alpha-like toxin Bom3	MGRDGYIAQPENCVYHCFPGSSGCDTLCKEKGATSGHCGFLPGSGVACWCNDLNPKN VPIVVGGEKCH
	P56678	Alpha-like toxin Lqh3	MVRDGYIAQPENCVYHCFPGSSGCDTLCKEKGATSGHCGFKVGHGLACWCNALPDNV GIIVEGEKCHS
	Q9NJC4	Chain (toxin BmKaTx17) [10–73] in toxin BmKaTx17	MLLMTGVESGRDAYIAKNYNCVYHCFRDDYCNGLTENGADSGYCYLAGKYGNACWC INLPDDKPIRIPGKCHRR
	Q4TUA4	Chain (alpha-toxin 4) [20–85] in alpha-toxin 4	MNLYLVFFSLALLMTGVESVRDGYIADDKNCAYFCGRNAYCDEECKKGAESGYCQWA GVYGNACWCYKLPDKVPIRVPGRGNGG
	P59863	Beta-toxin BotIT2	MDGYIKGYKGCKITCVINDDYCDTECKAEGGTGYCWKWGLACWCEDLPDEKRWKSE TNTC
	P60163	Toxin Cg2	MKDGVLVKNSTGCKYSCIENINDSHCNEECISSIRKSGSYGYCYFYCYCIGMPDSTQVYP IPGKTCSTE
	P60256	Toxin Boma6b	MVRDAYIAQNYNCVYDCARDAYCNELCTKNAGKSGHCEWFGPHGDACWCIDLNNVPI KVEGKCHRK
	O77091	Chain(beta-insect excitatory toxin BmK IT-AP) [19–90] in beta-insect excitatory toxin BmK IT-AP	MKFFLFLVIFPIMGVLGKKNGYAVDSSGKVAECLFNNYCNNECTKVYYADKGYCCLLK YCFGLADDKPVLDIWDSTKNYCDVQIIDL
	P21150	Toxin AaHIT4	MEHGYLLNKYTGCKVWCVINNEECGYLCNKRGGYGYCYFWKLACYCQGARKSELW NYKTNKCDL
	P80962	Beta-insect depressant toxin BaIT2	MDGYIRRRDGCKVSCFLFNGECDECKEAYGSGYCYCWWTGLACWCEGLPDDKTWKS ETNTCG
	P01485	Alpha-mammal toxin Bot3; chain (alpha-mammal toxin Bot3) [10–73] in alpha-mammal toxin Bot3	MLVMAGVESVKDGYIVDDRNCTYFCGRNAYCNEECTKLKGESGYCQWASPYGNACYC YKVPDHSVTKGPGRCN
	Q86BW9	Chain (Makatoxin-2) [20–83] in Makatoxin-2	MNLYLVISFALLMTSVESGRDAYIADSENCTYFCGSPNYCNDLCTENGAKSGYCQWAG RYGNACWCIDLDPKVPIRIPGPCRGR
	G4V3T9	Neurotoxin BmK AGAP- SYP2	MVKDGYIVDDKNCAYFCGRNAYCDEECKNGAESGYCQWAGVYGNACWCYKLPDKV PIRVPGRGNG
	P84614	Alpha-toxin Bs-Tx28	MGVRDAYIADDKNCVYTCGSNSYCNTECTKNGAESGYCQWFGRWGNGCWICLDPDKV PIRIPGKCR
	Q9BLM4	Toxin AahP1005; Chain (toxin AahP1005) [20–83] in toxin AahP1005	MNLYVMISLALLFMTGVESKKGDIYDDKNTFFCGRNAYCNDECKKGAESGYCQWA SPYGNACYCYKLPDRVSTKKKGCGNCR
	P86408	Neurotoxin MeuNaTx-1	MVRDGYIADDKNCAYFCGRNAYCDEECKKGAESGYCQWAGQYGNACWCYKLPDK VPIKVSCKCN
	P60255	Toxin Boma6a	MVRDAYIAQNYNCVYDCARDAYCNDLCTKNAGKSGYCEWFGPHGDACWCIDLNNV PIKVEGKCHRK
	P15225	Neurotoxin Os3	MGVRDGYIAQPHNCVYHCFPGSGGCDTLCKENGATGSSCFILGRGTACWCKDLPCR VGIVDGEKCH
	P45697	Alpha-like toxin BmK-M1; Chain (alpha-like toxin BmK- M1) [20–83] in alpha-like toxin BmK-M1	MNLYVMISFALLMTGVESVRDAYIAKPHNCVYECARNEYCNDLCTKNAGKSGYCQWV KGYNGCWICELPDNVPIRVPGKCHRR
	E4VP24	Chain [20–85] in sodium channel neurotoxin MeuNaTxalpha-1	MNSLVMISLALLVMTGVESVRDGYIADDKNCAYFCGRNAYCDEECKKGAESGYCQW AGQYGNACWCYKLPDKVPIKVSCKCNGR
	P55902	Alpha-insect toxin BotIT1	MVRDAYIAQNYNCVYFCMKDDYCNDLCTKNAGSSGYCQWAGKYGNACWCYALPDNV PIRIPGKCHS
	E7CAU3	Chain (neurotoxin BmK AGP- SYP1) [2–65] in neurotoxin BmK AGP- SYP1	MGRDAYIAQNYNCVYHCFRDDYCNGLTENGADSGYCYLAGKYGHACW CINLPDDKPIRIPGKCHRR
	Q11178	Toxin Td9	MIGMVAECKDGYLVGDDGCKMHCFTRPGHYCASECSRKVGKDGICYAW LACYCYNMPNWAPIWNSATNSCGKKGK
	A0A146CJ90	Chain [20–87] in Venom toxin meuNa32	MNLYLVISFALLVITGVESARDAYIAQNYNCVYFCLNPWSSYCDLCTKNAGK SGYQCIQFGYGNACWCIDLDPKVPIRIPGKCHFA

Coverage (%)	Measured MW (Da)	No. of peptides	No. of PSMs	No. of unique peptides	No. of protein groups	No. of AAs	calc.pl
98.46	6974.21	1	4	1	1	65	6.48
98.5	7012.14	1	1	1	1	67	6.71
98.52	7215.31	1	22	1	1	68	6.48
84	7062.13	1	1	1	1	75	7.58
77.64	7218.31	1	1	0	0	85	7.5
98.36	6564.78	1	1	1	1	61	4.84
88.4	6871.92	1	1	1	1	69	6.92
98.5	7307.23	1	4	1	1	67	7.2
80	7943.53	2	6	2	1	90	5.36
98.48	7791.58	1	6	1	1	66	8.46
100	6845.9	1	4	1	1	62	5.31
87.67	7289.18	1	5	1	1	73	7.53
75.29	7062.11	1	4	1	1	85	5.25
98.48	7289.18	1	6	1	1	66	5.31
98.48	7214.2	1	1	1	1	66	8.12
75.29	7316.26	1	3	1	1	85	8.46
98.46	7218.31	1	6	1	1	65	7.85
98.5	7221.18	1	12	1	1	67	7.09
98.52	6957.15	2	6	2	1	68	6.71
76.19	7429.4	1	4	1	1	84	7.88
77.64	7336.32	1	1	1	1	85	7.85
98.48	7345.15	1	2	1	1	66	7.55
98.5	7488.32	2	8	2	1	67	7.61
86.48	7076.01	1	2	1	1	74	7.84
78.16	7690.37	1	1	1	1	87	7.53

Table 2. Continued.

Category	Accession	Description	Identified Sequence
	P68410	Alpha-mammal toxin Ts2	MKEGYAMDHEGCKFSCFIRPAGFCDGYCKTHLKASSGYCAWPACYCYGV PDHIKWWDYATNKC
	P68726	Chain (Insect toxin 2–53) [22–82] in Insect toxin 2–53	MKLLLLLIVSASMLIESLVNADGYIKRRDGCKVACLGVNEGCDKECKAYGGSY GYCWTWGLACWCEGLPDDKTWKSETNTCGGKK
	Q1I163	Toxin Td8; chain (toxin Td8) [21– 83] in toxin Td8	MTRFVLFLSCFFLIGMVVECKDGYLVGGDGCKMHCFTRPGHYCASECSRVK GKDGYCAWLACYCYNMPNWAIPWNSATNRCRGRK
	P56569	Makatoxin-1	MGRDAYIADSENCTYTCALNPYCNDLCTKNGAKSGYCWAGRYGNACWC DLDPKVPIRISGSCR
	D8UWD3	Sodium channel neurotoxin MeuNaTxalpha-7	MARDGYIADKNCAYFCGRNAYCDEECKKGAESGYCQWAGQYGNACWC YKLDPKVPKIVSGKCNGR
	P0DMH9	Chain (alpha-toxin Bmalphatx47) [20–83] in alpha-toxin Bmalphatx47	MNVLIVISFALLMTGVQSGRDAYIADSENCTYTCALNPYCNDLCTKNGAKSG YCQWAGRYGNACWCIDLDPKVPIRISGSCRGR
	P01483	Neurotoxin Bot2	MGRDAYIAQPCNVYECANYSYNDLCTKNGAKSGYQWLGRWGNACYC IDLDPKVPIRIEGKCHF
	P17728	Chain (alpha-insect toxin LqhalT) [20–85] in alpha-insect toxin LqhalT	MNHLVMSIALLLLLLVESVRDAYIAKNYCNVYECFRDAYCNELCTKNGASS GYCQWAGRYGNACWCYALPDNPVIRVPGKCHRK
	P01496	Chain (toxin-3) [15–76] in toxin-5	MLVVVCLLTAGTEGKKDGYPVEYDNCAYICWNYDNAYCDKLCKDKKADSGY CYWVHILCYCYGLPDESEPTKTNGCKKSGKK
	Q1EG64	Chain [20–85] in sodium toxin peptide BmKTb'	MNVLVMSISFALLMTGVESARDAYIAQNYCNVYHARDAYCNELCTKNGAKS GSCPYLGEHKFACYCKDLPDNPVIRVPGKCNNG
	P01488	alpha-toxin Bot1	MGRDAYIAQPCNVYECANYSYNDLCTKNGATSGYQWLKGKYNACWC KDLPDNPVIRIPGKCHF
	P45698	Chain (neurotoxin BmK-M9) [15– 78] in neurotoxin BmK-M9	MISFALLMTGVESVRDAYIAKPENVCYHATNEGCNKLCTDNGAESGYCQW GGRYGNACWCIKLPDRVPIRVPGKCHR
	P83644	Toxin Lqh4	MGVRDYIADKNCVYTCGANSYCNTECTKNGAESGYCQWFGKYGNACWC IKLPDKVPIRIPGKCR
	P01487	Alpha-insect toxin Lqq3	MVRDAYIAKNYCNVYECFRDSYCNLCTKNGASSGYCQWAGKYGNACWC YALPDNPVIRVPGKCH
	H1ZZI7	Toxin Tpa6	MSIFIALALLLIGLEEAEARDGYPLSKNNNKIYCPDTCVCKDTCKNRASAP DGKCDGWNSCYCFKVPDHIPVWGDPTGKPCMT
	B8XGY6	Chain [20–85] in Putative alpha- toxin Tx17	MNVLILISLAVLLTSGVESVRDAYIAQNYCNVYTCFKDAYCNLCTKNGATSGY CQWVGKYNGCWYALPDNPVIRVPGKCHSR
	P81504	Insect toxin AaHIT5	MDGYIKRHDGCKVTCLINDNYCDETECKREGSGYCYSGVFACWCEGLPDD KAWKSETNTCD
	P68722	Chain (beta-insect excitatory toxin LqhlT1b) [19–88] in beta-insect excitatory toxin LqhlT1b	MKFFLLFLVLPIMGVLGKKNYAVDSKGKAPCEFLSNYCNNECTKVHYADK GYCCLLSYCFGLNDDKKVLEISDITTKYCYDFTIIN
	P60257	Toxin Boma6c	MVRDAYIAQNYCNVYTCFKDAHCNDLCTKNGASSGYCQWAGKYGNACWCY ALPDNPVIRIPGKCHRK
	M1J7U4	Putative sodium channel alpha- toxin Acra5	MVRDGYIMIKDTNCKFSCNIFKWEYCSPLCQSGAETGYCYNFGCWCLDL PDDVPVYGDGVICRTR
	Q9N682	Chain (neurotoxin BmK-M11) [20– 83] in neurotoxin BmK-M11	MNVLVMSISFALLMTGVESVRDAYIAKPENVCYHATNEGCNKLCTDNGAESG YCQWGGKYGNACWCIKLPDDVPVIRVPGKCHR
	P55903	beta-insect depressant toxin BotIT4	MDGYIRRRDGCKVSCFLGNEGCDKECKAYGSGYCWWTWGLACWCEGLPDD KTWKSETNTCG
	A0A0K0LBU9	Chain [20–83] in sodium channel blocker AbNaTx26	MRAALLAFSSLITGVLTGKSGYPTQHDGKIWCVFNFHFCSNYCETYGGSGYCYT WKLACWCDNIHDWWPTWSYATTKCRAK
	P0C910	Alpha-toxin Amm3	MGRDGYIVDTKNCVYHCYPPCDGLCKKNQAKSGSCGFLYPSGLACWCVALPENV PIKDPNDCHK
	P59360	Neurotoxin BmK-II	VRDAYIAKPHNCVYECARNEYCNLCTKDGAESGYCQWVGKYNGWCWCIPLDNPV PIRIPGNCH
	P81240	Insect toxin LqhlT5	MDGYIRGGDGCKVSCVIDHVFCNDNECKAAGSGYCYCWGWGLACWCEGLPADREW YETNTCG
	P01497	Chain (beta-insect excitatory toxin 1) [19–88] in beta-insect excitatory toxin 1	MKFFLLFLVLPIMGVFGKKNYAVDSSGKAPCEFLSNYCNNECTKVHYADKGYCCLL SCYCFGLNDDKKVLEISDITRKYCYDFTIIN
	V9P3B8	Chain [23–82] in Chain [23–82] in Meutoxin-3	MKILTVFMIFIANFLSMTQVFLSKDRFLLINGSYELCLYEENLDEDCERLCKEQNASDG FCRQPHCFCADMPDDYPTPTTR

Coverage (%)	Measured MW (Da)	No. of peptides	No. of PSMs	No. of unique peptides	No. of protein groups	No. of AAs	calc.pl
98.41	6655.84	1	6	1	1	63	7.61
71.76	6739.87	1	1	1	1	85	7.5
73.25	6986.05	1	3	1	1	86	8.34
98.46	7240.24	1	2	0	0	65	7.5
98.5	7295.24	1	1	1	1	67	8.1
75.29	7240.24	1	1	0	0	85	7.87
98.48	7240.24	1	19	1	1	66	7.55
77.64	7173.2	1	12	1	1	85	8.12
76.54	7105.03	1	20	1	1	81	7.49
77.64	7321.09	1	2	1	1	85	7.58
98.48	7074.14	1	3	1	1	66	6.92
81.01	7015.19	1	1	1	1	79	7.88
98.48	7155.25	1	3	1	1	66	8.1
98.5	6980.01	2	12	2	1	65	7.87
74.41	7059.12	1	2	1	1	86	5.38
77.64	7313.2	1	2	1	1	85	7.87
98.38	6894.89	1	8	1	1	62	4.83
79.54	7924.56	1	1	1	1	88	7.87
98.5	7308.21	2	14	2	1	67	8.31
98.52	7741.51	1	1	1	1	68	7.5
77.38	7179.21	2	2	2	1	84	7.09
100	6837.96	1	4	1	1	62	5.31
77.1	7505.2	1	1	1	1	83	8.31
98.46	7011.14	1	1	1	1	65	7.09
100	7431.33	2	14	2	1	65	7.09
100	6611.8	1	3	1	1	62	4.72
79.54	7928.54	1	10	1	1	88	7.53
73.17	7074.13	1	1	1	1	82	4.75

Table 2. Continued.

Category	Accession	Description	Identified Sequence
	Q8T3T0	Depressant insect toxin BmK ITa1	MKLFLLLLISASMLIDGLVNADGYIRGSNGCKVSLWNGECNKECGAYGASYGYCW TWGLACWCQGLPDDKTWKSESNTCGGKK
	Q9GQW3	Chain (toxin BmKaIT1) [20–83] in toxin BmKaIT1	MNYLVMSIFALLMTGVESVRDAYIAQNYNCVYHCARDAYCNELCTKNKAGSGSCP LGEHKFACYCKDLPDNPVPIRVPGKCHRR
	Q95WX6	Beta-insect depressant toxin BmKITb	MKLFLLLVISASMLIDGLVNADGYIRGSNGCKVSLWNGECNKECKAFGAYGYGYCW TWGLACWCQGLPDDKTWKSESNTCGGKK
	P0C5H1	Beta-toxin Isom1	MKKNGYAVDSSGKAPECLLSNYCNNECTKVHYADKGCCLLSCYCFGLSDDKKVLEIS DTRKKYCDYTIIN
	Q9GNG8	Toxin BmKaTX15	MNYLVFFSLALLVMTGVESVRDGYIADDKNCAYFCGRNAYCDECKKNGAESGYCQW AGVYGACWCYKLPDKVPIRVPGKCNNG
	M1JMR8	Sodium channel alpha-toxin Acra8	MVRDGYVDDKNCTFFCGRNAYCNDCKKGGESGYCQWASPYGNACWCYKLPDRV PIKEKGRGNR
	A0A0U4RDS7	Chain [20–87] in sodium channel toxin NaTx4	MNHLVMSLAFLFMTGVASVRDGYIAQPETCAYHCIPGSSGYCYLCKEKGESGHGCGW SGHGSAAWWCNDLPDKEGIIVDGKCTR
	P82814	Insect toxin BslT4	MDGYIKNGKCKVSCVINNVFCNSMCKSSSGSYGYCWSWGLACWCQGLPAKKWLY AATNTCG
	B8XGX9	Chain [20–87] in Putative alpha-toxin Tx2	MNYLIMISLALLMTGVESGTGVRDAYIADDKNCVYTCALNSYCNTECTKNGAESGYCQ WLGQYGNACWCIKLPDRVPIRVPGKCRG
	Q17254	Alpha-insect toxin Bot14	MSSLMISTAMKGKAPYRQVRDGYIAQPHNCAYHCLKISSGCDTLCKENGATSGHCGH KSGHGSACWCCKDLDPKVGIIHVHGEKCHR
KScTx	A0A059UI30	Chain (potassium channel toxin Meg-beta-KTx1) [28– 91] in potassium channel toxin Meg-beta-KTx1	MQRNLVLLFLGMVALSSCGLREKHFQKLKYAVPEGLRTIIQTAVHKLGTQFGCPA YQGYCDDHCQDIKKQEGFCHGFKCKGIPMGF
	Q9N661	Potassium channel toxin BmTXK-beta-2	MQRNLVLLFLGMVALSSCGLREKHFQKLKYAVPEGLRTIIQTAVHKLGTQFGCP AYQGYCDDHCQDIKKQEGFCHGFKCKGIPMGF
AMP	A0A0A1I6E7	AMP AcrAP1	MEIKYLLTVFLVLLIVSDHCQAFSLIPHAISGLISAFKGRKRDLGQIDFRFRNFRKR AELELLSKLPIY
Myotropic neuropeptide	F8THJ9	Putative orcokinin	MMFGIWLICGTAFFFCHVDAYLEYSNMAGYNALVRRRSMKQPSGRMFDNLGYNQE SLVKRNFDEIDNVGFNDFGPASRPGSGRSWFPKRNWELARYNLRRLVKRATQD ELMENKROELDEIDKSGFGFHKRNFDIDRSGFNDGKRSFDRFKLVRRADFN MQNIFWILGIVGICITAVQCDSEMISSRIDLTIRRYLYKARSVLDDLNNQDLTLHKRSC VLNLPGMDCYGDITGSGKDQDYWTSGRTPGKRRSYCSYLIGNSEECLETLQLKDDM TDFNSWNDKFRPGKK
Hypothetical secreted protein	F1CIZ9	Hypothetical secreted protein	

five of the entries were identified with 100% sequence coverage: neurotoxin BmK-II (P59360), beta-insect depressant toxin BotIT4 (P55903), beta-insect depressant toxin BaIT2 (P80962), insect toxin LqhIT5 (P81240), and insect toxin BsIT4 (P82814). These toxins were reported for the first time in this Moroccan venom, they corresponded to toxins already identified in other scorpion venom. The determined sequence of the neurotoxin BmK-II (P59360) showed 100% similarity with the database sequence, whereas the observed sequences of the other toxins showed methylation in the N-terminal part compared with sequences reported in Uniprot database (Fig. 4). Therefore, the other peptides corresponded approximately to toxins, previously identified in other scorpion species with a sequence identity ranging from 17% to 98% (Fig. S2).

Therefore, the detected peptides were divided into five categories on the basis of their molecular functions according to the UniProtKB database (<https://www.uniprot.org>); 63 neurotoxins acting on sodium channels

(NaScTxs), constitute 93% of the components and represent a MW from 6564.78 to 7943.53 Da; two neurotoxins acting on potassium channels (KScTxs) (2.94%, 2506.46–6889.3 Da); one antimicrobial peptide (AMP) (1.47%, 1959.13 Da); one myotropic neuropeptide (1.47%, 3112.45 Da); and one hypothetical secreted protein (1.47%, 3939.79 Da) (Fig. 5A).

Additionally, we have observed, that between these 68 peptides, 27 of them (40%) were detected as chains or fragments, for example, venom toxin meuNa32 (A0A146CJ90); potassium channel toxin Meg-beta-KTx1 (A0A059UI30); putative alpha-toxin Tx2 (B8XGX9); sodium channel toxin NaTx4 (A0A0U4RDS7); toxin BmKaIT1 (Q9GQW3); sodium channel blocker AbNaTx26 (A0A0K0LBU9); neurotoxin BmK-M11 (Q9N682); beta-insect excitatory toxin LqhIT1b (P68722); toxin-5 (P01496); toxin Td8 (Q1I163); alpha-like toxin BmK-M1 (P45697); toxin AahP1005 (Q9BLM4); makatoxin-2 (Q86BW9); and alpha-mammal toxin Bot3 (P01485) (Table 2).

Coverage (%)	Measured MW (Da)	No. of peptides	No. of PSMs	No. of unique peptides	No. of protein groups	No. of AAs	calc.pl
71.76	6632.71	1	18	1	1	85	6.38
75.29	7012.23	1	3	1	1	85	8.12
71.76	6775.93	1	4	1	1	85	7.85
98.59	7895.47	1	35	1	1	71	7.53
77.64	7211.14	1	1	1	1	85	6.4
98.5	7218.3	1	1	1	1	67	8.29
78.16	7243.29	2	4	2	1	87	7.66
100	6954.15	1	1	1	1	63	8.31
78.16	7394.28	1	3	1	1	87	7.5
78.82	7184.3	1	5	1	1	85	8.5
70.32	6889.3	1	9	1	1	91	8.76
25.27	2506.46	1	1	1	1	91	8.57
24.32	1959.13	1	1	1	1	74	9.31
16.96	3112.45	1	1	1	1	165	9.29
25.75	3939.79	1	1	1	1	132	7.99

Bottom-up proteomics

For the bottom-up workflow, two digest methods were performed: (a) in-solution digestion, the flow-through containing toxin < 30 kDa was directly reduced with DTT, alkylated with IAA, and digested with trypsin; and (b) in-gel digestion, the gel spot corresponding to peptides under 30 kDa (Fig. S1) was excised to small cubes, which after series of washings, were reduced, alkylated, and digested.

The results generated by the bottom-up approach using the in-gel digestion yielded the identification of 36 peptides, whereas 37 was the total of the identified peptide by in-solution digestion. The detected peptides showed similarity of sequences with peptides from other scorpion species, and with their sequence coverage ranging from 10.23% (P68721) to 86.15% (P01489) and from 8.75% (P0C294) to 92.86% (P80669) for the in-gel and in-solution digestions, respectively.

The identified categories of peptides using the in-gel digestion were as follows: 27 NaScTxS; seven KScTxS; and two ClTxS (Table 3). While, through the in-solution digestion, we identified in addition to 24 NaScTxS, eight KScTxS and three ClScTxS, one entry that shares 60% of similarity with neurotoxin Tx-2 (P83406) purified from *Hottentotta judaicus*, could correspond to a calcium channel activator 'CaScTx' scorpion. Besides neurotoxins, one amphipathic peptide was detected by this digestion method (Table 4).

According to the results, 23 of the entries were detected by both digestion methods (Tables 3 and 4). Thus, 14 peptides were identified only by the in-solution digestion method, for example, alpha-toxin Amm5 (P01482), alpha-mammal toxin Bot3 (P01485), potassium channel toxin alpha-KTx 9.3 (P80669), neurotoxin Tx-2 (P83406), neurotoxin P2 (P01498), and amphipathic peptide Tx348 (B8XH50). Otherwise, regarding the in-gel digestion results, 13 peptides were identified only by this method of digestion, for example,




Fig. 4. The detected amino acid sequences of the five toxins identified with 100% coverage by the top-down LC-MS/MS; neurotoxin BmK-II (P59360); beta-insect depressant toxin BalT2 (P80962); insect toxin BslT4 (P82814); insect toxin LqhIT5 (P81240); and beta-insect depressant toxin BotIT4 (P55903).




Fig. 5. (A) Relative abundance of the different peptide categories identified in reduced/alkylated *B. occitanus* venom filtrate by the top-down LC-MS/MS analysis. Peptides were divided on the basis of their molecular functions into: neurotoxins active on sodium channels (NaScTx), neurotoxins active on potassium channels (KScTx), myotrophic neuropeptide, AMP, and hypothetical secreted protein. (B) Relative abundance of the different peptide categories identified in reduced/alkylated and digested *B. occitanus* venom by bottom-up LC-MS/MS analysis. The peptides were divided on the basis of their molecular functions into: neurotoxins active on sodium channels (NaScTx), neurotoxins active on potassium channels (KScTx), neurotoxins active on chloride channels (ClScTx), neurotoxins active on calcium channels (CaScTx), toxin Acra, and amphipathic peptide.

potassium channel toxin alpha-KTx 9.11 (B3EWX9); sodium channel alpha-toxin Acra4 (M1JBC0); sodium channel alpha-toxin Acra8 (M1JMR8), alpha-toxin Ac3 (fragment) (D5HR52); and beta-insect depressant toxin BotIT5 (P55904).

Since the aim of using two methods of digestions was to identify the maximum of peptide, the data generated by bottom-up approaches using in-gel and in-solution digestions were then summarized in Table 5; the repeated molecules were deleted and thus allowed the detection of a total of 50 peptides, which were divided into different categories according to their molecular functions. The generated data from the bottom-up process confirmed that the family with the most diverse members in this venom is neurotoxins, with 31 NaScTx (62%, 4.3–10.2 kDa), 13 KScTx (26%, 2.9–10.4 kDa), three ClScTx (6%, 3.6–4 kDa), one CaScTx (2%, 2.9 kDa), and one toxin Acra (2%, 8.8 kDa).

In addition to these neurotoxins, we identified one amphipathic peptide (2%, 7.8 kDa) (Fig. 5B). Also, some peptides were detected as fragments (10% of

total): alpha-toxin Ac1 (D5HR50) and Ac3 (D5HR52); alpha-mammal toxin Bot3 (P01485); and neurotoxin 8 (P04098).

As we mentioned above, we aimed to gain a deeper understanding of the *B. occitanus* peptidome (under 30 kDa), so the molecular diversity of its toxins. In this context, we combined data from the top-down and bottom-up analyses and then analyzed the generated data to infer a global and comprehensive characterization of this venom.

According to this study, a total of 118 peptides were identified from *B. occitanus* venom; among them, 16 were identified by both approaches, for example, potassium channel toxin BmTXK-beta-2 (Q9N661); toxin AaHIT4 (P21150); and alpha-mammal toxin Bot3 (Fragment) (P01485).

Among the 102 identified peptides, the most representative category is neurotoxins, mainly NaScTx (77%), followed by KScTx (14%), ClScTx (3%), CaScTx (1%), and toxin Acra (1%). We also characterized other peptides with low percentage such as

Table 3. Bottom-up data generated from in-gel digestion of *B. occitanus* venom filtrate using nano-LC-MS/MS. Data sets generated from the mass spectrometer were analyzed by the PROTEOME DISCOVER 2.2 software, against UniProtKB/Swiss-Prot database.

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW [kDa]	calc. pI
NaScTx	<u>Q86SE0</u>	Toxin Aam2 OS = <i>Androctonus amoreuxi</i> PE = 1 SV = 1 - [SCX2_ANDAM]	198.74	24.42%	9	2	3	6	86	9.3	7.87
	<u>P21150</u>	Toxin AaHIT4 OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [SIX4_ANDAU]	85.81	29.23%	2	1	2	5	65	7.8	8.46
	<u>P13488</u>	Alpha-like toxin Bom3 OS = <i>Buthus occitanus mardochei</i> PE = 1 SV = 1 - [SCX3_BUTOM]	169.75	56.06%	2	3	3	15	66	6.9	6.71
	<u>P68721</u>	Beta-insect excitatory toxin LqhlT1a OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 3 SV = 1 - [SIX1A_LEIQH]	54.81	10.23%	2	1	2	3	88	9.9	8.09
	<u>P0DJH8</u>	Alpha-toxin Bu1 OS = <i>Buthacus macrocentrus</i> PE = 1 SV = 1 - [SCX1_BUTMA]	346.32	71.64%	1	3	5	7	67	7.5	8.48
	<u>P86406</u>	Neurotoxin MeuNaTx-6 OS = <i>Mesobuthus eupeus</i> PE = 1 SV = 1 - [SCXN6_MESEU]	134.56	15.15%	3	1	1	4	66	7.8	7.87
	<u>P83644</u>	Toxin Lqh4 OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 1 - [SCX4_LEIQH]	305.53	46.15%	8	1	3	7	65	7.2	8.1
	<u>P01489</u>	Alpha-toxin Lq4 OS = <i>Leiurus quinquestriatus quinquestriatus</i> PE = 1 SV = 1 - [SCX4_LEIQH]	531.95	86.15%	9	2	5	11	65	7.2	8.1
	<u>P01486</u>	Alpha-toxin Bot11 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SCXB_BUTOC]	106.19	35.38%	7	1	3	7	65	7.5	7.87
	<u>P60255</u>	Toxin Boma6a OS = <i>Buthus occitanus mardochei</i> PE = 3 SV = 1 - [SCXA_BUTOM]	65.84	15.15%	2	1	1	2	66	7.5	7.09
P17728	<u>P17728</u>	Alpha-insect toxin LqhaIT OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 2 - [SCXA_LEIQH]	174.28	31.76%	4	1	2	3	85	9.6	8.12
	<u>P04098</u>	Neurotoxin 8 (Fragment) OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SCX8_BUTOC]	202.83	72.22%	2	2	2	4	36	4.1	6.24
	<u>P55902</u>	Alpha-insect toxin BotIT1 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX1_BUTOC]	211.59	41.54%	2	1	2	4	65	7.3	7.55
	<u>P01488</u>	Alpha-toxin Bot1 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 2 - [SCX1_BUTOC]	136.52	20.00%	1	1	1	2	65	7.3	6.92

Table 3. (Continued).

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW [kDa]	calc. pI
	P81504	Insect toxin AaHIT5 OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [SIX5_ANDAU]	51.44	24.59%	1	1	1	2	61	6.9	4.83
	P59863	Beta-toxin BotIT2 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX2_BUTOC]	109.66	43.33%	1	2	2	3	60	6.9	4.84
	Q17254	Alpha-insect toxin Bot14 OS = <i>Buthus occitanus tunetanus</i> PE = 2 SV = 1 - [SCXE_BUTOC]	44.99	18.82%	1	1	1	3	85	9.2	8.5
	D5HR52	Alpha-toxin Ac3 (Fragment) OS = <i>Androctonus crassicauda</i> PE = 3 SV = 1 - [SCX3A_ANDCR]	139.86	63.77%	10	2	4	10	69	7.8	7.87
	P55904	Beta-insect depressant toxin BotIT5 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX5_BUTOC]	64.67	27.87%	21	2	2	9	61	6.8	5.31
	Q77091	Beta-insect excitatory toxin BmK IT-AP OS = <i>Mesobuthus martensii</i> GN = IT-AP PE = 1 SV = 1 - [SIXP_MESMA]	126.26	17.78%	8	2	2	4	90	10.2	5.36
	P59864	Beta-insect depressant toxin BotIT6 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX6_BUTOC]	32.37	11.29%	1	1	1	1	62	7.3	8.1
	P68723	Beta-insect excitatory toxin LqhIT1c OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 1 - [SIX1C_LEIQH]	182.31	11.36%	1	2	3	8	88	9.9	8.1
	P59360	Neurotoxin BmK-II OS = <i>Mesobuthus martensii</i> PE = 1 SV = 1 - [SCX2_MESMA]	48.26	15.63%	3	1	1	1	64	7.2	7.09
	P15224	Toxin Os1 OS = <i>Orthochirus scrobiculosus</i> PE = 1 SV = 1 - [SCX1_ORTSC]	39.14	19.70%	1	1	1	1	66	7.6	7.88
	D5HR50	Alpha-toxin Ac1 (Fragment) OS = <i>Androctonus crassicauda</i> PE = 2 SV = 1 - [SCX1A_ANDCR]	37.66	11.11%	2	1	1	2	81	8.7	7.55
	M1JMR8	Sodium channel alpha-toxin Acra8 OS = <i>Androctonus crassicauda</i> PE = 3 SV = 1 - [SCX8_ANDCR]	66.82	40.91%	3	2	3	5	66	7.5	8.29
	M1JBC0	Sodium channel alpha-toxin Acra4 OS = <i>Androctonus crassicauda</i> PE = 1 SV = 1 - [SCX4_ANDCR]	37.39	29.23%	1	1	2	4	65	7.1	8.31

Table 3. (Continued).

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW [kDa]	calc. pI
KScTx	P0C161	Potassium channel toxin alpha-KTx 2.8 OS = <i>Centruroides elegans</i> PE = 1 SV = 1 - [KAX28_CENEL]	45.57	17.95%	2	1	1	1	39	4.3	8.94
	<u>Q9NJC6</u>	Potassium channel toxin BmTXK-beta OS = <i>Mesobuthus martensii</i> PE = 2 SV = 1 - [KBX2_MESMA]	264.78	23.33%	2	1	2	6	90	10.4	8.82
	P59869	Potassium channel toxin alpha-KTx 5.4 OS = <i>Mesobuthus tamulus</i> PE = 1 SV = 1 - [KAX54_MESTA]	40.7	22.58%	2	1	2	2	31	3.5	8.02
	B8XH40	Potassium channel toxin BuTXK-beta OS = <i>Buthus occitanus israelis</i> PE = 2 SV = 1 - [KBX1_BUTOS]	298.65	42.86%	2	2	5	18	91	10.2	8.57
	<u>Q9N661</u>	Potassium channel toxin BmTXK-beta-2 OS = <i>Mesobuthus martensii</i> PE = 2 SV = 1 - [KBX1_MESMA]	230.62	42.86%	2	1	4	13	91	10.2	8.57
	B3EWX9	Potassium channel toxin alpha-KTx 9.11 OS = <i>Mesobuthus gibbosus</i> PE = 1 SV = 1 - [KAX9B_MESGB]	85.33	40.74%	4	1	1	2	27	2.9	5.01
	B8XH42	Potassium channel toxin alpha-KTx 16.6 OS = <i>Buthus occitanus israelis</i> PE = 2 SV = 1 - [KA166_BUTOS]	23.25	12.07%	1	1	1	1	58	6.5	8.12
	<u>P45639</u>	Chlorotoxin OS = <i>Leiurus quinquestriatus</i> quinquestriatus PE = 1 SV = 1 - [CTXL_LEIQU]	41.56	38.89%	1	1	1	3	36	4	8.13
	<u>P86436</u>	Chlorotoxin-like peptide OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [CTXL_ANDAU]	290.9	44.12%	1	1	1	7	34	3.6	8.34
	CIScTx										

Underlined peptide entries were identified by in-gel and in-solution digestion methods.

Table 4. Bottom-up data generated from in-solution digestion of *B. occitanus* venom filtrate using nano-LC-MS/MS. Data sets generated from the mass spectrometer were analyzed by the PROTEOME DISCOVER 2.2 software, against UniProtKB/Swiss-Prot database.

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW (kDa)	calc. pI
NaScTxs	<u>Q86SE0</u>	Toxin Aam2 OS = <i>Androctonus amoreuxi</i> PE = 1 SV = 1 - [SCX2_ANDAM]	250.68	24.42%	8	2	4	15	86	9.3	7.87
	<u>P21150</u>	Toxin AaHIT4 OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [SIX4_ANDAU]	192.23	30.77%	2	2	3	12	65	7.8	8.46
	<u>P01482</u>	Alpha-toxin Amm5 OS = <i>Androctonus mauretanicus</i> PE = 1 SV = 1 - [SCX5_ANDMA]	96.57	28.13%	1	1	1	2	64	7.3	7.5
	<u>P01481</u>	Alpha-mammal toxin Lq45 OS = <i>Leiurus quinquestriatus</i> PE = 1 SV = 1 - [SCX5_LEIOU]	77.71	25.00%	2	1	2	4	64	7.3	8.1
	<u>P13488</u>	Alpha-like toxin Bom3 OS = <i>Buthus occitanus mardochei</i> PE = 1 SV = 1 - [SCX3_BUTOM]	155.6	59.09%	2	2	4	18	66	6.9	6.71
	<u>P45698</u>	Neurotoxin BmK-M9 OS = <i>Mesobuthus martensii</i> PE = 1 SV = 1 - [SCX9_MESMA]	124.54	26.58%	11	1	3	15	79	8.8	7.88
	<u>P68721</u>	Beta-insect excitatory toxin Lq1T1a OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 3 SV = 1 - [SIX1A_LEIQH]	55.48	10.23%	2	1	2	4	88	9.9	8.09
	<u>P0DJH8</u>	Alpha-toxin Bu1 OS = <i>Buthacus macrocentrus</i> PE = 1 SV = 1 - [SCX1_BUTMA]	272.84	71.64%	1	2	4	9	67	7.5	8.48
	<u>P83644</u>	Toxin Lq4 OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 1 - [SCX4_LEIQH]	293.2	46.15%	7	1	3	10	65	7.2	8.1
	<u>P01489</u>	Alpha-toxin Lq4 OS = <i>Leiurus quinquestriatus quinquestriatus</i> PE = 1 SV = 1 - [SCX4_LEIQU]	569.9	90.77%	8	2	6	18	65	7.2	8.1
	<u>P01486</u>	Alpha-toxin Bot11 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SCXB_BUTOC]	76.63	35.38%	2	1	3	7	65	7.5	7.87
	<u>P60255</u>	Toxin Boma6a OS = <i>Buthus occitanus mardochei</i> PE = 3 SV = 1 - [SCXA_BUTOM]	46.01	15.15%	2	1	1	2	66	7.5	7.09

Table 4. (Continued).

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW (kDa)	calc. pI
	<u>P17728</u>	Alpha-insect toxin LqhαIT OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 2 - [SCXA_LEIQH]	369.61	51.76%	4	2	5	13	85	9.6	8.12
	<u>P04098</u>	Neurotoxin 8 (Fragment) OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SCX8_BUTOC]	536.34	77.78%	2	3	3	13	36	4.1	6.24
	<u>P55902</u>	Alpha-insect toxin Bot1T1 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX1_BUTOC]	296.35	61.54%	1	1	3	9	65	7.3	7.55
	<u>P01488</u>	Alpha-toxin Bot1 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 2 - [SCX1_BUTOC]	185.35	20.00%	1	1	1	3	65	7.3	6.92
	<u>P81504</u>	Insect toxin AaHIT5 OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [SIX5_ANDAU]	49.42	24.59%	1	1	1	2	61	6.9	4.83
	<u>P01485</u>	Alpha-mammal toxin Bot3 (Fragment) OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 2 - [SCX3_BUTOC]	436.17	61.11%	3	2	5	61	72	8.1	7.53
	<u>P59863</u>	Beta-toxin Bot1T2 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX2_BUTOC]	164.41	41.67%	1	2	2	4	60	6.9	4.84
	<u>Q17254</u>	Alpha-insect toxin Bot14 OS = <i>Buthus occitanus tunetanus</i> PE = 2 SV = 1 - [SCXE_BUTOC]	91.78	18.82%	1	1	1	9	85	9.2	8.5
	<u>O77091</u>	Beta-insect excitatory toxin BmK IT-AP OS = <i>Mesobuthus martensii</i> GN = IT-AP PE = 1 SV = 1 - [SIXP_MESMA]	50.93	17.78%	8	1	2	5	90	10.2	5.36
	<u>P59864</u>	Beta-insect depressant toxin Bot1T6 OS = <i>Buthus occitanus tunetanus</i> PE = 1 SV = 1 - [SIX6_BUTOC]	78.83	53.23%	1	2	3	7	62	7.3	8.1
	<u>P0C294</u>	Toxin Acra I-3 OS = <i>Androctonus crassicauda</i> PE = 2 SV = 1 - [TX13_ANDCR]	43.58	8.75%	1	1	1	1	80	8.8	8.25
	<u>P59360</u>	Neurotoxin BmK-II OS = <i>Mesobuthus martensii</i> PE = 1 SV = 1 - [SCX2_MESMA]	62.88	15.63%	3	1	1	2	64	7.2	7.09

Table 4. (Continued).

Category	Accession	Description	Score	Coverage	No. of proteins	No. of unique peptides	No. of peptides	No. of PSMs	No. of AAs	MW (kDa)	calc. pI
KScTxs	P0CC12	Potassium channel toxin alpha-KTx 8.5 OS = <i>Odontobuthus doriae</i> PE = 1 SV = 1 - [KAX85_ODODO]	86.25	48.28%	2	1	1	2	29	3.2	5.1
	P83407	Potassium channel toxin alpha-KTx 19.1 OS = <i>Mesobuthus martensii</i> PE = 1 SV = 1 - [KA191_MESMA]	82.22	32.26%	1	1	1	5	31	3.3	8.73
	Q95NU8	Potassium channel toxin alpha-KTx 17.1 OS = <i>Mesobuthus martensii</i> PE = 1 SV = 1 - [KA171_MESMA]	79.79	16.36%	1	1	1	5	55	6.2	8
	P80669	Potassium channel toxin alpha-KTx 9.3 OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 1 - [KAX93_LEIQH]	211.78	92.86%	3	2	2	9	28	3	6.98
Q9NJUC6		Potassium channel toxin BmTXK-beta OS = <i>Mesobuthus martensii</i> PE = 2 SV = 1 - [KBX2_MESMA]	135.05	27.78%	2	2	2	3	90	10.4	8.82
	Q9N661	Potassium channel toxin BmTXK-beta-2 OS = <i>Mesobuthus martensii</i> PE = 2 SV = 1 - [KBX1_MESMA]	96.9	42.86%	3	2	4	7	91	10.2	8.57
	P86399	Neurotoxin lamda-MeuTx OS = <i>Mesobuthus eupeus</i> PE = 1 SV = 2 - [TXL_MESEU]	262	25.00%	2	1	1	8	64	7.2	7.12
	P80670	Toxin GaTx2 OS = <i>Leiurus quinquestriatus hebraeus</i> PE = 1 SV = 1 - [KAX83_LEIQH]	86.02	48.28%	2	1	1	2	29	3.2	5.1
CaScTxs	P83406	Neurotoxin Tx-2 OS = <i>Buthotus judaicus</i> PE = 1 SV = 1 - [SCBT2_BUTJU]	287.63	60.71%	1	2	2	10	28	2.9	4.89
	P86436	Chlorotoxin-like peptide OS = <i>Androctonus australis</i> PE = 1 SV = 1 - [CTXL_ANDAU]	993.18	67.65%	1	3	3	65	34	3.6	8.34
CIScTxs	P45639	Chlorotoxin OS = <i>Leiurus quinquestriatus quinquestriatus</i> PE = 1 SV = 1 - [SCXL_LEIQU]	588.38	38.89%	1	2	2	38	36	4	8.13
	P01498	Neurotoxin P2 OS = <i>Androctonus mauretanicus</i> PE = 1 SV = 1 - [SCXP_ANDMA]	188.48	71.43%	1	2	2	5	35	3.7	7.88
Amphipathic peptide	B8XH50	Amphipathic peptide Tx348 OS = <i>Buthus occitanus israelis</i> PE = 2 SV = 1 - [NDB5R_BUTOS]	87.27	19.40%	4	1	1	1	67	7.8	9.19

Underlined peptide entries were identified by in-gel and in-solution digestion methods.

Table 5. List of the 50 peptides detected by the bottom-up analysis of the reduced/alkylated *B. occitanus* venom filtrate. Data sets generated from the mass spectrometer were analyzed by the PROTEOME DISCOVER 2.2 software, against UniProtKB/Swiss-Prot database.

Category	Accession	Description	MW (kDa)	Species	Digestion method
NaScTx	P86406	Neurotoxin MeuNaTx-6	7.8	<i>Mesobuthus eupeus</i>	In-gel digestion
	P59863	Beta-toxin BotIT2	6.9	<i>Buthus occitanus tunetanus</i>	Both
	D5HR52	Alpha-toxin Ac3 (Fragment)	7.8	<i>Androctonus crassicauda</i>	In-gel digestion
	P55904	Beta-insect depressant toxin BotIT5	6.8	<i>Buthus occitanus tunetanus</i>	In-gel digestion
	O77091	Beta-insect excitatory toxin BmK IT-AP	10.2	<i>Mesobuthus martensii</i>	Both
	P68723	Beta-insect excitatory toxin LqhIT1c	9.9	<i>Leiurus quinquestriatus hebraeus</i>	In-gel digestion
	P59360	Neurotoxin BmK-II	7.2	<i>Mesobuthus martensii</i>	Both
	P15224	Toxin Os1	7.6	<i>Orthochirus scrobiculosus</i>	In-gel digestion
	D5HR50	Alpha-toxin Ac1 (Fragment)	8.7	<i>Androctonus crassicauda</i>	In-gel digestion
	M1JMR8	Sodium channel alpha-toxin Acra8	7.5	<i>Androctonus crassicauda</i>	Both
	M1JBC0	Sodium channel alpha-toxin Acra4	7.1	<i>Androctonus crassicauda</i>	In-gel digestion
	Q86SE0	Toxin Aam2	9.3	<i>Androctonus amoreuxi</i>	Both
	P21150	Toxin AaHIT4	7.8	<i>Androctonus australis</i>	Both
	P01482	Alpha-toxin Amm5	7.3	<i>Androctonus mauretanicus</i>	In-solution digestion
				<i>mauretanicus</i>	
	P01481	Alpha-mammal toxin Lqq5	7.3	<i>Leiurus quinquestriatus quinquestriatus</i>	In-solution digestion
	P13488	Alpha-like toxin Bom3	6.9	<i>Buthus occitanus mardochei</i>	Both
	P45698	Neurotoxin BmK-M9	8.8	<i>Mesobuthus martensii</i>	In-solution digestion
	P68721	Beta-insect excitatory toxin LqhIT1a	9.9	<i>Leiurus quinquestriatus hebraeus</i>	Both
	P0DJH8	Alpha-toxin Bu1	7.5	<i>Buthacus macrocentrus</i>	Both
	P83644	Toxin Lqh4	7.2	<i>Leiurus quinquestriatus hebraeus</i>	Both
	P01489	Alpha-toxin Lqq4	7.2	<i>Leiurus quinquestriatus quinquestriatus</i>	Both
				<i>quinquestriatus</i>	
	P01486	Alpha-toxin Bot11	7.5	<i>Buthus occitanus tunetanus</i>	In-solution digestion
	P60255	Toxin Boma6a	7.5	<i>Buthus occitanus mardochei</i>	Both
	P17728	Alpha-insect toxin LqhalT	9.6	<i>Leiurus quinquestriatus hebraeus</i>	Both
	P04098	Neurotoxin 8 (Fragment)	4.1	<i>Buthus occitanus tunetanus</i>	Both
	P55902	Alpha-insect toxin BotIT1	7.3	<i>Buthus occitanus tunetanus</i>	Both
	P01488	Alpha-toxin Bot1	7.3	<i>Buthus occitanus tunetanus</i>	Both
	P81504	Insect toxin AaHIT5	6.9	<i>Androctonus australis</i>	Both
	P01485	Alpha-mammal toxin Bot3 (Fragment)	8.1	<i>Buthus occitanus tunetanus</i>	In-solution digestion
	P83406	Neurotoxin Tx-2	2.9	<i>Buthus occitanus tunetanus</i>	In-solution digestion
	Q17254	Alpha-insect toxin Bot14	9.2	<i>Buthus occitanus tunetanus</i>	Both
	P59864	Beta-insect depressant toxin BotIT6	7.3	<i>Buthus occitanus tunetanus</i>	In-solution digestion
	P0C294	Toxin Acra I-3	8.8	<i>Androctonus crassicauda</i>	In-solution digestion
KScTx	B3EWX9	Potassium channel toxin alpha-KTx 9.11	2.9	<i>Mesobuthus gibbosus</i>	In-gel digestion
	P0C161	Potassium channel toxin alpha-KTx 2.8	4.3	<i>Centruroides elegans</i>	In-gel digestion
	B8XH42	Potassium channel toxin alpha-KTx 16.6	6.5	<i>Buthus occitanus israelis</i>	Both
	P0CC12	Potassium channel toxin alpha-KTx 8.5	3.2	<i>Odontobuthus doriae</i>	In-solution digestion
	P59869	Potassium channel toxin alpha-KTx 5.4	3.5	<i>Mesobuthus tamulus</i>	In-gel digestion
	B8XH40	Potassium channel toxin BuTXK-beta	10.2	<i>Buthus occitanus israelis</i>	In-gel digestion
	Q95NJ8	Potassium channel toxin alpha-KTx 17.1	6.2	<i>Odontobuthus doriae</i>	In-solution digestion
	P83407	Potassium channel toxin alpha-KTx 19.1	3.3	<i>Mesobuthus martensii</i>	In-solution digestion
	P80669	Potassium channel toxin alpha-KTx 9.3	3	<i>Leiurus quinquestriatus hebraeus</i>	In-solution digestion
	P86399	Neurotoxin lamda-MeuTx	7.2	<i>Mesobuthus eupeus</i>	In-solution digestion
	Q9NJC6	Potassium channel toxin BmTXK-beta	10.4	<i>Mesobuthus martensii</i>	Both
	Q9N661	Potassium channel toxin BmTXK-beta-2	10.2	<i>Mesobuthus martensii</i>	Both
	P01498	Neurotoxin P2	3.7	<i>Androctonus mauretanicus</i>	In-solution digestion
				<i>mauretanicus</i>	
ClScTx	P86436	Chlorotoxin-like peptide	3.6	<i>Androctonus australis</i>	Both
	P45639	Chlorotoxin	4	<i>Leiurus quinquestriatus quinquestriatus</i>	Both
				<i>quinquestriatus</i>	
Amphipathic peptide	P80670	Toxin GaTx2	3.2	<i>Leiurus quinquestriatus hebraeus</i>	In-solution digestion
	B8XH50	Amphipathic peptide Tx348	7.8	<i>Buthus occitanus israelis</i>	In-solution digestion

Peptide entries in bold were identified by both top-down and bottom-up approaches.

AMPs (1%), amphipathic peptides (1%), hypothetical secreted proteins (1%), and myotropic neuropeptides (1%) (Fig. 6).

The majority of described peptides were identified for the first time in this Moroccan *B. occitanus*

scorpion venom. The identified peptides showed sequence similarities with toxins previously detected from several genera of scorpions (Fig. 7), principally *Mesobuthus* sp (30%), *Buthus* Sp (20%), and *Androctonus* sp (18%).




Fig. 6. Summary of the total peptides identified by top-down and bottom-up approaches. The 102 peptides were divided into neurotoxins, including NaScTx, KScTx, ClScTx, CaScTx and toxin Acra, amphipathic peptide, myotropic neuropeptide, AMPs, and hypothetical secreted protein.




Fig. 7. Percentage of *B. occitanus* peptides, which showed similarity of sequences with others from several scorpion genera.

Discussion

Envenomation following scorpion stings constitutes one of the most encountered emergencies in large parts of the world, especially in North Africa, where the data show the highest incidence and lethality [1]. Morocco is a country known for a high risk of envenomation owing to its huge and diversified scorpion fauna. Among the different scorpion species living in this country, the yellow scorpion *B. occitanus* is one of the most dangerous species with venom responsible for severe cases of envenomation.

Due to the limited knowledge about the composition and toxin arsenal of *B. occitanus* venom, we aimed in this study to elaborate the first exhaustive view of this scorpion venom peptidome and its molecular diversity, using mass spectrometry-based top-down and bottom-up approaches.

Top-down data sets showed that the venom of *B. occitanus* is very complex, counting around 200 MWs ranging from 1868 to 16 720 Da. A similar number of components have been revealed by previous studies [32–34], others showed fewer components, as well as *Leiurus abdullahbayrami* (45 masses) and *Opisthacanthus elatus* (106 masses) [35, 36], whereas some other scorpion venoms were more complex, such as the *Pandinus cavimanus* (390 masses) and *Centruroides limpidus* (395 masses) [37, 38]. Additionally, the repartition of MWs showed that < 1% were components with molecular masses < 2000 Da, 14% were those from 2000 to 5000 Da, 74% were those between 5000 and 8000 Da, and 10% were those over than 8000 Da, while the repartition of MW from the French *B. occitanus* scorpion venom showed an abundance of molecules ranging from 2000 to 3000 Da and those less than 2000 Da [39]. Most importantly, the whole sequences of five toxins were identified with 100% sequence coverage using the top-down approach. These neurotoxins were detected for the first time in this venom; they all belong to the NaScTx category and shared high similarities of sequence with toxins identified from other scorpion species: neurotoxin BmK-II (P59360), beta-insect depressant toxin BotIT4 (P55903), beta-insect depressant toxin BaIT2 (P80962), insect toxin LqhIT5 (P81240), and insect toxin BsIT4 (P82814). It is important to stress that the observed sequence of the P59360 entry with a MW of 7431.33 Da showed 100% similarity with the sequence of neurotoxin BmK-II isolated from the Chinese scorpion *Mesobuthus martensii*, this neurotoxin is active in mammal and insect Nav channel [40]. In contrast, the detected sequence of the P81240 entry (6611.8 Da)

showed the presence of methionine in the N-terminal compared with the database sequence of the Insect toxin LqhIT5, an excitatory insect beta-toxin from the *Leiurus hebraeus* scorpion [41]. Similar to the P82814 entry (6954.15 Da), in which the observed sequence corresponds 100% to the insect toxin BsIT4, a depressant insect beta-toxins was isolated from *Hottentotta tamulus indicus* [42]. Also, the observed sequence of the peptide corresponding to the depressant toxin BotIT4 (6837.96 Da) presents methionine in N-terminal compared with the database sequence. This toxin, identified for the first time from the Tunisian *Buthus tunetanus* [43], showed also 100% sequence identity with the P80962 entry (6845.9 Da), referred to the beta-insect depressant toxin BaIT2 isolated from the *Buthacus arenicola* scorpion [44]. The high similarity of the amino acid sequence, in both detected depressant toxins and in the other peptides is commonly observed in scorpion toxins.

Interestingly, the combined top-down and bottom-up data sets of *B. occitanus* venom provide the identification of 102 different peptides, whereas 147 proteins were characterized from the yellow Brazilian scorpion *Tityus serrulatus*, 60 of which were detected by the top-down approach [45]. The major representative category of components identified in our venom was neurotoxins, mainly NaScTx (77%), these neurotoxins are abundant in species from the Buthidae family [38,46,47] and less representative in scorpions from the non-Buthidae family [33,48,49]. Those toxins are the ones responsible for envenomation symptoms [39]; their high content in the *B. occitanus* venom could explain the involvement of this scorpion in lethal cases of envenoming in the country.

Between the entries corresponding to NaScTx, there are alpha-like toxins, this type of toxins had been already identified in several *Buthus* sp; yet, the alpha-toxin Bot1 (P01488) has never been found in other Moroccan *Buthus* subspecies except from *Buthus mardochei* [39,50–53], but identified herein with a high sequence coverage (98.48% on top-down data set). We should mention also that we identified for the first time, in this scorpion venom, peptides corresponding to atypical NaScTx, as well as makatoxin-1, fragment from makatoxin-2, toxin Cg2, chain [20–87] in venom toxin meuNa32, and AaHIT4 toxin (which could bind on receptor site 3 or 4 of sodium channel) [33].

Besides NaScTx and KScTx (14%), ClScTx (3%) were identified, these categories of peptides showed activities against autoimmune disease and cancers, respectively [54–58]; also, we identified one entry that shared 60% of similarity with neurotoxin Tx-2 (P83406), a calcium channel activator identified for the

first time from the *Buthotus judaicus*, this category of toxins was identified in few scorpion species, for example, *Parabuthus transvaalicus* (Kurt toxin) and *Parabuthus granulatus* (Kurt toxin-like I) but never been detected in a Moroccan scorpion venom [59, 60]. And last but not least, peptides referring to toxin Acra category have also been screened in *B. occitanus* venom, these toxins probably acting on ion channels.

Some peptides with antibacterial activities were also found, for example, amphipathic peptide (B8XH50) and AMP AcrAPI (A0A059UI30); this category was commonly present in scorpion venom due to its role in the protection of venom glands and its involvement in the neurotoxic effects [61–65]. Additionally, other components were identified with a low percentage, such as orokinin, a myotropic neuropeptide identified from crustaceans, insects, and arachnids [17, 66], and hypothetical secreted proteins, which are proteins with unknown activities. Finally, we notice that some of the detected toxins were identified as fragments and chains, which may be due to the proteolysis of toxins. This process seems to be a usual PTM in scorpion and snake venoms, whereas its biological pertinence remains obscure [17, 45].

This study decrypted the peptidome arsenal of the Moroccan *B. occitanus* scorpion venom through proteomic view without the *de novo* sequence annotation. These findings constitute a step forward to a 'deeper' understanding of this scorpion venom; nevertheless, complete identification of this complex matrix is still a challenging task, especially with the lack of a specific database and/or a complete sequenced genome of this venom.

Conclusion

Herein, we reported the first proteomic study of the Moroccan *B. occitanus* scorpion peptidome, using mass spectrometry-based top-down and bottom-up venomic approaches. The combination of these approaches allowed the identification of 102 components classified, with approximation, on different categories, mainly neurotoxins (96%), including NaScTx (77%), KScTx (14%), ClScTx (3%), CaScTx (1%), and toxin Acra (1%). We also identified AMPs (1%), amphipathic peptides (1%), hypothetical secreted proteins (1%), and myotropic neuropeptides (1%). This study constitutes for sure a step forward to a deeper understanding of the *B. occitanus* venom; nevertheless, complete identification of this complex matrix is still a challenging task, especially with the lack of a specific database and a complete sequenced genome.

Acknowledgements

We declare that this study received financial support from the French government through the grant No 870634D.

Conflict of interest

The authors declare no conflict of interest.

Data Accessibility

All generated data during this study are included in this article.

Author contributions

NO and JCR conceived the research. KD and CM performed experiments. KD and CM analyzed the data. KD interpreted data and wrote the manuscript. AL, BD, and SC participated in writing. JMS and RC reviewed the manuscript. NO designed the project, supervised the study, and reviewed the manuscript. All authors read and approved the final version for publication.

References

- Chippaux JP (2012) Emerging options for the management of scorpion stings. *Drug Des Devel Ther* **6**, 165–173.
- Sunagar K, Undheim EA, Chan AH, Koludarov I, Muñoz-Gómez SA, Antunes A and Fry BG (2013) Evolution stings: the origin and diversification of scorpion toxin peptide scaffolds. *Toxins* **12**, 2456–2487.
- Possani LD, Becerril B, Delepierre M and Tytgat J (1999) Scorpion toxins specific for Na⁺ channels. *Eur J Biochem* **2**, 287–300.
- Rodríguez de la Vega RC and Possani LD (2005) Overview of scorpion toxins specific for Na⁺ channels and related peptides: biodiversity, structure–function relationships and evolution. *Toxicon* **8**, 831–844.
- Almaaytah A, Zhou M, Wang L, Chen T, Walker B and Shaw C (2012) Antimicrobial/cytolytic peptides from the venom of the North African scorpion, *Androctonus amoreuxi*: biochemical and functional characterization of natural peptides and a single site–substituted analog. *Peptides* **2**, 291–299.
- Cao L, Dai C, Li Z, Fan Z, Song Y, Wu Y, Cao Z and Li W (2012) Antibacterial activity and mechanism of a scorpion venom peptide derivative *in vitro* and *in vivo*. *PLoS One* **7**, e40135.
- El Oufir R (2019) Piqures et Envenimations Scorpioniques (PES) Centre Anti Poison du Maroc

- (CAPM). Rapports général et spécifiques. Revue toxicologie Maroc **43**:11.
- 8 Touloun O, Slimani T and Boumezzough A (2001) Epidemiological survey of scorpion envenomation in Southwestern Morocco. *J Venom Anim Toxins Incl Trop Dis* **7**, 99–218.
 - 9 Ghalim N, Ghalim N, Sebti F, El-Hafny B, Lazar N, Moustani R and Heikel J (2000) Scorpion envenomation and serotherapy in Morocco. *Am J Trop Med Hyg* **2**, 277–283.
 - 10 Aboumaâd B, Lahssaini M, Tiger A and Benhassain SM (2014) Clinical comparison of scorpion envenomation by *Androctonus mauritanicus* and *Buthus occitanus* in children. *Toxicon* **90**, 337–343.
 - 11 Oukkache N, El Jaoudi R, Ghalim N, Chgoury F, Bouhaouala B, Mdaghri NE and Sabatier JM (2014) Evaluation of the lethal potency of scorpion and snake venoms and comparison between intraperitoneal and intravenous injection routes. *Toxins* **6**, 1873–1881.
 - 12 Daoudi K, Chgoury F, Rezzak M, Bourouah O, Boussadda L, Soukri A, Sabatier JM and Oukkache N (2017) Consequences of *Androctonus mauretanicus* and *Buthus occitanus* scorpion venoms on electrolyte levels in rabbits. *Heliyon* **1**, e00221.
 - 13 Emerich BL, De Lima ME, Martin-Eauclaire M and Bougis PE (2018) Comparative analyses and implications for antivenom serotherapy of four Moroccan scorpion *Buthus occitanus* venoms: subspecies tunetanus, paris, malhommei, and mardochei. *Toxicon* **149**, 26–36.
 - 14 Pimenta AM, Stöcklin R, Favreau P, Bougis PE and Martin-Eauclaire MF (2001) Moving pieces in a proteomic puzzle: mass fingerprinting of toxic fractions from the venom of *Tityus serrulatus* (Scorpiones, Buthidae). *Rapid Commun Mass Spectrom* **17**, 1562–1572.
 - 15 Batista CV, D'Suze G, Gómez-Lagunas F, Zamudio FZ, Encarnación S, Sevcik C and Possani LD (2006) Proteomic analysis of *Tityus discrepans* scorpion venom and amino acid sequence of novel toxins. *Proteomics* **12**, 3718–3727.
 - 16 Favreau P, Menin L, Michalet S, Perret F, Cheneval O, Stöcklin M, Bulet P and Stöcklin R (2006) Mass spectrometry strategies for venom mapping and peptide sequencing from crude venoms: case applications with single arthropod specimen. *Toxicon* **6**, 676–687.
 - 17 Rates B, Ferraz KK, Borges MH, Richardson M, De Lima ME and Pimenta AM (2008) *Tityus serrulatus* venom peptidomics: assessing venom peptide diversity. *Toxicon* **5**, 611–618.
 - 18 Verano-Braga T, Rocha-Resende C, Silva DM, Ianzer D, Martin-Eauclaire MF, Bougis PE, de Lima ME, Santos RA and Pimenta AM (2008) *Tityus serrulatus* Hypotensins: a new family of peptides from scorpion venom. *Biochem Biophys Res Commun* **3**, 515–520.
 - 19 Hempel BF, Damm M, Mrinalini, Göçmen B, Karış M, Nalbantsoy A, Kini RM and Süßmuth RD (2020) Extended snake venomomics by top-down in-source decay: investigating the newly discovered Anatolian meadow viper subspecies, *Vipera anatolica senliki*. *J Proteome Res* **4**, 1731–1749.
 - 20 Compton PD and Kelleher NL (2012) Spinning up mass spectrometry for whole protein complexes. *Nat Methods* **11**, 1065–1066.
 - 21 Smith LM and Kelleher NL (2013) Consortium for top down proteomics. Proteoform: a single term describing protein complexity. *Nat Methods* **3**, 186–187.
 - 22 Mayne J, Ning Z, Zhang X, Starr AE, Chen R, Deeke S, Chiang CK, Xu B, Wen M, Cheng K *et al.* (2016) Bottom-up proteomics (2013–2015): keeping up in the era of systems biology. *Anal Chem* **1**, 95–121.
 - 23 Fornelli L, Toby TK, Schachner LF, Doubleday PF, Szrentić K, DeHart CJ and Kelleher NL (2018) Top-down proteomics: where we are, where we are going? *J Proteomics* **175**, 3–4.
 - 24 Melani RD, Skinner OS, Fornelli L, Domont GB, Compton PD and Kelleher NL (2016) Mapping proteoforms and protein complexes from king cobra venom using both denaturing and native top-down proteomics. *Mol Cell Proteomics* **7**, 2423–2434.
 - 25 Petras D, Heiss P, Harrison RA, Süßmuth RD and Calvete JJ (2016) Top-down venomomics of the East African green mamba, *Dendroaspis angusticeps*, and the black mamba, *Dendroaspis polylepis*, highlight the complexity of their toxin arsenals. *J Proteomics* **146**, 148–164.
 - 26 Melani RD, Nogueira FCS and Domont GB (2017) It is time for top-down venomomics. *J Venom Anim Toxins Incl Trop Dis* **23**, 44.
 - 27 Trevisan-Silva D, Bednaski AV, Fischer JSG, Veiga SS, Bandeira N, Guthals A, Marchini FK, Leprevost FV, Barbosa VC, Senff-Ribeiro A *et al.* (2017) A multi-protease, multi-dissociation, bottom-up-to-top-down proteomic view of the *Loxosceles intermedia* venom. *Sci Data* **4**, 170090.
 - 28 Estrada-Gomez S, Cardoso FC, Vargas-Muñoz LJ, Quintana-Castillo JC, Arenas Gómez CM, Pineda SS and Saldarriaga-Cordoba MM (2019) Venomic, transcriptomic, and bioactivity analyses of pamphobeteus verdolaga venom reveal complex disulfide-rich peptides that modulate calcium channels. *Toxins* **9**, 496.
 - 29 Ghezellou P, Garikapati V, Kazemi SM, Strupat K, Ghassempour A and Spengler B (2019) A perspective view of top-down proteomics in snake venom research. *Rapid Commun Mass Spectrom* **1**, 20–27.
 - 30 Oukkache N, Chgoury F, Lalaoui M, Cano AA and Ghalim N (2013) Comparison between two methods of scorpion venom milking in Morocco. *J Venom Anim Toxins Incl Trop Dis* **1**, 5.

- 31 Shevchenko A, Wilm M, Vorm O and Mann M (1996) Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal Chem* **5**, 850–858.
- 32 Miyashita M, Otsuki J, Hanai Y, Nakagawa Y and Miyagawa H (2007) Characterization of peptide components in the venom of the scorpion *Liocheles australasiae* (Hemiscorpiidae). *Toxicon* **3**, 428–437.
- 33 Luna-Ramírez K, Quintero-Hernández V, Vargas-Jaimes L, Batista CV, Winkel KD and Possani LD (2013) Characterization of the venom from the Australian scorpion *Urodacus yaschenkoi*: Molecular mass analysis of components, cDNA sequences and peptides with antimicrobial activity. *Toxicon* **63**, 44–54.
- 34 Cid-Urbe JI, Santibáñez-López CE, Meneses EP, Batista CVF, Jiménez-Vargas JM, Ortiz E and Possani LD (2018) The diversity of venom components of the scorpion species *Paravaejovis schwenkmeyeri* (Scorpiones: Vaejovidae) revealed by transcriptome and proteome analyses. *Toxicon* **151**, 47–62.
- 35 Erdeş E, Doğan TS, Coşar I, Danişman T, Kunt KB, Seker T, Yücel M and Ozen C (2014) Characterization of *Leiurus abdullahbayrami* (Scorpiones: Buthidae) venom: peptide profile, cytotoxicity and antimicrobial activity. *J Venom Anim Toxins Incl Trop Dis* **1**, 48.
- 36 Estrada-Gómez S, Vargas Muñoz LJ, Saldarriaga-Córdoba M and Quintana Castillo JC (2016) Venom from *Opisthacanthus elatus* scorpion of Colombia, could be more hemolytic and less neurotoxic than thought. *Acta Trop* **153**, 70–78.
- 37 Diego-García E, Peigneur S, Clynen E, Marien T, Czech L, Schoofs L and Tytgat J (2012) Molecular diversity of the telson and venom components from *Pandinus cavimanus* (Scorpionidae Latreille 1802): transcriptome, venomomics and function. *Proteomics* **2**, 313–328.
- 38 Cid-Urbe JI, Meneses EP, Batista CVF, Ortiz E and Possani LD (2019) Dissecting toxicity: the venom gland transcriptome and the venom proteome of the highly venomous scorpion *Centruroides limpidus* (Karsch, 1879). *Toxins* **5**, 247.
- 39 Martin-Eauclaire MF, Bosmans F, Céard B, Diochot S and Bougis PE (2014) A first exploration of the venom of the *Buthus occitanus* scorpion found in southern France. *Toxicon* **79**, 55–63.
- 40 Ji YH, Mansuelle P, Terakawa S, Kopeyan C, Yanaihara N, Hsu K and Rochat H (1996) Two neurotoxins (BmK I and BmK II) from the venom of the scorpion *Buthus martensi* Karsch: purification, amino acid sequences and assessment of specific activity. *Toxicon* **9**, 987–1001.
- 41 Moskowitz H, Herrmann R, Jones AD and Hammock BD (1998) A depressant insect-selective toxin analog from the venom of the scorpion *Leiurus quinquestriatus* hebraeus-purification and structure/function characterization. *Eur J Biochem* **1**, 44–49.
- 42 Ali SA, Stoeva S, Grossmann JG, Abbasi A and Voelter W (2001) Purification, characterization, and primary structure of four depressant insect-selective neurotoxin analogs from scorpion (*Buthus sindicus*) venom. *Arch Biochem Biophys* **2**, 197–206.
- 43 Borchani L, Stankiewicz M, Kopeyan C, Mansuelle P, Kharrat R, Cestèle S, Karoui H, Rochat H, Pelhate M and el Ayeb M (1997) Purification, structure and activity of three insect toxins from *Buthus occitanus tunetanus* venom. *Toxicon* **3**, 365–382.
- 44 Cestèle S, Kopeyan C, Oughideni R, Mansuelle P, Granier C and Rochat H (1997) Biochemical and pharmacological characterization of a depressant insect toxin from the venom of the scorpion *Buthacus arenicola*. *Eur J Biochem* **1–2**, 93–99.
- 45 Verano-Braga T, Dutra AA, León IR, Melo-Braga MN, Roepstorff P, Pimenta AM and Kjeldsen F (2013) Moving pieces in a venom puzzle: unveiling post-translationally modified toxins from *Tityus serrulatus*. *J Proteome Res* **7**, 3460–3470.
- 46 DeBin JA, Maggio JE and Strichartz GR (1993) Purification and characterization of chlorotoxin, a chloride channel ligand from the venom of the scorpion. *Am J Physiol* **264** (2 Pt 1), C361–C369.
- 47 Zeng XC, Peng F, Luo F, Zhu SY, Liu H and Li WX (2001) Molecular cloning and characterization of four scorpion K(+) toxin-like peptides: a new subfamily of venom peptides (alpha-KTx14) and genomic analysis of a member. *Biochimie* **9**, 883–889.
- 48 Abdel-Rahman MA, Quintero-Hernandez V and Possani LD (2013) Venom proteomic and venomous glands transcriptomic analysis of the Egyptian scorpion *Scorpio maurus palmatus* (Arachnida: Scorpionidae). *Toxicon* **74**, 193–207.
- 49 Quintero-Hernández V, Ramírez-Carreto S, Romero-Gutiérrez MT, Valdez-Velázquez LL, Becerril B, Possani LD and Ortiz E (2015) Transcriptome analysis of scorpion species belonging to the Vaejovis genus. *PLoS One* **2**, e0117188.
- 50 Volkova TM, Garsia AF, Telezhinskaia I N, Potapenko NA and Grishin EV (1985) Neurotoxins from the venom of the Central Asian scorpion *Buthus eupeus*. *Bioorg Khim* **11**, 1445–1456.
- 51 Vargas O, Martin MF and Rochat H (1987) Characterization of six toxins from the venom of the Moroccan scorpion *Buthus occitanus mardochei*. *Eur J Biochem* **3**, 589–599.
- 52 Koo GC, Blake JT, Talento A, Nguyen M, Lin S, Sirotna A, Shah K, Mulvany K, Hora D Jr, Cunningham P *et al.* (1997) Blockade of the voltage-gated potassium channel Kv1.3 inhibits immune responses *in vivo*. *J Immunol* **11**, 5120–5128.
- 53 Liu ZR, Ye P and Ji YH (2011) Exploring the obscure profiles of pharmacological binding sites on voltage-

- gated sodium channels by BmK neurotoxins. *Protein Cell* **6**, 437–444.
- 54 Lyons SA, O'Neal J and Sontheimer H (2002) Chlorotoxin, a scorpion-derived peptide, specifically binds to gliomas and tumors of neuroectodermal origin. *Glia* **2**, 162–173.
 - 55 Sidach SS and Mintz IM (2002) Kurtotoxin, a gating modifier of neuronal high- and low-threshold Ca channels. *J Neurosci* **6**, 2023–2034.
 - 56 Varga Z, Gurrola-Briones G, Papp F, Rodríguez de la Vega RC, Pedraza-Alva G, Tajhya RB, Gaspar R, Cardenas L, Rosenstein Y, Beeton C *et al.* (2012) Vm24, a natural immunosuppressive peptide, potently and selectively blocks Kv1.3 potassium channels of human T cells. *Mol Pharmacol* **3**, 372–382.
 - 57 Yue PJ, He L, Qiu SW, Li Y, Liao YJ, Li XP, Xie D and Peng Y (2014) OX26/CTX-conjugated PEGylated liposome as a dual-targeting gene delivery system for brain glioma. *Mol Cancer* **13**, 191.
 - 58 Misra SK, Ye M, Kim S and Pan D (2015) Defined nanoscale chemistry influences delivery of peptidotoxins for cancer therapy. *PLoS One* **6**, e0125908.
 - 59 Olamendi-Portugal T, García BI, López-González I, Van Der Walt J, Dyason K, Ulens C, Tytgat J, Felix R, Darszon A and Possani LD (2002) Two new scorpion toxins that target voltage-gated Ca²⁺ and Na⁺ channels. *Biochem Biophys Res Commun* **4**, 562–568.
 - 60 Carballar-Lejarazú R, Rodríguez MH, de la Cruz Hernández-Hernández F, Ramos-Castañeda J, Possani LD, Zurita-Ortega M, Reynaud-Garza E, Hernández-Rivas R, Loukeri T, Lycett G *et al.* (2008) Recombinant scorpine: a multifunctional antimicrobial peptide with activity against different pathogens. *Cell Mol Life Sci* **19**, 3081–3092.
 - 61 Pascual N, Castresana J, Valero ML, Andreu D and Bellés X (2004) Orcokininins in insects and other invertebrates. *Insect Biochem Mol Biol* **11**, 1141–1146.
 - 62 Hernández-Aponte CA, Silva-Sanchez J, Quintero-Hernández V, Rodríguez-Romero A, Balderas C, Possani LD and Gurrola GB (2011) Vejovine, a new antibiotic from the scorpion venom of *Vaejovis mexicanus*. *Toxicon* **1**, 84–92.
 - 63 Parente AMS, Daniele-Silva A, Furtado AA, Melo MA, Lacerda AF, Queiroz M, Moreno C, Santos E, Rocha HAO, Barbosa EG *et al.* (2018) Analogs of the scorpion venom peptide stigmurin: structural assessment, toxicity, and increased antimicrobial activity. *Toxins* **4**, 161.
 - 64 Das Neves RC, Mortari MR, Schwartz EF, Kipnis A and Junqueira-Kipnis AP (2019) Antimicrobial and antibiofilm effects of peptides from venom of social wasp and scorpion on multidrug-resistant *Acinetobacter baumannii*. *Toxins* **4**, 216.
 - 65 Amorim-Carmo B, Daniele-Silva A, Parente AMS, Furtado AA, Carvalho E, Oliveira JWF, Santos ECG, Silva MS, Silva SRB, Silva-Júnior AA *et al.* (2019) Potent and broad-spectrum antimicrobial activity of analogs from the scorpion peptide stigmurin. *Int J Mol Sci* **3**, 623.
 - 66 Hofer S, Dirksen H, Tollbäck P and Homberg U (2005) Novel insect orckokinins: characterization and neuronal distribution in the brains of selected dicondylar insects. *J Comp Neurol* **1**, 57–71.

Supporting information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Fig. S1. SDS/PAGE profile of the < 30 kDa filtrate of *Buthus occitanus* venom. Molecular weight markers (MM) are indicated in kDa. Proteins/Peptides were stained with Coomassie Brilliant Blue R (InstantBlue, Expedion, CA, USA). Stained bands corresponding to proteins/peptides with mass < 30 kDa were manually excised into equal small cubes of 1 mm³ and subjected to a nanoLC-MS/MS analysis.

Fig. S2. Detected amino acid sequences of the 68 peptides identified by Top-down approach.