

## Benthic nutrients and oxygen fluxes at the water sediment interface in a pearl farming atoll (Ahe, Tuamotu, French Polynesia)

Christian Grenz, Martine Rodier, Claire Seceh, David Varillon, Gabriel Haumani, Christel Pinazo

### ▶ To cite this version:

Christian Grenz, Martine Rodier, Claire Secch, David Varillon, Gabriel Haumani, et al.. Benthic nutrients and oxygen fluxes at the water sediment interface in a pearl farming atoll (Ahe, Tuamotu, French Polynesia). Marine Pollution Bulletin, 2021, 173, pp.112963. 10.1016/j.marpolbul.2021.112963 . hal-03360097

### HAL Id: hal-03360097 https://amu.hal.science/hal-03360097

Submitted on 30 Sep 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

### **Marine Pollution Bulletin**

# Benthic nutrients and oxygen fluxes at the water sediment interface in a pearl farming atoll (Ahe, Tuamotu, French Polynesia) --Manuscript Draft--

Manuscript Number:	MPB-D-21-00945R1						
Article Type:	VSI: MANA						
Keywords:	Benthic nutrient fluxes; sediment respiration; phytobenthic production; Atoll lagoon; pearl farming; French Polynesia						
Corresponding Author:	Christian Grenz, PhD, HDR Aix Marseille University Marseille, FRANCE						
First Author:	Christian Grenz, PhD, HDR						
Order of Authors:	Christian Grenz, PhD, HDR						
	Rodier Martine						
	Seceh Claire						
	Varillon David						
	Haumani Gabriel						
	Pinazo Christel, Assistant Professor						
Abstract:	Benthic exchanges of oxygen and nutrient at the sediment-water interface were investigated under light and dark conditions at 5 sites in a sub-tropical atoll. Mean oxygen fluxes were - $1316.5 \pm 242.0 \ \mu mol m \ -2 h \ -1$ and mean effluxes of oxygen under light conditions were $2231.7 \pm 626.4 \ \mu mol m \ -2 h \ -1$ , presumably due to microphytobenthos present at the sediment-water interface. The consequences of this high related productivity was a systematic consumption of nutrients (DIN, PO 4 and Si(OH) 4) during almost all light incubations, contrasting with the effluxes of nutrients during dark incubations. Our results suggest that sediments were net autotrophic and the oxygen balance in favor of microbenthic productivity were high ( $3423 \pm 1192 \ \mu mol m \ -2 h \ -1$ ) which emphasize the role of microphytobenthos in maintaining the oxygen reservoir in tropical lagoons.						

Benthic Nutrients and oxygen fluxes at the water sediment interface in a pearl farming atoll (Ahe, Tuamotu, French Polynesia) Grenz Christian<sup>a,\*</sup>, Rodier Martine<sup>a,b</sup>, Seceh Claire<sup>a</sup>, Varillon David<sup>c</sup>, Haumani Gabriel<sup>d</sup>, Pinazo Christel<sup>a</sup> б <sup>a</sup> Aix-Marseille Université, Université Toulon, CNRS/INSU, IRD, Mediterranean Institute of Oceanography MIO UM 110, Marseille, France <sup>b</sup> Institut de Recherche pour le Développement, UMR 241 EIO (Université de la Polynésie française, IRD, ILM, Ifremer), B.P. 6570, 98702 Faa'a, Tahiti, French Polynesia <sup>c</sup> Institut de Recherche pour le Développement, US 191 IMAGO, BP A5, 98848 Nouméa cedex, New-Caledonia, France <sup>d</sup> Direction des Ressources Marines, Antenne de Takapoto, BP 20, 98713 Papeete, French Polynesia \* Corresponding author. Email addresses: christian.grenz@mio.osupytheas.fr (C. Grenz), martine.rodier@ird.fr (M. Rodier), claire.seceh@mio.osupytheas.fr (C. Seceh), david.varillon@ird.fr (D. Varillon), gabriel.haumani@gmail.com (G Haumani), christel.pinazo@mio.osupytheas.fr (C. Pinazo). Keywords: Benthic nutrient fluxes, sediment respiration, phytobenthic production, Atoll lagoon, Pearl farming, French Polynesia ABSTRACT Benthic exchanges of oxygen and nutrient at the sediment-water interface were investigated under light and dark conditions at 5 selected sites in a sub-tropical atoll. Mean oxygen fluxes were -  $1316.5 \pm$ 242.0  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> and mean effluxes of oxygen under light conditions were 2231.7 ± 626.4  $\mu$ mol m<sup>-2</sup>  $h^{-1}$ , presumably due to microphytobenthos present at the sediment-water interface. The consequences of this high related productivity was a systematic consumption of nutrients (DIN, PO<sub>4</sub> and Si(OH)<sub>4</sub>) during almost all light incubations, contrasting with the effluxes of nutrients during dark incubations. Our results suggest that the sediments were net autotrophic and the oxygen balance in favor of microbenthic production when compared to community demand. Diurnal rates of gross benthic primary productivity were high (3423  $\pm$  1192  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup>) which emphasize the role of microphytobenthos in maintaining the oxygen reservoir in tropical lagoons. 1. Introduction Sediment compartment contributes intensely to the biogeochemical cycling of organic matter through mineralization of settled particles driven by living benthic macro and microorganisms (Aller & Cochran, 2019; Middelburg et al., 1993; Middelburg, 2018). These processes have a profound effect on the local and global cycling of many elements (Burdige, 2006) especially in coastal areas due the shallowness of the water column and the high primary production compared to open ocean (Kopp et al., 2015). Benthic pelagic coupling processes play a major role in the balance between carbon preservation and 

remineralization (Ehrnsten et al., 2019). Moreover, most of the microbial degradation and nutrient recycling occurs at the sediment-water interface where benthic microbes are confined and gather most valuable electron acceptors for organic matter degradation (Canfield et al., 2005). As a consequence, recycled nutrients can return to surface waters via mixing processes (Caffrey et al., 2002) and supply potentially the pelagic N and P demand for phytoplankton production (Grenz et al., 2003; Boynton et al., 2018). 

As the pelagic plankton represents the main portion of the black pearl oyster (*Pinctada margaritifera*) diet for juveniles and adults, productivity of oyster farming depends on pelagic phytoplankton biomass. Pelagic primary production was shown to be nitrogen limited in Ahe atoll (Charpy et al., 2012; Pagano et al., 2017). Gaertner-Mazouni et al. (2012) estimated that benthic nitrogen fluxes could contribute up to 28% of the pelagic nitrogen demand. 

Due to the shallowness of coastal environments, benthic primary production may be significant at some sites, even where only small amounts of light reach the sediment surface (Jahnke et al. 2000). Microphytobenthos (MPB), a mixture of benthic unicellular algae and cyanobacteria (MacIntyre et al, 1996) significantly contributes to primary production, and it is responsible for more than 50% of the total primary production in some shallow waters (Underwood & Kromkamp, 1999; Underwood, 2002). Indeed, benthic primary production can considerably moderate the sediment-water fluxes of nutrients (Sundbäck et al., 1991) and compete with the macroalgal mats for the sediment nutrient pool (Sundbäck and Miles, 2002). In some sub-tropical lagoons, photosynthetic oxygen production by MPB enhance oxygen availability and increase oxygen penetration into the sediment favouring coupled nitrification-denitrification within the sediments (An and Joye 2001). 

Sub-tropical atoll lagoons are generally highly productive ecosystem compared to the surrounding oligotrophic ocean (Harcher 1997). Their shallowness and the very high amount of light at sediment surface may facilitate the microphytobenthic primary production and nutrient regeneration and accentuate the asymmetric light-nutrient competition between benthic and pelagic primary producers in freshwater (Jäger and Diehl, 2014) and marine (Chatterjee et al., 2013) systems. 

The aim of this study was to quantify spatial patterns in oxygen and nutrient exchanges at the sediment
 water interface and the role of MPB in nutrient cycling in a sub-tropical lagoon submitted to intense
 black pearl oyster cultivation in French Polynesia.

### 40<br/>41692. Materials and Methods

### <sup>42</sup><sub>43</sub> 70 *2.1. Context, site description and water column sampling*

This work is a contribution of the MANA project which aim is to produce new knowledge and develop numerical tools designed to support management decisions in the context of exploited atoll lagoons affected by the impact of climate change. Ahe atoll like other lagoons of the Central Pacific Ocean is dedicated to aquaculture and specifically the culture of the black lip oyster Pinctada margaritifera that produces the prized black pearls.

Ahe atoll located 456 km North-East from Tahiti, is one of the 76 coral atolls of the Tuamotu Archipelago (French Polynesia) and, like most of these, is characterized by an internal lagoon. Measuring around 20 km in length and up to 12 km across, the lagoon covers an area of 138 km<sup>2</sup> bordered by a land area of around 12 km<sup>2</sup>. Like all the atolls of the region, Ahe atoll rises sharply from the seabed (2700 m depth) and is connected to the open ocean by a 200 m wide pass (Tiareroa pass) located on the north-western side of the atoll rim. The mean depth is around 40 m and water renewal time was estimated at 252 days (Dumas et al., 2012). Ahe atoll is exposed year round to easterly trade winds and tidal amplitudes are small (Andréfouët et al, 2012a) but exhibits some seasonal variations (Bouvy et al., 2012; Charpy et al., 2012; Lefèbvre et al., 2012; Thomas et al., 2010). The general climate 

is characterized by a wet and a dry season and dominant trade winds (Dutheil et al., 2020). Our experiments took place at the beginning of the hot season during a period of moderate trade winds.

Five study sites (Fig. 1) were studied during a field campaign in Ahe lagoon between 27 November and 7 December 2017, chosen to reflect a range of different bathymetric (lesser than 20 m depth) and environmental conditions related to pearl culture. At each station a CTD profiler (SBE 19+, SeaBird) was used to record a vertical profile from surface to the bottom of temperature, salinity and PAR (Photosynthetic Available Radiation). Bottom water samples for nutrient analyses were sampled with a 5L Niskin bottle operated by the divers at the beginning of each incubation.



Fig. 1. Location of the sampling stations (black dots) in Ahe atoll (Tuamotu Archipelago) during the field trip in Nov – Dec 2017. Concessions for oyster breeding (dark grey) and spat collection (light grey) were redrawn from Direction des Ressources Marines (Pu Fa' Ahuta Moana). Tiareroa pass (North) and the village (South) are shown along the atoll rim.

2.2 Sediment core sampling and analyses

55 116 Sediments were sampled in each station by scuba divers using 3 cores (20 cm high, diameter 3 cm) and 56 117 the upper cm of the core was analysed for water content, porosity and pigments. A first surface level of samples was wet and dry weighted for water content. Porosity was calculated using the classical formulation of Berner (1980):

WW - DW1.035 POR = -WW – DW (DW -FW+1.036

Where WW is the sample wet weight, DW dry weight, FW empty flask weight, and considering a water density of 1.036 (mean salinity 36.3) and a bulk density of 2.56 g cm<sup>-3</sup>. 

Pigments were determined on a second 1 cm thick sediment sample, measured fluorometrically before and after acidification (methanol extraction) and using a Turner Design fluorometer equipped with the 10 125 module No 7200-040 (Le Bouteiller et al., 1992). A third surface level (3 cm thick) was used to extract under pressure the interstitial water. Water samples were immediately poisoned by HgCl2 (at 6g/l) and stored in an icebox and analysed while back at lab. Concentrations of nitrate and nitrite (NOx), 14 128 phosphate and silicates were determined by colorimetric methods using a segmented-flow Auto-Analyzer (AA3, SEAL corporation, Mequon, WI, USA), as described in Raimbault et al. (1990) and Aminot and Kérouel (2007). The detection limits for NOx, PO<sub>4</sub> and NH<sub>4</sub> were 0.05, 0.02, and 0.01 µmol L<sup>-1</sup> respectively.



Fig. 2. Clear and opaque benthic chambers placed on the sediment. Control chamber (bottom right).

#### 2.3 Fluxes at the sediment water interface

Benthic primary production and respiration as well as fluxes of dissolved nutrients were measured using opaque and transparent benthic chambers. We used similar devices as Barranguet et al (1996) slightly adapted to host new equipment. Briefly, the 37 cm diameter chambers (Fig. 2) are hemispherical domes made of polymethyl methacrylate isolating a water volume of 13.3 L over a sediment area of 0.108 m<sup>2</sup>. The chambers are clipped on cylindrical PVC frames equipped with an O-46 147 ring to avoid any leaking. The 10 cm high frames are inserted into the sediment depending on the toughness of the bottom. On top of the chamber, watertight housing hosts a battery powered motor and is coupled to a magnetic stir bar to homogenize the enclosed volume, preventing the formation of 50 150 a gradients within the chamber (Tengberg et al., 2004). During deployment, divers control visually that **151** no resuspension is artificially created. A diffusive oxygen probe (SDOT – NKE equipped with an Aanderaa 3835 Optode) is inserted in a lateral sampling port for dissolved Oxygen (DO) and temperature measurements. A second sampling port at the opposite position is used by divers to **154** withdraw water sub-samples (120 ml syringe equipped with a 10 cm silicone tube) during the in situ incubations. When a sample is withdrawn, it fills up with external water that remains isolated from the incubated water, so that it avoids a pressure change inside the chamber. The port holes are sealed with stoppers between the samplings. Control chambers equipped with a round plate isolating the chamber space from the bottom are used in parallel as a "blank" correction and to separate water-

column processes from benthic fluxes. As suggested by Foster et al. (2016), between 2 and 3 1 160 observational chambers and a control chamber (clear and opaque respectively) were used during every sampling occurrence as well as recordings of ambient conditions (outside the chambers).

DO content and temperature were continuously recorded at a 1 min frequency during 4 h cycles. The oxygen concentrations in the chambers never fell below 85% (respiration) or above 125% (production) of the initial O<sub>2</sub> content. Nutrient sampling was performed by divers every 1 h, during the first experiment to test the linear uptake or release of nutrient from the sediments. During the following experiments, enclosed water was sampled at the beginning and the end of each 4 h-incubation period. This start-end strategy was favored in order to reduce the number of samples (Dalsgaard et al., 2000). 

<sub>13</sub> 168 Fluxes at the sediment water interface were calculated by regressing the change of oxygen 14 169 concentrations in the chambers against time ( $\mu$ mol O<sub>2</sub> m<sup>-2</sup> h<sup>-1</sup>). A correction for pelagic processes was <sup>15</sup> 170 applied by subtracting the changes calculated for the control chambers. The equation used was as follows:

$$\frac{\left(\frac{\Delta C_{1}}{\Delta t} Vol1\right) - \left(\frac{\Delta C_{2}}{\Delta t} Vol2\right)}{Surf}$$

22<sup>-</sup> 173 Where  $\Delta C/\Delta t$  represent the uptake or release of O<sub>2</sub> and nutrients in the chambers (µmol L<sup>-1</sup> h<sup>-1</sup>), Vol the total volume of the chambers (L<sup>1</sup>) and Surf the area of the sediment enclosed (m<sup>2</sup>). Subscript 1 and **174** <sup>24</sup> 175 2 correspond to observational chambers and control chambers, respectively. The volume of the PVC base frames was added to the volume of the dome for Vol1. The volumes were corrected to take **177** account of the space occupied by the equipment (mixing device and oxygen probes). Once the syringe 28 178 retrieved on board, water samples were filtrated on GFF filters and immediately poisoned by HgCl<sub>2</sub> (at 6g/l) and stored in an icebox for later analysis of nutrients using an autoanalyser (same methods as for incubation samples - see above 2.2). 

#### 3. Results

#### 3.1 Environmental settings

Our study period was characterized by a hot season during the austral winter in 2017, with only few <sub>38</sub> 184 rain events observed. Table 2 shows the main environmental conditions during the sampling period. **185** The temperature of the water column was vertically homogenous between 28.4 and 28.9 °C (mean <sup>40</sup> 186  $28.6 \pm 0.2$ °C) and showed a slight increase at the south-western part of the lagoon. Bottom water nutrient concentrations were low and typical of oligotrophic conditions. Oxygenation of the water masses was efficient with O<sub>2</sub> saturation reaching between 83 and 97 %. Light conditions at the bottom 44 189 were optimal, between 17 and 42 % of the surface irradiance depending on depth and cloud cover. 

#### 46 190 Table 1

Physical and chemical features of the bottom waters measured at the five sampling sites in the Ahe lagoon. DO and PAR correspond to values measured outside the chambers and averaged over the incubation period.

94										
95	Statior	ns T (°C)	Depth (m)	Nutrients (µmol L <sup>-1</sup> )			DO (µmol L <sup>-1</sup> )	DO % sat	PAR (%)	
96				$NH_4$	NOx	$PO_4$	Si(OH)4			
	LO	28.7	9	0.07	0.29	0.29	1.41	233.5	96.7	42
	L1	28.9	13	0.05	0.13	0.24	1.36	206.2	85.8	17
	L8	28.4	15	0.05	0.22	0.35	1.29	210.5	86.7	30
	L4	28.5	18	0.03	0.33	0.35	1.68	213.9	88.1	30
	L10	28.4	15	0.08	0.59	0.51	1.53	202.9	83.0	34

Sediments were characterized by fine grained white sands with a mean water content of 15.8 % and a 1 203 porosity of 0.32. Nutrients in interstitial waters were high for NH<sub>4</sub> (between 30.6 and 85.1  $\mu$ mol L<sup>-1</sup>) and PO<sub>4</sub> (8.1 to 30.5  $\mu$ mol L<sup>-1</sup>) and low for NOx (0.7 to 1.4  $\mu$ mol L<sup>-1</sup>). Silicates showed intermediate values (from 1.6 to 10.0  $\mu$ mol L<sup>-1</sup>). Sediment pigments (chlorophyll a and phaeopigments) were comparable between stations L1, L8 and L10 and higher for station L4 (Table 2).

#### Table 2

Sediment characteristics in the first sediment layer at sampling sites L1 to L10. Water Content (WC)

in %, porosity (Poro) dimensionless, interstitial nutrients in µmol L<sup>-1</sup>, sediment chlorophyll a (SCH) 

and phaeopigments (SPH) in mg  $m^{-2}$  and %CH is the percentage of Chla over total pigments.

Interstitial nutrient were extracted from a 0-3 cm sediment layer and 0-1 cm for the other variables.

213	Stations	WC	Poro	Interst	Interstitial nutrients (µmol.L <sup>-1</sup> )					%CH
214				$NH_4$	NOx	PO <sub>4</sub>	Si(OH)4			
215	L1	16.7	0.33	68.2	1.2	30.5	7.2	2.99	2.87	51
216	L8	16.5	0.33	65.1	1.4	8.1	10.0	2.79	2.66	51
217	L4	14.8	0.30	85.1	ND	ND	ND	11.30	5.42	67
218	L10	15.1	0.30	30.6	0.7	13.3	1.6	2.21	3.83	37
219	Station LO samples were lost. ND: not determined.									

Station LO samples were lost. ND: not determined.

3.2 Sediment oxygen exchanges and nutrient fluxes at the sediment-water interface

The use of dark and transparent benthic chambers allows to infer both sediment oxygen demand (SOD) and light O<sub>2</sub> fluxes (LOF). Results (Fig. 3) showed the variability observed between the different stations. Station 8 showed the highest rates in SOD and LOF, -1776.5 (sd 169.9) and 3303.2 (sd 534.0) µmol m<sup>-2</sup> h<sup>-1</sup> respectively. Stations L4 and L10 showed similar values around -1307.2 and 1057.8 µmol  $m^{-2}$  h<sup>-1</sup> for SOD respectively, and 1934.2 and 2003.6 µmol m<sup>-2</sup> h<sup>-1</sup> for LOF respectively. Stations LO and L1 showed the lowest values of 1167.7  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> for mean LOF, but different values for SOD (228.9 and 1124.6 µmol m<sup>-2</sup> h<sup>-1</sup>). Unfortunately we got only one replicat for both clear and dark chambers at station LO. Excluding this station we calculated a mean SOD of -1316.5  $\pm$  242.0  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> and a mean LOF of 2231.7  $\pm$  626.4  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup>.



Fig. 3. Sediment oxygen fluxes measured during light (light grey) and dark (dark grey) incubations at the different stations (vertical bars indicate sd for triplicates).

**237** 

**239** 

**236** 

**238** 

58 240 



Nutrient fluxes showed also a large variability between replicate chambers and stations (Fig. 4). 1 274 Concerning nitrogen, dark chambers showed generally an efflux directed towards the water column. Conversely, clear chambers showed an influx at all stations. DIN fluxes representing the sum of NH<sub>4</sub> and NOx fluxes were largely driven by NOx fluxes because these fluxes were an order of magnitude higher than NH<sub>4</sub> fluxes. Highest DIN effluxes in dark chambers were recorded at stations L0, L1 and L8 б (138.5, 49.7 and 92.8 µmol m<sup>-2</sup> h<sup>-1</sup> respectively) whereas stations L4 and L10 showed fluxes close to zero. Highest DIN influxes in clear chambers were recorded at L1 and L8 (-88.4 and 119.9 µmol m<sup>-2</sup> h<sup>-1</sup> respectively) whereas these were close to zero at LO and slightly negative for L4 and L10 (-11.3 and -10 281 28.3  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> respectively). It is noteworthy that the variability between replicate chambers is elevated enabling any statistical significant comparison test. 

<sup>13</sup> 283 For PO<sub>4</sub> fluxes (Fig. 5A), the same observations were made with a general efflux in the dark chambers and an influx in the clear chambers. As for N-nutrients, the huge variability enables to test **285** statistically the difference between the stations. The range of values we found is between 0.5 and 4.9  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> in the dark chambers and between -0.2 and -3.8  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> in the clear chambers.

19 287 Finally, silicate fluxes (Fig. 5B) were positive most of the time in the dark chambers (efflux) and negative (influx) in clear chambers. Again the inter-chamber variability was high but the general trend observed for the dark chambers was a higher efflux of silicates at station 8 (94.9 µmol m<sup>-2</sup> h<sup>-1</sup>) and a lower Si flux for all other stations (-7.3, 15.4, 10.6 and 16.5  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> respectively for stations L0, L1, L4 and L10). **290** In the clear chambers the Si influxes were also highly variable, and showed the same trend with a highest influx observed for station L8 (-49.1  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup>) and the lowest at the other stations (-20.8, 3.7, -8.3 and 11.9  $\mu$ mol m<sup>-2</sup> h<sup>-1</sup> respectively for stations L0, L1, L4 and L10). 

#### 4. Discussion

Ahe atoll is one of the most thoroughly studied lagoon of the Tuamotu Archipelago (Andréfouët et al., 2012b). Nevertheless, nutrient fluxes and SOD measurements at the sediment water interface are poorly documented in Ahe atoll. Gaertner-Mazouni et al. (2012) measured nutrient fluxes at 4 stations in October 2010. Lacoste & Gaertner-Mazouni (2016) supplemented the data of 2010 by new ones obtained at 3 similar stations in May 2013. To our knowledge no published data are available regarding SOD or microphytobenthic production in this system.

41 302 Nutrient fluxes we measured in the dark chambers were in the same range of variation than those published previously by Lacoste & Gaertner-Mazouni (2016) in the case of ammonia and slightly higher for NOx. In New Caledonia lagoon, Boucher and Clavier (1990) and Grenz et al. (2010) observed a 45 305 similar range of nutrient fluxes, with a prevalence of fluxes oriented towards the water column indicating a net source of nutrients for the pelagic compartment. Through mineralization processes of organic matter at the sediment interface, the release of dissolved nitrogen concurrently to nitrification processes should lower the accumulation of NH<sub>4</sub> and raise NOx which in turn diffused to the overlaying **309** waters. Dunn et al. (2012) measured SOD and DIN effluxes during light and dark incubations in a subtropical coastal lagoon (Coombabah Lake, Australia). They concluded that rates of total nitrate reduction were relatively low, ranging from 3 to 26  $\mu$ mol N m<sup>-2</sup> h<sup>-1</sup> and exhibited only minor seasonal **312** variations. Finally, they stated that dissimilatory nitrate reduction to ammonium (DNRA) was the <sup>55</sup> 313 dominant pathway for nitrate reduction and denitrification processes seemed a less effective process probably because of the high amount of oxygen available (Sørensen, 1978). Same conclusions were <sub>58</sub> 315 made by Sundbäck et al. (1991) who performed a laboratory continuous-flow study with sediments 59 316 from SE Kattegat maintained at 15°C. In our case we measured lower concentrations of NOx compared to ammonium in the interstitial water in the sub-surface sediment layer. 

Silicate and phosphate fluxes in dark were in the same order of magnitude than those found by 1 319 Gaertner-Mazouni et al. (2012) for the same lagoon and as for Terminos lagoon in Mexico during the dry season (Grenz et al., 2019). Silicate release by sediments is a fast reaction possibly resulting from dissolution of biogeogenic Si from freshly deposited diatom frustules (Willey, 1978, Kamatani, 1982). 

During light incubations DIN, PO<sub>4</sub> and silicate fluxes were oriented towards the sediment. This could be related to the photosynthetic processes of the microphytobenthos at the sediment water interface. The mean DIN/PO<sub>4</sub> flux ratio was estimated around 16.6 during day light which is close to the Redfield ratio(16/1) underlying a potential sink of nutrients driven by microphytobenthic production. During <sub>11</sub> 326 the dark incubation this ratio increased to 45.6 showing that the proportion of nitrogen produced by 12 327 remineralization is higher than the phosphate produced. The loss of phosphate can be due to other processes like phosphorous adsorption by iron or manganese oxides (Sarrazin et al., 1988, Jensen and Thamdrup, 1993, Van Cappellen and Wang 1996) and/or reduced dissimilatory sulfate reduction (Pan 16 330 et al., 2020). Conversely, an increase in the nitrogen pool, due to a reduced assimilation of N by microphytobenthos in the dark, can enhance the available N in the interstitial waters and subsequently modify the solute exchange rates between the sediment and the overlaying water column. <sub>20</sub> 333 Nevertheless firm conclusions cannot be drawn as the calculated ratios were highly variable between the individual flux measurements (Ratio confidence intervals of  $\pm$  144.8 and  $\pm$  57.0, respectively for 21 334 light and dark incubations). 

#### Table 3 Example of benthic oxygen consumptions (min max) measured in subtropical shallow aquatic systems.

Site	Latitude	O₂ (µm	ol m <sup>-2</sup> h <sup>-1</sup> )	Author
Shing-Mun River, Hong Kong	22°23N	1188	18988	Chen et al., 2000
Tolo Habor	22°25N	607	1653	Chau, 2002
New Caledonia lagoon	22°22N	550	1570	Grenz et al., 2003
Nichupte, Mexico	21°70N	250	12875	Valdes-Lozano et al., 2006
Goa, W Coast India	15°27N	1460	4120	Pratihary et al., 2009
Pompey Reef, Australia	21°00S	821	3400	Alongi et al., 2011
Cochin Backwater System	9°49N	995	2267	Abhilash et al., 2012
Semariang Batu River, Malaysia	1°37N	982	14803	Ling et al., 2009
Ahe lagoon	15°00N	1150	3300	This study

The sediment oxygen demand (SOD) is a good measure for total system mineralization as oxygen in 44 350 sediments is utilized both directly for organic matter respiration (aerobic mineralization) and for <sup>45</sup> 351 reoxidation of reduced substances formed by anoxic mineralization (Rasmussen and Jorgensen, 1992). Our SOD values recorded in Ahe lagoon were in the range of observations made in New Caledonia lagoon (Clavier et al., 2005, Grenz et al., 2010) and other subtropical systems (Table 3). Temperature 49 354 combined with organic matter influx to the sediments control benthic mineralization logically driving coastal tropical lagoons in the upper range of SOD values, as demonstrated by the maximum values reported from shallow rivers in Hong Kong and Malaysia (Chen et al. 2000, Ling et al. 2009), and from **357** coastal lagoons near Cancun (Valdes-Lozano et al. 2006). Gaertner-Mazouni et al. (2012) measured low <sup>54</sup> 358 organic matter content in Ahe lagoon sediment, underneath oyster cultures and at a control station (around 3 %). This low values indicated that organic matter deposition produced by oyster biodeposition or natural sedimentation from the pelagic compartment is not stored in the sediment **361** they sampled but either transported and deposited elsewhere, or metabolized by the local benthic community. The low organic content can also be related to the dilution by terrigenous particles but in 

the case of atoll lagoons with low terrestrial inputs, this process should be restricted in space as in Tahiti lagoon (Frouin 2000). The low range in temperature variation encountered during our measurements precludes any temperature dependency related to our SOD estimates but the elevated temperature of the bottom waters suggests optimal conditions for microbial-mediated benthic degradation processes (Hicks et al., 2018).

б 368 Chlorophyll pigments indicated densities of live microphytobenthos while phaeopigments, the amount 7 8 369 of decaying microalgal material present in the sediments. We found pigment concentrations in the 9 370 upper sediment layers close to the lowest values reported for Nouméa lagoon (Garrigue, 1993) and for 10 <sub>11</sub> 371 Tikehau atoll, French Polynesia (Charpy and Charpy-Roubaud, 1990) but equivalent to the 12 372 concentrations measured in Terminos lagoon at stations distant from river discharge and urbanized 13 373 areas (Grenz et al., 2019). A mean ratio of 1:1 was calculated between Chla versus phaeopigments 14 374 indicating a microphytobenthos in relative healthy conditions. 15

16 375 Heil et al.(2004) showed for coral reef sediments that benthic microalgal biomass was strongly 17 18 376 correlated with oxygen production by sediments. Moreover they showed that microphytobenthos 19 377 dominated over pelagic biomass by a factor of around 100, when integrated through the water depth. 20 378 Taking the pelagic Chla measurements reported for the same cruise by Rodier et al. (2021), i.e. 21 379 between 2.3 and 5.0 mg m<sup>-2</sup>, we found about the same biomasses in the surficial sediments, underlying 22 23 **380** the importance of this compartment for the ecosystem.

24 25 **381** Incubations in clear chambers enable to quantify the exchanges of nutrients under light conditions but 26 382 also the microphytobenthos production during the diurnal period. The net oxygen production (NOP) 27 383 was always higher than SOD suggesting that the sediments were net autotrophic and the oxygen 28 384 balance in favor of microbenthic production when compared to community demand. This has been 29 30 385 observed in New Caledonia lagoon by Clavier et al. (2005), but not by Dunn et al. (2012). These later 31 386 authors observed that oxygen fluxes under light or dark conditions were consistently directed towards 32 387 the sediment. In our case, light O<sub>2</sub> fluxes were positive and diurnal rates of gross benthic primary 33 productivity ( $\Delta O_2$  = difference between the light and dark  $O_2$  fluxes) were high (3423 ± 1192  $\mu$ mol m<sup>-2</sup> 34 388 <sup>35</sup> 389  $h^{-1}$ ). Our values are close to the gross productivities observed by Dunn et al. (2012) at 2 stations in 36 390 summer and autumn, but lower than the gross productivity measured by Eyre et al. (2011) in southern 37 391 Moreton Bay, Australia, especially in summer. To our knowledge, there is no data available on benthic 38 39 **392** microalgal composition for Ahe lagoon. This lack of data should be addressed in future studies as 40 393 benthic microalgae are key components of coral reef ecosystems (Heil et al., 2004) and probably also 41 394 in atoll lagoons. 42

43 395 As already stated by Gaertner-Mazouni et al. (2012), the impact of oyster culture on the sediments of 44 396 Ahe lagoon is limited, due to a low sedimentation rate of organic matter underneath the spat collectors 45 46 397 and breading lines. Our data on nutrient fluxes confirm the order of magnitude already published but 47 398 at a larger scale. The impact of oyster culture on sediment metabolism seems to be reduced compared 48 399 to other areas where intense shellfish aquaculture at high stocking density disturb ecological systems 49 50 400 (see review of Dumbauld et al., 2009), moreover in poorly flushed bays. This has some direct 51 401 implications in the management of pearl oyster farming, especially in the context of the assimilative 52 402 capacity for organic loading to the sediment and the subsequent benthic metabolism (Ferreira et al., 53 403 2018). 54

Finally, benthic gross productivity/respiration (p/r), a measure of the balance between autotrophic production and heterotrophic respiration in the benthic communities, were almost > 1 (between 1.2 and 1.7) underlying the net autotrophic character at the sediment-water interface and the prominent role of the sediment compartment in the lagoon wide oxygen balance, in the tropics.

61 62

63 64 65

1

25

### <sup>1</sup>/<sub>2</sub> 409 **5. Conclusion**

3 410 This study investigated the nutrient and oxygen fluxes at the sediment water of Ahe atoll in the 4 5 411 Tuamotu Archipelago (French Polynesia). The use of clear and dark chambers incubated in situ allowed 6 412 to quantify simultaneously sediment oxygen demand (SOD) and microphytobenthic (MPB) production 7 413 as net oxygen production (NOP). Baring in mind the high variability encountered and the lack of spatial 8 9 414 replicates, our results showed some interesting points.

10 415 NOP were consistently directed towards the water column at all sampling stations whereas SOD
 11 416 logically pointed to the opposite direction, with a lower rate. Gross oxygen production (GOP) may
 13 417 contribute to the reported high oxygen saturation rates. The ratio between GOP and SOD over 1
 14 418 underlined the autotrophic state of the sediment water interface where light conditions are optimal.

<sup>15</sup> 419 We observed nutrient effluxes during almost all dark incubations and a sink during light conditions due 16 420 to the influence of the photosynthetic activity of MPB, reducing the availability of recycled nutrients 17 18 421 for the pelagic demand. The relatively stable environmental conditions over the year, compared to 19 **422** temperate regions, enable our results to be broadened on other temporal scales and extended to 20 423 similar systems. These new data on mineralization and production rates at the sediment-water 21 424 interface will be used in the frame of numerical modeling as foreseen in the MANA project 22 23 **425** (MANagement of Atolls - Seceh et al., 2021). <sup>24</sup> 426

# <sup>26</sup> 427 Acknowledgments<sup>27</sup>

428
 428
 428
 429
 429
 429
 429
 430
 430
 431
 431
 432
 431
 433
 434
 435
 435
 436
 436
 437
 438
 439
 430
 430
 430
 430
 431
 431
 431
 432
 433
 434
 435
 435
 436
 436
 437
 438
 439
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430
 430

# 3334 432 References

Abhilash, K.R., Raveendran, T.V., Mol, V.P.L., Deepak, M.P., 2012. Sediment oxygen demand in Cochin
 backwaters, a tropical estuarine system in the south-west coast of India. Marine Environmental
 Research 79, 160–166.

- 40
   41
   436 Aller, R.C., Cochran, J.K., 2019. The Critical Role of Bioturbation for Particle Dynamics, Priming
   42
   437 Potential, and Organic C Remineralization in Marine Sediments: Local and Basin Scales. Front. Earth
   438 Sci. 7, 157. doi: 10.3389/feart.2019.00157
- 45 439 Alongi, D., Trott, L., Mohl, M., 2011. Strong tidal currents and labile organic matter stimulate benthic
   46 440 decomposition and carbonate fluxes on the southern Great Barrier Reef shelf. Continental Shelf
   48 441 Research 31, 1384-1395.
- 49
  50
  442 Aminot, A., Kérouel, R., 2007. Dosage automatique des nutriments dans les eaux marines: méthodes
  51
  443 en flux continu (Editions Quae).
- An, S., Joye S., B., 2001. Enhancement of coupled nitrification-denitrification by benthic
   445 photosynthesis inshallow estuarine sediments. . Limnol. Oceanogr. 46(1), 62–74.
- Andréfouët, S., Ardhuin, F., Quefeullou, P., Le Gendre, R., 2012a. Island shadow effects and the wave
   climate of the Western Tuamotu Archipelago (French Polynesia) inferred from altimetry and
   numerical model data. Mar. Pollut. Bull. 65, 415–424.
- 60 61
- 62 63

- Andréfouët, S., Charpy, L., Lo-Yat, A., Lo, C., 2012b. Recent research for pearl oyster aquaculture
   1 450 management in French Polynesia. Mar. Pollut. Bull. 65, 407–414.
  - 451 https://doi.org/10.1016/j.marpolbul.2012.06.021.

Barranguet, C., M.R. Plante-Cuny, Alivon E., 1996. Microphytobenthos production in the Gulf of Fos,
French Mediterranean coast. Hydrobiologia 333, 181-193.

- Berner, R.A., 1980. Early Diagenesis a Theoretical Approach. Princeton University Press, Princeton,
  New York.
- 456 Boucher, G., Clavier, J. 1990. Contribution of benthic biomass to overall metabolism in New 457 Caledonian lagoon sediments. Marine Ecology Progress series 64, 271-280.
- <sup>4</sup> 458 Bouvy, M., Dupuy, C., Pagano, M., Barani, A., Charpy, L., 2012. Do Human Activities Affect the
   <sup>5</sup> 459 Picoplankton Structure of the Ahe atoll Lagoon (Tuamotu Archipelago, French Polynesia)? Mar.
   <sup>7</sup> 460 Pollut. Bull. 65, 516-524.
- Boynton, W.R., Ceballos, M.A.C., Bailey, E.M., Hodgkins, C.L S., Humphrey, J.L., Testa, J.M., 2018.
   Oxygen and Nutrient Exchanges at the Sediment-Water Interface: a Global Synthesis and Critique of
   Estuarine and Coastal Data. Estuaries and Coasts 41, 301–333. https://doi.org/10.1007/s12237-017 0275-5
- <sup>4</sup> 465 Burdige, D., 2006. Geochemistry of Marine Sediments. PRINCETON; OXFORD: Princeton University
   <sup>6</sup> 466 Press, 610 pp. doi:10.2307/j.ctv131bw7s
- 467 Caffrey, J. M., Harrington, N., Ward, B., 2002. Biogeochemical processes in a small California estuary.
   468 1. Benthic fluxes and pore water constituents reflect high nutrient freshwater inputs. Marine Ecology
   469 Progress Series 233, 39–53. doi:10.3354/meps233039
- <sup>2</sup> 470 Canfield, D.E., Kristensen, E., Thamdrup, B., 2005. In: Canfield, D.E., Kristensen, E., Thamdrup, B.
   <sup>3</sup> 471 (Eds.), Aquatic Geomicrobiology. Academic Press, 656 pp.
- <sup>5</sup> 472 Charpy, L., Charpy-Roubaud, C.J., 1990. Trophic structure and productivity of the lagoonal
   7 473 communities of Tikehau atoll (Tuamotu Archipelago, French Polynesia). Hydrobiology 207, 43–52.
- 474 Chatterjee, A., Klein, C., Naegelen, A., Claquin, P., Masson, A., Legoff, M., Amice, E., L'Helguen, S.,
   475 Chauvaud, L., Leynaert, A., 2013. Comparative dynamics of pelagic and benthic micro-algae in a
   476 coastal ecosystem. Estuarine, Coastal and Shelf Science 133, 67-77. 10.1016/j.ecss.2013.08.015.
- <sup>3</sup> 477 Chau, K.W., 2002. Field measurements of SOD in a land-locked embayment in Hong Kong. Advances
   <sup>4</sup> 478 in Environmental Research 6, 135-142.
- 479 Chen, G.H., Leong, I.M., Liu, J., Huang, J.C., Lo, I.M.C., Yen, B.C., 2000. Oxygen deficit determinations
   480 for a major river in eastern Hong Kong, China. Chemosphere 41, 7-13.
- Clavier, J., Boucher, G., Chauvaud, L., Fichez, R., Chifflet, S., 2005. Benthic response to ammonium
   pulses in a tropical lagoon: Implications for coastal environmental processes. Journal of Experimental
   Marine Biology and Ecology, 231-241. 10.1016/j.jembe.2004.11.004.
- <sup>4</sup> 484 Dalsgaard, T., Nielsen, L.P., Brotas, V., Viaroli, P., Underwood, G.J.C., Nedwell, D.B., Sundbäck, K.,
   <sup>6</sup> 485 Rysgaard, S., Miles, A., Bartoli, M., Dong, L., Thornton, D.C.O., Ottosen, L.D.M., Castaldelli, G.,
   <sup>7</sup> 486 Risgaard-Petersen, N., 2000. Protocol Handbook for NICE-Nitrogen Cycling in Estuaries: A Project
   <sup>8</sup> 487 under the EU Research Programme: Marine Science and Technology (MAST III). Ministry of

488 Environment and Energy National Environmental Research Institute, Denmark<sup>©</sup> Department of Lake 1 489 and Estuarine Ecology. 2

3 490 Dumas, F., Le Gendre, R., Thomas, Y., Andréfouët, S., 2012. Tidal flushing and wind driven circulation 4 491 of Ahe atoll lagoon (Tuamotu Archipelago, French Polynesia) from in situ observations and numerical 5 492 modelling. Mar. Pollut. Bull. 65, 425–440. 6

493 Dunn, R.J.K.K., Welsh, D.T., Jordan, M., Waltham, N.J., Lemckert, C.J., Teasdale, P.R., 2012. Benthic 9 494 metabolism and nitrogen dynamics in a sub-tropical coastal lagoon: microphytobenthos stimulate 10 495 nitrification and nitrate reduction through photosynthetic oxygen evolution. Estuar. Coast Shelf Sci. 11 496 113, 272-282. 12

13 497 Dutheil, C., Andréfouët, S., Jullien, S., Le Gendre, R., Aucan, J., Menkès, C., 2020. Characterization of 14 <sub>15</sub> 498 south central Pacific Ocean wind regimes in present and future climate for pearl farming application. Marine Pollution Bulletin 160, 111584. 16 **499** 

Eyre, B.D., Ferguson, A.J.P., Webb, A., Maher, D., Oakes, J.M., 2011. Metabolism of different benthic 18 500 <sup>19</sup> 501 habitats and their contribution to the carbon budget of a shallow oligotrophic sub-tropical coastal 20 502 system (southern Moreton Bay, Australia). Biogeochemistry 102, 87–110. 21 22 **503** https://doi.org/10.1007/s10533-010-9424-7.

- 23 <sub>24</sub> 504 Ferreira, J.G., Corner, R.A., Moore, H., Service, M., Bricker, S., Rheault, R., 2018. Ecological carrying 25 505 capacity for shellfish aquaculture –Sustainability of naturally occurring filter-feeders and cultivated 26 506 bivalves. Journal of Shellfish Research 37: 1-18. 27
- <sup>28</sup> 507 Frouin, P., 2000. Effects of anthropogenic disturbances of tropical soft-bottom benthic communities. 29 508 Marine Ecology Progress Series 194: 39-53. 30
- 31 509 Gaertner-Mazouni, N., Lacoste, E., Bodoy, A., Peacock, L., Rodier, M., Langlade, M.-J., Orempuller, J., 32 33 510 Charpy, L., 2012. Nutrient fluxes between water column and sediments: potential influence of the 34 511 pearl oyster culture. Mar. Pollut. Bull. 65 : 500-505. https://doi.org/10.1016/j.marpolbul.2012.02.013 <sup>35</sup> 512
- <sup>36</sup> 513 Grenz, C., Denis, L., Boucher, G., Chauvaud, L., Clavier, J., Fichez, R., Pringault, O., 2003. Spatial 37 514 variability in sediment oxygen consumption under winter conditions in a lagoonal system in New 38 39 515 Caledonia (South Pacific). Journal of Experimental Marine Biology and Ecology 285-286, 33-47.
- 40 41 516 Heil, C.A., Chaston, K., Jones, A., Bird, P., Longstaff, B., Costanzo, S., Dennison, W.C., 2004. Benthic <sup>42</sup> 517 microalgae in coral reef sediments of the southern Great Barrier Reef, Australia. Coral Reefs 23, 336-43 518 343. 44
- 45 519 Hicks, N., Liu, X., Gregory, R., Kenny, J., Lucaci, A., Lenzi, L., Paterson, D.M., Duncan, K.R., 2018. 46 <sub>47</sub> 520 Temperature Driven Changes in Benthic Bacterial Diversity Influences Biogeochemical Cycling in 48 521 Coastal Sediments. Frontiers in Microbiology 9, 1730. 10.3389/fmicb.2018.01730. 49
- 50 **522** Holmes, R.M., Aminot, A., Kérouel, R., Hooker, B.A., Peterson, B.J., 1999. A simple and precise <sup>51</sup> **523** method for measuring ammonium in marine and freshwater ecosystems. Can. J. Fish. Aquat. Sci. 56, 52 524 1801-1808. 53
- 54 525 Jäger, C.G., Diehl, S., 2014. Resource competition across habitat bounderies; asymmetric interactions 55 56 **526** between benthic and pelagic producers. Ecological Monographs 84 (2), 287-302.

57 58 **527** Jahnke, R. A., Nelson, J.R., Marinelli, R.L., Eckman, J.E., 2000. Benthic flux of biogenic elements on the 59 528 Southeastern US continental shelf: Influence of pore water advective transport and benthic <sup>60</sup> 529 microalgae. Cont. Shelf Res. 20, 109-127. 61

63 64

62

65

7

8

Jensen, H. S., Thamdrup, B., 1993. Iron-bound phosphorus in marine sediments as measured by 530 1 531 bicarbonate-dithionite extraction. Hydrobiologia 253, 47-59. 2 3 532 Kamatani, A., 1982. Dissolution rates of silica from diatoms decomposing at various temperatures. 4 533 Mar. Biol. 68, 91–96. 5 6 534 Kopp, D., Lefebvre, S., Cachera, M., Villanueva, M.C., Ernande, B., 2015. Reorganization of a marine 7 535 trophic network along an inshore-offshore gradient due to stronger pelagic-benthic coupling in 8 9 536 coastal areas. Prog. Oceanogr. 130, 157–171. https://doi.org/10.1016/j.pocean.2014.11.001. 10 11 **537** Lacoste, E., Gaertner-Mazouni, N., 2016 Nutrient regeneration in the water column and at the 12 538 sediment-water interface in pearl oyster culture (Pinctada margaritifera) in a deep atoll lagoon (Ahe, 13 539 French Polynesia). Estuarine, Coastal and Shelf Science 182, 304-309. 14 <sub>15</sub> 540 https://doi.org/10.1016/j.ecss.2016.01.037. 16 <sub>17</sub> 541 Le Bouteiller, A., Blanchot, J., Rodier, M., 1992. Size distribution patterns of phytoplankton in the western Pacific: towards a generalization for the tropical open ocean. Deep Sea Research Part A. 18 **542** 19 543 Oceanographic Research Papers 39 (5), 805-823. 20 <sup>21</sup> 544 Ling, T.K., Chiat-Siew, N., Lee, N., Buda, D., 2009. Oxygen demand of the sediment from the 22 545 Semariang Batu River. Malaysia World Applied Sciences Journal 7(4), 440-447. 23 24 546 MacIntyre, H. L., Geider, R. J., Miller, D., 1996. Microphytobenthos: the ecological role of the "secret 25 26 **547** garden" of unvegetated, shallow water marin habitats. I. Distribution, abundance and primary 27 548 production. Estuaries 19, 186–201. doi: 10.2307/1352224 28 29 549 Middelburg, J.J., Vlug, T., Jaco, F., van der Nat, W.A., 1993. Organic matter mineralization in marine <sup>30</sup> 550 systems. Global Planet. Change 8 (1), 47-58. 31 <sup>32</sup> 551 Middelburg, J.J, 2018. Reviews and syntheses: to the bottom of carbon processing at the seafloor. 33 552 Biogeosciences 15, 413-427. 10.5194/bg-15-413-2018 34 35 553 Pan, F., Guo, Z., Cai, Y., Fu, Y., Wu, J., Wang, B., Liu, H., Gao, A., 2020. Cyclical patterns and 36 (im)mobilization mechanisms of phosphorus in sediments from a small creek estuary: Evidence from 37 554 38 555 in situ monthly sampling and indoor experiments. Water Research 171, 115479. 39 556 10.1016/j.watres.2020.115479. 40 41 557 Pratihary, A.K., Naqvi, S.W.A., Naik, H., Thorat, B.R., Narvenkar, G., Manjunatha, B.R., Rao, V.P., 2009. 42 558 Benthic fluxes in a tropical estuary and their role in the ecosystem. Estuarine Coastal Shelf Science 43 44 559 85(3), 387-398. 45 46 560 Raimbault, P., Slawyk, G., Coste, B., Fry, J., 1990. Feasibility of using an automated colorimetric <sup>47</sup> 561 procedure for the determination of seawater nitrate in the 0 to 100nM range: examples from field 48 562 and culture. Mar. Biol. 104, 347-351. 49 50 563 Rasmussen, H., Jorgensen, B.B., 1992. Microelectrode studies of seasonal oxygen uptake in a coastal 51 <sub>52</sub> 564 sediment: role of molecular diffusion. Mar. Ecol. Prog. Ser. 81, 289–303. 53 <sub>54</sub> 565 Rodier, M., Pinazo, C., Seceh, C., Varillon, D., 2021. Pelagic stocks and carbon and nitrogen uptake in 55 **566** a pearl farming atoll (Ahe, French Polynesia). Marine Pollution Bulletin 167, 112352. 56 57 **567** Sarrazin, G., Charpy-Roubaud, C., Charpy, L., 1988. Early diagenesis of organic matter in the <sup>58</sup> 568 sediments of the central basin of Tikehau lagoon-reef (Tuamotu Archipelago - French Polynesia). 59 569 Proceedings of the 6th Int Coral Reef Symposium Vol 3: 373-378. 60 61 62 14 63 64 65

- Seceh, C., Pinazo, C, Rodier, M., Lajaunie-Salla, K., Mazoyer, C., Grenz, C., Le Gendre, R., 2021. 1 571 Biogeochemical model of nitrogen cycling in a South Pacific coral atoll with pearl farming: Ahe (French Polynesia). Marine Pollution Bulletin 169, 112526. Sørensen, J., 1978. Capacity for denitrification and reduction of nitrate to ammonia in a coastal marine sediment. Appl. Environ. Microbiol. 35, 301–305. Sundbäck, K., Enoksson, V., Granéli, W., Pettersson, K., 1991. Influence of sublittoral 9 576 microphytobenthos on the oxygen and nutrient flux between sediment and water: a laboratory 10 577 continuous-flow study. Mar Ecol Prog Ser 74, 262–279. 12 578 Sundbäck, K., Miles, A., 2002. Role of microphytobenthos and denitrification for nutrient turnover in embayments with floating macroalgal mats: A spring situation. Aquatic Microbial Ecology 30, 91-101. <sub>15</sub> 580 10.3354/ame030091. <sub>17</sub> 581 Underwood, G.J.C., Kromkamp, J., 1999. Primary production by phytoplankton and microphytobenthos in estuaries In: Nedwell DB, Raffaelli DG (eds) Advances in ecological research, **582** Vol 29. Academic Press, San Diego, CA, 93 –153. <sup>21</sup> **584** Underwood, G.J.C., 2002. Adaptations of tropical marine microphytobenthic assemblages along a gradient of light and nutrient availability in Suva Lagoon, Fiji. Eur J Phycol 37, 449-462. Valdes-Lozano, D.S., Chumacero, M., Real, E., 2006. Sediment oxygen consumption in a developed **587** coastal lagoon of the Mexican Caribbean. Indian Journal of Marine Sciences 35, 227–234.
- Van Cappellen, P. and Wang, Y. 1996. Cycling of iron and manganese in surface sediments: A general
   theory for the coupled transport and reaction of carbon, oxygen, nitrogen, sulfur, iron and
   manganese. American Journal of Science 296, 197-243.
- <sup>32</sup> 591 Willey, J. D., 1978. Release and uptake of dissolved silica in seawater by marine sediments. Marine
   <sup>33</sup> 592 Chemistry 7, 53-65. https://doi.org/10.1016/0304-4203(78)90042-7.

### Credit author statement

Grenz Christian Conceptualization, Investigation, Supervision, Writing-Original Draft
Rodier Martine : Conceptualization, Investigation, Writing-Review & Editing
Claire Seceh : Visualization, Writing-Review & Editing
Varillon David : Resources, Investigation
Haumani Gabriel : Resources, Investigation

Pinazo Christel : Conceptualization, Investigation, Writing-Review & Editing

### **Declaration of interests**

 $\boxtimes$  The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

□The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: