



Gemella massiliensis sp. nov., a new bacterium isolated from the human sputum

Maxime Descartes Mbogning Fonkou, Cheikh Ibrahima Lo, Zouina Mekhalif, Melhem Bilen, Enora Tomei, Edmond Kuete Yimagou, Grégory Dubourg, Didier Raoult, Florence Fenollar, Pierre-Edouard Fournier

► To cite this version:

Maxime Descartes Mbogning Fonkou, Cheikh Ibrahima Lo, Zouina Mekhalif, Melhem Bilen, Enora Tomei, et al.. *Gemella massiliensis* sp. nov., a new bacterium isolated from the human sputum. *Archives of Microbiology*, 2021, 203 (9), pp.5817-5823. <10.1007/s00203-021-02493-2>. <hal-03421043>

HAL Id: hal-03421043

<https://amu.hal.science/hal-03421043v1>

Submitted on 6 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



Gemella massiliensis sp. nov., a new bacterium isolated from the human sputum

Maxime Descartes Mbogning Fonkou^{1,2} · Cheikh Ibrahima Lo^{2,3} · Zouina Mekhalif^{1,2} · Melhem Bilen¹ · Enora Tomei^{1,2} · Edmond Kuete Yimagou^{1,2} · Grégory Dubourg^{1,2} · Didier Raoult^{1,2,4} · Florence Fenollar^{2,3} · Pierre-Edouard Fournier^{2,3}

Received: 18 February 2021 / Revised: 28 June 2021 / Accepted: 14 July 2021 / Published online: 21 August 2021
© The Author(s) 2021

Abstract

Thanks to its ability to isolate previously uncultured bacterial species, culturomics has dynamized the study of the human microbiota. A new bacterial species, *Gemella massiliensis* Marseille-P3249^T, was isolated from a sputum sample of a healthy French man. Strain Marseille-P3249^T is a facultative anaerobe, catalase-negative, Gram positive, coccus, and unable to sporulate. The major fatty acids were C_{16:0} (34%), C_{18:1n9} (28%), C_{18:0} (15%) and C_{18:2n6} (13%). Its 16S rRNA sequence exhibits a 98.3% sequence similarity with *Gemella bergeri* strain 617-93^T, its phylogenetically closest species with standing in nomenclature. Its digital DNA–DNA hybridization (dDDH) and OrthoANI values with *G. bergeri* of only 59.7 ± 5.6% and 94.8%, respectively. These values are lower than the thresholds for species delineation (> 70% and > 95%, respectively). This strain grows optimally at 37 °C and its genome is 1.80 Mbp long with a 30.5 mol% G + C content. Based on these results, we propose the creation of the new species *Gemella massilienis* sp. nov., strain Marseille-P3249^T (= CSUR P3249 = DSMZ 103940).

Keywords *Gemella massiliensis* sp. nov. · Respiratory microbiota · Taxono-genomics · Bacteria

Abbreviations

| | |
|--------------|--|
| DSMZ | Deutsche Sammlung von Mikroorganismen und Zellkulturen |
| CSUR | Collection de Souches de l'Unité des Rickettsies |
| MALDI-TOF MS | Matrix-assisted laser desorption ionization time-of-flight |
| MIC | Minimal inhibitory concentration |

dDDH
COG

DNA–DNA hybridization
Clusters of orthologous groups

Introduction

The genus *Gemella* has been described for the first time by Berger 1961 (Berger). Members of this genus are usually Gram-positive, coccoid-shaped, facultatively anaerobic, and do not produce any catalase activities (Collins 2006). The first species of this genus such as *Gemella morbillorum* and *Gemella haemolysans* are commensals of mucous membranes of humans but are sometimes responsible for human infections (Kilpper-Bälz and Schleifer 1988).

Gemella bergeri and *Gemella sanguinis* were recovered from human clinical specimens (Collins et al. 1998a, b), whereas *Gemella palaticanis* was isolated from a dog (Collins et al. 1999). Although the pathogenicity of members of this genus is not yet proven, it seems likely that they are also residents of the mucous membranes.

During a project on the human microbiota, we studied sputum samples by culturomics as previously described (Lagier et al. 2018) which allowed us to isolate a new

Communicated by Erko Stackebrandt.

✉ Pierre-Edouard Fournier
pierre-edouard.fournier@univ-amu.fr

¹ Aix Marseille Univ, IRD, AP-HM, MEPHI, Marseille, France

² Institut Hospitalo-Universitaire Méditerranée-Infection, 19-21 Boulevard Jean Moulin, 13385 Marseille Cedex 05, France

³ Aix Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

⁴ Special Infectious Agents Unit, King Fahd Medical Research Center, King Abdulaziz University, Jeddah, Saudi Arabia

bacterial strain belonging to the phylum *Firmicutes*. Herein, we report a taxonogenomic description (Fournier et al. 2015) of *Gemella massiliensis* sp. nov., which is previously announced by our research group (Fonkou et al. 2018).

Materials and methods

Growth conditions

A bacterial strain was isolated from a sputum sample from a healthy Frenchman by culturomics to explore the human microbiome. The study was approved by the ethics committee of the Institut Federatif de Recherche IFR48 under the number 09-022 and then the patient gave his formal agreement by signing the informed consent. Thus optimal growth conditions of strain Marseille-P3249 were evaluated using various culture conditions. Culture assays were done at 28, 37, 45 and 55 °C under anaerobic (GENbag anaer, bioMérieux), microaerophilic (GENbag Microaer, bioMérieux) and aerobic conditions. Tolerance to acidity and halotolerance were evaluated independently with growth assays at pH 6, 6.5, 7 and 8.5 and by using 0, 5, 10, 50, 75 and 100 g/L NaCl concentrations, respectively.

Morphological, biochemical and antibiotic susceptibility analysis

The main biochemical features of strain Marseille-P3249^T were tested using API strips (ZYM, 50CH and 20A (bioMérieux, France)). Motility and Gram stain were checked using a DM1000 photonic microscope (Leica Microsystems, Nanterre, France). Additionally, sporulation was evaluated after exposing a bacterial suspension to a 20 min heat shock at 80 °C. Cell morphology images were obtained using a scanning electron (SEM) microscope (TM4000 Plus, Hitachi High-Technologies Corp., Tokyo, Japan).

Cellular fatty acid methyl ester (FAME) analyses were performed with GC/MS with 10 mg of bacterial biomass per tube. GC/MS and FAME analyses were performed as previously reported (Elsawi et al. 2017).

The minimal inhibitory concentrations (MIC) of strain Marseille-P3249 were evaluated using Etest (bioMérieux) for benzylpenicillin, amoxicillin, cefotaxime, ceftriaxone, imipenem, erythromycin, daptomycin, amikacin, rifampicin, minocycline, teicoplanin, vancomycin, metronidazole, and colistin.

DNA extraction and genome sequencing

A total of 82.1 ng/μL of genomic DNA (gDNA) were extracted from strain Marseille-P3249 as previously described (Elsawi et al. 2017). gDNA was sequenced using

the MiSeq technology (Illumina Inc, San Diego, CA, USA) with the Mate-pair strategy and were run and barcoded with 11 additional projects using the Nextera Mate-Pair sample prep kit (Illumina) as formerly described (Elsawi et al. 2017). The DNA fragment size ranged from 1.5 kb up to 11 kb with an optimal size of 6.29 kb. No size selection was done and 177.24 ng of tagged fragments were circularized. The circularized DNAs were sheared mechanically to smaller fragments with an optimal size at 1393 bp on the Covaris device S2 in T6 tubes (Covaris, Woburn, MA, USA). Using a high sensitivity bioanalyzer LabChip (Agilent Technologies Inc, Santa Clara, CA, USA), the library profile was visualized with a final concentration of 15.59 nmol/L. The latter were normalized at 2 nM and pooled with other samples, and finally diluted to 15 pM. Automated cluster generation and sequencing run were performed in a single 2 × 251-bp run. Total information of 9.5 Gb was obtained from a 1050 K/mm² cluster density with a cluster passing quality control filters of 92.5% (18,644,000 passing filter paired-reads). Within this run, the index representation for strain Marseille-P3249^T was determined to 4.67%. The 870,362 paired reads were trimmed, assembled, annotated and analyzed.

Genome-to-Genome Distance Calculator (<http://ggdc.dsmz.de>) was used for digital DNA–DNA hybridization (dDDH) estimates with confidence intervals under recommended settings (Formula 2, BLASTP).

Phylogenetic analysis

For phylogenetic analyses, 16S rRNA gene sequences of closely related species were recovered from the Genbank database (<https://www.ncbi.nlm.nih.gov/genbank/>). Muscle was used for sequence alignment and phylogenetic inferences were generated using the approximately-maximum-likelihood method within the FastTree software (Edgar 2004; Price et al. 2009). In addition, a phylogenetic tree based on housekeeping genes such as *groES*, *groEL*, *recA*, *gyrA*, and *rpoB* was performed using iTOL software online (<https://itol.embl.de/>). Genes are extracted from annotated genomic sequences and then concatenated for each strain.

Results and discussion

Strain identification

MALDI-TOF MS failed to identify strain Marseille-P3249^T. Therefore, 16S rRNA gene sequencing was performed and using a blast comparison against the NCBI nucleotide database, strain Marseille-P3249^T exhibited a 98.3% sequence similarity with *Gemella bergeri* strain 617-93^T, being the phylogenetically closest species with standing

in nomenclature (Fig. 1) (Collins et al. 1998a). Thus, and according to Kim et al., this strain may be classified within a new bacterial species within the *Gemella* genus as it exhibits more than 1.35% sequence divergence with its phylogenetically closest species with a validly published name (Kim et al. 2014). Furthermore, the MLSA tree performed with concatenated genes shows that *G. massiliensis* strain Marseille-P3249^T is positioned within the *Gemella* species but is clearly distinct from them on a single branch (Fig. 2).

General characteristics of strain Marseille-P3249

Cells from strain Marseille-P3249^T were Gram-positive cocci. Colonies grew in optimally at 37 °C in aerobic conditions with pH range between 6 and 8.5 and NaCl concentrations below 50 g/L and measured from 0.5 to 1.2 mm in diameter after 24 h of incubation. Cells were not motile and non-spore forming with a mean diameter of 0.78 µm. They metabolize D-fructose, amygdalin, and L-sorbose possessed enzymes such as esterase, leucine arylamidase, and naphthol-AS-BI-phosphohydrolase. Biochemical criteria of

strain Marseille-P3249^T are compared with those of closely related species in standing in nomenclature (Table 1).

The major fatty acids were hexadecanoic acid (34%), 9-Octadecenoic acid (28%), octadecanoic acid (15%) and 9,12-octadecadienoic acid (13%). A wide variety of other fatty acids were described but present with low amounts (Table 2).

Strain Marseille-P3249 exhibited MICs with benzylpenicillin (0.012 µg/mL), amoxicillin (0.016 µg/mL), cefotaxime (0.016 µg/mL), ceftriaxone (0.016 µg/mL), imipenem (0.016 µg/mL), erythromycin (0.19 µg/mL), daptomycin (> 6 µg/mL), amikacin (0.125 µg/mL), rifampicin (0.03 µg/mL), minocycline (0.64 µg/mL), teicoplanin (0.032 µg/mL), vancomycin (0.75 µg/mL), metronidazole (> 256 µg/mL) and colistin (> 256 µg/mL).

Genome characteristics of strain Marseille-P3249

The genome was 1,804,813 bp long with a 30.5 mol% G + C content (Fig. 3). It is composed of 7 scaffolds (composed of 8 contigs). Of the 1727 predicted genes, 1677 were

Fig. 1 16S rRNA gene sequence phylogenetic analysis highlighting the position of strain Marseille-P3249 relative to other species. This tree is formally already published but it was remade with slight changes (Fonkou et al. 2018). Sequence alignment and phylogenetic inferences were obtained using the maximum likelihood method within MEGA 7 software. The scale bar represents a 2% sequence divergence using 1000 replicates. GenBank accession numbers are indicated in parenthesis





Fig. 2 Neighbour-joining tree displaying the relationships among species of the genus *Gemella* based on concatenated *groES*, *groEL*, *recA*, *gyrA*, and *rpoB* sequences

Table 1 Differential characteristics of *Gemella massiliensis* strain Marseille-P3249^T (GMA), *Gemella bergeri* 617-93^T (GBE) (Collins et al. 1998a), *Gemella assaccharolytica* EU427463^T (GAS) (Ulger-Toprak et al. 2010), *Gemella cuniculi* AJ251987^T (GCU) (Hoyleset al. 2000), *Gemella morbillorum* L14327^T (GMO) (Kilpper-Bälz and Schleifer 1988), and *Gemella sanguinis* Y13364^T (GSA) (Collins et al. 1998b)

| Properties | GMA | GBE | GAS | GCU | GMO | GSA |
|------------------------|---------------|-------------------|-------------------|--------------------|-------------------|-------------------|
| Cell diameter (μm) | 0.78 | Na | 0.5 | Na | 0.3–0.8 | Na |
| Oxygen requirement | Fa | Fa | Fa | Fa | Fa | Fa |
| Gram stain | + | + | V | + | + | + |
| Endospore formation | – | – | – | Na | Na | – |
| Production of | | | | | | |
| Alkaline phosphatase | – | – | – | + | Na | + |
| Catalase | – | – | – | – | – | – |
| Urease | – | – | – | – | Na | – |
| β-galactosidase | – | – | Na | Na | Na | – |
| N-acetyl-β-glucosamine | – | Na | Na | – | Na | – |
| L-Arabinose | – | – | – | – | Na | – |
| D-Ribose | – | – | – | – | Na | – |
| D-Mannose | – | Na | – | Na | + | Na |
| D-Mannitol | – | – | – | + | + | + |
| D-glucose | – | + | – | + | + | + |
| D-fructose | + | – | + | Na | – | Na |
| D-maltose | – | W | – | V | + | + |
| D-lactose | – | – | – | – | – | V |
| G + C content (mol%) | 30.5 | 30.3 | 26.7 | 28.9 | 30.8 | 29.7 |
| Habitat | Sputum sample | Clinical specimen | Clinical specimen | Abcess of a rabbit | Clinical specimen | Clinical specimen |

Fa facultative anaerobic; Na data not available; V variable; W weakly positive

Table 2 Cellular fatty acid composition (%)

| Fatty acids | Name | Mean relative % ^a |
|---------------------------|------------------------------|------------------------------|
| C _{16:0} | Hexadecanoic acid | 34.1 ± 0.3 |
| C _{18:1n9} | 9-Octadecenoic acid | 27.6 ± 0.2 |
| C _{18:0} | Octadecanoic acid | 14.8 ± 0.1 |
| C _{18:2n6} | 9,12-Octadecadienoic acid | 12.5 ± 0.4 |
| C _{18:1n7} | 11-Octadecenoic acid | 2.3 ± 0.1 |
| C _{18:1n5} | 13-Octadecenoic acid | 2.1 ± 0.1 |
| C _{14:0} | Tetradecanoic acid | 1.2 ± 0.1 |
| C _{17:0} | Heptadecanoic acid | TR |
| C _{15:0} | Pentadecanoic acid | TR |
| C _{15:0 anteiso} | 12-methyl-tetradecanoic acid | TR |
| C _{16:1n7} | 9-Hexadecenoic acid | TR |

TR trace amounts < 1%

^aMean peak area percentage

protein-coding genes and 50 were RNAs (5 genes were 5S rRNA, 2 genes were 16S rRNA, 2 genes were 23S rRNA, and 41 genes were tRNA genes). A total of 1 276 genes (76.09%) were assigned a putative function (by cogs or by NR blast). Twenty-six genes were classified as ORFans

(1.55%). The remaining genes were annotated as hypothetical proteins (304 genes (18.13%)). The distribution of genes into COG functional categories is detailed in supplementary Table S1.

Genome comparison

The draft genome sequence of strain Marseille-P3249^T was larger than those of *Gemella cuniculi* DSM 15828^T, *Gemella sanguinis* ATCC 700632^T and *Gemella haemolysans* ATCC 10379^T, but smaller than those of *Gemella asaccharolytica* WAL 1945J^T, *Gemella bergeri* 617-93^T and *Gemella morbillorum* NCTC11323^T (Table 3).

Additionally, the G + C content of strain Marseille-P3249^T is smaller than those of *G. asaccharolytica* WAL 1945J^T, *G. cuniculi* DSM 15828^T, *G. sanguinis* ATCC 700632^T and *G. bergeri* 617-93^T, but larger than those of *G. morbillorum* NCTC11323^T and *G. haemolysans* ATCC 10379^T. In the same way, the gene content of strain Marseille-P3249^T was compared with the closely related *Gemella* species.

Strain Marseille-P3249^T shared the highest number of orthologous proteins with *G. cuniculi* (1039). Furthermore,

Fig. 3 Graphical circular map of the chromosome. From outside to the center: genes on the forward strand colored by COG categories (only genes assigned to COG), genes on the reverse strand colored by COG categories (only gene assigned to COG), RNA genes (tRNAs green, rRNAs red), GC content and GC skew




Table 3 Genome information of the species involved in the genomic comparative analyses

| Species | Size (Mb) | GC (%) | Gene content |
|--|-----------|--------|--------------|
| <i>Gemella asaccharolytica</i> WAL 1945J ^T | 1.28 | 26.6 | 1251 |
| <i>Gemella bergeri</i> 617-93 ^T | 1.60 | 30.3 | 1524 |
| <i>Gemella morbillorum</i> NCTC11323 ^T | 1.75 | 30.7 | 1622 |
| <i>Gemella massiliensis</i> Marseille-P3249 ^T | 1.80 | 30.5 | 1677 |
| <i>Gemella cuniculi</i> DSM 15828 ^T | 1.86 | 28.9 | 1687 |
| <i>Gemella haemolysans</i> ATCC 10379 ^T | 1.91 | 30.8 | 1710 |
| <i>Gemella sanguinis</i> ATCC 700632 ^T | 1.90 | 29.6 | 1861 |

this bacterium shared 1031, 1032, 1054, and 778 orthologous proteins with *G. haemolysans*, *G. morbillorum*, *G. sanguinis* and *G. asaccharolytica*, respectively. Strain Marseille-P3249^T exhibited the highest OrthoANI values of 94.8% with *G. bergeri* and 70.3% as the lowest value *G. asaccharolytica* (Fig. 4). dDDH values obtained during analysis were not exceeded 59.7% between *G. massiliensis* strain Marseille-P3249 and *G. bergeri* (Table 4). Based on

DDH values below 70%, the recommended threshold for delineating a new species (Wayne 1988; Tindall et al. 2010), we consider this strain Marseille-P3249 to be a new species of the genus *Gemella*.

Description of *Gemella massiliensis* sp. nov.

Gemella massiliensis (*mas.si.li.en'sis*. L. fem. adj, *massiliensis*, pertaining to Massilia, the Latin name of the city of Marseille, where this bacterium was discovered). Strain Marseille-P3249^T is a facultative anaerobic bacterium but grows optimally at 37 °C under aerobic conditions. Using a 50 CH strip, this strain exhibits positive reactions for D-fructose, amygdaline, and L-sorbose. Positive reactions are also observed for esterase (C4), esterase lipase (C8), leucine arylamidase, phosphatase acid, and naphthol-AS-BI-phosphohydrolase. In addition, using an API 20A (bioMérieux), positive reactions are observed for esculin ferric citrate only. The genome is 1.80 Mbp with 30.5 mol% G + C content.

The type strain Marseille-P3249^T (=CSURP3249=DSM103940) was isolated from the sputum sample of a healthy French man.

The 16S rRNA and whole-genome sequences of *G. massiliensis* sp. nov., were deposited in EMBL-EBI

Fig. 4 Heatmap generated with OrthoANI values calculated using the OAT software between *Gemella massiliensis* sp. nov., strain Marseille-P3249 and other closely related species with standing in nomenclature




Table 4 Digital DNA–DNA hybridization values (%) obtained by strain Marseille-P3249^T with other closely-related species using the GGDC formula 2 software (dddH estimates based on identities/HSP length)

| | GMA | GAS | GCU | GHA | GMO | GSA | GBE |
|-----|------|-------------|-------------|-------------|-------------|-------------|-------------|
| GMA | 100% | 21.3 ± 4.7% | 22.6 ± 4.7% | 21.7 ± 4.7% | 22.1 ± 4.7% | 21.9 ± 4.7% | 59.7 ± 5.6% |
| GAS | | 100% | 23.4 ± 4.7% | 23.2 ± 4.7% | 22.4 ± 4.7% | 21.6 ± 4.6% | 21.0 ± 4.7% |
| GCU | | | 100% | 21.8 ± 4.7% | 22.0 ± 4.7% | 22.1 ± 4.7% | 22.7 ± 4.7% |
| GHA | | | | 100% | 22.9 ± 4.7% | 23.5 ± 4.8% | 22.1 ± 4.7% |
| GMO | | | | | 100% | 23.0 ± 4.8% | 21.9 ± 4.7% |
| GSA | | | | | | 100% | 21.9 ± 4.8% |
| GBE | | | | | | | 100% |

GMA *Gemella massiliensis* Marseille-P3249^T; GAS *Gemella asaccharolytica* strain KA00071^T; GCU *Gemella cuniculi* DSM 15828^T; GHA *Gemella haemolysans* strain NCTC10459^T; GMO *Gemella morbillorum* strain NCTC11323^T; GSA *Gemella sanguinis* strain FDAARGOS 742^T; GBE *Gemella bergeri* 617-93^T

under accession numbers LT628479 and FQLS00000000, respectively.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00203-021-02493-2>.

Acknowledgements The authors thank Ludvine Brechard for sequencing the genome and Aurelia Caputo for submitting the genomic sequence to GenBank. We thank also Jaishriram Rathored and Mamadou Beye for their assistance in genomic analysis and Frédéric Cadoret for ensuring the deposit of the strain in different collections.

Author contributions MDMF, MB and CIL drafted the manuscript and analyzed the data. MDMF, ZM, EK and ET performed the technical characterization on strain Marseille-P3249. PEF and DR conceived the study. CIL, GD, FF, and PEF revised the manuscript and participated in its design and coordination. All authors read and approved the final manuscript.

Funding This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the National Research Agency under the program «Investissements d'avenir», reference ANR-10-IAHU-03, the Région Provence Alpes Côte d'Azur, and European funding FEDER PRIM1.

Declarations

Conflict of interest Prs Fournier and Raoult are co-founders of the Techno Jouvence startup. The techno Jouvence startup had no role in this study.

Informed consent The volunteer has given freely his authorization by signed and informed consent for advanced studies to be done on the collected sample. In addition, all the methods used in this study were performed in accordance with relevant guidelines and regulations conformed to the Declaration of Helsinki.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Berger U (1961) A proposed new genus of gram-negative cocci: *Gemella*. Int J Syst Evol Microbiol 11:17–19. <https://doi.org/10.1099/0096266X-11-1-17>
- Collins MD (2006) The genus *Gemella*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H, Stackebrandt E (eds) The prokaryotes, 3rd edn. Springer, New York, pp 511–518
- Collins MD, Hutson RA, Falsen E et al (1998a) *Gemella bergeriae* sp. nov., isolated from human clinical specimens. J Clin Microbiol 36:1290–1293
- Collins MD, Hutson RA, Falsen E et al (1998b) Description of *Gemella sanguinis* sp. nov., isolated from human clinical specimens. J Clin Microbiol 36:3090–3093
- Collins MD, Rodriguez Jovita M, Foster G et al (1999) Characterization of a *Gemella*-like organism from the oral cavity of a dog: description of *Gemella palaticanis* sp. nov. Int J Syst Bacteriol 49(Pt 4):1523–1526. <https://doi.org/10.1099/00207713-49-4-1523>
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Elsawi Z, Togo AH, Beye M et al (2017) *Hugonella massiliensis* gen. nov., sp. nov., genome sequence, and description of a new strictly anaerobic bacterium isolated from the human gut. Microbiologyopen. 6(4):e00458. <https://doi.org/10.1002/mbo3.458>
- Fonkou MDM, Bilén M, Cadoret F et al (2018) “*Enterococcus timonensis*” sp. nov., “*Actinomyces marseillensis*” sp. nov., “*Leptotrichia massiliensis*” sp. nov., “*Actinomyces pacensis*” sp. nov., “*Actinomyces oralis*” sp. nov., “*Actinomyces culturomici*” sp. nov. and “*Gemella massiliensis*” sp. nov., new bacterial species isolated from the human respiratory microbiome. New Microbes New Infect 22:37–43. <https://doi.org/10.1016/j.nmni.2017.12.005>
- Fournier P-E, Lagier J-C, Dubourg G, Raoult D (2015) From culturomics to taxonomogenomics: a need to change the taxonomy of prokaryotes in clinical microbiology. Anaerobe 36:73–78. <https://doi.org/10.1016/j.anaerobe.2015.10.011>
- Hoyle L, Foster G, Falsen E, Collins MD (2000) Characterization of a *Gemella*-like organism isolated from an abscess of a rabbit: description of *Gemella cunicula* sp. nov. Int J Syst Evol Microbiol 50(Pt 6):2037–2041. <https://doi.org/10.1099/00207713-50-6-2037>
- Kilpper-Bälz R, Schleifer KH (1988) Transfer of *Streptococcus morbillorum* to the Genus *Gemella* as *Gemella morbillorum* comb. nov. Int J Syst Evol Microbiol 38:442–443. <https://doi.org/10.1099/00207713-38-4-442>
- Kim M, Oh H-S, Park S-C, Chun J (2014) Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. Int J Syst Evol Microbiol 64:346–351. <https://doi.org/10.1099/ijs.0.059774-0>
- Lagier J-C, Dubourg G, Million M et al (2018) Culturing the human microbiota and culturomics. Nat Rev Microbiol 16:540–550. <https://doi.org/10.1038/s41579-018-0041-0>
- Price MN, Dehal PS, Arkin AP (2009) FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. Mol Biol Evol 26:1641–1650. <https://doi.org/10.1093/molbev/msp077>
- Tindall BJ, Rosselló-Móra R, Busse H-J et al (2010) Notes on the characterization of prokaryote strains for taxonomic purposes. Int J Syst Evol Microbiol 60:249–266. <https://doi.org/10.1099/ijs.0.016949-0>
- Ulger-Toprak N, Summanen PH, Liu C et al (2010) *Gemella asaccharolytica* sp. nov., isolated from human clinical specimens. Int J Syst Evol Microbiol 60:1023–1026. <https://doi.org/10.1099/ijs.0.001966-0>
- Wayne LG (1988) International Committee on Systematic Bacteriology: announcement of the report of the ad hoc Committee on Reconciliation of Approaches to Bacterial Systematics. Zentralbl Bakteriol Mikrobiol Hyg A 268:433–434

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.