



HAL
open science

Association between mortality and highly antimicrobial-resistant bacteria in intensive care unit-acquired pneumonia

Ines Lakbar, Sophie Medam, Romain Ronflé, Nadim Cassir, Louis Delamarre, Emmanuelle Hammad, Alexandre Lopez, Alain Lepape, Anaïs Machut, Mohamed Boucekine, et al.

► To cite this version:

Ines Lakbar, Sophie Medam, Romain Ronflé, Nadim Cassir, Louis Delamarre, et al.. Association between mortality and highly antimicrobial-resistant bacteria in intensive care unit-acquired pneumonia. *Scientific Reports*, 2021, 11 (1), 10.1038/s41598-021-95852-4 . hal-03429611

HAL Id: hal-03429611

<https://amu.hal.science/hal-03429611v1>

Submitted on 28 Jun 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



OPEN

Association between mortality and highly antimicrobial-resistant bacteria in intensive care unit-acquired pneumonia

Ines Lakbar^{1,2,26}, Sophie Medam^{1,26}, Romain Ronflé¹, Nadim Cassir³, Louis Delamarre^{1,2}, Emmanuelle Hammad¹, Alexandre Lopez^{1,3}, Alain Lepape^{4,5,8}, Anaïs Machut^{5,7}, Mohamed Boucekine⁶, Laurent Zieleskiewicz¹, Karine Baumstarck⁶, Anne Savey^{5,7,8}, Marc Leone^{1,3,9}✉ & REA RAISIN Study Group*

Data on the relationship between antimicrobial resistance and mortality remain scarce, and this relationship needs to be investigated in intensive care units (ICUs). The aim of this study was to compare the ICU mortality rates between patients with ICU-acquired pneumonia due to highly antimicrobial-resistant (HAMR) bacteria and those with ICU-acquired pneumonia due to non-HAMR bacteria. We conducted a multicenter, retrospective cohort study using the French National Surveillance Network for Healthcare Associated Infection in ICUs (“REA-Raisin”) database, gathering data from 200 ICUs from January 2007 to December 2016. We assessed all adult patients who were hospitalized for at least 48 h and presented with ICU-acquired pneumonia caused by *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, or *A. baumannii*. The association between pneumonia caused by HAMR bacteria and ICU mortality was analyzed using the whole sample and using a 1:2 matched sample. Among the 18,497 patients with at least one documented case of ICU-acquired pneumonia caused by *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, or *A. baumannii*, 3081 (16.4%) had HAMR bacteria. The HAMR group was associated with increased ICU mortality (40.3% vs. 30%, odds ratio (OR) 95%, CI 1.57 [1.45–1.70], $P < 0.001$). This association was confirmed in the matched sample (3006 HAMR and 5640 non-HAMR, OR 95%, CI 1.39 [1.27–1.52], $P < 0.001$) and after adjusting for confounding factors (OR ranged from 1.34 to 1.39, all $P < 0.001$). Our findings suggest that ICU-acquired pneumonia due to HAMR bacteria is associated with an increased ICU mortality rate, ICU length of stay, and mechanical ventilation duration.

Hospital-acquired pneumonia (HAP) is a common condition that is responsible for a large proportion of hospital-acquired infections, reaching 22% of cases in the United States and 15.6% of cases in France^{1,2}. In intensive care units (ICUs), HAP refers to both healthcare-associated pneumonia and ventilator-associated pneumonia³. The attributable mortality of ventilator-associated pneumonia has been extensively evaluated in recent studies^{4–6}, although this has led to conflicting results because of confounding biases⁷. Likewise, the attributable mortality of ICU-acquired pneumonia—that is, healthcare-associated pneumonia diagnosed after a 48 h stay in the

¹Department of Anesthesiology and Intensive Care Unit, Aix Marseille University, Assistance Publique Hôpitaux Universitaires de Marseille, Nord Hospital, Marseille, France. ²Department of Anesthesiology and Intensive Care Unit, University hospital of Toulouse, Toulouse, France. ³MEPHI, IHU Méditerranée Infection, Aix Marseille Université, Marseille, France. ⁴Intensive Care Unit, Centre Hospitalier Lyon Sud, Pierre Bénite, Hospices Civils de Lyon, France. ⁵Rea-Raisin study group (National network for Healthcare-Associated Infection surveillance in ICU, Marseille, France. ⁶APHM, EA 3279 CERESS, School of Medicine, La Timone Medical Campus, Health Service Research and Quality of Life Center, Aix Marseille Université, Marseille, France. ⁷Infection Control & Prevention, Hôpital Henry Gabrielle, Hospices Civils de Lyon, Saint Genis Laval, France. ⁸PHE3ID, Centre International de Recherche en Infectiologie, INSERM U1111, CNRS Unité Mixte de Recherche 5308, ENS de Lyon, Université Claude Bernard Lyon 1, Saint Genis Laval, France. ⁹Service d’anesthésie et de réanimation, Chemin des Bourrely, Hôpital Nord, 13015 Marseille, France. ²⁶These authors contributed equally: Ines Lakbar and Sophie Medam. *A list of authors and their affiliations appears at the end of the paper. ✉email: marc.leone@ap-hm.fr

ICU—remains difficult to accurately assess. However, these infections most likely have a detrimental effect on the outcomes of patients in the ICU^{1,8–10}.

In ICU-acquired pneumonia, the resistance level of the causative microorganism may also affect outcome^{11–13}. Investigating the relationship between antibiotic resistance and clinical outcome is challenging, as it is difficult to discriminate the confounders and determinants of this relationship. In addition, patients at the highest risk of death are also likely to be those at the highest risk of infection by highly antimicrobial-resistant (HAMR) bacteria¹⁴. Therefore, this study aimed to compare ICU mortality rates between patients who developed ICU-acquired pneumonia caused by HAMR bacteria and those who developed ICU-acquired pneumonia caused by non-HAMR bacteria among the following causative pathogens: *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, and *A. baumannii*.

The first aim of our study was to compare the ICU mortality rates in patients who developed ICU-acquired pneumonia due to HAMR and non-HAMR bacteria. The secondary aims were to compare the durations of ICU stay and mechanical ventilation between these two groups.

Material and Methods

We performed a retrospective, observational 9-year study using the REA-RAISIN database (from January 2007 to December 2016), a French national surveillance network for healthcare-associated infections in ICUs (the surveillance period was of 6 months from 2007 to 2014 and then surveillance became continuous as of 2015)¹⁵. The number of ICUs contributing to the database increased between 2007 and 2016, varying from 165 to 200. All the patients or their relatives were informed that their data would be used anonymously unless they disagreed with being included. This study was approved by the French Commission Nationale Informatique et Liberté (CNIL No. 588909) and Institutional Review Board (IRB No. 00009118). All included patients were followed up until death or discharge from the ICU.

The inclusion criteria were admission to an ICU for at least 48 h and diagnosis of ICU-acquired pneumonia. The diagnosis criteria for pneumonia, following the European Centre for Disease Prevention and Control (ECDC) definition, were one (if the patient had no past medical history of cardiac or pulmonary disease) or two chest radiographs showing pulmonary infiltrates and (1) at least one of the following clinical signs: hyperthermia (> 38 °C) or a leukocyte count less than 4,000 cells/mm³ or greater than 12,000 cells/mm³; (2) at least one of the following clinical criteria: onset of purulent secretions or changes in characteristics; suggestive auscultation, cough, dyspnea, or tachypnea; low oxyhemoglobin saturation; or increased pulmonary oxygen consumption; and (3) microbiological confirmation by a positive culture from directed bronchoalveolar lavage or from tracheal secretions or by an alternative method¹⁶. Probable cases of ICU-acquired pneumonia defined as positive according to the radiological, biological, and clinical criteria but with no positive microbiology were excluded. For each patient, we considered only the first episode of ICU-acquired pneumonia. The minimum delay between ICU admission and the onset of ICU-acquired pneumonia was 48 h. The REA-Raisin database allowed the registration of up to two causative pathogens per infection, and resistance profiles were reported only if the causative pathogens were one of the following: *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, or *A. baumannii*.

Data collection. Demographic and clinical data, including clinical and microbiological assessments from the electronic medical charts were analyzed. We extracted age, gender, Simplified Acute Physiology Score (SAPS) 2 at ICU admission, administration of antibiotic treatment within 48 h before or after ICU admission, length of ICU stay, patient's provenance before ICU admission (in-hospital patient or out-hospital patient, hospitalizations occurring before the stay of interest were not recorded), immunodeficiency according to Acute Physiology and Chronic Health Evaluation II score, medical or surgical origin of patients, use of mechanical ventilation, and trauma diagnosis.

Definition of HAMR status. Each episode of ICU-acquired pneumonia was microbiologically confirmed to identify the causal pathogens. ICU-acquired pneumonia due to HAMR bacteria was defined as the identification of at least one antimicrobial-resistant bacterium in a clinical sample, as reported in Table 1¹⁵.

Statistical methods. To assess the association between HAMR bacteria and ICU mortality, analyses were conducted in two steps: (1) on the whole sample and (2) on a 1:2 matched sample (at least one control). Matching was based on seven factors: sex, age (within 5 years), SAPS 2 (within 10 points), antibiotic treatment at admission (yes–no), category of patient (medical vs. surgical), mechanical ventilation and type of pathogen. The last factor was determined as follows: when pneumonia was caused by a single pathogen (n = 15,717) or two identical pathogens (n = 991), patients were matched based on the same causative pathogen; for pneumonia caused by two different pathogens (n = 2096), a Delphi review was performed by the authors and a panel of experts to determine on which pathogens the matching should be based. The results of the Delphi review are provided in the Supplementary Material. Patients were matched using the %match SAS macro¹⁷, which implements an optimal matching algorithm¹⁸. The optimal algorithm sorts cases and controls, identifies all pairs that satisfy the specified distance measures, and then selects the set of pairs that minimizes the total distance between all pairs.

For each sample, patients with pneumonia due to HAMR bacteria were compared to patients with pneumonia due to non-HAMR according to the main characteristics. To assess the link between HAMR status and ICU mortality, comparisons based on socio-demographic, clinical, and hospital data between survivors and non-survivors were performed (1) on the whole sample using chi² tests or Student's t tests according to the nature of the variable and (2) on the matched sample using conditional logistic regression¹⁹, taking into account the matched procedure. Odds ratios (OR) with a 95% confidence interval (CI) were estimated. Multivariate models were assessed to confirm the effect of HAMR status on ICU mortality after adjusting for the main confounding

Resistance classification 2007-2010: the profile considered to be HAMR in our analysis is framed in a box											
	0	1	2	3							
<i>Staphylococcus aureus</i>	OXA-S	OXA-NS	OXA-NS, genta-NS	GLY-NS							
<i>Enterobacteriace</i>	AMP-S	AMP-NS, CTX-S	CTX-NS (ESBL)	CTX-NS (no ESBL)							
<i>Acinetobacter baumannii</i>	-	CAZ-S	CAZ-NS	-							
<i>Pseudomonas aeruginosa</i> <i>Burkholderia cepacia</i> <i>Stenotrophomonas maltophilia</i>	ticar-S	ticar-NS, CAZ-S	CAZ-NS	-							
Resistance classification 2011-2015: the profile considered to be HAMR in our analysis is framed in a box											
<i>Staphylococcus aureus</i>	OXA-S, VAN-S	OXA-NS	VAN-NS	-							
<i>Enterobacteria</i>	CTX-S, IMP-S	CTX-NS non ESBL	CTX-NS ESBL, IMP-S	IMP-NS							
<i>Acinetobacter baumannii</i>	CAZ-S, IMP-S	CAZ-NS, IMP-S	CAZ-S, IMP-NS	CAZ- NS, IMP- NS							
<i>Pseudomonas aeruginosa</i>	CAZ-S, IMP-S	CAZ- NS, IMP-S	CAZ-S, IMP- NS	CAZ- NS, IMP- NS							
Resistance classification 2016: the profile considered to be HAMR in our analysis is framed in a box											
	OXA	AMP	GLY	AMC	C3G	PTZ	CAZ	CAR	COL	ESBL	PanR
<i>Staphylococcus aureus</i>	X		X								X
<i>Enterobacteria</i>				X	X			X		X	X
<i>Pseudomonas aeruginosa</i>						X	X	X	X		X
<i>Acinetobacter baumannii</i>							X	X	X		X

Table 1. Evolution of the classification of the antimicrobial resistance by year and by micro-organisms in the REA-RAISIN database. NS = Non-susceptible S = susceptible OXA: oxacillin (or methicillin), AMP: ampicillin (or amoxicillin), GLY: glycopeptide (vancomycin or teicoplanin), AMC: amoxicillin-clavulanic acid, ticar: ticarcillin, C3G: 3rd generation cephalosporins = cefotaxime (or ceftriaxone) PTZ: piperacillin-tazobactam, CAZ: ceftazidime, CAR: carbapenem = imipenem or doripenem or meropenem, IMP: imipenem, VAN: vancomycin, COL: colistin, ESBL: extended-spectrum beta-lactamase-producing, PANR: non susceptible to all tested agents, HAMR: highly antimicrobial-resistant.

factors (1) with the whole sample using logistic regression (adjustment for sex, age, SAPS 2, antibiotic at admission, immunodeficiency status, mechanical ventilation, traumatic situation, provenance, category of patients, delay of pneumonia) and (2) with the matched sample using a generalized linear model, PROC GLIMMIX SAS (adjustment for provenance, immunodeficiency, trauma, delay of pneumonia). The link between HAMR status and ICU mortality was also assessed in predefined subgroups: men and women, younger (< 65 years) and older (≥ 65 years) patients, medical and surgical patients, mechanical ventilation and no mechanical ventilation, and patients with or without antibiotics at ICU admission. A two-sided p-value of less than 0.05 was considered to indicate statistical significance. All statistical analyses were conducted using IBM SPSS Statistics for Windows (Version 21.0. Armonk, NY: IBM Corp) and SAS 9.4 (SAS Institute).

Results

For the study period of 9 years, the database contained 355,116 patients, of which 30,561 (8.6%) developed at least one episode of ICU-acquired pneumonia. A total of 25,096 patients had a documented infection, and for 18,529, a bacteria profile of the isolated strains corresponding to *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, or *A. baumannii* was available. Of these 18,529 patients, the vital status of 18,497 was available at the time of discharge from the ICU. A flowchart of the study is displayed in Fig. 1, and the patient features are presented in Table 2.

Of the 18,497 included cases of infection, 3081 (17%) were infected with HAMR bacteria and 15,416 (83%) with non-HAMR bacteria (details about the pathogens are provided in the Supplemental Material). The ICU mortality rate was 32%, representing 5872 patients aged 68 ± 13 years with an average SAPS 2 of 55 ± 18 . The average ICU length of stay was 33 ± 26 days. Invasive mechanical ventilation was required in 18,109 (98%) patients for a duration of 28 ± 25 days. Of note, 11,512 (62%) patients received antibiotics within 48 h of admission. The reasons for ICU admission were medical (67%) and surgical (33%).




Figure 1. Selection of the final study group ($n = 18,772$) from 355,116 patients hospitalized ≥ 48 h in French ICUs, 2007–2016.

For several sociodemographic and clinical variables, there were significant differences between patients with ICU-acquired pneumonia due to HAMR bacteria and those with ICU-acquired pneumonia due to non-HAMR bacteria (Table 2). Therefore, 5640 non-HAMR ICU-acquired pneumonia cases were matched with 3006 ICU-acquired cases of pneumonia caused by HAMR bacteria. Details are provided in Table 2.

In the whole sample, HAMR group and non-HAMR group were associated with 40.3% and 30.0% ICU mortality rates, respectively (differential 10, odds ratio (OR) and 95% confidence interval (CI) 1.57[1.45–1.70], $P < 0.001$). Age, sex, provenance, immunosuppression, ICU length of stay, and HAMR status were associated with ICU mortality (Table 3). HAMR status was still associated with ICU mortality (1) in the matched sample (OR 95%, CI 1.39 [1.27–1.52], $P < 0.001$) (Table 3); (2) after adjusting for the main confounding factors (ORs ranged from 1.34 to 1.39, all p -values < 0.001) (Table 4); and (3) in prespecified subgroups: females *versus* males, age below 65 years *versus* above (or equal) 65 years, antibiotic at ICU admission *versus* no-antibiotic at ICU admission, medical patient *versus* surgical patient, mechanical ventilation *versus* no mechanical ventilation, and in-hospital patient *versus* out-hospital patient (Fig. 2).

The mean durations of ICU length of stay (37 ± 26 days *versus* 33 ± 26 days, $P < 0.001$) and mechanical ventilation (31 ± 26 days *versus* 27 ± 24 days, $P < 0.001$) were higher in the HAMR group than in the non-HAMR group. The delay between ICU admission and the pneumonia onset differed between HAMR group and non-HAMR group from 16.0 days \pm 12.4 to 14.1 days \pm 14.4, respectively ($P < 0.001$).

Discussion

According to our results, developing ICU-acquired pneumonia due to HAMR bacteria was an independent risk factor for ICU mortality. To our knowledge, our study included one of the largest cohorts to assess the association between infection due to HAMR bacteria and ICU mortality.

Lambert et al. published the largest prospective European study ($n = 119,699$ patients, of whom 8525 were diagnosed with HAP)²⁰. They concluded that the effect of antimicrobial resistance on mortality was modest. In the same line, Paramythiotou et al. concluded that a direct association between infections caused by Gram-negative

	Patients with vital status at discharge						
	Whole sample			P-value	Matched sample		P-value
	Non HAMR	HAMR	Non HAMR		HAMR		
	n = 15,416	n = 3081	n = 5640	n = 3,006			
Age (years)							
M ± SD	62.2 ± 16.4	64.0 ± 14.9	<0.001	64.7 ± 13.6	64.2 ± 14.7	–	
m [IQR]	65 [53–75]	66 [56–75]		67 [58–75]	66 [56–75]		
Sex							
Female	4318 (28.0)	842 (27.3)	0.442	1506 (26.7)	807 (26.8)	–	
Male	11,098 (72.0)	2239 (72.7)		4134 (73.3)	2199 (73.2)		
Provenance							
Inpatient	6897 (44.9)	1677 (54.5)	<0.001	2735 (48.6)	1642 (54.7)	<0.001	
Outpatient	8474 (55.1)	1398 (45.5)		2897(51.4)	1361 (45.3)		
Type							
Medical	10,170 (66.1)	2216 (72.2)	<0.001	4083 (72.4)	2174 (72.3)	–	
Surgery	5216 (33.9)	854 (27.8)		1556 (27.6)	831 (27.7)		
SAPS 2 at admission							
M ± SD	50.5 ± 18.2	52.4 ± 18.2	<0.001	51.3 ± 15.4	52.2 ± 17.6	–	
m [IQR]	49 [37–62]	50 [39–64]		50 [40–60]	50 [39–64]		
Immunosuppression							
No	12,816 (86.2)	2357 (78.4)	<0.001	4673 (85.0)	2310 (78.5)	<0.001	
Yes	2058 (13.8)	650 (21.6)		821 (14.9)	632 (21.5)		
Mechanical ventilation							
No	319 (2.1)	58 (1.9)	0.505	62 (1.1)	38 (1.3)	–	
Yes	15,089 (97.9)	3020 (98.1)		5578 (98.9)	2968 (98.7)		
Mechanical ventilation (days)							
M ± SD	27.3 ± 24.5	31.5 ± 25.9	<0.001	29.2 ± 25.1	31.5 ± 25.8	<0.001	
m [IQR]	21 [12–35]	24 [15–41]		22 [13–37]	24 [15–41]		
Trauma patients							
No	12,967 (84.3)	2881(93.6)	<0.001	4990 (88.5)	2818 (93.7)	<0.001	
Yes	2416 (15.7)	197 (6.4)		646 (11.5)	188 (6.3)		
Antibiotic treatment at admission							
No	6137 (40.2)	683 (22.3)	<0.001	1283 (22.8)	668 (22.3)	–	
Yes	9131 (59.8)	2381 (77.7)		4345 (77.2)	2331 (77.7)		
ICU duration (days)							
M ± SD	32.9 ± 26.2	37.2 ± 26.7	<0.001	35.1 ± 27.6	37.3 ± 26.6	<0.001	
m [IQR]	26 [16–41]	30 [19–48]		28 [18–43]	30 [19–48]		
Delay between ICU admission and event							
M ± SD	12.8 ± 12.3	15.9 ± 12.3	<0.001	14.1 ± 14.4	16.0 ± 12.4	<0.001	
m [IQR]	10 [6–16]	13 [8–19]		11 [7–17]	13 [8–19]		
Delay between pneumonia and ICU discharge (days)							
M ± SD	21.1 ± 21.7	22.3 ± 22.1	<0.001	22.0 ± 22.3	22.3 ± 22.1	0.391	
m [IQR]	15 [8–26]	16 [8–29]		15 [9–27]	16 [8–29]		
ICU mortality							
Survivors	10,785 (70.0)	1840 (59.7)	<0.001	3799 (67.4)	1794 (59.7)	<0.001	
Non-survivors	4631 (30.0)	1241 (40.3)		1841 (32.6)	1212 (40.3)		

Table 2. Patient characteristics and comparison according to the HAMR status. *ICU* intensive care unit, *HAMR* high antimicrobial resistance, *SAPS2* simplified acute physiology score. M ± SD: mean ± standard deviation; m [IQR]: median [interquartile range]. *with vital status not missing.

resistant bacteria and ICU mortality was not confirmed²¹. However, most studies were performed in single centers and included small numbers of patients. In addition, most were characterized by a high degree of heterogeneity that prevented definitive conclusions from being made²¹. Three studies have suggested that antibiotic resistance led to an increase in crude mortality, even after adjusting for two of them^{12,14,22}. Here, we found an association between ICU mortality and the occurrence of an infection due to HAMR bacteria.


The effect of bacterial resistance on patient outcomes can be explained by three determinants. First, patients infected by HAMR bacteria are more likely to receive an inadequate empirical antimicrobial therapy²³. As there is

	Whole sample*			
	Survivors	Non-survivors	OR [95%CI]**	P-value
	n = 12,625	n = 5872		
Age (years)				
M ± SD	60.0 ± 16.8	67.8 ± 13.4	1.03 [1.03–1.04]	<0.001
m [IQR]	62 [51–73]	70 [60–78]		
Sex				
Female	3507 (27.8)	1653 (28.2)	0.98 [0.91–1.05]	0.599
Male (1)	9118 (72.2)	4219 (71.8)		
Provenance				
Inpatient	5537 (44.0)	3037 (51.9)	0.72 [0.68–0.77]	<0.001
Outpatient (1)	7055 (56.0)	2817 (48.1)		
Type				
Medical	8012 (63.6)	4374 (74.8)	0.58 [0.55–0.63]	<0.001
Surgery (1)	4592 (36.4)	1477 (25.2)		
SAPS 2 at admission				
M ± SD	48.7 ± 18.0	55.2 ± 18.1	1.02 [1.01–1.02]	
m [IQR]	47 [36–60]	54 [42–67]		
Immunosuppression				
No	10,655 (87.3)	4518 (79.6)	1.79 [1.62–1.92]	<0.001
Yes (1)	1547 (12.7)	1161 (20.4)		
ICU duration (days)				
M ± SD	34.5 ± 25.9	31.7 ± 27.3	1.00 [0.99–1.00]	<0.001
m [IQR]	28 [18–43]	25 [15–40]		
Delay between ICU admission and event				
M ± SD	12.8 ± 10.9	14.4 ± 14.8	1.01 [1.00–1.01]	<0.001
m [IQR]	10 [6–16]	11 [7–18]		
Delay between event and ICU discharge				
M ± SD	22.7 ± 21.8	18.3 ± 21.5	0.99 [0.98–0.99]	<0.001
m [IQR]	16 [9–29]	12 [6–23]		
Mechanical ventilation				
No	317 (2.5)	60 (1.0)	2.49 [1.88–3.29]	<0.001
Yes (1)	12,301 (97.5)	5808 (99.0)		
Mechanical ventilation duration (days)				
M ± SD	27.9 ± 24.9	28.3 ± 24.8	1.00 [1.00–1.00]	0.344
m [IQR]	21 [12–35]	22 [13–36]		
Trauma patients				
No	10,436 (82.8)	5412 (92.4)	0.39 [0.35–0.43]	<0.001
Yes (1)	2169 (17.2)	444 (7.6)		
Antibiotic treatment at ICU admission				
No	4912 (39.2)	1908 (32.9)	1.31 [1.23–1.40]	<0.001
Yes (1)	7612 (60.8)	3900 (67.1)		
Bacteria feature				
Non HAMR	10,785 (85.4)	4631 (78.9)	1.57 [1.45–1.70]	<0.001
HAMR (1)	1840 (14.6)	1241 (21.1)		
Matched sample*				
	Survivors	Non-survivors	OR [95%CI]**	P-value
	n = 5593	n = 3053		
Bacteria feature				
Non HAMR	3799 (67.4)	1794 (59.7)	1.39 [1.27–1.52]	<0.001
HAMR (1)	1841 (32.6)	1212 (40.3)		

Table 3. Factors associated with ICU mortality on the whole sample and the matched sample (univariate analysis). *ICU* intensive care unit, *HAMR* highly antimicrobial resistant *SAPS2* simplified acute physiology score. *M ± SD*: mean ± SD; *m [IQR]*: median [interquartile range]; *OR [95%CI]*: odd ratio [95% confidence interval]. *with vital status not missing; **OR is provided for the modality (1).

Whole sample				Matched sample		
Model adjusted for...		OR [95%CI]	P-value	Model adjusted for...		P-value
1	Sex, age, SAPS2	1.48 [1.36–1.61]	<0.001	1	Provenance	1.39 [1.27–1.53] <0.001
2	Sex, age, SAPS2, antibiotic at admission,	1.45 [1.34–1.58]	<0.001	2	Immunosuppression	1.37 [1.24–1.51] <0.001
3	Sex, age, SAPS2, antibiotic at admission, immunodeficiency, mechanical ventilation	1.41 [1.29–1.53]	<0.001	3	Provenance, immunosuppression	1.36 [1.23–1.50] <0.001
4	Sex, age, SAPS2, antibiotic at admission, immunodeficiency, mechanical ventilation, traumatic situation, provenance, category of patient	1.36 [1.25–1.48]	<0.001	4	Provenance, immunosuppression, trauma	1.34 [1.21–1.48] <0.001
5	Sex, age, SAPS2, antibiotic at admission, immunodeficiency, mechanical ventilation, traumatic situation, provenance, delay of pneumonia	1.35 [1.24–1.47]	<0.001	5	Provenance, immunosuppression, trauma, delay of pneumonia	1.33 [1.20–1.46] <0.001

Table 4. Highly antimicrobial resistant status and ICU mortality (variate analysis). SAPS2 simplified acute physiology score.



Inf 65y: sub-group below 65 years of age,
 Sup 65y: subgroup greater than or equal to 65 years of age
 HAMR: highly antimicrobial resistant

Figure 2. Risk of mortality associated with HAMR status by subgroup.

an association between the adequateness of the empirical antimicrobial therapy and survival, this hypothesis may explain the increased ICU mortality that we reported here²⁴. Second, an increased virulence has been suspected in some resistant bacteria²⁵, as suggested by a murine model of infection due to *P. aeruginosa*²⁶. The authors found that the acquisition of antibiotic resistance improved the fitness of the bacteria and thus promoted its survival and virulence. However, the higher virulence of HAMR bacteria remains unlikely as conversely, other studies described a loss of potency and virulence in specific bacteria-antibiotic pairs^{27,28}. The third determinant relates to host factors and co-morbidities. A frail patient has a higher risk of recurrent hospitalizations, exposure to antibiotics, and thus colonization and infection by HAMR bacteria²⁹. Moreover, the effects of antibiotics themselves could be deleterious, as suggested previously³⁰.

Our study has several limitations. First, this was a retrospective analysis of a large database. Hence, our choices for the statistical approach could be a matter of debate. However, our results were confirmed using several statistical approaches. Second, the definition of ICU-acquired pneumonia relies on each on-site physician, with different sampling techniques, without external confirmation, while the diagnosis of HAP remains challenging³¹. Third, only *S. aureus*, *Enterobacteriaceae*, *P. aeruginosa*, and *A. baumannii* pneumonia were included in the analysis, thus excluding, notably, streptococci and other gram-negative bacteria. Finally, there was no mention in the database of the antimicrobial therapy, specifically the adequacy and delay of the empirical treatment. As discussed above, this is a major determinant of mortality in these patients^{23,24}. However, in our study, the ICU-acquired pneumonia due to HAMR bacteria occurred later than those due to non-HAMR bacteria, suggesting an increased number of late-onset pneumonia in the HAMR group. Following international guidelines, the patients with late-onset pneumonia are more prone to receive broad-spectrum antibiotics than those with early-onset pneumonia³¹. Notably, 62.2% of the patients included in our study received antibiotics at admission in the ICU. This finding is in line with the rates recently reported in an international observational 24-h point prevalence among 15,202 patients³². The selection of our population was based on voluntary participation in the network and a duration of ICU stay of at least 48 h (for a reduced surveillance workload). Thus, our findings may not be reflective of the entire ICU patient population, as they may pertain specifically to patients exposed to infections acquired in the ICU. Finally, the definition of antimicrobial-resistant bacteria evolved during the study period, which could have affected our findings, despite efforts to ensure comparability across the nine years. As ecology

and therapies have evolved over the years with the arrival of new molecules in our therapeutic arsenal, the classification of resistances, established prospectively by the designers of the database, has also evolved towards a more recent and precise definition in 2016 which is more consistent with recent guidelines^{1,31}.

Conclusion

In conclusion, the findings of our study suggest that ICU-acquired pneumonia due to HAMR bacteria was associated with an increased ICU mortality rate, duration of ICU stay, and mechanical ventilation duration. However, the reasons behind this association remain to be elucidated.

Data availability

This study was approved by the French Commission Nationale Informatique et Liberté (CNIL No. 588909) and Institutional Review Board (IRB No. 00009118). Our study has no attached data.

Received: 26 April 2021; Accepted: 26 July 2021

Published online: 13 August 2021

References

- Kalil, A. C. *et al.* Management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the infectious diseases society of America and the American thoracic society. *Clin. Infect. Dis.* **63**, e61–e111 (2016).
- Daniau, C. *et al.* Infections associées aux soins en établissement de santé: résultats de l'enquête nationale de prévalence 2017, france/healthcare-associated infections in healthcare facilities: results of french national point prevalence survey. *Science* **16**, 412–423 (2017).
- Timsit, J.-F., Esaied, W., Neuville, M., Bouadma, L. & Mourvillier, B. Update on ventilator-associated pneumonia. *F1000Res* **6**, 2061 (2017).
- Timsit, J.-F., Zahar, J.-R. & Chevret, S. Attributable mortality of ventilator-associated pneumonia. *Curr. Opin. Crit. Care* **17**, 464–471 (2011).
- Bekaert, M. *et al.* Attributable mortality of ventilator-associated pneumonia: a reappraisal using causal analysis. *Am. J. Respir. Crit. Care Med.* **184**, 1133–1139 (2011).
- Nguile-Makao, M. *et al.* Attributable mortality of ventilator-associated pneumonia: respective impact of main characteristics at ICU admission and VAP onset using conditional logistic regression and multi-state models. *Intensive Care Med.* **36**, 781–789 (2010).
- Papazian, L., Klompas, M. & Luyt, C.-E. Ventilator-associated pneumonia in adults: a narrative review. *Intensive Care Med.* <https://doi.org/10.1007/s00134-020-05980-0> (2020).
- Torres, A. *et al.* International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia. *Eur. Respir. J.* **50**, 1–26 (2017).
- Ibn, S. W. *et al.* A comparison of the mortality risk associated with ventilator-acquired bacterial pneumonia and nonventilator ICU-acquired bacterial pneumonia. *Crit. Care Med.* **47**, 345–352 (2019).
- Melsen, W. G. *et al.* Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies. *Lancet. Infect. Dis* **13**, 665–671 (2013).
- Damas, P. *et al.* Severity of ICU-acquired pneumonia according to infectious microorganisms. *Intensive Care Med.* **37**, 1128–1135 (2011).
- Razazi, K. *et al.* Frequency, associated factors and outcome of multi-drug-resistant intensive care unit-acquired pneumonia among patients colonized with extended-spectrum β -lactamase-producing Enterobacteriaceae. *Ann. Intensive Care* **7**, 1–7 (2017).
- Zahar, J.-R. *et al.* Is Methicillin resistance associated with a worse prognosis in *Staphylococcus aureus* ventilator-associated pneumonia? *Clin. Infect. Dis.* **41**, 1224–1231 (2005).
- Bottazzi, A. *et al.* Mortality attributable to different Klebsiella susceptibility patterns and to the coverage of empirical antibiotic therapy: a cohort study on patients admitted to the ICU with infection. *Intensive Care Med.* **44**, 1709–1719 (2018).
- Lepape, A., MacHut, A. & Savey, A. National network REA-RAISIN of adult intensive care acquired infections methods and main results. *Med. Intensive Reanim.* **27**, 197–203 (2018).
- Plachouras, D., Lepape, A. & Suetens, C. ECDC definitions and methods for the surveillance of healthcare-associated infections in intensive care units. *Intensive Care Med.* **44**, 2216–2218 (2018).
- Bergstralh, E. J. & Kosanke, J. L. Computerized matching of cases to controls. *Tech. Rep.* **56**, 1–28 (1995).
- Rosenbaum, P. R. Optimal matching for observational studies. *J. Am. Stat. Assoc.* **84**, 1024–1032 (1989).
- Hosmer, D. W., Lemeshow, S. & Sturdivant, R. X. *Applied Logistic Regression* (Wiley, 2013).
- Lambert, M. L. *et al.* Clinical outcomes of health-care-associated infections and antimicrobial resistance in patients admitted to European intensive-care units: a cohort study. *Lancet. Infect. Dis* **11**, 30–38 (2011).
- Paramythiotou, E. & Routsis, C. Association between infections caused by multidrug-resistant gram-negative bacteria and mortality in critically ill patients. *World J. Crit. Care Med.* **5**, 111 (2016).
- Barbier, F. *et al.* Colonization and infection with extended-spectrum β -lactamase-producing Enterobacteriaceae in ICU patients: What impact on outcomes and carbapenem exposure? *J. Antimicrob. Chemother.* **71**, 1088–1097 (2016).
- Rottier, W. C., Ammerlaan, H. S. M. & Bonten, M. J. M. Effects of confounders and intermediates on the association of bacteraemia caused by extended-spectrum β -lactamase-producing enterobacteriaceae and patient outcome: a meta-analysis. *J. Antimicrob. Chemother.* **67**, 1311–1320 (2012).
- Rhodes, A. *et al.* Surviving Sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med.* **43**, 304–377 (2017).
- Guillard, T., Pons, S., Roux, D., Pier, G. B. & Skurnik, D. Antibiotic resistance and virulence: understanding the link and its consequences for prophylaxis and therapy. *BioEssays News Rev. Mol. Cell. Dev. Biol.* **38**, 682–693 (2016).
- Roux, D. *et al.* Fitness cost of antibiotic susceptibility during bacterial infection. *Sci. Transl. Med.* **7**, 29 (2015).
- Hraiech, S. *et al.* Impaired virulence and fitness of a colistin-resistant clinical isolate of *Acinetobacter baumannii* in a rat model of pneumonia. *Antimicrob. Agents Chemother.* **57**, 5120–5121 (2013).
- Andersson, D. I. & Hughes, D. Antibiotic resistance and its cost: is it possible to reverse resistance? *Nat. Rev. Microbiol.* **8**, 260–271 (2010).
- Giarratano, A., Green, S. E. & Nicolau, D. P. Review of antimicrobial use and considerations in the elderly population. *Clin. Interv. Aging* **13**, 657–667 (2018).
- Jensen, J. U. *et al.* Procalcitonin-guided interventions against infections to increase early appropriate antibiotics and improve survival in the intensive care unit: a randomized trial. *Crit. Care Med.* **39**, 2048–2058 (2011).
- Torres, A. *et al.* International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia. *Eur. Respir. J.* **50**, 64 (2017).

32. Vincent, J.-L. *et al.* Prevalence and outcomes of infection among patients in intensive care units in 2017. *JAMA* **323**, 1478 (2020).

Acknowledgements

The authors are indebted toward the physicians and nurses who took care of patients and provided the individual data in each center.

Author contributions

Conceptualization: I.L., S.M., R.R., A.L., M.L. Data curation: I.L., K.B., M.B., L.D. Formal analysis: K.B., M.B. Supervision: L.Z., M.L. Writing – original draft: I.L., S.M., M.L. Writing – review & editing: N.C., E.H., A.L., A.L., A.M., A.S., M.L. All authors reviewed the manuscript.

Funding

Department of anesthesiology and intensive care unit of Nord Hospital, Marseille, France.

Competing interests

IL, SM, RR, LD, EH, AL, LZ, MB, NC, AS, AM and KB have no conflict of interest to disclose. ML served as speaker for MSD, Pfizer and as consultant for Amomed, Aguettant and Gilead AL served as consultant for Fresenius.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-95852-4>.

Correspondence and requests for materials should be addressed to M.L.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

REA RAISIN Study Group

Serge Alfandari¹⁰, Sébastien Bailly¹¹, Odile Bajolet¹², Olivier Baldesi¹³, Anne Berger-Carbonne¹⁴, Pierre-Edouard Bollaert¹⁵, Cedric Bretonniere¹⁶, Céline Chatelet¹⁷, Philippe Corne¹⁸, Isabelle Durand-Joly¹⁹, Arnaud Friggeri²⁰, Gaëlle Gasan¹⁷, Rémy Gauzit²¹, Marine Giard⁷, Caroline Landelle¹¹, Thierry Lavigne²², Didier Lepelletier¹⁶, Pierre-François Perrigault¹⁸, Santiago Picos²³, Marie-Aline Robaux²⁴, Vincent Stoeckel²⁵, Jean-François Timsit²¹ & Philippe Vanhems⁷

¹⁰Infection Control, Hospital of Tourcoing, Tourcoing, France. ¹¹University Hospital of Grenoble, Grenoble, France. ¹²University Hospital of Reims, Reims, France. ¹³Hospital of Aix-en-Provence, Aix-en-Provence, France. ¹⁴Santé Publique France, Paris, France. ¹⁵University Hospital of Nancy, Nancy, France. ¹⁶University Hospital of Nantes, Nantes, France. ¹⁷Hospital of Lens, Lens, France. ¹⁸University Hospital of Montpellier, Montpellier, France. ¹⁹Hospital of Dunkerque, Dunkerque, France. ²⁰Hospices Civils de Lyon, Civils de Lyon, France. ²¹Assistance Publique des hôpitaux de Paris, Paris, France. ²²University Hospital of Strasbourg, Strasbourg, France. ²³Hospital of Draguignan, Draguignan, France. ²⁴Hospital of Bayonne, Bayonne, France. ²⁵Hospital of Chalons en Champagne, Chalons en Champagne, France.