



Effects of solar park construction and solar panels on soil quality, microclimate, CO₂ effluxes, and vegetation under a Mediterranean climate

Quentin Lambert, Armin Bischoff, Sixtine Cueff, Alexandre Cluchier, Raphaël Gros

► To cite this version:

Quentin Lambert, Armin Bischoff, Sixtine Cueff, Alexandre Cluchier, Raphaël Gros. Effects of solar park construction and solar panels on soil quality, microclimate, CO₂ effluxes, and vegetation under a Mediterranean climate. *Land Degradation and Development*, 2021, 32 (18), pp.5190-5202. <10.1002/ldr.4101>. <hal-03440295>

HAL Id: hal-03440295

<https://amu.hal.science/hal-03440295v1>

Submitted on 5 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

i. Manuscript Title

**Effects of solar park construction and solar panels on soil
quality, microclimate, CO₂ effluxes and vegetation under
Mediterranean climate**

ii. Authors

Quentin Lambert^{1*}, Armin Bischoff², Sixtine Cueff³, Alexandre Cluchier⁴, Raphael
Gros¹

iii. Author's institutional affiliations

1 Aix Marseille Université, Université d'Avignon, IRD, CNRS, Institut Méditerranéen de
Biodiversité et d'Écologie marine et continentale (IMBE), Campus l'Etoile, Av. Escadrille
Normandie Niémen, 13397, Marseille, Cedex 20, France

2 Aix Marseille Université, Université d'Avignon, IRD, CNRS, Institut Méditerranéen de
Biodiversité et d'Écologie marine et continentale (IMBE), IUT Avignon, Agroparc, BP
61207, 84911 Avignon Cedex 9, France

3 Current address : INRAE, AgroParisTech, UMR ECOSYS, 78850 Thiverval-Grignon, France4
ECO-MED Ecologie & Médiation, Tour Méditerranée, 65 avenue Jules Cantini, 13298
Marseille cedex 20, France

4 ECO-MED Ecologie & Médiation, Tour Méditerranée, 65 avenue Jules Cantini, 13298
Marseille cedex 20, France

iv. Corresponding author: qntnlmbrt@gmail.com

v. Short informative containing containing the major key words.

Effects of solar parks on soil quality, CO₂ effluxes and vegetation under Mediterranean climate

Solar parks are expanding in Europe, but their impact on soil and vegetation is not well studied yet. In the present study including three parks in the Mediterranean region, we show that the construction of solar parks reduces the physical quality of the soil altering main soil functions. Moreover, the presence of solar panels decreases CO₂ emissions and temperature but does not change the structure of plant communities.

vi. Short running title

Effects of solar parks on soil quality, CO₂ effluxes and vegetation

vii. Acknowledgements

Funding was provided by the French Agency for Environmental Transition (ADEME) via a PhD grant to Quentin Lambert and the project PIESO (agreement N°1405C0035). We thank Pierre Illac and Marine David (Total Quadran) for their logistic support. We further thank Mathilde Dionisi, Jean Bigotte, Xavier Fortuny, Lisa Foli for technical assistance in the lab and in the field.

viii. Abstract and Keywords

Abstract:

Solar energy is increasingly used to produce electricity in Europe, but the environmental impact of constructing and running solar parks (SP) is not yet well studied. Solar park construction requires partial vegetation removal and soil leveling. Additionally, solar panels may alter soil microclimate and functioning. In our study of three French Mediterranean solar parks, we analysed 1) effects of solar park construction on soil quality by comparing solar park soils with those of semi-natural land cover types (pinewood and shrubland) and abandoned croplands

(former vineyards) and 2) the effect of solar panels on soil microclimate, CO₂ effluxes and vegetation. We measured 21 soil properties of physical, chemical, and microbiological soil quality in one solar park and its surroundings to calculate integrated indicators of soil quality. We surveyed soil temperature and moisture, CO₂ effluxes and vegetation below and outside solar panels of three solar parks. Soil aggregate stability was reduced by SP construction resulting in a degradation of soil physical quality. Soil chemical quality and a general indicator of soil quality were lower in anthropogenic (SP and abandoned vineyards) than in semi-natural (pinewood and shrubland) land cover types. However, differences between abandoned vineyards representing the pre-construction land cover type and solar parks were not significant. Solar panels reduced the soil temperature by 10% and soil CO₂ effluxes by 50% but did not affect early successional plant communities. Long-term monitoring is needed to evaluate the effects of solar panels on vegetation.

Keywords: renewable energy, soil functions, land cover, microclimate, soil respiration, plant communities

ix. Main text

1. Introduction

The use of solar energy to produce electricity is increasingly common in Europe and requires large areas in order to be cost-effective (Murphy *et al.*, 2015 ; Ong *et al.*, 2013). Solar park construction involves clearing and grading the soil surface, burying of electric cables, vegetation removal and soil compaction increasing runoff and erosion. Grading, compaction, and erosion change the physical and chemical properties of the soil and thus reduce its quality. Since solar park construction destroys the vegetation and affects the soil, a careful analysis of the environmental impact of solar parks is needed (Armstrong *et al.*, 2016; Hernandez *et al.*, 2015).

Although soil quality is an important indicator of ecosystem functioning, the effect of solar park construction on soil quality has not yet been reported elsewhere. After the installation of solar panels, the vegetation is regularly mown or grazed limiting vegetation height to prevent shading of panels. The solar panels also change the microclimate such as temperature, humidity, solar radiation (Tanner *et al.*, 2020; Armstrong *et al.*, 2016). Such changes in microclimate may affect soil processes and plant communities under panels, in particular in the European Mediterranean with high solar irradiation compared to temperate regions.

Soil quality is “the capacity of a specific kind of soil to function, within natural or managed ecosystem boundaries, to sustain plant and animal productivity, maintain or enhance water and air quality and support human health and habitation” (Karlen *et al.*, 1997). Three soil quality indicator groups are commonly used: physical, chemical and biological soil properties (Bünemann *et al.*, 2018 ; Costantini *et al.*, 2016 ; Maurya *et al.*, 2020). Physical properties, such as bulk density and texture influence water holding capacity and plant communities by modulating root growth (Scarpore *et al.* 2019 ; Lampurlanés, Cantero-Martínez 2003). Chemical properties such as inorganic N, total C and pH control plant nutrition and microbiological activity. Biological indicators include the activity of decomposers such as invertebrates or microorganisms. These organisms control organic matter decomposition and nutrient cycling (Maurya *et al.*, 2020).

Velasquez *et al.* (2007) developed a single general indicator of soil quality (GISQ) that integrates a set of physical, chemical, and biological soil properties. Such soil properties are chosen and measured to evaluate multifaceted aspects of soil functions and further combined to calculate sub-indicators of physical quality, chemical fertility, and biological functioning. The GISQ combines the sub indicators to provide a global assessment of soil quality based on soil

ecosystem services and facilitates the comparison of soils between different sites/habitats. In a comparative study on four land use types, (Raiesi & Salek- Gilani, 2020) showed, using an adapted GISQ, that soil quality was 1.5 times lower in anthropogenic than in natural soils. Joimel *et al.* (2016) observed a decrease in soil physico-chemical quality along an anthropization gradient from forest to urban soils whereas Joimel *et al.* (2017) did not find any difference in biological quality of these soils. The construction of solar parks on natural and semi-natural land use types (e.g. forest, shrubland and abandoned vineyards) may reduce soil quality and affect ecosystem functions such as infiltration and storage of water, fertility and plant reestablishment, soil organic matter and nutrient cycling (Khare & Goyal, 2013; Romero-Díaz *et al.*, 2017; Rutgers *et al.*, 2009; Scarpore *et al.*, 2019; Yin *et al.*, 2020).

Plant communities and soil functioning may also be affected by changes in microclimate under solar panels. Solar panels reduce solar radiation, air humidity and soil temperatures, but in winter, soil temperatures are generally higher under panels (Armstrong *et al.*, 2016). Adeh *et al.* (2018) reported highest soil moisture and local heterogeneity of soil water conditions under solar panels. Such changes in microclimate may alter plant community composition and soil respiration that can be measured as CO₂ release. Mediterranean plant communities are dominated by heliophilous plants (Bagella & Caria, 2012). The reduction of solar radiation under solar panels may thus result in a plant community shift towards shade-tolerant species. Seed germination of Mediterranean species may be limited by light reduction (Gresta *et al.*, 2010), and the mortality of heliophilous plants increases in competition to shade-tolerant species. (Novara *et al.*, 2012; de Dato *et al.*, 2010). The change in air and soil microclimate under panels reduced the soil respiration under temperate oceanic climate (Armstrong *et al.*, 2016). Under Mediterranean climate with higher annual temperatures and summer drought, changes in microclimate under

solar panels may be higher resulting in a strong disturbance of seasonal soil respiration dynamics (González-Ubierna & Lai, 2019). Plant communities contribute to soil CO₂ release by respiration of roots and rhizosphere microorganisms (Raich & Tufekciogul, 2000) but also by changes in soil structure (Yang *et al.*, 2009; Zou *et al.*, 2005). Furthermore, plants are the principal carbon source of decomposer microorganisms (Wall *et al.*, 2012). Thus, solar panels may also change soil conditions indirectly by a shift in plant community composition since plants are very sensitive to change in microclimate.

The aims of our study were to assess 1) the effect of solar park construction on soil quality in comparing solar parks with semi-natural land cover types (pinewood and shrubland) and abandoned vineyards and 2) the effects of solar panels on soil microclimate, CO₂ effluxes and vegetation under Mediterranean climate. We expected that 1) solar park construction reduces physical, chemical, and biological soil quality, 2) solar panels change soil microclimate and plant community composition, and 3) solar panels change soil respiration according to the season.

2. Material and Methods

2.1. Study sites

Two studies were conducted in three solar parks (SP) located in Southern France (La Calade, Pouzols-Minervois and Roquefort des Corbières, Figure 1A.) with a distance of 10 to 30 km from one another (Table 1). These SP were constructed in 2011, 2014 and 2016, respectively, covered between 8.5 and 16 Ha and used ground-fixed photovoltaic (PV) systems carrying the solar panels at a fixed inclination. The solar panels are aligned to form rows (height of 0.6 m min and 2 m max) exposed to the South, with a gap of 4 m between rows. The study region is

characterized by typical Mediterranean climate with summer drought and mild, wet winters. The SP are mainly bordered by pinewood (*Pinus halepensis*), shrublands and vineyards. The soils of the SPs are characterized by carbonatic pedofeatures (i.e. fine calcareous silty clay soil).

2.2. Sampling designs

Study 1: effect of solar park construction

To study the effects of solar park construction on soil quality, four sampling plots (50 × 2 m) separated by 150 m were randomly chosen within the SP at Roquefort des Corbières (inter-rows between solar panels). In the surroundings of the SP, we randomly selected four sampling plots (100 m²) for each of the three dominant habitat types in the study region: pinewood, shrubland and abandoned vineyards (Figure 1B). These three land cover types are also representative of regional land use (forestry, ancient pastures and viticulture), and the studied SP was constructed on an abandoned vineyard. The plots were at least 200 m apart from each other and not more than 400 m from the SP. The pinewood is essentially composed of *Pinus halepensis* and *Quercus coccifera*. The shrublands are dominated by *Quercus coccifera*, *Pistacia lentiscus*, *Rosmarinus officinalis*, *Myrtus communis*, *Genista scorpius*, *Brachypodium retusum* and *Cistus monspelliensis*. The vineyards had been abandoned for five years before solar park construction and were dominated by grapevine, *Anisantha rubens*, *Dittrichia viscosa* and *Lysimachia arvensis*. In March 2016, ten soil samples were randomly collected (10 cm depth) within each plot, mixed to one composite sample per plot. Composite samples were sieved (mesh size: 2 mm) prior to analyses. An aliquot of samples was air-dried (1 week, 30 °C). For each sample, another aliquot was stored at 4 °C for microbial analyses.

Study 2: effect of solar panels

To study the effect of solar panels on soil respiration, temperature, and moisture and on plant communities, we randomly selected within each of the three SP four sampling plots (50 × 2 m) below the solar panels, both separated by at least 100 m, and four adjacent sampling plots (50 × 2 m) in the inter-rows between the solar panel.

2.3. Measurements of soil physico-chemical and microbiological quality

Soil physical properties

Water content (g.kg⁻¹) was determined after drying samples (24 hours, 105 °C). Water holding capacity (WHC) was analyzed according to Saetre (1998) but using a modified protocol. 10 g of dried soil were weighted in a PVC cylinder and saturated with water. WHC was defined as the water content remaining in the soil after 12 h (4 °C). The different soil fractions (*i.e.* sand, silt, clay) were determined using the Robinson's pipette method (Olmstead *et al.*, 1930) after organic matter removal by oxidation with H₂O₂ (30%, 48 hours). Bulk density (BD) was determined by measuring dried soil mass sampled in a Siegrist's cylinder. According to Huang *et al.*, (2004), a value of 2.65 g.cm⁻³ was assumed for real soil density (RD). Soil porosity was calculated using the following equation.

$$\text{Soil porosity} = 100 \times \frac{\text{RD} - \text{BD}}{\text{RD}} \text{ (Equation. 1)}$$

Mean weight diameter (MWD) of soil aggregates was measured according to Kemper and Rosenau (1986).

177
178 Soil chemical properties
179 The soil pH was measured in distilled water and KCL (1M) (Aubert, 1978). Total Carbon (TC)
180 and Nitrogen (TN) content were determined by combustion in an elemental analyzer CN FlashEA
181 1112 (ThermoFisher) (NF ISO 10694, NFISO 13878). Calcium carbonate (CaCO_3) content was
182 measured using a Bernard calcimeter (Müller & Gastner, 1971) and the percentage of C in
183 CaCO_3 (C- CaCO_3) was determined as: $\text{C-CaCO}_3 = 11.991 / 100 \times \text{CaCO}_3$. Inorganic nitrogen
184 (NH_4^+ and NO_3^-) was extracted in KCL solution (1 M) and analysed calorimetrically using the
185 nitroprusside-salicylate and nitrosalicylic acid method according to Mulvaney (1996) and Keeney
186 and Nelson (1983), respectively.

187
188 Soil microbiological properties
189 Microbial Biomass (MB) was measured using substrate induced respiration (SIR) rates
190 (Anderson and Domsch, 1978). Basal respiration was determined without adding glucose to
191 calculate the metabolic quotient $q\text{CO}_2$ (the ratio of basal respiration to microbial biomass), which
192 is a sensitive ecophysiological indicator of soil stress (Anderson, 2003). Three enzyme activities
193 (*i.e.* fluorescein diacetate hydrolase, phosphatase and tyrosinase) involved in carbon and
194 phosphorous cycles were assessed (n=3 per sample) to determine the catabolic potential of
195 microbial communities. Fluorescein diacetate hydrolase (FDase, U.g^{-1} dry weight) was measured
196 according to Green *et al.* (2006), phosphatase (U.g^{-1} dry weight) according to Tabatabai and
197 Bremner (1969) and the activity of tyrosinase ($\mu\text{mol.min}^{-1}.\text{g}^{-1}$ dry weight) according to Saiya-
198 Cork *et al.* (2002).

199

200 2.4. Measurements of solar panel effects on soil moisture, temperature and *in situ* respiration

201 Soil respiration, temperature and moisture were recorded in March and June 2017 in each

202 sampling plot of the study on solar panel effects. *In situ* CO₂ release (g.CO₂.m⁻¹.h⁻¹) from soils,

203 plant roots, soil organisms and chemical oxidation of C compounds was measured after removal

204 of aboveground vegetation, using a portable gas analyser (IRGA, EGM-4, PP-system). The

205 device was connected to a closed soil respiration chamber (SRC-1, PP systems Massachusetts,

206 USA). To prevent leakage of CO₂ when placing the chamber on the soil, a PVC tube (10 cm x 11

207 cm) was buried 1 cm deep into the soil prior to measurements. Soil temperature was recorded in a

208 depth of 7 cm using the soil temperature probe (STP-1, PP-system) connected to the

209 respirometer. Soil moisture was recorded on four points at 7 cm depth using a portable time-

210 domain reflectometry (TDR) device (Delta-T Devices, ML2 Theta Probes).

211

212 2.5. Measurements of solar panel effects on vegetation

213 In the sampling plots of the study on solar panel effects, vegetation surveys were carried out in

214 2016 and 2017. Three rectangular sub-plots of 10 m² (2 m × 5 m) were placed at the ends and the

215 center of each plot. Percentage cover of all occurring vascular plant species was estimated as the

216 vertical projection of aboveground plant organs. A ratio of shadow-tolerant (sciophilous) to hemi-

217 heliophilous and heliophilous plant species (Julve, 2020) was calculated .

218

219 2.6. Statistical analyses

We calculated a General Indicator of Soil Quality (GISQ) according to Velazquez *et al.* (2007). Information on 21 variables of physical, chemical, and microbiological soil properties was used to create three sub-indicators related to main soil functions: 1) physical properties that determine the infiltration and storage of water, 2) chemical properties that affect fertility and plant reestablishment in solar parks, 3) microbiological properties that drive soil organic matter decomposition and nutrient cycling. For each group of variables (physical, chemical and microbiological), a principal component analysis (PCA) with data scaled to unit of variance was run using “FactoMineR” (Husson *et al.*, 2020) and “Factoextra” (Kassambara & Mundt, 2020) packages. The Kaiser–Meyer–Olkin measure (>0.50). Bartlett’s test of sphericity (0.05) were used to test the sampling adequacy of variables included in the PCA. Variables with communality values <0.05 were removed and main components were identified using the latent root criterion (eigenvalues >1.0). A synthetic index of quality for each group of variables at a plot *i* (Iq_i) was calculated as the sum of *n* variables (v_i) multiplied by their respective weight (w_i) in the determination of axes 1 and 2 of the PCA (Equation 2.).

$$Iq_i = \sum_{i=1}^n w_i v_i \text{ (Equation 2.)}$$

The values of Iq_i were then reduced to a common range (0.1 to 1.0) using a homothetic transformation to obtain the sub-indicators of physical, chemical and microbiological soil quality (hereafter pSQ, cSQ and mSQ respectively, Equation 3.). In this equation, *a* is the maximal and *b* the minimal Iq value for the plot *i*.

$$p, c \text{ or } mSQ = 0.1 + \frac{Iq_i - b}{a - b} \times 0.9 \text{ (Equation 3.)}$$

Finally, a PCA was run with the 3 sub-indicators. The GISQ was obtained by summing the products of the respective contributions of variables to factors 1 and 2 by the % inertia explained by factors, respectively. Finally, the sum of these products gave the following formula for the GISQ (Equation 4.):

$$GISQ = 0.29 pSIq + 0.28 cSIq + 0.33 mSIq \text{ (Equation 4.)}$$

All data were analysed using R software (3.6.1, R Core Team, 2020). Effects of land cover type on soil physical, chemical and microbiological properties, sub-indicators of soil quality and GISQ were assessed using one way-analysis of variance (ANOVA). In the case of a significant land cover type effect, a Tukey HSD post hoc test was used to analyze differences between specific land cover types. To analyze the effect of solar panels on soil temperature, water content, CO₂ effluxes and vegetation, linear mixed-effect models (LMMs) (R package “lme4”) were applied including month and treatment (below vs outside panels) as fixed factors and solar park identity as random factor. Assumptions of ANOVA were checked by Shapiro-Wilks test for normality and by Levene test for homoscedasticity. When necessary, data were transformed using the “bestNormalize” package (Peterson, 2019) to meet these assumptions. Effects of solar panels on plant communities were visualized by non-metric multidimensional scaling (nMDS) based on the Bray-Curtis dissimilarity. Differences in plant community composition were tested using permutational multivariate analysis (PERMANOVA) in R package “vegan” (Oksanen *et al.*, 2019). A subsequent pairwise post hoc test was conducted to analyze differences between factor levels. False discovery rate (fdr) was used to correct for multiple comparisons.

3. Results

3.1. Effects of solar park construction on soil properties

Seven of the eight tested physical soil properties were significantly different between land cover types (Table 2). Among these seven properties, only two showed a significant difference between the two semi-natural (pinewood and shrubland) land cover types and the SP. Soil water content was 5.5% lower in the SP ($p < 0.01$) than in shrubland. The mean weight diameter of aggregates was twice as high in abandoned vineyards as in SP, and three times higher in pinewood and shrubland than in SP ($p < 0.001$). Organic carbon was about 2.5 times higher in semi-natural than in anthropogenic land cover types ($p < 0.001$). Sand and silt content, soil porosity and bulk density were significantly different between abandoned vineyards and pinewood (Table 2). Silt content and soil porosity were 1.4 and 1.3 times lower in abandoned vineyards than in pinewood, respectively. Sand content and BD were about 1.5 times higher in abandoned vineyards than in pinewood. Pinewood and shrubland showed similar physical properties without significant differences.

For most soil chemical properties, SP showed significant differences to pinewood and shrubland but not to abandoned vineyards (Table 2). Total carbon contents were 1.44 times higher in semi-natural land cover types than in anthropogenic land cover types ($p < 0.01$). Organic carbon contents were about 2.76 times higher in semi-natural land cover types than in anthropogenic land cover types ($p < 0.01$). Total nitrogen (TN) content showed the same pattern. TN was twice as high in pinewood and shrubland as in the SP and abandoned vineyards ($p < 0.001$). The water pH

284 ranged between 8.02 and 8.11 and showed a small but significant difference between shrubland
285 and abandoned vineyards.

286 Two microbiological properties were significantly different between land cover types (Table 2).
287 Land cover type had a significant influence on basal respiration ($p < 0.03$) being two times lower
288 in the SP and abandoned vineyards than in forest and shrubland. The FDAse activity was two
289 times higher in shrubland and pinewood than in the SP and abandoned vineyards. Microbial
290 biomass was twice as low (marginally significant) in SP and abandoned vineyards as in the semi-
291 natural land cover types.

292 3.2. Effects of solar park construction on soil quality

293 The first two axes of the PCA run on physical properties explained 80.56 % of the total variance
294 (see A.1.A). The semi-natural land cover types are separated from the anthropogenic soils along
295 the first axis. Silt, water content, water holding capacity and mean weight diameter of aggregates
296 had the highest score on the first PCA axis, while soil porosity was strongly correlated with the
297 second axis (see A.1.A). The highest physical quality index of 0.92 was measured in pinewood
298 and shrubland being two times higher than in abandoned vineyards ($p < 0.001$). The pSQ (Figure
299 2A) was two times and four times lower in SP than in the abandoned vineyards and semi-natural
300 land cover types, respectively ($p < 0.01$).

301 The first two axes of the PCA used to calculate the sub-indicator of soil chemical quality (cSQ)
302 explained 80.98% of the total variance (see A.1.B). The semi-natural land cover types are
303 separated from the anthropogenic soils along the first axis. Total carbon, organic carbon, total
304 nitrogen and ammonium were most correlated with the first axis and nitrate with the second axis

(see A.1.B). With a value of 0.18, the cSQ was four times lower in the SP and abandoned vineyards than in shrubland and pinewood ($p < 0.001$, Figure 2B).

The first two axes of the PCA used to calculate the sub-indicator of soil microbiological quality explained 80.28 % of the total variance (see A.1.C). Basal respiration, microbial biomass, FDAse and phosphatase were highly correlated with the first PCA axis, while the qCO_2 was correlated with the second one (see A.1.C). The mSQ was not significantly different between land cover types ($p = 0.74$) (Figure 2C).

The General Indicator of Soil Quality was four times lower in the SP and abandoned vineyards than in the pinewood and shrubland ($p < 0.001$).

3.3. Effects of solar panels on soil temperature, water content and *in situ* CO_2 effluxes.

Soil temperature and water content were significantly different between months ($p < 0.05$; Figure 3A and 3B). Solar panels significantly decreased soil temperature in March and June (Figure 3A) but did not affect soil water content ($p = 0.79$). Soil CO_2 effluxes did not change between months but were twice as high outside solar panels than below solar panels ($p < 0.001$).

3.3. Effects of solar panels on plant communities

Neither the species richness nor the total cover of plant community was significantly affected by the solar panels (Table 4). A marginally significant difference was detected for the ratio ‘Sciaphile: Heliophile plants’, being higher below than outside solar panels. The NMDS and PEMANOVA did not reveal any significant solar panel effect on plant community composition ($p = 0.3461$, Figure 4). However, community composition was significantly different between

the solar parks ($p < 0.001$). No significant difference was detected between observation years (data not shown).

4. Discussion

Solar Park (SP) construction reduced physical and chemical soil quality compared with semi-natural land cover types (forest and shrubland) but not biological soil quality. A change in soil temperature and CO₂ effluxes also demonstrated a negative solar panel effect on soil microclimate and functioning. However, in early stages of plant succession following solar park construction, plant community composition below and outside solar panels was not significantly different.

4.1 Effects of solar park construction on soil quality

Soil quality assessments require the measurement of a wide range of physical, chemical, and biological properties involving a high complexity of potential analyses (Maurya *et al.*, 2020). In this study, we assessed soil quality using a multi-proxy approach including 21 soil properties. The reduction of the number of variables using PCA to group these properties allows an integrated evaluation of soil quality based on their main functions, such as infiltration and storage of water, soil fertility, plant reestablishment and soil organic matter and nutrient cycling. We found that two of three integrated sub-indicators and the general indicator of soil quality were lower in SP than in the semi natural land cover types.

Among the physical soil properties, the aggregate MWD was 1.5 times lower in the SP than in the semi-natural land cover types (Table 2). A low MWD may result in a low aggregate stability.

Similarly, Kabir *et al.*, (2017) showed that the MWD decreases in anthropogenic soils associated with a degraded vegetation. In our study, soil levelling and vegetation removal prior to SP construction may have decreased soil organic matter (SOM) content reducing MWD. By binding colloids and stabilizing soil structure, SOM plays a key role in soil physical properties and nutrient cycling (Six *et al.*, 2004). Telak and Bogunovic (2020) showed a decrease in SOM and MWD in a vineyard of Croatia after intensive and frequent tillage. Such mechanical disturbance for many years may have affected soil structure of the vineyard on which the studied SP (Roquefort des Corbières) was constructed. Accordingly, overall physical soil quality of SP was lower than that of abandoned vineyards which was in turn lower than that of semi-natural land cover types. The physical soil quality index (Figure 2A) revealed that the construction of a SP increased the degradation of the physical soil quality of habitats already degraded by land management (abandoned vineyards). In particular, the stability of the soil, key factor of soil functioning, was lower in SP than in abandoned vineyards. Moreover, a lower SOM affects microbial activity and production of mucus resulting in a decrease of aggregate MWD and thus a soil more sensitive to erosion (Blavet *et al.*, 2009; Le Bissonnais *et al.*, 2018). The soil levelling and vegetation removal during SP construction may have increased surface runoff and soil erosion (Rabaia *et al.*, 2021).

In contrast to our expectations, the SP construction did neither increase soil bulk density nor decrease porosity compared to the abandoned vineyards (Table 2). In our study, soil bulk density was found to be lower in abandoned vineyards than in semi-natural land cover types. The past land management of vineyards may have degraded these properties before SP construction limiting effects of construction work. It is well known that the use of agricultural machinery considerably increases the soil bulk density (Dunjó *et al.*, 2003) . Under similar Mediterranean

climate and land use history in Spain, the bulk density was 30% higher in vineyards than in pinewood (Dunjó *et al.* 2003). Such changes in soil physical properties can reduce the infiltration and storage of water. To improve the process of water infiltration and the capacity of water storage, a decompaction of the soil surface may be useful in particular if soils were already compacted by previous land management. Such decompaction also facilitates the germination and establishment of plant species (Bassett *et al.*, 2005). The resulting improved revegetation may limit erosion and protect functions supported by soil physical quality (Beatty *et al.*, 2017).

Soil chemical properties, such as total and organic carbon and total nitrogen are directly linked to soil fertility and plant growth (Liu *et al.*, 2014). In our study, these properties showed lower values in anthropogenic soils than in semi-natural land cover types (Table 2). Joimel *et al.* (2016) obtained similar results along a gradient from natural to anthropogenic habitats in which total carbon and nitrogen decreased significantly from forests to vineyards. Soil disturbance such as soil tillage in vineyards or construction activities increases mineralization of organic matter reducing organic C and N (Brantley & Young, 2010). Accordingly, Choi *et al.* (2020) found a significantly lower C and N content in SP than in grassland soil. In our study, SP construction did not reduce neither C and N content nor soil chemical quality compared to degraded vineyard soil (Table 2). Consequently, the construction of SP did not further degrade the soil chemical properties. The low index of chemical quality (Figure 2B) is thus most likely related to the past vineyard management. Low C and N content suggest that nutrient cycling was lower in vineyards and SP soils and that several SP soil functions (carbon sequestration, soil structure, biological processes) were hampered compared to semi-natural soils. Appropriate site selection may limit such a loss of key soil functions by SP construction (Choi *et al.* 2020).

Soil microorganisms (*i.e.* bacteria and fungi) contribute actively to soil nutrient cycling (Schimel and Schaeffer, 2012). Thus, their genetic and physiologic characteristics are important indicators of ecosystem functioning such as nutrient cycling (Ranjard *et al.*, 2011). Microbiological soil properties showed differences between land cover types for fluorescein diacetate hydrolase (FDAse) activity and basal respiration. FDAse is an appropriate proxy to evaluate soil microbial activities because the ubiquitous esterase enzymes (e.g. lipase, protease) are involved in the hydrolysis of FDA (Schnürer & Rosswall, 1982). In our study, the FDAse was two times lower in anthropogenic soils (Table 2) suggesting a reduction of microbiological activity and nutrient cycling. Soil basal respiration showed the same pattern confirming a degradation of soil functions compared to semi-natural soils (Sparling 1997). Under a similar Mediterranean climate in Italy, basal respiration was three to four times lower in vineyards than in coniferous forests, mixed forests and shrublands (Marzaioli *et al.* (2010). The decrease in soil respiration may be related to the lower organic carbon and nitrogen content or the decrease in microbial biomass (Fernandes *et al.*, 2005). Previous vineyard management, in particular soil tillage, may have severely decreased microbial biomass resulting in low microbial activity. Five years abandonment were probably not sufficient to entirely reestablish microbial communities (Quintana *et al.*, 2020). Despite the reduction of microbial properties in abandoned vineyards and solar park soils, the microbiological soil quality index (mSiQ) was not significantly different between land cover types (Figure 2C). Other microbiological properties (BM and phosphatase) mainly contributing to the first PCA axis were hardly affected by land cover type and thus overruled significant response variables in mSiQ calculation.

As a consequence of lower physical and chemical sub-indicators, the general indicator of soil quality was about three times lower in SP compared to semi-natural land cover types (Figure 2D).

The key processes involved in degradation of soil quality were soil tillage, partial topsoil removal increasing erosion (Quinton *et al.*, 2010) and organic matter mineralization. Reduced organic matter content and increase of soil compaction decrease water holding capacity (Mujdeci *et al.*, 2017) and soil stability (Simansky *et al.*, 2013). Soil restoration by revegetation may improve soil physical and microbiological qualities of solar parks (Hernandez *et al.*, 2019). Revegetation may increase the stability of aggregates by increasing root biomass and the production of binding agents (Erktan *et al.*, 2016). The root exudates can also stimulate microbial biomass and activity and thus improve nutrient cycling (Eisenhauer *et al.*, 2010 ; Feng *et al.*, 2019).

4.2 Effects of solar panels on soil microclimate, vegetation, and *in-situ* CO₂ effluxes

Climatic conditions influence both soil microbial activities (Shao *et al.*, 2018) and plant communities (García- Fayos & Bochet, 2009). In our study, solar panels reduced the soil temperature in spring and in summer by about 5 °C (Figure 3A). Similarly, Armstrong *et al.* (2016) found a soil temperature reduction of 2 °C under solar panels during the summer (UK). The lower temperature under solar panels was the direct effect of shading although night temperatures may be higher (Tanner *et al.*, 2020). Solar panels also intercept precipitation, and Tanner *et al.* (2020) found a significant reduction in soil humidity under solar panels in the Mojave desert. However, we did not find any significant soil humidity difference under solar panels and outside (Figure 3B). The result may be explained by a lower evapotranspiration limiting humidity losses during drought periods as suggested by Tanner *et al.* (2020). Mediterranean vegetation is dominated by heliophilous plants (Bagella & Caria, 2012). So we expected that light reduction by solar panels strongly affects plant communities. However, we did not find a significant effect of solar panels on plant community composition and structure (Figure

440 4). The effect of solar panels on the ratio of shadow-tolerant to heliophilous species was only
441 marginally significant and no influence on plant species richness was detected (Table 3). Other
442 studies showed, however, a reduction in plant cover and species richness under solar panels
443 resulting from lower germination and higher mortality (Armstrong *et al.* 2016). Due to light
444 limitation, heliophilous plants are expected to be less competitive under solar panels (Chen *et al.*,
445 2004). However, protection against strong solar radiation and drought during Mediterranean
446 summer may have compensated for reduction of light and precipitation in our study. Accordingly,
447 Tanner *et al.* (2020) observed that in a desert plant richness was marginally greater under their
448 solar panels than in the control. In our study, the absence of a solar panel effect on the vegetation
449 (Figure 4, table 3) may also be explained by the low age of our solar parks limiting differential
450 effects on the vegetation. In early successional stages, the vegetation is dominated by ubiquitous
451 annual species germinating and developing under a great variety of environmental conditions.
452 Responses to the specific microclimate under solar panels may be slow in Mediterranean
453 vegetation types (Coiffait-Gombault *et al.*, 2012; Kinzig *et al.*, 1999). Long-term monitoring is
454 required to finally evaluate the influence of solar panels on plant communities.

455 A lower soil respiration is an indicator of lower litter decomposition and nutrient cycling
456 suggesting that these ecosystem functions may be reduced under solar panels (Incerti *et al.*,
457 2011). In our study, soil respiration was highly affected by solar panels (Figure 3C). Half of the
458 CO₂ fluxes from soils are produced by heterotrophic organisms (Bond-Lamberty *et al.*, 2004).
459 Heterotrophic respiration, but also plant respiration, are driven by environmental factors, mainly
460 temperature and moisture, explaining the strong influence of solar panels (Francioni *et al.*, 2020,
461 Moinet *et al.*, 2019). Accordingly, Armstrong *et al.* (2016) found a reduction of soil CO₂ effluxes
462 under solar panels in May and June. We already detected a reduced CO₂ efflux from March

onwards suggesting that the warm Mediterranean spring results in earlier temperature differences between soils under and outside solar panels compared to temperate oceanic climate. This early decrease in temperature under solar panels compared to controls outside solar panels may have reduced microbial activity and thus heterotrophic respiration. The reduction of CO₂ effluxes under solar panels may also be the result of light reduction reducing plant growth and root respiration.

4.4 Conclusions

Physical, chemical, and general soil quality indexes were lower in a solar park than in semi-natural land cover types. Clearing and grading the soil surface during solar park construction resulted in a strong degradation of soil physical quality, especially of soil structure, but did neither disturb soil chemical quality nor global quality compared to abandoned vineyards. These results suggest that solar parks should be preferably constructed on anthropogenic soils or that construction must be accompanied by environmental compensation measures and/or ecological restoration. At our Mediterranean study sites, solar panels reduced soil temperature from spring onwards. Neither light nor spring temperature reduction under solar panels altered plant communities in early stages of plant succession but reduced CO₂ effluxes. Our study demonstrated that solar park construction and solar panels changed soil quality and microclimate to a magnitude known to affect key soil functions. Long-term monitoring including different seasons is required to evaluate the final response of soil properties and vegetation to solar panels.

x. Reference

- Adeh EH, Selker JS, Higgins CW. 2018. Remarkable agrivoltaic influence on soil moisture, micrometeorology and water-use efficiency. *Plos One* **13**: e0203256. DOI: 10.1371/journal.pone.0203256
- Anderson JP, Domsch KH. 1978. A physiological method for the quantitative measurement of microbial biomass in soils. *Soil biology and biochemistry* **10**: 215–221
- Anderson T-H. 2003. Microbial eco-physiological indicators to asses soil quality. *Agriculture, Ecosystems & Environment* **98**: 285–293. DOI: 10.1016/S0167-8809(03)00088-4
- Armstrong A, Ostle NJ, Whitaker J. 2016. Solar park microclimate and vegetation management effects on grassland carbon cycling. *Environmental Research Letters* **11**: 074016. DOI: 10.1088/1748-9326/11/7/074016
- Aubert G. 1978. *Methodes d'analyses des sols: documents de travail tous droits reserves*. Centre régional de documentation pédagogique
- Bagella S, Caria MC. 2012. Diversity and ecological characteristics of vascular flora in Mediterranean temporary pools. *Comptes Rendus Biologies* **335**: 69–76. DOI: 10.1016/j.crvi.2011.10.005
- Beatty B, Macknick J, McCall J, Braus G, Buckner D. 2017. *Native Vegetation Performance under a Solar PV Array at the National Wind Technology Center*. National Renewable Energy Lab. (NREL), Golden, CO (United States). DOI: 10.2172/1357887
- Blavet D, De Noni G, Le Bissonnais Y, Leonard M, Maillo L, Laurent JY, Asseline J, Leprun JC, Arshad MA, Roose E. 2009. Effect of land use and management on the early stages of soil water erosion in French Mediterranean vineyards. *Soil and Tillage Research* **106**: 124–136. DOI: 10.1016/j.still.2009.04.010
- Bond-Lamberty B, Wang C, Gower ST. 2004. A global relationship between the heterotrophic and autotrophic components of soil respiration? *Global Change Biology* **10**: 1756–1766. DOI: 10.1111/j.1365-2486.2004.00816.x
- Botta GF, Rivero D, Tourn M, Melcon FB, Pozzolo O, Nardon G, Balbuena R, Becerra AT, Rosatto H, Stadler S. 2008. Soil compaction produced by tractor with radial and cross-ply tyres in two tillage regimes. *Soil and Tillage Research* **101**: 44–51. DOI: 10.1016/j.still.2008.06.001
- Brantley ST, Young DR. 2010. Shrub expansion stimulates soil C and N storage along a coastal soil chronosequence. *Global Change Biology* **16**: 2052–2061. DOI: 10.1111/j.1365-2486.2009.02129.x
- Bünemann EK, Bongiorno G, Bai Z, Creamer RE, De Deyn G, de Goede R, Flesskens L, Geissen V, Kuyper TW, Mäder P, Pulleman M, Sukkel W, van Groenigen JW, Brussaard L. 2018.

- 519 Soil quality – A critical review. *Soil Biology and Biochemistry* **120**: 105–125. DOI:
520 10.1016/j.soilbio.2018.01.030
- 521 Choi CS, Cagle AE, Macknick J, Bloom DE, Caplan JS, Ravi S. 2020. Effects of Revegetation on
522 Soil Physical and Chemical Properties in Solar Photovoltaic Infrastructure. *Frontiers in*
523 *Environmental Science* **8**. DOI: 10.3389/fenvs.2020.00140
- 524 Coiffait-Gombault C, Buisson E, Dutoit T. 2012. Are old Mediterranean grasslands resilient to
525 human disturbances? *Acta Oecologica* **43**: 86–94. DOI: 10.1016/j.actao.2012.04.011
- 526 Costantini EAC, Branquinho C, Nunes A, Schwilch G, Stavi I, Valdecantos A, Zucca C. 2016.
527 Soil indicators to assess the effectiveness of restoration strategies in dryland ecosystems.
528 *Solid Earth* **7**: 397–414. DOI: <https://doi.org/10.5194/se-7-397-2016>
- 529 de Dato GD, De Angelis P, Sirca C, Beier C. 2010. Impact of drought and increasing
530 temperatures on soil CO₂ emissions in a Mediterranean shrubland (gariga). *Plant and Soil*
531 **327**: 153–166. DOI: 10.1007/s11104-009-0041-y
- 532 Dunjó G, Pardini G, Gispert M. 2003. Land use change effects on abandoned terraced soils in a
533 Mediterranean catchment, NE Spain. *CATENA* **52**: 23–37. DOI: 10.1016/S0341-
534 8162(02)00148-0
- 535 Erktan A, Cécillon L, Graf F, Roumet C, Legout C, Rey F. 2016. Increase in soil aggregate
536 stability along a Mediterranean successional gradient in severely eroded gully bed
537 ecosystems: combined effects of soil, root traits and plant community characteristics.
538 *Plant and Soil* **398**: 121–137. DOI: 10.1007/s11104-015-2647-6
- 539 Feng C, Ma Y, Jin X, Wang Z, Ma Y, Fu S, Chen HYH. 2019. Soil enzyme activities increase
540 following restoration of degraded subtropical forests. *Geoderma* **351**: 180–187. DOI:
541 10.1016/j.geoderma.2019.05.006
- 542 Fernandes SAP, Bettiol W, Cerri CC. 2005. Effect of sewage sludge on microbial biomass, basal
543 respiration, metabolic quotient and soil enzymatic activity. *Applied Soil Ecology* **30**: 65–
544 77. DOI: 10.1016/j.apsoil.2004.03.008
- 545 Francioni M, Trozzo L, Toderi M, Baldoni N, Allegranza M, Tesei G, Kishimoto-Mo AW, Foresi
546 L, Santilocchi R, D'Ottavio P. 2020. Soil Respiration Dynamics in *Bromus erectus*-
547 Dominated Grasslands under Different Management Intensities. *Agriculture* **10**: 9. DOI:
548 10.3390/agriculture10010009
- 549 García- Fayos P, Bochet E. 2009. Indication of antagonistic interaction between climate change
550 and erosion on plant species richness and soil properties in semiarid Mediterranean
551 ecosystems. *Global Change Biology* **15**: 306–318. DOI: [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2486.2008.01738.x)
552 2486.2008.01738.x
- 553 González-Ubierna S, Lai R. 2019. Modelling the effects of climate factors on soil respiration
554 across Mediterranean ecosystems. *Journal of Arid Environments* **165**: 46–54. DOI:
555 10.1016/j.jaridenv.2019.02.008

- 556 Green VS, Stott DE, Diack M. 2006. Assay for fluorescein diacetate hydrolytic activity:
557 Optimization for soil samples. *Soil Biology and Biochemistry* **38**: 693–701. DOI:
558 10.1016/j.soilbio.2005.06.020
- 559 Gresta F, Cristaudo A, Onofri A, Restuccia A, Avola G. 2010. Germination response of four
560 pasture species to temperature, light, and post-harvest period. *Plant Biosystems* **144**: 849–
561 856. DOI: 10.1080/11263504.2010.523549
- 562 Hernandez RR, Armstrong A, Burney J, Ryan G, Moore-O’Leary K, Diédhiou I, Grodsky SM,
563 Saul-Gershenz L, Davis R, Macknick J, Mulvaney D, Heath GA, Easter SB, Hoffacker
564 MK, Allen MF, Kammen DM. 2019. Techno–ecological synergies of solar energy for
565 global sustainability. *Nature Sustainability* **2**: 560–568. DOI: 10.1038/s41893-019-0309-z
- 566 Hernandez RR, Hoffacker MK, Murphy-Mariscal ML, Wu GC, Allen MF. 2015. Solar energy
567 development impacts on land cover change and protected areas. *Proceedings of the*
568 *National Academy of Sciences* **112**: 13579–13584. DOI: 10.1073/pnas.1517656112
- 569 Huang Q, Akinremi OO, Rajan RS, Bullock P. 2004. Laboratory and field evaluation of five soil
570 water sensors. *Canadian Journal of Soil Science*. DOI: 10.4141/S03-097
- 571 Husson F, Josse J, Le S, Mazet J. 2020. *FactoMineR: Multivariate Exploratory Data Analysis*
572 *and Data Mining*.
- 573 Incerti G, Bonanomi G, Giannino F, Rutigliano FA, Piermatteo D, Castaldi S, De Marco A,
574 Fierro A, Fioretto A, Maggi O, Papa S, Persiani AM, Feoli E, De Santo AV, Mazzoleni S.
575 2011. Litter decomposition in Mediterranean ecosystems: Modelling the controlling role
576 of climatic conditions and litter quality. *Applied Soil Ecology* **49**: 148–157. DOI:
577 10.1016/j.apsoil.2011.06.004
- 578 Joimel S, Cortet J, Jolivet CC, Saby NPA, Chenot ED, Branchu P, Consalès JN, Lefort C, Morel
579 JL, Schwartz C. 2016. Soil biological quality: A new concept. *Science of The Total Environment* **545–546**: 40–47. DOI:
580 10.1016/j.scitotenv.2015.12.035
- 581 Joimel S, Schwartz C, Hedde M, Kiyota S, Krogh PH, Nahmani J, Pérès G, Vergnes A, Cortet J.
582 2017. Urban and industrial land uses have a higher soil biological quality than expected
583 from physicochemical quality. *Science of The Total Environment* **584–585**: 614–621.
584 DOI: 10.1016/j.scitotenv.2017.01.086
- 585 Julve P. 2020. Baseflor. Index botanique, écologique et chorologique de la flore de France.
- 586 Kabir EB, Bashari H, Mosaddeghi MR, Bassiri M. 2017. Soil aggregate stability and organic
587 matter as affected by land-use change in central Iran. *Archives of Agronomy and Soil*
588 *Science* **63**: 1823–1837. DOI: 10.1080/03650340.2017.1308492
- 589 Karlen DL, Mausbach MJ, Doran JW, Cline RG, Harris RF, Schuman GE. 1997. Soil Quality: A
590 Concept, Definition, and Framework for Evaluation (A Guest Editorial). *Soil Science*
591 *Society of America Journal* **61**: 4–10. DOI:
592 <https://doi.org/10.2136/sssaj1997.03615995006100010001x>

- 593 Kassambara A, Mundt F. 2020. *factoextra: Extract and Visualize the Results of Multivariate*
594 *Data Analyses*.
- 595 Keeney D, Nelson D. 1983. Nitrogen –inorganic forms. *Methods of Soil Analysis. Part 2.*
596 *Chemical and Microbiological Properties*. John Wiley & Sons, Ltd, i–xxiv. DOI:
597 10.2134/agronmonogr9.2.2ed.frontmatter
- 598 Kemper WD, Rosenau RC. 2018. Aggregate Stability and Size Distribution. *Methods of Soil*
599 *Analysis*. John Wiley & Sons, Ltd, 425–442. DOI: 10.2136/sssabookser5.1.2ed.c17
- 600 Khare P, Goyal DK. 2013. Effect of high and low rank char on soil quality and carbon
601 sequestration. *Ecological Engineering* **52**: 161–166. DOI: 10.1016/j.ecoleng.2012.12.101
- 602 Kinzig AP, Levin SA, Dushoff J, Pacala S. 1999. Limiting Similarity, Species Packing, and
603 System Stability for Hierarchical Competition- Colonization Models. *The American*
604 *Naturalist* **153**: 371–383. DOI: 10.1086/303182
- 605 Lampurlanés J, Cantero- Martínez C. 2003. Soil Bulk Density and Penetration Resistance under
606 Different Tillage and Crop Management Systems and Their Relationship with Barley
607 Root Growth. *Agronomy Journal* **95**: 526–536. DOI:
608 <https://doi.org/10.2134/agronj2003.5260>
- 609 Le Bissonnais Y, Prieto I, Roumet C, Nespoulous J, Metayer J, Huon S, Villatoro M, Stokes A.
610 2018. Soil aggregate stability in Mediterranean and tropical agro-ecosystems: effect of
611 plant roots and soil characteristics. *Plant and Soil* **424**: 303–317. DOI: 10.1007/s11104-
612 017-3423-6
- 613 Liu Z, Zhou W, Shen J, Li S, Ai C. 2014. Soil quality assessment of yellow clayey paddy soils
614 with different productivity. *Biology and Fertility of Soils* **50**: 537–548. DOI:
615 10.1007/s00374-013-0864-9
- 616 Marzaioli R, D’Ascoli R, De Pascale RA, Rutigliano FA. 2010. Soil quality in a Mediterranean
617 area of Southern Italy as related to different land use types. *Applied Soil Ecology* **44**: 205–
618 212. DOI: 10.1016/j.apsoil.2009.12.007
- 619 Maurya S, Abraham JS, Somasundaram S, Toteja R, Gupta R, Makhija S. 2020. Indicators for
620 assessment of soil quality: a mini-review. *Environmental Monitoring and Assessment* **192**:
621 604. DOI: 10.1007/s10661-020-08556-z
- 622 Moinet GYK, Midwood AJ, Hunt JE, Rumpel C, Millard P, Chabbi A. 2019. Grassland
623 Management Influences the Response of Soil Respiration to Drought. *Agronomy-Basel* **9**:
624 124. DOI: 10.3390/agronomy9030124
- 625 Mujdeci M, Simsek S, Uygur V. 2017. The Effects of Organic Amendments on Soil Water
626 Retention Characteristics Under Conventional Tillage System. *Fresenius Environmental*
627 *Bulletin* **26**: 4075–4081

- 628 Müller G, Gastner M. 1971. The 'Karbonat-Bombe', a simple device for the determination of
629 carbonate content in sediment, soils, and other materials. *Neues Jahrbuch für*
630 *Mineralogie-Monatshefte* **10**: 466–469
- 631 Mulvaney RL. 1996. Nitrogen—Inorganic Forms. *Methods of Soil Analysis*. John Wiley & Sons,
632 Ltd, 1123–1184. DOI: 10.2136/sssabookser5.3.c38
- 633 Murphy DJ, Horner RM, Clark CE. 2015. The impact of off-site land use energy intensity on the
634 overall life cycle land use energy intensity for utility-scale solar electricity generation
635 technologies. *Journal of Renewable and Sustainable Energy* **7**: 033116. DOI:
636 10.1063/1.4921650
- 637 Novara A, Armstrong A, Gristina L, Semple KT, Quinton JN. 2012. Effects of soil compaction,
638 rain exposure and their interaction on soil carbon dioxide emission. *Earth Surface*
639 *Processes and Landforms* **37**: 994–999. DOI: 10.1002/esp.3224
- 640 Olmstead LB, Alexander LT, Middleton HE. 1930. *A pipette method of mechanical analysis of*
641 *soils based on improved dispersion procedure*.
- 642 Ong S, Campbell C, Denholm P, Margolis R, Heath G. 2013. *Land-Use Requirements for Solar*
643 *Power Plants in the United States*. National Renewable Energy Lab. (NREL), Golden,
644 CO (United States). DOI: 10.2172/1086349
- 645 Quinton JN, Govers G, Van Oost K, Bardgett RD. 2010. The impact of agricultural soil erosion
646 on biogeochemical cycling. *Nature Geoscience* **3**: 311–314. DOI: 10.1038/ngeo838
- 647 Rabaia MKH, Abdelkareem MA, Sayed ET, Elsaid K, Chae K-J, Wilberforce T, Olabi AG. 2021.
648 Environmental impacts of solar energy systems: A review. *Science of The Total*
649 *Environment* **754**: 141989. DOI: 10.1016/j.scitotenv.2020.141989
- 650 Raich JW, Tufekciogul A. 2000. Vegetation and soil respiration: Correlations and controls.
651 *Biogeochemistry* **48**: 71–90. DOI: 10.1023/A:1006112000616
- 652 Raiesi F, Salek- Gilani S. 2020. Development of a soil quality index for characterizing effects of
653 land-use changes on degradation and ecological restoration of rangeland soils in a semi-
654 arid ecosystem. *Land Degradation & Development* **31**: 1533–1544. DOI:
655 <https://doi.org/10.1002/ldr.3553>
- 656 Ranjard L, Dequiedt S, Jolivet C, Saby NPA, Thioulouse J, Harmand J, Loisel P, Rapaport A,
657 Fall S, Simonet P, Joffre R, Bouré NC-P, Maron P-A, Mougél C, Martin MP, Toutain B,
658 Arrouays D, Lemanceau P. 2011. Biogeography of Soil Microbial Communities: A
659 Review and a Description of the Ongoing French National Initiative. *Sustainable*
660 *Agriculture Volume 2* 857–865. DOI: 10.1007/978-94-007-0394-0_37
- 661 Romero-Díaz A, Ruiz-Sinoga JD, Robledano-Aymerich F, Brevik EC, Cerdà A. 2017. Ecosystem
662 responses to land abandonment in Western Mediterranean Mountains. *CATENA* **149**:
663 824–835. DOI: 10.1016/j.catena.2016.08.013

664 Rutgers M, Schouten AJ, Bloem J, Eekeren NV, Goede RGMD, Akkerhuis GAJMJ, Wal AV der,
665 Mulder C, Brussaard L, Breure AM. 2009. Biological measurements in a nationwide soil
666 monitoring network. *European Journal of Soil Science* **60**: 820–832. DOI:
667 <https://doi.org/10.1111/j.1365-2389.2009.01163.x>

668 Saetre P. 1998. Decomposition, Microbial Community Structure, and Earthworm Effects Along a
669 Birch–Spruce Soil Gradient. *Ecology* **79**: 834–846. DOI: 10.1890/0012-
670 9658(1998)079[0834:DMCSAE]2.0.CO;2

671 Saiya-Cork KR, Sinsabaugh RL, Zak DR. 2002. The effects of long term nitrogen deposition on
672 extracellular enzyme activity in an *Acer saccharum* forest soil. *Soil Biology and*
673 *Biochemistry* **34**: 1309–1315. DOI: 10.1016/S0038-0717(02)00074-3

674 Scarpore FV, van Lier Q de J, de Camargo L, Pires RCM, Ruiz-Correa ST, Bezerra AHF, Gava
675 GJC, Dias CTS. 2019. Tillage effects on soil physical condition and root growth
676 associated with sugarcane water availability. *Soil & Tillage Research* **187**: 110–118. DOI:
677 10.1016/j.still.2018.12.005

678 Schimel J, Schaeffer SM. 2012. Microbial control over carbon cycling in soil. *Frontiers in*
679 *Microbiology* **3**. DOI: 10.3389/fmicb.2012.00348

680 Schnürer J, Rosswall T. 1982. Fluorescein Diacetate Hydrolysis as a Measure of Total Microbial
681 Activity in Soil and Litter. *Applied and Environmental Microbiology* **43**: 1256–1261

682 Shao P, He H, Zhang X, Xie H, Bao X, Liang C. 2018. Responses of microbial residues to
683 simulated climate change in a semiarid grassland. *Science of the Total Environment* **644**:
684 1286–1291. DOI: 10.1016/j.scitotenv.2018.07.055

685 Simansky V, Bajcan D, Ducsay L. 2013. The effect of organic matter on aggregation under
686 different soil management practices in a vineyard in an extremely humid year. *Catena*
687 **101**: 108–113. DOI: 10.1016/j.catena.2012.10.011

688 Six J, Bossuyt H, Degryze S, Denef K. 2004. A history of research on the link between
689 (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil and Tillage Research*
690 **79**: 7–31. DOI: 10.1016/j.still.2004.03.008

691 Sparling G. 1997. Soil microbial biomass, activity and nutrient cycling as indicators of soil
692 health. *Biological indicators of soil health* 97–119

693 Tabatabai MA, Bremner JM. 1969. Use of p-nitrophenyl phosphate for assay of soil phosphatase
694 activity. *Soil biology and biochemistry* **1**: 301–307

695 Tanner KE, Moore- O’Leary KA, Parker IM, Pavlik BM, Hernandez RR. 2020. Simulated solar
696 panels create altered microhabitats in desert landforms. *Ecosphere* **11**: e03089. DOI:
697 10.1002/ecs2.3089

698 Team RC. 2020. R: A language and environment for statistical computing.

- 699 Telak LJ, Bogunovic I. 2020. Tillage-induced impacts on the soil properties, soil water erosion,
700 and loss of nutrients in the vineyard (Central Croatia). *Journal of Central European*
701 *Agriculture* **21**: 589–601. DOI: 10.5513/JCEA01/21.3.2810
- 702 Wall DH, Bardgett RD, Behan-Pelletier V, Herrick JE, Jones TH, Six J, Strong DR, Putten WH
703 van der, Ritz K. 2012. *Soil Ecology and Ecosystem Services*. OUP Oxford
- 704 Yang L, Liu N, Ren H, Wang J. 2009. Facilitation by two exotic Acacia: *Acacia auriculiformis*
705 and *Acacia mangium* as nurse plants in South China. *Forest Ecology and Management*
706 **257**: 1786–1793. DOI: 10.1016/j.foreco.2009.01.033
- 707 Yin R, Kardol P, Thakur MP, Gruss I, Wu G-L, Eisenhauer N, Schaedler M. 2020. Soil
708 functional biodiversity and biological quality under threat: Intensive land use outweighs
709 climate change. *Soil Biology & Biochemistry* **147**: 107847. DOI:
710 10.1016/j.soilbio.2020.107847
- 711 Zou CB, Barnes PW, Archer S, McMurtry CR. 2005. Soil moisture redistribution as a mechanism
712 of facilitation in savanna tree–shrub clusters. *Oecologia* **145**: 32–40. DOI:
713 10.1007/s00442-005-0110-8
- 714

715 xi. Table

716 Table 1 Environmental and technical characteristics of solar parks.

	La Calade	Pouzols- Minervois	Roquefort des Corbières
Altitude (m)	77	100	62
Slope (%)	5	5	5
Temperature (annual mean, °C)	15.5	13.6	15.5.
Precipitation (annual mean, mm)	557	648	557
Sunshine duration (annual mean, hours)	2465	2119	2324
Soil texture	Loamy soil	Loamy soil	Loamy soil
Land cover	Shrubland	Abandoned	Abandoned
before construction		Vineyards and shrublandsQ	Vineyards
Commissioning of the SP	2011	2014	2016
Maximum power (Kwc)	5102	4950	11152
Solar panel soil localisation		Ground-fixed	
Solar panel cell technologies		Crystalline	
A Solar panel power (Wc)	180	250	260
Number of solar panels	26856	19800	46473
Area of the SP (ha)	8.5	10.7	16

717

Table 2: Soil physical, chemical, and microbiological properties in each type of land cover. Mean values with standard errors in parentheses. Different letters indicate significant differences between land cover types (significant P-values in bold). BD: bulk density, WC: water content; WHC : water holding capacity; MWD : mean weight diameter; OC : organic carbon; TC: Total carbon, TN: total nitrogen; BR : basal respiration; MB : microbial biomass; qCO₂ : metabolic quotient; FDAse : Fluorescein diacetate hydrolase.

Properties		Pinewood	Shrubland	Abandoned Vineyards	Solar park	p value
Physical	Sand (%)	35.13 (5.07) ^a	45.91 (8.96) ^{ab}	47.78 (4.34) ^b	42.68 (2.58) ^{ab}	0.04
	Silt (%)	47.32 (8.63) ^a	35.81 (6.67) ^{ab}	33.16 (2.97) ^b	35.97 (1.03) ^{ab}	0.03
	Clay (%)	17.54 (4.70)	18.28 (6.33)	19.06 (1.37)	21.35 (1.57)	0.59
	BD (g.cm ⁻³)	1.11 (0.18) ^a	1.13 (0.17) ^a	1.47 (0.10) ^b	1.32 (0.13) ^{ab}	0.02
	WC (%)	19.55 (3.78) ^{ab}	22.14 (3.15) ^a	16.67 (2.01) ^{ab}	16.36 (0.44) ^b	0.03
	WHC (%)	65.66 (12.46)	70.81 (15.30)	51.93 (13.03)	59.24 (9.06)	0.23
	Porosity (%)	58.15 (6.95) ^a	57.54 (6.38) ^a	44.69 (3.92) ^b	50.19 (4.88) ^{ab}	0.02
	MWD (mm)	2626.47 (260.47) ^a	2618.40 (223.73) ^a	1593.30 (194.09) ^b	879.22 (271.43) ^c	<0.001
Chemical	OC (%)	4.92 (0.62) ^a	4.13(0.70.64) ^a	1.46(0.19) ^b	1.61 (0.17) ^b	<0.001
	TC (%)	8.59 (0.35) ^a	8.07 (1.28) ^a	5.63 (0.46) ^b	5.93 (0.71) ^b	<0.001

TN (%)	0.22 (0.06) ^a	0.20 (0.07) ^a	0.09 (0.03) ^b	0.10 (0.02) ^b	<0.01
Soil pH in water	8.03 (0.04) ^{ab}	8.02 (0.03) ^b	8.11 (0.05) ^a	8.06 (0.04) ^{ab}	0.03
Soil pH in KCl	7.45 (0.04)	7.48 (0.06)	7.52 (0.06)	7.49 (0.05)	0.38
Nitrate ($\mu\text{g } \mu\text{g N-NO}_3^+ \cdot \text{g}^{-1}$)	1.34 (0.32)	1.06 (0.80)	0.72 (0.26)	1.71 (0.61)	0.12
Ammonium ($\mu\text{g N-NH}_4^+ \cdot \text{g}^{-1}$)	2.90 (0.17) ^a	2.92 (0.19) ^a	2.65 (0.14) ^b	2.65 (0.04) ^b	0.03
Microbiological	BR ($\mu\text{g C-CO}_2 \cdot \text{g}^{-1} \cdot \text{h}^{-1}$)	1.28 (0.33) ^a	1.31 (0.61) ^a	0.61 (0.36) ^b	0.03
	MB ($\mu\text{g C-CO}_2 \cdot \text{g}^{-1} \cdot \text{h}^{-1}$)	0.40 (0.10)	0.37 (0.15)	0.24 (0.12)	0.07
	qCO ₂	3.20 (0.26)	3.48 (0.56)	3.33 (2.68)	0.99
	FDAse ($\text{u} \cdot \text{g}^{-1}$)	0.0007 (0.0001) ^{ab}	0.0008 (0.0003) ^a	0.0004 (0.0001) ^b	0.02
	Tyrosinase ($\text{u} \cdot \text{g}^{-1}$)	0.0526 (0.0144)	0.0321 (0.0061)	0.0504 (0.0084)	0.11
	Phosphatase ($\text{u} \cdot \text{g}^{-1}$)	0.0067 (0.0004)	0.0058 (0.0015)	0.0046 (0.0024)	0.28

Table 3: Effects of solar panels on plant communities. Mean values with standard errors in parentheses.

Parameters	Below solar panels	Outside solar panels	p
Species richness	12.56 (5.92)	13.25 (5.49)	0.29
Total cover (%)	351.2 (165.12)	379.97 (183.31)	0.22
Hemi-heliophilous: Heliophilous ratio	0.12 (0.14)	0.10 (0.10)	0.09

xii. Figure legend:

Figure 1: (A) Position of the three solar parks with 1 : Pouzols-Minervois, 2 : Roquefort-des-Corbières 3: La Calade (A) (B) The Roquefort-des-Corbières solar park in detail with surrounding land use types and sampling points.

Figure 2: Sub-indicators of soil physical (A), chemical (B), and microbiological (C) quality and general soil quality indicator (D) for different types of land cover. Error bars are means \pm standard error. Different letters indicate significant differences ($p < 0.05$).

Figure 3: Soil temperature (A), water content (B) and CO₂ effluxes in March (black bars) and June (grey bars) below and outside solar panels. Error bars are means \pm standard error; different capital and lowercase letters indicate significant differences under and outside panels in March and June, respectively.

Figure 4: NMDS plot with polygons indicating the plant species composition of the three solar parks under (hatched polygon) and outside (solid polygon) solar panels, NMDS stress: 0.084. Different letters indicate significant differences after pairwise post hoc test ($p < 0.05$).





