



HAL
open science

Tectonics from topography : insights from high-resolution hillslope morphology analysis

Vincent Godard, Jean-Claude Hippolyte, Edward Cushing, Nicolas Espurt, Jules Fleury, Olivier Bellier, Vincent Ollivier, Team Aster

► To cite this version:

Vincent Godard, Jean-Claude Hippolyte, Edward Cushing, Nicolas Espurt, Jules Fleury, et al.. Tectonics from topography : insights from high-resolution hillslope morphology analysis. EGU General Assembly 2020, 2020, VIenna, Austria. 10.5194/egusphere-egu2020-2985 . hal-03541364

HAL Id: hal-03541364

<https://amu.hal.science/hal-03541364>

Submitted on 24 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

EGU2020-2985

<https://doi.org/10.5194/egusphere-egu2020-2985>

EGU General Assembly 2020

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



Tectonics from topography : insights from high-resolution hillslope morphology analysis

Vincent Godard¹, Jean-Claude Hippolyte¹, Edward Cushing², Nicolas Espurt¹, Jules Fleury¹, Olivier Bellier¹, Vincent Ollivier³, and Aster Team¹

¹Aix-Marseille Univ, CNRS, IRD, INRA, Coll France, CEREGE, Aix-en-Provence, France (godard@cerege.fr)

²PSE-ENV/SCAN, Institut de Radioprotection et de Surete Nucleaire, Fontenay-aux-Roses, France

³Aix-Marseille Univ, CNRS, Minist Culture, LAMPEA, Aix-en-Provence, France

The spatial distribution of tectonic uplift is often investigated using river profiles, as fluvial gradient is predicted to be strongly dependent on rock uplift. A similar response is expected from hillslope morphology which is also dependent on the relative base-level lowering rate. However, the reduced sensitivity of near-threshold hillslopes and the limited availability of high resolution topographic data has often been a major limitation for their use to investigate the distribution of tectonic activity.

Here we combined high-resolution analysis of hillslope morphology and cosmogenic nuclide-derived denudation rates to constrain the distribution of rock uplift across a thrust system at the Southwestern Alpine front in France. Our study is located in the Valensole Mio-Pliocene basin, where a series of folds and thrusts has deformed a plateau surface. Using a 1-m LiDAR Digital Terrain Model, we analyzed the morphology of hillslopes and extracted proxies for the relative spatial variations in denudation such as hilltop curvature (C_{HT}) and non-dimensional erosion rates (E^*). We observed systematic variation of these metrics coincident with the location of a major underlying thrust system identified by seismic surveys. Using a simple deformation model, the inversion of the E^* pattern allows us to constrain the geometry of a blind thrust, which is consistent with available geological and geophysical data.

We also sampled clasts from eroding conglomerate at several hilltop locations for ^{10}Be and ^{26}Al concentration measurements. Calculated hilltop denudation rates range from 40 to 120 mm/ka. These denudation rates appear to be correlated with E^* and C_{HT} extracted from the morphological analysis, and are used to derive absolute estimates for the fault slip rate. This high resolution hillslope analysis allows us to resolve short wavelength variations in rock uplift that would not be possible to unravel using commonly used channel profile-based methods.