

FP09.05 Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study

L. Mezquita, F. Barlesi, G. Ielsch, J. Merlio, D. Debieuvre, J. Mosser, C. Ricordel, L'Houcine Ouafik, I. Rouquette, I. Monnet, et al.

▶ To cite this version:

L. Mezquita, F. Barlesi, G. Ielsch, J. Merlio, D. Debieuvre, et al.. FP09.05 Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study. Journal of Thoracic Oncology, 2021, 16 (3), pp.S214. 10.1016/j.jtho.2021.01.122 . hal-03603784

HAL Id: hal-03603784 https://amu.hal.science/hal-03603784

Submitted on 10 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

and 11+ drinks per week (Table). Those with FH of LC were more likely to be obese regardless of demographic factors, and less likely to exercise 2+ times per week (Table). Those with FH of LC were more likely to have received 10+ imaging scans and to have been exposed to hydrocarbons compared to those without FH (Table). There were no reported differences in any other occupational or home exposures, including asbestos and radon in those with v. without FH of LC (Table). **Conclusion:** FH of LC is associated with higher-risk behaviors that increase the likelihood of developing both lung and other cancers. As such, FH can be used in cancer prevention and control programs to identify higher-risk patients and educate patients that FH of lung cancer is associated with risk-increasing behaviors that are nonetheless modifiable. **Keywords:** Family History, Risky Behaviors, Prevention

FP09.05

Driver Oncogenic Alterations and Indoor Radon in NSCLC Patients From the IFCT Biomarker Cohort: Bioradon France Study

L. Mezquita,¹ F. Barlesi,² G. Ielsch,³ J. Merlio,⁴ D. Debieuvre,⁵ J. Mosser,⁶ C. Ricordel,⁷ L. Ouafik,⁸ I. Rouquette,⁹ I. Monnet,¹⁰ F. Escande,¹¹ A. Langlais,¹² F. Morin,¹² C. Vignaud,³ E. Auclin,¹³ J.C. Benitez,² D. Planchard,² K. Leuraud,³ D. Laurier,³ B. Besse,² V. Westeel¹⁴ ¹Hospital Clinic, Medical Oncology Department, Barcelona; Medical Oncology Department, Gustave Roussy Cancer Campus, Villejuif, France, Spain/ES, ²Medical Oncology Department, Gustave Roussy Cancer Campus, Villejuif, France, Villejuif/FR, ³Institute for Radiological Protection and Nuclear Safety – Irsn, France, Paris/FR, ⁴University Hospital and University Bordeaux, Bordeaux/france, Bordeaux/FR, ⁵Groupe Hospitalier Régional Mulhouse Sud Alsace (Ghrmsa), Hôpital Emile Muller, Mulhouse Cedex/france, Mulhouse Cedex/FR, ⁶CHU Rennes, Rennes/france, Rennes/FR, ⁷Centre Hospitalier Régional Et Universitaire de Rennes, Rennes/france, Rennes/FR, ⁸Aix Marseille Univ, Aphm, Cnrs, Inp, Inst Neurophysiopathol, Marseille, France, Faculté de Médecine Secteur Nord, Service de Transfert D'Oncologie Biologique, Marseille, France, Marseille/france, Marseille/FR, ⁹Oncopole - CHU Toulouse, Toulouse/france, Toulousse/FR, ¹⁰Chi Creteil, Creteil/france, Creteil/FR, ¹¹Chru Lille, Lille/france, Lille/FR, ¹²Ifct, Paris/france, Paris/FR, ¹³Hopital Européen Georges Pompidou, Paris/FR, ¹⁴University Hospital of Besancon, Besançon/france, Besançon/FR

Introduction: Radon is a radioactive gas, considered as the leading cause of lung cancer in non-smokers. In a previous work, we studied the correlation between the estimation of radon concentration from the French Indoor Radon Map (*Institut de Radioprotection et de Sûreté Nucléaire*, IRSN) and the regional prevalence of driver alterations in a cohort of 116.424 NSCLC patients in France (*Mezquita et al, WCLC 2018*). The prevalence of driver oncogene alterations was significantly

higher in high-radon areas but clinical data were not available. We aim to confirm this hypothesis in an annotated database of NSCLC patients with matched molecular data available for adjustment (Barlesi et al, Lancet 2016). Methods: Retrospective assessment of patients with NSCLC tested for EGFR/BRAF/HER2/KRAS mutations (m) and ALK fusion from the 28 Platform led by the National Cancer Institute between Apr.2012 and Apr.2013, and included in the Biomarkers France dataset. We studied the association between the prevalence of driver oncogenic alterations (EGFR/ALK/BRAF/HER2/KRAS) and the radon mean concentration in the area where the patient was born according to the IRSN Map. Adjustment on age, gender and smoking was performed. Results: Out of 17664 patients, we analyzed 3994 with birthplace available: 63% males, 82% smokers, with a median age of 64 years [18-94]. Lung cancer tumors were mostly adenocarcinoma (76%), followed by other histologies (18%) and squamous (6%). By molecular alterations: 468 tumors harbor EGFRm (12%), 129 ALK (3%), 89 BRAFm (2%), 32 HER2m (1%), 985 KRASm (25%); 2273 wildtype or harbor other non-driver alterations (control; 57%). Adenocarcinoma histology (83.7% vs. 80.2%, p=0.0034), and nonsmoker habit (19.5% vs. 16.5%, p=0.0251) were more common in radon high-risk group (comparatively at the low risk group). The mean radon concentration by birthplace was 74.36 Bq/m³ ±53.28SD [range 16.6-622.3], and by molecular groups: EGFRm 72.49 Bq/m³ \pm 47.98 SD [16.6-461.4], ALK 80.24 Bq/m³ ±55.22SD [19.3-384.7], BRAFm 73.22 Bq/m³ ±47.86SD [19.3-319.3], *HER2m* 72.74 Bq/m³ ±39.51SD Bq/m³ [27.8-231.3], KRASm 71.79 Bq/m³ ±53.32SD [16.6-576.8] and control group 75.67 Bq/m³ \pm 54.5SD [16.6-622.3] (*p*=0.20). The prevalence of driver alterations was higher in high-radon areas (table 1; p=0.0472); but no significant difference was observed after adjustment on age, gender and smoking. Conclusion: We observed a higher prevalence of driver oncogenic alterations in NSCLC patients born in high radon areas; but no significant difference was observed after adjustment on age, gender and smoker. This study warrants further research on radon gas and driver oncogenes. Keywords: radon, lung cancer, driver oncogenic alterations

FP10 SMALL CELL LUNG CANCER/NET

FP10.01

Survival in Advanced SCLC: Projected Impact of Immuno-Oncology-Associated Durable Response on Population Health Gains in US



J. Gordon,¹ P. Mcewan,¹ R. Young,¹ P. Kirkness,¹ J. Penrod,² Y. Yuan^{2 1}Health Economics and Outcomes Research Ltd., Cardiff/GB, ²Bristol-Mysers Squibb, Princeton/NJ/US

			Low radon <50 Bq/m3	High radon ≥50 Bq/m3	P value	P value adjusted*
EGFR	Mutation	N (%)	155 (11.4%)	313 (13.2%)	0.1218	0.3024
	Control (1)	N (%)	1199 (88.6)	2059 (86.8)		
ALK	Fusion	N (%)	42 (3.4%)	87 (4.1%)	0.3272	0.4708
	Control (1)	N (%)	1199 (96.6)	2059 (95.9)		
BRAF	Mutation	N (%)	32 (2.6%)	57 (2.7%)	0.8708	0.9865
	Control (1)	N (%)	1199 (97.4)	2059 (97.3)		
HER2	Mutation	N (%)	7 (0.6%)	25 (1.2%)	0.0880	0.1781
	Control (1)	N (%)	1199 (99.4)	2059 (98.8)		
KRAS	Mutation	N (%)	375 (31.3%)	610 (29.6%)	0.3227	0.3478
	Control (2)	N (%)	824 (68.7)	1449 (70.4)		
DRIVER	Positive	N (%)	236 (16.4%)	482 (19%)	0.0472	0.2128
	Control (1)	N (%)	1199 (83.6)	2059 (81)		

(1) population no EGFRm, noBRAFm, no HER2m and no ALKr. (2) population no EGFRm, noBRAFm, no HER2m, no ALKr and no KRASm. *adjustment on age, gender and smoking