



**HAL**  
open science

# Phylogenetic relationship between the endosymbiont “*Candidatus RIESIA pediculicola*” and its human louse host

Alissa Hammoud, Meriem Louni, Dorothée Missé, Sébastien Cortaredona,  
Florence Fenollar, Oleg Mediannikov

► **To cite this version:**

Alissa Hammoud, Meriem Louni, Dorothée Missé, Sébastien Cortaredona, Florence Fenollar, et al..  
Phylogenetic relationship between the endosymbiont “*Candidatus RIESIA pediculicola*” and its human  
louse host. *Parasites & Vectors*, 2022, 15 (1), pp.73. 10.1186/s13071-022-05203-z . hal-03663771

**HAL Id: hal-03663771**

**<https://amu.hal.science/hal-03663771>**

Submitted on 13 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH

Open Access



# Phylogenetic relationship between the endosymbiont “*Candidatus RIESIA pediculicola*” and its human louse host

Alissa Hammoud<sup>1,2</sup>, Meriem Louni<sup>1,2,3\*</sup>, Dorothée Missé<sup>4</sup>, Sébastien Cortaredona<sup>1,5</sup>, Florence Fenollar<sup>1,5</sup> and Oleg Mediannikov<sup>1,2\*</sup>

## Abstract

**Background:** The human louse (*Pediculus humanus*) is a haematophagous ectoparasite that is intimately related to its host. It has been of great public health concern throughout human history. This louse has been classified into six divergent mitochondrial clades (A, D, B, F, C and E). As with all haematophagous lice, *P. humanus* directly depends on the presence of a bacterial symbiont, known as “*Candidatus RIESIA pediculicola*”, to complement their unbalanced diet. In this study, we evaluated the codivergence of human lice around the world and their endosymbiotic bacteria. Using molecular approaches, we targeted lice mitochondrial genes from the six diverged clades and *Candidatus RIESIA pediculicola* housekeeping genes.

**Methods:** The mitochondrial cytochrome *b* gene (*cytb*) of lice was selected for molecular analysis, with the aim to identify louse clade. In parallel, we developed four PCR primer pairs targeting three housekeeping genes of *Candidatus RIESIA pediculicola*: *ftsZ*, *groEL* and two regions of the *rpoB* gene (*rpoB-1* and *rpoB-2*).

**Results:** The endosymbiont phylogeny perfectly mirrored the host insect phylogeny using the *ftsZ* and *rpoB-2* genes, in addition to showing a significant co-phylogenetic congruence, suggesting a strict vertical transmission and a host-symbiont co-speciation following the evolutionary course of the human louse.

**Conclusion:** Our results unequivocally indicate that louse endosymbionts have experienced a similar co-evolutionary history and that the human louse clade can be determined by their endosymbiotic bacteria.

**Keywords:** Human lice, *Candidatus RIESIA pediculicola*, Co-evolution, Mitochondrial genes, Housekeeping genes, Phylogenetic analysis

## Background

The human louse, *Pediculus humanus* (Phthiraptera: Anoplura), has been a great public health concern throughout human history. It is one of the most ancient haematophagous ectoparasites and intimately related to its host [1]. Two ecotypes can infest *Homo sapiens*: *Pediculus humanus corporis* and *Pediculus humanus*

*capitis*. *Pediculus h. corporis*, known as the body louse, infests people living in poor hygienic conditions and is the principal vector of *Rickettsia prowazekii* (epidemic typhus agent), *Borrelia recurrentis* (relapsing fever agent), *Bartonella quintana* (trench fever agent) [2, 3] and probably *Yersinia pestis* (pandemic plague agent) [4]. *Pediculus h. capitis*, known as the head louse, has a widespread infestation rate regardless of the hygiene conditions [5, 6]. However, its capacity to be a potential vector of disease remains poorly understood [7]. The genetic diversity of human lice has been extensively

\*Correspondence: louni.meriem@yahoo.fr; oleguss1@gmail.com

<sup>1</sup> Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, 13005 Marseille, France

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

investigated using mitochondrial (mt) genes [cytochrome *b* (*cytb*), cytochrome oxidase subunit 1 (*cox1*) and 12S ribosomal RNA (*12S*)], allowing their classification into six divergent clades that may be grouped in three sister groups (A–D, B–F and C–E), each exhibiting a specific geographical distribution [8]. Clade A is the most prevalent, with a worldwide distribution, while clade D is only found in central Africa. Clade B is reported on all continents, while the genetically close clade F has a geographically restricted distribution and has been recently reported in South America. In addition, clade C has been identified in lice from central Africa and Asia, whereas clade E is reported in central and west Africa [6–9].

Like all haematophagous lice, *P. humanus* directly depends on the presence of endosymbiotic bacteria to supplement its unbalanced diet and metabolic integration [9]. Symbiosis is a common and widespread phenomenon that has a major effect on the biology of haematophagous arthropods. This interaction encompasses a whole range of possible symbiotic associations, ranging from strict parasitism to obligate mutualism [10]. Body and head lice host the same primary endosymbiotic bacterium, *Candidatus* *Riesia pediculicola* [9], which is classified in the genus *Candidatus* *Riesia* (class *Gamma-Proteobacteria*, family *Enterobacteriaceae*) [11, 12]. The symbionts are transovarially transmitted to progeny and are housed in the mycetome, which is localized on the ventral side of the louse midgut. Migration is initiated by a stimulating factor associated with the adult moult. In females, the symbionts migrate to the lateral oviducts at the beginning of oogenesis, and in male adults, the stomach disc will degenerate over time [13–15]. A genomic study of the human body louse and its primary endosymbiont has provided new insights into *Candidatus* *Riesia pediculicola* [11]. This bacterium has a small genome (<600 genes) containing a panel of genes encoding for the synthesis of essential B-group vitamins that are crucial to the host's diet [16, 17]. Indeed, the symbiont supplements the host's diet with thiamine, riboflavin, niacin, pantothenic acid, pyridoxine, biotin and folate (B1, B2, B3, B5, B6, B7 and B9 vitamins, respectively) [16–18]. Removal of the mycetome from *Pediculus* females leads to their death a few days later, as well as to the production of deformed eggs [10, 16]. It is a distinct possibility that the development of transgenic lineages of host and symbiont genes will facilitate our understanding of host–symbiont function and integration [19].

Human and chimpanzee lice (*Pediculus schaeffi*) diverged from a common ancestor, as did their human and chimpanzee hosts (*Pan troglodytes*), respectively, sometime between approximately five and seven million years ago. Interestingly, *Candidatus* *Riesia pediculicola* shared a common ancestor with the *P. schaeffi*

endosymbiont (*Candidatus* *Riesia pediculischaeffi*) roughly 5.4 million years ago [17, 18]. The evaluation of this co-evolutionary association between lice and their endosymbionts might provide new insights into human evolution [17]. Also, phylogenetic studies have shown a higher sequence similarity between clade A head and body lice endosymbionts than between clade A and clade B head lice endosymbionts. These results suggest that the endosymbionts co-evolved with their hosts' clades [17].

The aim of the present study was to establish a co-evolutionary relationship between the endosymbiotic bacteria and their human lice hosts from different clades using molecular approaches. We investigated mt genes from the six divergent clades of human lice and the housekeeping genes of *Candidatus* *Riesia pediculicola* in order to determine the louse clade using its endosymbiont bacteria population.

## Methods

### Lice selection and DNA extraction

From among the human lice collection of the IHU Mediterannée Infection laboratory, we selected 126 head and body lice that had been collected from around the world to perform the molecular study (Additional file 1: Table S1). These specimens had been collected in dry tubes, transported to our laboratory and frozen at – 20 °C for subsequent analysis. Prior to DNA extraction, each louse was externally decontaminated as previously reported [20]. Each specimen was cut longitudinally, and one half was frozen for subsequent analysis. DNA was extracted using a DNA extraction kit (QIAamp Tissue Kit; Qiagen, Hilden, Germany), using the EZ1 instrument in accordance with the manufacturer's protocol.

### Lice genotypic status

#### Haplogroup identification using qPCR assays

To identify the lice mt clades, DNA samples were subjected to clade-specific quantitative duplex real-time PCR (qPCR) targeting a portion of the *cytb* gene [21]. Each duplex is specific to clades A–D and B–C, noting that the B–C duplex also amplifies clade E lice, classified as a sub-sister clade within clade C lice. DNA amplification was performed as described previously [22]. Lice with a known clade were used as a positive control, while the master mixtures served as negative controls.

#### Haplotype identification using standard PCR and sequencing

Based on the qPCR results, we randomly selected 46 lice specimens encompassing the full range of clade diversity for phylogenetic analysis. DNA samples were subjected to standard PCR, targeting a 347-bp fragment of the *cytb* gene [23]. The final reaction volume (25 µl) consisted of 12.5 µl Amplitaq gold master mixes, 0.5 µM of each

primer, 5 µl DNA template and water. *cytb* amplification was performed in the Applied Biosystems 2720 Thermal Cycler (Applied Biosystems, Thermo Fisher Scientific, Waltham, MA, USA) with the following thermal cycling profile: 1 cycle at 95 °C for 15 min; then 40 cycles of 1 min at 95 °C, 30 s at 56 °C and 1 min at 72 °C; followed by a final extension step for 5 min at 72 °C. Successful amplification was validated by electrophoresis in an 1.5% agarose gel. Amplicons were then purified on NucleoFast 96 PCR plates (Macherey–Nagel EURL, Hoerdt, France) according to the manufacturer’s instructions and sequenced using the Big Dye Terminator Cycle Sequencing Kit (Thermo Fisher Scientific) with an Applied Biosystems automated sequencer.

**Candidatus Riesia pediculicola housekeeping gene analysis**

**Primer design**

In order to investigate the genotypic profile of the endosymbiotic bacteria, four standard PCR systems were designed targeting three *Candidatus Riesia pediculicola* housekeeping genes: *ftsZ*, *groEL* and two regions of *rpoB*. Four genomes of *Candidatus Riesia pediculicola* belonging to clade A and B lice deposited in the GenBank database (accession numbers CP012841, CP012843, CP012845, CP001085) [17, 18] were aligned using Muscle in MEGA7 software [24] and screened for conserved and discriminative genes.

Based on the variability in the available *Riesia* genomes, three housekeeping genes were selected as candidates for primer design. In order to find a suitable and conserved region for primers, two sequence fragments (approx. 100 bp) separated by a minimum of 500 bp for each gene were submitted to Primer3 software v. 0.4.0 (<http://primer3.ut.ee/>). The melting temperature of each primer was tested using the free online software programme Oligo Analyser 3.1 (<https://eu.idtdna.com/>

[calc/analyzer](#)) [25]. Designed primers (Table 1) were then tested in silico, using the NCBI BLAST nucleotide sequence similarity tool (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

**Housekeeping gene amplification**

Prior to endosymbiotic DNA amplification, the designed primers were tested against a panel of negative controls consisting of the DNA of various bacterial species and arthropods (Additional file 2: Table S2). Once validated, 73 samples were randomly selected from the 126 lice to amplify *ftsZ*, *groEL* and the two *rpoB* regions of the endosymbiont (Table 1). Standard PCRs were performed as described for *cytb*, and amplicons were visualized on a 1.5% agarose gel.

**Data analysis**

**Phylogenetic analysis**

In total, 28 lice specimens harbored common sequences for the *cytb* and the endosymbionte genes, but only 21 specimens had good quality sequences and were chosen for the phylogenetic analysis. The obtained *cytb* sequences were combined and compared with the worldwide *cytb* data previously reported and deposited in the GenBank database [8, 26]. Alignments were performed using MEGA7.0.26 software, and a maximum-likelihood (ML) tree was constructed using the Kimura2-parameter model under 1000 bootstrap replicates [27]. *Candidatus Riesia pediculicola* sequences were combined with data reported previously by Boyd et al. [16, 17] and were analysed as described for the *cytb* gene. Specifically, we used an Orthologous Average Nucleotide Identity Tool (OAT) [28] to define the overall similarities between the published *Candidatus Riesia pediculicola* genomes reported previously by Boyd et al. [17]. We also performed a Procrustes Approach to Cophylogenetic (PACo) analysis

**Table 1** Details of designed primers for *Candidatus Riesia pediculicola* gene amplification

Endosymbiont of <i>Pediculus humanus</i>	Target gene	Primer sequences (5’–3’)	Tm	Fragment length (bp)
<i>Candidatus Riesia pediculicola</i>	<i>ftsZ</i>	<i>ftsZ</i> -196F_GGGAATTTCTGATCTTCTCTGCG	56 °C	454
		<i>ftsZ</i> -650R_CTTTACTGGATGCTTTTGGYGC		
	<i>groEL</i>	<i>groEL</i> -560F_GATAGAGGTTATCTGTCTCC	50 °C	631
		<i>groEL</i> -1191R_GCAGCTCKAGTAGCATGTA		
	<i>rpoB</i>	<i>rpoB</i> -865F_ACCTGGTGATAAATCGTCTCC	54 °C	677
		<i>rpoB</i> -1542R_GAAAGAATCGTTCAGAAAGATCGG		
<i>rpoB</i> -2619F_CAGCCATCTCTCCGATTGAACG				
		<i>rpoB</i> -3085R_CGATGGAAAGCTAATTTGTTTCAGC	56 °C	466

Tm, Melting temperature

[29] using R [30] with 10,000 permutations, to test the dependence of the endosymbionte phylogeny on that of the louse clades by superimposing principal coordinates generated from the phylogenetic distance matrixes of *Candidatus* *Riesia pediculicola* and *P. humanus*.

**Results**

**Lice clade identification**

In this study, we collected 126 head lice and body lice worldwide (Additional file 1: Table S1) and selected 110 (87%) and 16 (13%), respectively, from these two populations for clade determination. qPCR assay of 125 of the 126 specimens showed that 76 (61%) of the lice belonged to the most prevalent clade, clade A, with various origins, including West Africa (17/76, 22.4%), Central Africa (10/76, 13.2%), North Africa (15/76, (9.7%), South Asia (24/76, 31.6%) and Europe (2/76, 2.6%), as well as reared lice used in the study as reference (8/76, 10.5%). Of the remaining lice specimens tested, 8/125 (6.3%) were genotyped as clade D; all these specimens were collected in the African continent, with six (75%) and two (25%) collected from Central and West Africa, respectively. Further, 39/125 (31.2%) of the analysed specimens belonged to clades C/E, with worldwide distribution: 21/39 (53.8%) and 11/39 (28.2%) specimens with Clades C/E were collected from West and Central Africa, respectively, and 7/39 (18%) were collected from Europe. Finally, only 2/125 (1.5%) samples belonged to clade B; these were collected in South Asia (Pakistan).

Standard PCR and sequencing of the *cytb* gene of 46/126 (36.5%) samples revealed that 12/46 (26.1%) belonged to clade A, 6/46 (13%) belonged to clade D, 2/46 (4.3%) belonged to clade B and only 1/46 (2.2%) belonged to the novel clade, clade F. In addition, 4/46 (8.7%) specimens belonged to clade C, while 21/46 (45.7%) of lice belonged to clade E (Table 2). Phylogenetical analysis was performed on 21 representative samples of all lice clades and origins.

**Phylogenetic relationship between the endosymbiont and their lice host**

To determine the phylogenetic profile of *Candidatus* *Riesia pediculicola*, we selected 73/126 (57.9%) lice specimens for DNA amplification and sequencing of three housekeeping genes (*ftsZ*, *groEL* and *rpoB*). For *ftsZ*, *groEL* and the first and second regions of the *rpoB*, we were able to generate all four fragments in 46/73 (63%) selected louse sequences: 65/73 (89%), 62/73 (84.9%), 60/73 (82.2%) and 57/73 (78.1%) sequences, respectively (Table 1). However, 27/46 (58.7%) of the endosymbionts were in clade A lice collected worldwide [7/27 (26%) North Africa; 16/27 (59.3%) South Asia; 4/27 (14.7%) Orlando strain]. Of these 46 sequences, four (8.6%) belonged to clade D lice and originated from West and Central Africa; two (4.4%) specimens were clade B endosymbionts collected in South Asia (Pakistan); one (2.2%) belonged to the novel clade F collected in Europe

**Table 2** Results of mitochondrial analysis and *Candidatus* *Riesia pediculicola* housekeeping genes of human lice samples

Origin	Country	Number of lice	Clade mitochondrial analysis		<i>Candidatus</i> <i>Riesia pediculicola</i> housekeeping gene analysis (number of lice–qPCR/clade type)			
			Number of lice-qPCR/clade type	Number of lice-standard PCR/clade type	<i>ftsZ</i> gene	<i>groEL</i> gene	<i>rpoB</i> -1 gene	<i>rpoB</i> -2 gene
West Africa	Senegal	10	3/A; 2/D; 5/C	2/A; 2/D; 4/E	2/A; 2/D; 4/E	2/A; 2/D; 3/E	2/D; 1/E	2/A 2/D
	Guinea	6	6/C	6/E	6/E	6/E	6/E	6/E
East Africa	Ethiopia	3	3/C	2/C	3/C	3/C	2/C	2/C
Central Africa	Gabon	1	1/C	1/C	1/C	1/C	1/C	1/C
	Democratic Republic of the Congo	6	4/D; 2/C	4/D; 2/E	4/D; 2/E	3/D; 2/E	3/D; 1/E	3/D; 2/E
	Congo Brazzaville	3	3/C	NT	NT	NT	3/C	NT
North Africa	Algeria	3 <sup>a</sup>	3/A	3/A	2/A	2/A	1/A	2/A
	Morocco	6	6/A	2/A	6/A	6/A	6/A	6/A
South Asia	India	13	13/A	2/A	11/A	10/A	12/A	11/A
	Pakistan	12	11/A; 2/B	2/A; 2/B	10/A; 2/B	11/A; 2/B	11/A; 2/B	11/A; 2/B
Europe	France	3	2/C	1/C; 1/F	2/C; 1/F	2/C; 1/F	1/C; 1/F	1/F
USA	Orlando strain	7 <sup>a</sup>	7/A	2/A	6/A	6/A	6/A	6/A

NT, Not tested

<sup>a</sup> Body lice



(France); and four (8.7%) were clade C endosymbionts collected in East Africa. Finally, for its sister clade E, 8/46 (17.4%) specimens were collected from West and Central Africa (Table 2).

Maximum likelihood phylogenetic trees were constructed for the housekeeping genes, including 21 samples already analysed for the *cytb* gene. Interestingly, louse endosymbionts from each mt host clade clustered in a separate group (Fig. 1a, d), while the phylogeny based on the endosymbiotic *ftsZ* and the second region of *rpoB* (*rpoB-2*) genes followed that of *cytb*, which is not the case for the *groEL* and first region of *rpoB* (*rpoB-1*) genes (Fig. 1b, c). To further investigate the present congruent phylogenies, we proceeded to concatenate *ftsZ* and *rpoB-2* sequences (848 bp of the final fragment), and a phylogenetic tree was generated and compared to that of *cytb*. We noted that the concatenated 848-bp fragment of different *Candidatus* *Riesia pediculicola* harboured a higher bootstrap value and grouped in clades which were almost perfect when compared to the one gene-based tree. In addition, the clustering of sister clades was also visible for the B–F and C–E clades, mimicking the mt phylogeny of *P. humanus* (Fig. 2). Furthermore, PACo analysis showed a significant co-phylogenetic congruence between *Candidatus* *Riesia pediculicola* and *P. humanus* phylogenies across all clades. These results indicated a sum of squared residuals ( $m^2$ ) of 0.38 ( $P < 0.001$ ) (Fig. 3).


The degree of genomic similarity of the deposited endosymbiont genomes of clade A and B lice [17] strongly support our results. A higher similarity (99.97%) was observed between *Candidatus* *Riesia pediculicola* of clade A head and body lice, with a lower similarity (97.85%) observed between clade A and B *P. h. capitis* specimens (Additional file 3: Figure S1).

## Discussion


In this study, we highlighted the codivergence of the endosymbiont *Candidatus* *Riesia pediculicola* and the mt clades of their *P. humanus* host. Louse endosymbionts were first reported in the 1920s, and they were subsequently successfully characterized in various histological, embryonic, experimental and nutritional studies [10]. This obligate intracellular primary endosymbiont is fully and uniquely attached to its host [16]. Its major role resides in providing essential B vitamins that are crucial for the survival of the louse and is lacking in the delivered blood meals [12, 31]. This bacterium has never been isolated in pure axenic culture, but studies employing molecular techniques suggest a polyphyletic origin [32]. Endosymbiotic microorganisms are generally associated with diverse arthropods, such as *Buchnera* of aphids, *Carsonella* of psyllids, *Portiera* of whiteflies, *Sulcia* of many homopterans, *Baumannia* of sharpshooters,

*Blochmannia* of carpenter ants and *Nardonella* of weevils, as well as with bloodsucking insects, such as *Wigglesworthia* of tsetse flies [33, 34]. Their endosymbiont phylogeny generally mirrors the host phylogeny, indicating a stable and intimate host–symbiont association over time. This is also the case of the *Nycteribiidae* family of bat flies, which is involved in ectoparasitic blood-feeding on bats: phylogeny of the endosymbiont *Candidatus* *Aschnera chinzeii* clades shows co-speciation over the evolutionary course of the *Nycteribiidae* family [35]. Allen et al. dated the divergence between the *Riesia* and *Arsenophonus* (endosymbiont of *Lipoptena cervi*) clades at 13–25 million years ago [36]. Furthermore, 16S rDNA sequences confirmed a strict coevolution between the endosymbionts of *Anoplura* (i.e. *Haematopinus* sp. of ungulates, *Solenoptes* sp. of cattle, *Pediculus* sp. of hominids and *Polyplax* sp. of rodents) and *Rhyncophthirinan* (*Haematomyzus* sp. of Asian elephants) genera, with the endosymbiont sequences forming five separate monophyletic branches, each composed from only one louse genus [37]. Finally, the gene content of the *Columbicola wolffhuegeli* (endosymbiont of Pied Imperial Pigeon louse) was so similar to the gene content of the *Candidatus* *Riesia pediculicola*, based on the phylogenetic tree, that the human head louse and *C. wolffhuegeli* acquired their endosymbionts independently [38]. These findings suggest that every louse group has its own endosymbiont.


We demonstrated the evolutionary phylogenetic relationship that links *Candidatus* *Riesia pediculicola* to their host mt clades. Individual phylogenetic trees based on *ftsZ*, *rpoB-2* and their concatenated genes enabled an identical endosymbiotic clusterization depending on human lice clades. The endosymbiont phylogeny perfectly mirrored the host insect phylogeny, suggesting strict vertical transmission and host–symbiont co-speciation during the evolutionary course of the human louse. These data will allow the classification of human louse clades through their endosymbiotic bacteria based on the *ftsZ* and *rpoB-2* genes. The slight discordance of *rpoB-1*- and, in particular, *groEL*-based trees (Fig. 1b, c) with host *cytb*-based topologies may be due to the lack of sufficient informative characters. The concatenated tree of the four gene fragment analysed, however, is in perfect agreement with the host tree (Fig. 2). We further investigated this phenomenon by constructing a ML phylogenetic tree of human, gorilla and chimpanzee lice and their endosymbionts; however, the paucity of available sequences did not allow us to conclude if there is a probable gene tree conflict (Additional file 4: Figure S2). While we observed a clear characterization of the lice clades for the first region of the *rpoB-1* gene, differences regarding clades E and C were observed. The slightly incongruent trees obtained within the same gene can be explained by a difference in



**Fig. 1** ML phylogenetic trees of the *Pediculus humanus* endosymbiont housekeeping genes, distributed in six divergent clades. **a** ML tree of a 454-bp fragment of the *ftsZ* gene, **b** ML tree of a 631-bp fragment of the *groEL* gene, **c, d** ML tree of the 677- and 466-bp fragments of the *rpoB* gene (*rpoB*-1 and *rpoB*-2, respectively). Phylogenetic inference was conducted in MEGA 7 using the ML method under the Kimura 2-parameter with 1000 bootstrap replicates. Abbreviations: ML, Maximum likelihood



**Fig. 2** Phylogenetic tree of the *cytb* mitochondrial gene of *P. humanus* (a) projected with the concatenated *ftsZ* and *rpoB* (*rpoB-2*) genes (848 bp) of *Candidatus Riesia pediculicola* (b), showing the relationship and co-evolution of the endosymbiont dependently on the mitochondrial clades of their host. Phylogenetic inference was conducted in MEGA 7 using the ML method under the Kimura 2-parameter with 1000 bootstrap replicates



**Fig. 3** Procrustes superimposition plot of *Candidatus Riesia pediculicola* and *P. humanus*. The extended principal coordinate matrices (X and Y) are centred by mean column vectors and subjected to Procrustes analysis. The configuration of the endosymbiont (dots) has been rotated and scaled to fit the lice configuration (arrowheads). Asterisk indicates lice samples, underlining indicates body lice. Abbreviations: r, Reference sequences from GenBank; see phylogenetic trees in Figs. 1 and 2 for sample abbreviations



the mutation level between these two regions. Our findings need further investigation by sequencing and analysing the endosymbionts' whole genomes within all human lice clades to better establish the evolutionary time courses within their hosts.

## Conclusion

Based on phylogenetic and genomic analyses, we have highlighted the co-evolutionary relationship between *Candidatus* *Riesia pediculicola* and their host mt clades. Our results unequivocally indicate that louse endosymbionts have experienced a similar co-evolutionary history and that human lice clades can be determined by their endosymbiotic bacteria based on their *ftsZ* and *rpoB* housekeeping genes. In future studies, further robust phylogenetic examination of all endosymbiont genome clades will be fundamental to a better understanding of the evolution of *Candidatus* *Riesia pediculicola* depending on the mt divergence of their hosts. However, it is crucial to isolate and identify this bacterium in order to evaluate the effectiveness of a drug treatment targeting the louse endosymbiont.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13071-022-05203-z>.

**Additional file 1: Table S1.** Results of mitochondrial analysis of human lice samples and *Candidatus* *Riesia pediculicola* housekeeping gene analysis.

**Additional file 2: Table S2.** Negative control DNA used for sensitivity and specificity determination of designed oligonucleotides.

**Additional file 3: Figure S1.** Heatmap generated according to OrthoANI values calculated using Orthologous Average Nucleotide Identity Tool (OAT) software (<https://www.ezbiocloud.net/tools/orthoani>) to measure the overall similarity between the genomes of *Candidatus* *Riesia* sp. strains and other related members of the *Riesia* genus. *Abbreviations:* BL, body lice; HapA, *Candidatus* *Riesia pediculicola* from *P. humanus* clade A; HapB, *Candidatus* *Riesia pediculicola* from *P. humanus* clade B; HL, head lice

**Additional file 4: Figure S2.** ML phylogenetic tree of primates lice (left) and their endosymbionts "*Ca. Riesia* sp." (right). Phylogenetic inference was conducted in MEGA 7 using the maximum likelihood method under the Kimura 2-parameter with 1000 bootstrap replicates.

## Acknowledgements

We thank the IHU Méditerranée-Infection for its support.

## Authors' contributions

AH, ML and OM designed the study. AH performed the experiments. AH, ML, SC and OM analysed the data. AH and ML wrote the manuscript. AH, ML, DM, FF and OM edited and revised the manuscript. All authors read and approved the final manuscript.

## Funding

This study was supported by the Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, the National Research Agency under the "Investissements d'avenir" programme, reference ANR-10-IHU-03, the Région Provence-Alpes-Côte d'Azur and European ERDF PRIM1 funding.

## Availability of data and materials

The sequences of the *Candidatus* *Riesia pediculicola* housekeeping genes referred to in this article and the attributed GenBank accession numbers are as follows. *ftsZ* gene: clade A (Orlando-MW588459, Senegal-MW588463), clade D (Congo-MW588466, Senegal-MW588464), clade B (Pakistan-MW588462), clade F (France-MW588460), clade C (France-MW58846) and clade E (Congo-MW588465, Guinea-MW588458). The second region of the endosymbiont *rpoB* gene: clade A (Orlando-MW588485, India-MW588486, Algeria-MW588487), clade D (Senegal-MW588488, Congo-MW588489); clade B (Pakistan-MW588491), clade F (France-MW588490), clade C (Ethiopia-MW588492) and clade E (Congo-MW588494, Guinea-MW588493).

## Declarations

### Competing interests

The authors declare that they have no competing interests. The funders had no role in the design of the study; in the collection, analysis, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

### Author details

<sup>1</sup>Institut Hospitalo-Universitaire (IHU) Méditerranée Infection, 13005 Marseille, France. <sup>2</sup>Institut de Recherche pour le Développement (IRD), Assistance Publique-Hôpitaux de Marseille (APHM), Microbes Evolution Phylogeny and Infections (MEPHI), Aix-Marseille University, 13005 Marseille, France. <sup>3</sup>Department of Biology, Faculty of Sciences, M'Hamed Bougara University, 35000 Boumerdès, Algeria. <sup>4</sup>IRD, CNRS, MIVEGEC, Université Montpellier, 34394 Montpellier, France. <sup>5</sup>Institut de Recherche pour le Développement (IRD), Assistance Publique-Hôpitaux de Marseille (APHM), Vectors Infections Tropicales and Mediterranean (VITROME), Aix-Marseille University, 13005 Marseille, France.

Received: 10 September 2021 Accepted: 14 February 2022

Published online: 05 March 2022

## References

1. Reed DL, Clayton DH. Genetic analysis of lice supports direct contact between modern and archaic humans. *PLoS Biol.* 2004;2:e340.
2. Raoult D, Roux V. The body louse as a vector of reemerging human diseases. *Clin Infect Dis.* 1999;29:888–911.
3. Light JE, Toups MA, Reed DL. What's in a name: the taxonomic status of human head and body lice. *Mol Phylogenet Evol.* 2008;47:1203–16.
4. Barbieri R, Drancourt M, Raoult D. The role of louse-transmitted diseases in historical plague pandemics. *Lancet Infect Dis.* 2021;21:e17–25.
5. Amanzougaghene N, Fenollar F, Raoult D, Mediannikov O. Where are we with human lice? A review of the current state of knowledge. *Front Cell Infect Microbiol.* 2020;9:474.
6. Light JE, Allen JM, Long LM, Carter TE, Barrow L, Suren G, et al. Geographic distributions and origins of human head lice (*Pediculus humanus capitis*) based on mitochondrial data. *J Parasitol.* 2008;94:1275–81.
7. Ashfaq M, Prosser S, Nasir S, Masood M, Ratnasingham S, Hebert PDN. High diversity and rapid diversification in the head louse, *Pediculus humanus* (Pediculidae: Phthiraptera). *Sci Rep.* 2015;5:14188.
8. Amanzougaghene N, Fenollar F, Davoust B, Djossou F, Ashfaq M, Bitam I, et al. Mitochondrial diversity and phylogeographic analysis of *Pediculus humanus* reveals a new Amazonian clade "F". *Infect Genet Evol.* 2019;2019:1–8.
9. Boyd BM, Reed DL. Taxonomy of lice and their endosymbiotic bacteria in the post-genomic era. *Clin Microbiol Infect.* 2012;18:324–31.
10. Sasser D, Epis S, Pajoro M, Bandi C. Microbial symbiosis and the control of vector-borne pathogens in tsetse flies, human lice, and triatomine bugs. *Pathog Glob Health.* 2013;107:285–92.
11. Perotti MA, Allen JM, Reed DL, Braig HR, Perotti MA, Allen JM, et al. Host-symbiont interactions of the primary endosymbiont of human head and body lice. *FASEB J.* 2007;21:1058–66.
12. Sasaki-Fukatsu K, Koga R, Nikoh N, Yoshizawa K, Kasai S, Mihara M, et al. Symbiotic bacteria associated with stomach discs of human lice. *AEM.* 2006;72:7349–52.

13. Eberle MW, McLean DL. Initiation and orientation of the symbiote migration in the human body louse *Pediculus humanus* L. *J Insect Physiol.* 1982;28:417–22.
14. Eberle MW, McLean DL. Observation of symbiote migration in human body lice with scanning and transmission electron microscopy. *Can J Microbiol.* 1983;29:755–62.
15. Buchner P. Endosymbiosis of animals with plant microorganisms. Interscience publishers. 1965; 909
16. Kirkness EF, Haas BJ, Sun W, Braig HR, Perotti MA, Clark JM, et al. Genome sequences of the human body louse and its primary endosymbiont provide insights into the permanent parasitic lifestyle. *Proc Natl Acad Sci USA.* 2010;107:12168–73.
17. Boyd BM, Allen JM, Nguyen N-P, Vachaspati P, Quicksall ZS, Warnow T, et al. Primates, lice and bacteria: speciation and genome evolution in the symbionts of hominid lice. *Mol Biol Evol.* 2017;34:1743–57.
18. Boyd BM, Allen JM, de Crécy-Lagard V, Reed DL. Genome sequence of *Candidatus Riesa pediculischaeffi*, endosymbiont of chimpanzee lice, and genomic comparison of recently acquired endosymbionts from human and chimpanzee lice. *G3 (Bethesda).* 2014;4:2189–95.
19. Gerardo NM, Wilson ACC. The power of paired genomes. *Mol Ecol.* 2011;20:2038–40.
20. La Scola B, Raoult D. *Acinetobacter baumannii* in human body louse. *Emerg Infect Dis.* 2004;10:1671–3.
21. Amanzougaghene N, Akiana J, Ndombe GM, Davoust B, Nsana NS, Parra H-J, et al. Head lice of Pygmies reveal the presence of relapsing fever *Borrelia* in the Republic of Congo. *PLoS Negl Trop Dis.* 2016;10:e0005142
22. Amanzougaghene N, Fenollar F, Sangaré AK, Sissoko MS, Doumbo OK, Raoult D, et al. Detection of bacterial pathogens including potential new species in human head lice from Mali. *PLoS ONE.* 2017;12:e0184621.
23. Li W, Ortiz G, Fournier P-E, Gimenez G, Reed DL, Pittendrigh B, et al. Genotyping of human lice suggests multiple emergences of body lice from local head louse populations. *PLoS Negl Trop Dis.* 2010;4:e641.
24. Kumar S, Stecher G, Tamura K. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 2016;33:1870–4.
25. Owczarzy R, Tataurov AV, Wu Y, Manthey JA, McQuisten KA, Almbrazi HG, et al. IDT SciTools: a suite for analysis and design of nucleic acid oligomers. *Nucleic Acids Res.* 2008;36:W163–9.
26. Hammoud A, Louni M, Baldé MC, Beavogui AH, Gautret P, Raoult D, et al. Molecular characterization and genetic diversity of haplogroup E human lice in Guinea, West Africa. *Microorganisms.* 2021;9:257.
27. Efron B, Tibshirani R. Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy. *Statist Sci.* 1986;55:54–75.
28. Lee I, Kim YO, Park SC, Chun J. OrthoANI: an improved algorithm and software for calculating average nucleotide identity. *Int J Syst Evol Microbiol.* 2016;66:1100–3.
29. Balbuena JA, Míguez-Lozano R, Blasco-Costa I, Moreau CS. PACo: a novel Procrustes application to cophylogenetic analysis. *PLoS ONE.* 2013;8:e61048.
30. Hutchinson MC, Cagua EF, Balbuena JA, Stouffer DB, Poisot T, Fitzjohn R. PACo: implementing Procrustes approach to cophylogeny in R. *Methods Ecol Evol.* 2017;8:932–40.
31. Douglas AE. The B vitamin nutrition of insects: the contributions of diet, microbiome and horizontally acquired genes. *Curr Opin Insect Sci.* 2017;23:65–9.
32. Kikuchi Y. Endosymbiotic bacteria in insects: their diversity and culturability. *Microb Environ.* 2009;24:195–204.
33. Manzano-Marín A, Ocegüera-Figueroa A, Latorre A, Jiménez-García LF, Moya A. Solving a bloody mess: B-vitamin independent metabolic convergence among gammaproteobacterial obligate endosymbionts from blood-feeding arthropods and the leech *Haementeria officinalis*. *Genome Biol Evol.* 2015;7:2871–84.
34. Xiong H, Campelo D, Pollack RJ, Raoult D, Shao R, Alem M, et al. Second-generation sequencing of entire mitochondrial coding-regions (approx. 15.4 kb) holds promise for study of the phylogeny and taxonomy of human body lice and head lice. *Med Vet Entomol.* 2014;28:40–50.
35. Hosokawa T, Nikoh N, Koga R, Satō M, Tanahashi M, Meng X-Y, et al. Reductive genome evolution, host–symbiont co-speciation and uterine transmission of endosymbiotic bacteria in bat flies. *ISME J.* 2012;6:577–87.
36. Allen JM, Light JE, Perotti MA, Braig HR, Reed DL, Tregenza T. Mutational meltdown in primary endosymbionts: selection limits Muller's Ratchet. *PLoS ONE.* 2009;4:e4969.
37. Hypša V, Křížek J. Molecular evidence for polyphyletic origin of the primary symbionts of sucking lice (Phthiraptera, Anoplura). *Microb Ecol.* 2007;54:242–51.
38. Alickovic L, Johnson KP, Boyd BM. The reduced genome of a heritable symbiont from an ectoparasitic feather feeding louse. *BMC Ecol Evo.* 2021;21:108.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

