



HAL
open science

Introduction of the SARS-CoV-2 Beta variant from Comoros into the Marseille geographical area

van Thuan Hoang, Loutfia Assoumani, Jérémy Delerce, Linda Houhamdi, Marielle Bedotto, Jean-Christophe Lagier, Matthieu Million, Anthony Levasseur, Pierre-Edouard Fournier, Bernard La Scola, et al.

► To cite this version:

van Thuan Hoang, Loutfia Assoumani, Jérémy Delerce, Linda Houhamdi, Marielle Bedotto, et al.. Introduction of the SARS-CoV-2 Beta variant from Comoros into the Marseille geographical area. *Travel Medicine and Infectious Disease*, 2022, 46, pp.102277. 10.1016/j.tmaid.2022.102277. hal-03678697

HAL Id: hal-03678697

<https://amu.hal.science/hal-03678697>

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

TITLE PAGE

Full-length title:

Introduction of the SARS-CoV-2 Beta variant from Comoros into the Marseille geographical area

Short title (for the running head):

SARS-CoV-2 Beta variant introduction in Marseille

Author list: Van Thuan Hoang^{1,2,3}, Loutfia Assoumani¹, Jérémy Delerce¹, Linda Houhamdi^{1,5}, Marielle Bedotto¹, Jean-Christophe Lagier^{1,4,5}, Matthieu Million^{1,4,5}, Anthony Levasseur^{1,4,5}, Pierre-Edouard Fournier^{1,4,5}, Bernard La Scola^{1,4,5}, Didier Raoult^{1,4,5}, Philippe Gautret^{1,2,5} *, Philippe Colson^{1,4,5} *

Affiliations: ¹ IHU-Méditerranée Infection, 19-21 boulevard Jean Moulin, 13005 Marseille, France; ² Aix Marseille Univ, IRD, SSA, VITROME, Marseille, 27 boulevard Jean Moulin, 13005 Marseille, France; ³ Thai Binh University of Medicine and Pharmacy, P. Kỳ Bá, Thái Bình, Thai Binh, Vietnam; ⁴ Aix Marseille Univ, IRD, MEPHI, 27 boulevard Jean Moulin, 13005 Marseille, France; ⁵ Assistance Publique-Hôpitaux de Marseille, 264 rue Saint-Pierre, 13385 Marseille.

***Corresponding authors:** Philippe Colson and Philippe Gautret, Institut Hospitalo-Universitaire Méditerranée Infection, 19-21 Boulevard Jean Moulin 13385 Marseille Cedex 05, France. Phone: + 33 (0) 4 13 73 24 01. Fax: + 33 (0) 4 13 73 24 02. E-mail addresses: philippe.colson@univ-amu.fr; philippe.gautret@ap-hm.fr

Key words: SARS-CoV-2; COVID-19; Beta variant; Comoros; travel; epidemic.

Word count: abstract: 200; text: 1,683; **Figures:** 2; **Table:** 0; **References:** 27

Supplementary Material: **Supplementary Figure:** 1; **Supplementary Table:** 1

ABSTRACT

Background: We describe the epidemiology of the first cases diagnosed in our institute of infections with the Beta variant and how this variant was imported to Marseille.

Methods: The SARS-CoV-2 Beta variant was identified based on analyses of sequences of viral genomes or of a spike gene fragment obtained by next-generation sequencing using Illumina technology, or a real-time reverse-transcription-PCR (qPCR) specific of the Beta variant.

Results: The first patient diagnosed as infected with the SARS-CoV-2 Beta variant was sampled on 15 January 2021. Twenty-nine patients were diagnosed in January 2021 (two weeks). Fifteen (52%) patients were of Comorian nationality. Eight (28%) had travelled abroad, including six who had returned from Comoros. Phylogeny based on SARS-CoV-2 genomes from 11 of these patients and their best BLAST hits from the GISAID database showed that seven patients, including the four returning from Comoros, were clustered with 27 other genomes from GISAID that included the six first Beta variant genomes described in Comoros in January 2021.

Conclusions: Our analyses highlight that, as for the case of other SARS-CoV-2 variants that have been diagnosed in Marseille, the Beta variant was imported to Marseille through travel from abroad. It had limited spread in our geographical area.

TEXT

INTRODUCTION

Since its emergence, SARS-CoV-2 has spread worldwide and has led to a COVID-19 pandemic [1]. Due to the considerable variability and spread of this virus, multiple variants have emerged that have been and continue to be responsible for distinct epidemics that occurred concurrently or successively [2]. At the University Hospital Institute (IHU) - Méditerranée Infection, in Marseille, southern France, we described 10 variants as early as during the summer of 2020 by performing genomic surveillance using next-generation sequencing and variant specific qPCR [2,3]. In addition, we observed differences in clinical severity between some of these variants [4]. However, it took until December 2020 for such variants to be considered as being worthy of detection and monitoring, when the emergence of another variant, currently named Alpha (or B.1.1.7 or 20I), was described in the United Kingdom and then spread worldwide [5]. Subsequent SARS-CoV-2 variants have been increasingly scrutinised through genomic surveillance, and three other major variants were first described in South Africa (Beta, or B.1.351 or 20H), Brazil (Gamma, or P.1 or 20J), and India (Delta, or B.1.617 or 21A) [6]. The Beta variant, first described in late 2020 [7,8], was classified by the World Health Organization as a variant of concern, and spread to 105 countries worldwide with >25,000 sequences published [9].

At the IHU Méditerranée Infection, we previously reported how several variants had been imported from abroad, including the Marseille-1 variant, of sub-Saharan origin, from North Africa by boat [10]; the Delta variant from India through a sailor who was to embark on a boat in Marseille [11]; and the Marseille-501 variant from Mayotte, a French overseas department in the

Comoros archipelago [12]. Here we describe the epidemiology of the first cases which were diagnosed by our institute as being infected with the Beta variant in January 2021, and how this variant was imported to Marseille.

PATIENTS AND METHODS

Identification of the Beta variant was performed using different techniques, on viral RNA extracted from 200 µL of nasopharyngeal swab fluid collected from patients to diagnose SARS-CoV-2 infection by real-time reverse-transcription-PCR (qPCR) [2]. Extracted viral RNA was either reverse-transcribed using SuperScript IV (ThermoFisher Scientific) prior to cDNA second strand synthesis with Klenow Fragment DNA polymerase (New England Biolabs, Beverly, MA, USA), or processed for obtention of cDNA according to the COVIDSeq protocol (Illumina Inc.). In most cases, SARS-CoV-2 genotyping consisted of next-generation sequencing of viral genomes, or of a spike gene fragment (2,129 nucleotides corresponding to positions 21,296-23,424 of the genome of the Wuhan-Hu-1 isolate GenBank Accession no. NC_045512.2) as previously described [12]. Next-generation sequencing was performed using the Illumina technology, with the Nextera XT paired-end procedure on a MiSeq instrument (Illumina Inc., San Diego, CA, USA) or with the Illumina COVIDSeq protocol on a NovaSeq 6000 instrument (Illumina Inc), as previously described [2,10]. Sequencing reads generated by next-generation sequencing were assembled by mapping on the SARS-CoV-2 genome GenBank Accession no. NC_045512.2 (Wuhan-Hu-1 isolate) with the CLC Genomics workbench v.7 software (<https://digitalinsights.qiagen.com/>) or with the Minimap2 software [13]. Assembled sequences were analysed for their classification using the Nextclade online tool (<https://clades.nextstrain.org/>) [14], the Pangolin online tool ([4](https://cov-</p></div><div data-bbox=)

lineages.org/pangolin.html) [15], and an in-house script written in Python as previously described [2]. SARS-CoV-2 genome sequences obtained in the present study have been deposited on the GISAID sequence database (<https://www.gisaid.org/>) [17] (Supplementary Table S1). Alternative SARS-CoV-2 genotype identification used an in house-designed qPCR system specific of the Beta variant, as previously described [2]. This qPCR system targets the envelope gene and uses forward primer C_SA_3_MBF: TGAATTGCAGACACCTTTTGA, reverse primer C_SA_3_MBR: CAACCCTTGGTTGAATAGTCTTG, and probe C_SA_3_MBP: TGACATCTTCAATGGGGAATGT. Phylogenetic analyses were performed based either on SARS-CoV-2 genome sequences when available, or on spike gene sequences. Phylogeny reconstructions were performed using the MEGAX software v.10.2.6 (<http://www.megasoftware.net/>) [16] with the neighbour-joining method and the Kimura 2-parameter method. The five best BLAST hits from the GISAID sequence database (<https://www.gisaid.org/>) [17] for each SARS-CoV-2 genome analysed here, as well as the six first Beta variant genomes detected in Comoros in January 2021 [18] were incorporated into phylogenies. SARS-CoV-2 culture was performed by inoculating nasopharyngeal samples on Vero E6 cells, as previously described [19].

RESULTS

The first SARS-CoV-2-infected patient to be diagnosed with the Beta variant was sampled on 15 January 2021. As of 31 August 2021, 611 patients had been diagnosed as being infected with this variant through routine SARS-CoV-2 genotyping (Figure 1). Twenty-nine patients were diagnosed in January 2021 (over a two-week period), from whom the Beta variant was identified by genome or spike gene next-generation sequencing in 11 and 16 cases,

respectively, and by qPCR specific of this variant, in two cases. Seventeen patients were male and 12 were female (sex ratio= 1.4). The median age was 37 years (interquartile= [27 - 50], range= 10-62 years). Additional epidemiological or clinical data were available for 26 of the 29 patients. They were tested because they had respiratory symptoms or were contact cases of patients diagnosed with SARS-CoV-2-diagnosed patients. Fifteen (52%) patients were of Comorian origin (Supplementary Table S1). Eight (28%) had travelled abroad, including six who had returned from Comoros (all were of Comorian nationality), and two who had returned from Italy and United Arab Emirates. Four of the six people returning from Comoros transited through Tunisia. Seven of the eight travellers (six returning from Comoros and one from United Arab Emirates) were diagnosed over a three-day period, between 16-18 January 2021, which corresponded to the first evidence of the introduction of the Beta variant in Marseille. Time between return to France and diagnosis ranged from 0 to 4 days for the eight patients who travelled abroad and from 0 to 1 day for the six patients who returned from Comoros (Supplementary Table S1). Among those patients who had not travelled, two had participated in mass events (a wedding and a religious festival), eleven were contact cases, and no specific exposure was reported in five symptomatic patients. SARS-CoV-2 culture was performed for a respiratory sample collected from twelve patients and was positive in all cases, a cytopathic effect being observed in a mean time of 5.7 ± 4.2 days (range, 2-16 days).

The phylogenetic tree based on SARS-CoV-2 genomes showed that seven of the genomes that were obtained in our institute from the 10 patients, including the four from patients returning from Comoros, were clustered (with a bootstrap value of 75%) with 27 other genomes from GISAID, which included the first six Beta variant genomes detected in January 2021 and described in Comoros [17], 16 genomes obtained from patients in Mayotte, a French overseas

department in the Comoros archipelago, and two genomes obtained from patients in Reunion Island, another French overseas department in the Indian Ocean (Figure 2). Three other genomes that were obtained in our institute, including one from a patient returning from Italy and one from a patient returning from Dubai, were clustered with genomes obtained from patients in Luxembourg, France, Belgium, the Netherlands and Israel, while the last genome was not robustly clustered with another genome, although it was also most closely related to genomes from Comoros. The phylogenetic tree based on the SARS-CoV-2 spike gene fragment showed lower bootstrap values than that based on genomes. Nonetheless, sequences from all fifteen patients of Comorian nationality including those from all six patients returning from Comoros were clustered (bootstrap value, 54%) with all other sequences from patients sampled in Comoros islands (Supplementary Figure S1).

DISCUSSION

Our analyses highlight that, as is the case of the other SARS-CoV-2 variants that we have diagnosed in Marseille [2, 10-12], the Beta variant was imported to Marseille through travel from abroad, primarily from the Comoros. This coincided with the initial detection of genomes of the Beta variant in Comoros in January 2021, which comprised two clusters with the most recent common ancestor dating to 30 October 2020, and which were close to viruses identified in Mayotte, part of the Comoros archipelago [18]. In the present study, sources other than Comoros, including return from Italy, United Arab Emirates and unknown sources, contributed to the early emergence or spread of the Beta variant in our geographical area.

Marseille includes a large community of more than 50,000 people originating from the Comoros [20, 21]. Comorians often travel between Comoros and Marseille, which can spread

infectious diseases. The transmission of several infectious agents between Comoros and Marseille has previously been described [22-24]. Furthermore, Marseille is France's second largest city and is located on the Mediterranean coast. Daily passenger traffic takes place by air and by boat, especially between Marseille and African countries [22].

The Beta variant, along with other N501Y variants including the Alpha, Gamma, and Marseille-501 variants, was part of the third period of high SARS-CoV-2 incidence in Marseille [4]. It has been described as being more infectious than previous SARS-CoV-2 variants [25], and as displaying decreased sensitivity to neutralising antibodies, including from the convalescent plasma of South African patients [26]. However, following its introduction, it had limited spread in our geographical area, with only 611 cases as of 31 August 2021, compared to 9,436 cases (15 times more) with the Alpha variant, first detected in late December 2020; 6,742 cases (11 times more) with the Marseille-4 variant (or B.1.160 lineage according to the Pangolin classification) that we first detected in July 2020; and 6,079 cases (10 times more) with the Delta variant, that we first detected in April 2021. The extent of the Beta variant in the Marseille area was, in contrast, higher than that of the Gamma variant (87 cases as of 31 August 2021). The causes of such differences in the spread of SARS-CoV-2 variants on the local and national scales merit further study. These causes may include viral features but also various epidemiological parameters, including the frequency of events introducing these variants from abroad and the effect of the co-circulation of several variants during the same period of time.

Travels abroad during the SARS-CoV-2 pandemic, including after the reopening of international borders, create favourable conditions for the spread of SARS-CoV-2 variants on a global scale [2,25]. Overall, our data exemplify that such viral variants can rapidly spread internationally through travel, thus causing distinct epidemics according to geographic location

and over time. They also show that there are multiple travel routes and that closing national borders is complicated. Epidemiological and genomic surveillance is needed to decipher the dynamic of these variants and their epidemiological and clinical specificities.

Author contributions

Van Thuan Hoang: Methodology, Investigation, Formal analysis, Writing - original draft, review & editing; **Loutfia Assoumani:** Methodology, Investigation, Formal analysis;

Jérémy Delerce: Methodology, Investigation; **Linda Houhamdi:** Methodology, Investigation;

Marielle Bedotto: Methodology, Investigation; **Jean-Christophe Lagier:** Investigation;

Matthieu Million: Investigation; **Anthony Levasseur:** Methodology, Validation;

Pierre-Edouard Fournier: Methodology, Validation; **Bernard La Scola:** Methodology, Investigation; **Didier Raoult:** Conceptualization, Formal analysis, Writing - review & editing, Supervision; **Philippe Gautret:** Conceptualization, Methodology, Investigation, Formal analysis, Validation, Writing - original draft, review & editing, Supervision; **Philippe Colson:** Conceptualization, Methodology, Investigation, Formal analysis, Validation, Writing - original draft, review & editing, Supervision.

Ethics

This study has been approved by the ethics committee of our institution (N°2020-016-03).

Access to the patients' biological and registry data issued from the hospital information system

was approved by the data protection committee of Assistance Publique-Hôpitaux de Marseille (APHM) and was recorded in the European General Data Protection Regulation registry under number RGPD/APHM 2019-73.

Funding

This work was supported by the French Government under the “Investments for the Future” program managed by the National Agency for Research (ANR), Méditerranée-Infection 10-IAHU-03 and was also supported by Région Provence Alpes Côte d’Azur and European funding FEDER PRIMMI (Fonds Européen de Développement Régional-Plateformes de Recherche et d’Innovation Mutualisées Méditerranée Infection), FEDER PA 0000320 PRIMMI.

Conflicts of Interest

The authors declare no conflict of interest. Didier Raoult has been a consultant for Hitachi High-Technologies Corporation, Tokyo, Japan from 2018 to 2020. He is a scientific board member of Eurofins company and a founder of a microbial culture company (Culture Top). The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

REFERENCES

1. European Centre for Disease Prevention and Control. COVID-19 situation update worldwide, as of week 36, updated 16 September 2021. Available at: <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases> (Accessed on September 21, 2020)
2. Colson P, Fournier PE, Chaudet H, Delerce J, Giraud-Gatineau A, Houhamdi L, et al. Analysis of SARS-CoV-2 variants from 24,181 patients exemplifies the role of globalisation and zoonosis in pandemics. *MedRxiv* 2021, <https://doi.org/10.1101/2021.09.10.21262922>.
3. Colson P, Levasseur A, Delerce J, Chaudet H, Bossi V, Ben Khedher M, et al. Dramatic increase in the SARS-CoV-2 mutation rate and low mortality rate during the second epidemic in summer in Marseille. *IHU Preprints* 2020, doi: <https://doi.org/10.35088/68c3-ew82>.
4. Dao TL, Hoang VT, Nguyen NN, Delerce J, Chaudet H, Levasseur A, et al. Clinical outcomes in COVID-19 patients infected with different SARS-CoV-2 variants in Marseille, France. *Clin Microbiol Infect* 2021;27:1516.e1-1516.e6.
5. Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. *Science* 2021;372:eabg3055.
6. Oude Munnink BB, Worp N, Nieuwenhuijse DF, Sikkema RS, Haagmans B, Fouchier RAM, et al. The next phase of SARS-CoV-2 surveillance: real-time molecular epidemiology. *Nat Med* 2021;27:1518-1524.
7. Fink S. South Africa announces a new coronavirus variant. *The New York Times* 2020. Available at: <https://www.nytimes.com/2020/12/19/world/south-africa-announces-a-new-coronavirus-variant.html> (Accessed on September 21, 2021)
8. Tegally H, Wilkinson E, Giovanetti M, Iranzadeh A, Fonseca V, Giandhari J, et al. Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature* 2021;592:438–43.
9. Cov-lineages.org. Report B.1.351 variant. Available at: https://cov-lineages.org/global_report_B.1.351.html (Accessed on September 21, 2021)
10. Colson P, Levasseur A, Gautret P, Fenollar F, Thuan Hoang V, Delerce J, et al. Introduction into the Marseille geographical area of a mild SARS-CoV-2 variant originating from sub-Saharan Africa: An investigational study. *Travel Med Infect Dis* 2021;40:101980.
11. La Scola B, Lavrard P, Fournier P-E, Colson P, Lacoste A, Raoult D. SARS-CoV-2 variant from India to Marseille: The still active role of ports in the introduction of epidemics. *Travel Med Infect Dis* 2021;42:102085.

12. Colson P, Levasseur A, Delerce J, Pinault L, Dudouet P, Devaux C, et al. Spreading of a new SARS-CoV-2 N501Y spike variant in a new lineage. *Clin Microbiol Infect* 2021;27:1352.e1-1352.e5.
13. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 2018 Sep 15;34(18):3094-3100.
14. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics* 2018;34:4121-3.
15. Rambaut A, Holmes EC, O'Toole A, Hill V, McCrone JT, Ruis C, et al. A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic epidemiology. *Nat Microbiol* 2020;5:1403-1407.
16. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics Analysis across computing platforms. *Mol Biol Evol* 2018;35:1547-1549.
17. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from vision to reality. *Euro Surveill* 2017;22:30494.
18. Agoti CN, Githinji G, Mohammed KS, Lambisia AW, de Laurent ZR, Mburu MW, et al. Detection of SARS-CoV-2 variant 501Y.V2 in Comoros Islands in January 2021. *medRxiv* 2021.04.08.21254321; doi: <https://doi.org/10.1101/2021.04.08.21254321>.
19. Wurtz N, Penant G, Jardot P, Duclos N, La Scola B. Culture of SARS-CoV-2 in a panel of laboratory cell lines, permissivity, and differences in growth profile. *Eur J Clin Microbiol Infect Dis* 2021;40:477-484.
20. Marseille Population 2021 (Demographics, Maps, Graphs) n.d. <https://worldpopulationreview.com/world-cities/marseille-population> (Accessed September 21, 2021).
21. Katibou A. Les migrations Comoriennes en France. Available at : <https://www.recherches-internationales.fr/RI90/RI90-Katibou.pdf> (Accessed September 21, 2021).
22. Griffiths KM, Savini H, Brouqui P, Simon F, Parola P, Gautret P. Surveillance of travel-associated diseases at two referral centres in Marseille, France: a 12-year survey. *J Travel Med* 2018;25:tay007.
23. Parola P, Gazin P, Pradines B, Parzy D, Delmont J, Brouqui P. Marseilles: a surveillance site for malaria from the Comoros Islands. *J Travel Med* 2004;11:184-6.
24. Gautret P, Simon F, Askling HH, Bouchaud O, Leparç-Goffart I, Ninove L, et al. Dengue type 3 virus infections in European travellers returning from the Comoros and Zanzibar, February-April 2010. *Eurosurveillance* 2010;15:19541.
25. Dao TL, Hoang VT, Colson P, Lagier JC, Million M, Raoult D, Levasseur A, Gautret P. SARS-CoV-2 Infectivity and Severity of COVID-19 According to SARS-CoV-2 Variants:

Current Evidence. *J Clin Med* 2021;10:2635.

26. Wibmer CK, Ayres F, Hermanus T, Madzivhandila M, Kgagudi P, Oosthuysen B, et al. SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19 donor plasma. *Nat Med* 2021;27:622-5.
27. Lemey P, Ruktanonchai N, Hong SL, Colizza V, Poletto C, Van den Broeck F, et al. Untangling introductions and persistence in COVID-19 resurgence in Europe. *Nature* 2021;595:713-7.

FIGURE LEGENDS

Figure 1. Weekly distribution of patients with the Beta variant

Figure 2. Phylogenetic analysis based on SARS-CoV-2 genomes

The SARS-CoV-2 phylogenetic tree is based on the SARS-CoV-2 genomes. The five best BLAST hits from the GISAID sequence database (<https://www.gisaid.org/>) [16] for SARS-CoV-2 genome analysed here (labelled with a white bold font and a black background), as well as the six first Beta variant genomes detected in Comoros in January 2021 [17] (labelled with a blue bold font and a grey background) and the genome of the Wuhan-Hu-1 isolate (GenBank Accession no. NC_045512.2) were incorporated into the phylogeny reconstruction. SARS-CoV-2 genomes obtained from patients sampled in the Comoros archipelago are labelled with a blue bold font. Evolutionary history was inferred using MEGAX software (<http://www.megasoftware.net/>) [15] using the neighbour-joining method and the Kimura 2-parameter method. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1,000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree; the scale bars indicate the number of nucleotide substitutions per site. Bootstrap values > 50% are indicated on the tree.

Fig. 1




Fig. 2

