

Multi-elemental compound-specific isotope analysis of pesticides for source identification and monitoring of degradation in soil: a review

Patrick Höhener, Delphine Guers, Ouassim Boukaroum, Laure Malleret, Fabrice Martin-Laurent, Jérémy Masbou, Sylvain Payraudeau, Gwenaël Imfeld

▶ To cite this version:

Patrick Höhener, Delphine Guers, Ouassim Boukaroum, Laure Malleret, Fabrice Martin-Laurent, et al.. Multi-elemental compound-specific isotope analysis of pesticides for source identification and monitoring of degradation in soil: a review. Environmental Chemistry Letters, 2022, 20 (3927–3942), 10.1007/s10311-022-01489-8. hal-03760066

HAL Id: hal-03760066 https://amu.hal.science/hal-03760066

Submitted on 15 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Multi-elemental compound-specific isotope analysis of pesticides for		
2	source identification and monitoring of degradation in soil: a review		
3			
C			
4	Patrick Höhener ^{1*} , Delphine Guers ¹ , Laure Malleret ¹ , Ouassim Boukaroum ¹ , Fabrice Martin-		
5	Laurent, Jeremy Masbou, Sylvain Payraudeau, Gwenaer Imreid		
6			
7	¹ Aix Marseille University – CNRS, UMR 7376, Laboratory of Environmental Chemistry, Marseille,		
8	France		
9	² INRAE, Institut Agro, Université de Bourgogne, Université de Bourgogne Franche-Comté,		
10	Agroécologie, Dijon, France		
11	³ Université de Strasbourg, CNRS, ENGEES, Institut Terre et Environnement de Strasbourg (ITES,		
12	UMR 7063), France		
13	*Corresponding author: patrick.hohener@univ-amu.fr		
14	All E-Mails of authors:		
15	delphineguers04@gmail.com, mohamed.boukaroum@etu.univ-amu.fr, laure.malleret@univ-		
16	amu.fr, fabrice.martin@inrae.fr, masbou@unistra.fr, sylvain.payraudeau@engees.unistra.fr,		
17	imfeld@unistra.fr		
18			
19	Revised version Env Chem Letters ECLE-D-22-00326		
10			
20			
21			
22			
23	Key Words		

24 Persistent Pollutants; Pesticides; Fate; Stable isotopes; Soil; Monitoring

25 ABSTRACT

The transfer of pesticides from agricultural soils to food and drinking water is a huge environmental problem in many countries. Robust approaches to identify, characterize and quantify the dissipation of pesticides in soils at the field scale are scarce. A relatively mature tool to study industrial pollutants in groundwater is multi-elemental compound-specific isotope analysis (ME-CSIA) which may help to follow-up pesticide residues in soils. Direct adaptation to soils is challenging because pesticide concentrations are generally low, and soil constituents may hamper chromatographic separation and accurate analysis of pesticide residues.

33

34 We reviewed recent literature on ME-CSIA as an approach to improve the evaluation of sources 35 and transformations of synthetic pesticides in agricultural soils. First, we discuss current extraction 36 and purification techniques for pesticides in soils. Then, the analytical techniques for reliable 37 measurements of the stable isotope composition of the pesticides are examined. Five hundred and 38 fortyseven isotopic compositions for seventy-one active molecules and issued from different pesticide manufacturers were retrieved and listed in a database, for the following isotopes: ¹³C 39 (61%), ¹⁵N (14%), ³⁷Cl (14%), and ²H (11%). This unique database reflects the overall isotopic 40 41 variability and trackability of pesticide residues in the environment. Since the environmental fate of 42 pesticides is governed by multiple transformation processes, including bio- and photodegradation, 43 and other abiotic degradations, isotope fractionation values were retrieved from the peer-reviewed 44 literature. This results in a catalogue of two hundred seventy fractionation values for thirty-three 45 compounds to guide the interpretation of isotopic fractionation in the environment, including legacy 46 pesticides such as atrazine, lindane, dichlorodiphenyltrichloroethane, chlordecone and 47 organophosphorus compounds. Specific transformation processes may be identified and quantified using the Rayleigh concept for isotope fractionation during reaction. The review article summarizes 48 49 existing modelling approaches and future trends to predict the reactive transport of pesticides in 50 soils and associated hydrological systems.

51 **1. Introduction**

52 Soil contamination by organic compounds such as pesticides, hydrocarbons or chlorinated 53 solvents is a widespread problem occurring on agricultural soils, urban and industrial sites. In 54 particular, an estimated 1 to 2.5 million tons of active pesticide ingredients were used annually in 55 2012, mainly in agriculture (Fenner et al. 2013). In principle, pesticides are only registered for use if 56 they are demonstrated not to persist too long in the environment beyond their intended period of 57 action. However, many pesticides persist for long periods in soil, mainly in the form of non-58 extractable residues. On the one hand, pesticides leaching from the soil into surface and 59 groundwater may cause severe deterioration of aquatic ecosystems (Brock et al. 2009). On the 60 other hand, uptake of recalcitrant pesticides into crops can lead to human health risks and 61 economic losses for farmers and distributors. Additionally, pesticide residues can impact soil living 62 organisms with consequences on soil ecosystemic services. Over the last decades, efforts to 63 reduce the use of pesticides have mostly failed, while global warming is expected to expand the 64 prevalence of pests in the future and increase the societal dependence on chemical crop protection (Stokstad, 2018). The continuous and partially increasing use of synthetic pesticides in 65 66 agriculture jeopardizes the ecological and chemical status of soil and aquatic ecosystems, 67 threatening human health (Mateo-Sagasta et al. 2017). In the future, agroecology and organic 68 farming may globally increase and clean and green chemistry in industry may give hope for change 69 in the conception of modern pesticides. But the issue of the persistence in soils of historical and 70 current pesticides and other pollutants still needs to be solved.

71 The recurrent detection and persistence of pesticides in aquatic ecosystems also emphasizes 72 difficulties to extrapolate pesticide testing in the laboratory or in lysimeter to field conditions 73 (Fenner et al. 2013). Previous research postulated several reasons for the slow degradation of 74 pesticides in soils, including: i) the overall low bioavailability, due to sorption and sequestration of 75 pesticides into soil organic matter (Allen-King et al. 2002), and diffusion-limited desorption from soil 76 aggregates or from minerals (Grathwohl and Reinhard 1993), ii) the poor ability of microorganisms to degrade at low concentrations (Roch and Alexander 1997), iii) the absence of adapted 77 78 microorganisms with genes encoding for degrading enzymes (Vogel 1996), iv) the inability of 79 microorganisms to degrade due to insufficient available nutrients (N. C source) for co-metabolism. 80 and v) combinations of either reasons. For example, the soil fumigant 1,2-dibromoethane was 81 found in agricultural soils up to 19 years after its last reported application (Steinberg et al. 1987). 82 while half-life at 20°C in laboratory is 70 days (Lewis et al. 2016). In contrast to freshly added 1,2-83 dibromoethane, residual 1,2-dibromoethane remains recalcitrant to both mobilization (desorption 84 into air and water) and microbial degradation, while diffusion-limited desorption was the main 85 process driving 1,2-dibromoethane removal (Steinberg et al. 1987).

Back in 1962, the publication of the book "Silent spring" by Rachel Carson (1962) revealed for the
first time how dangerous persistent organic pesticides in soils can be for man and nature. Since

then, huge advancements in analytical chemistry have been made to identify, follow-up and predict
pesticides in soils and to evaluate their degradation.

90 Some bioreporter bacteria, such as e.g., luminescent PAH degraders, have been developed in the 91 past to identify less bioavailable pollutants such as hydrocarbons. However, such approaches are 92 restricted to a narrow range of pollutants and proved to be generally inefficient in the case of 93 strongly adsorbing soils. Often, conventional methods cannot resolve whether pollutants are truly 94 recalcitrant, or degradation is just too slow for being monitored. The occurrence of biodegradation 95 of persistent pollutants in the soil is of fundamental importance to improve their monitoring, to evaluate their potential ecotoxicological effects, and to predict ensuing risks following their 96 97 exposure.

98 One of the latest important analytical developments was the coupling of gas/liquid-chromatography 99 to continuous-flow isotope ratio mass spectrometry to study various stable isotopes ratios specific 100 to each pesticide molecule. When at least two different isotope ratios are analyzed, one speaks of 101 multi-elemental compound-specific isotope analysis (ME-CSIA; Elsner and Imfeld 2016; Vogt et al. 102 2016). Starting from the early 2000s, ME-CSIA has successfully been applied to some common 103 pollutants in many different environments (Lichtfouse, 2000), but especially in groundwater. The 104 Environmental Protection Agency of the United States of America (US-EPA) published in 2008 a 105 auideline for the application of CSIA to groundwater contamination (Hunkeler et al. 2008). More 106 recently, current research on ME-CSIA was reviewed in two short review articles on hydrocarbons 107 (Vogt et al. 2016) and on pesticides (Elsner and Imfeld 2016).

108 In pesticide-contaminated soils, ME-CSIA is much more difficult to apply compared to surface or 109 groundwater applications mainly due to three reasons. First, soil is a complex matrix and contains 110 compounds, such as humic and fulvic acids, which may hamper the extraction (Kalathoor et al. 111 2015), the chromatographic separation and the identification of pollutants due to the so-called 112 matrix effect (Bakkour et al. 2018). This is generally less of a problem in groundwater which is 113 often a well-filtered matrix. Second, pesticides in soil water are usually in the microgram-per-liter 114 range or below it, while industrial pollutants in groundwater often exceed the mg/L range. In 115 addition, pesticides in soil cannot be as easily concentrated as aqueous samples. Third, pesticides 116 in agricultural soils differ from fuel compounds and solvents by their polarity and by a generally 117 larger molecular mass, limiting measurement of stable isotope fractionation using traditional 118 techniques.

119 Nevertheless, in the last years many studies revealed that ME-CSIA can not only be applied to 120 several pesticides such as s-triazine herbicides (Meyer et al. 2008; Reinnicke et al. 2012), 121 phenoxyacid herbicides (Milosevic et al. 2013), substituted phenylureas such as isoproturon 122 (Penning and Elsner 2007), glyphosate (Kujawinski et al. 2013), bentazone (Reinnicke et al. 2010), 123 organophosphorus insecticides (Wu et al. 2014), hexachlorocyclohexanes (Badea et al. 2011), 124 DDT (Holmstrand et al. 2004) and chlordecone (Chevallier et al. 2018) but also to transformation 125 products amino methyl phosphonic acid (AMPA, Kujawinski et al. 2013) and 2,6-126 Dichlorobenzamide (Reinnicke et al. 2012). Analytical developments were made toward novel 127 sample clean-up techniques, using for instance preparative HPLC (Schreglmann et al. 2013) or 128 molecularly imprinted polymers (Bakkour et al. 2018), and novel injection systems including pre-129 concentration, like solid-phase microextraction (SPME) or large-volume injections. While all these 130 studies brought a proof of concept for soil ME-CSIA to the scientific literature, they were 131 sometimes restricted to artificially contaminated laboratory systems (e.g., soil microcosms or 132 lysimeters) and higher concentrations, while field studies remain scarce (Alvarez-Zaldivar et al. 133 2018).

134 The purpose of this review article is thus to describe recent developments of ME-CSIA for 135 pesticides in soils and to provide an overview on what ME-CSIA offers to better understand 136 sources and the fate of pesticides in soils. The underlying hypothesis of this review is that ME-137 CSIA bears the potential to identify, characterize, quantify and predict, when combined with 138 reactive transport models, in situ transformation of pesticides in soils, even if no transformation 139 products are found. A list of isotopic compositions of active molecules from available pesticide 140 formulations, analytical standards and major degradation products is provided to gain insight in the 141 overall variability of molecules from different sources. Furthermore, stable isotope fractionation 142 processes were compiled in another list to help interpret the fate of pesticides in soils. The 143 knowledge gathered in this article will help not only to take actions to assist the self-restoration of 144 soils (e.g., by nutrient addition or plant introduction) and evaluate its resilience, but also to provide 145 an expected timeframe to manage pesticide ecotoxicity when remediation is not feasible, as e.g. 146 for Chlordecone in the French West-Indies (Chevallier et al, 2018).

147 2. Analysis of soils

148 2.1 Extraction and clean-up for ME-CSIA of pesticide residues in soils: analytical challenges 149 and prospects

150 Challenges. The extension of ME-CSIA of pesticide residues in soils faces three main analytical 151 challenges limiting the evaluation of pesticide degradation in soils: (i) the need of a sufficient mass 152 of analyte for ME-CSIA combined with low pesticide concentrations (sub-ug/q_{soil}) following 153 degradation, ageing or leaching, (ii) the wide range of pesticide physico-chemical characteristics 154 (polar to apolar, non-volatile to volatile), and (iii) the co-enrichment of non-volatile soil components 155 leading to the so-called 'matrix effect' during chromatographic separation. As for water samples, 156 the development of appropriate sample preparation procedures, including optimized robust 157 extraction and cleanup, is a fundamental key to maximize the analytical performance for precise 158 and sensitive ME-CSIA without altering the isotope ratio of the target compounds (Torrento et al. 159 2019).

160 The low concentrations of pesticides typically found in soils, the chemical properties of pesticides 161 and the complexity of soil matrices, impose a significant pre-concentration step (from 1000 to 162 100,000-fold to allow >10-15 ng C or Cl, >30 ng N or H on column) depending on the elemental 163 contribution to the molar mass of the target analyte(s), as well as its solubility, polarity and 164 volatility. In addition, the selectivity and efficiency of pesticide extraction largely varies according to 165 the soil moisture, pH, texture and the amount of organic matter content controlling the amount of 166 available pesticide in soils. For instance, pesticides typically sorb to soils with high clay and organic 167 matter content and low water content (e.g. (Correia-Sa et al. 2012). Pesticide sequestration and 168 extraction also depends on pesticide ageing, i.e., the pesticide physical and chemical distribution 169 and diffusion over time in soil pore water, minerals and organic matter.

170 Extraction. Pesticides from different families were already extracted from diverse soils for CSIA in 171 several recent analytical (lvdra et al. 2014; Wu et al. 2019a), laboratory (Masbou et al. 2018; Liu et 172 al. 2020; Liu et al. 2021; Perez-Rodriguez et al. 2021; Torrento et al. 2021) and field studies 173 (Alvarez-Zaldivar et al. 2018; Wu et al. 2019a). Both classical and modern methods of pesticide 174 extraction, described in several leading reviews (e.g. (Tadeo et al. 2012; Bruzzoniti et al. 2014), 175 were adapted for the target analytes. Extraction methods include quick, easy, cheap, effective, rugged and safe procedure (QuEChERS), accelerated solvent extraction, ultrasonic assisted 176 177 extraction (USE) and focused ultrasonic extraction (FUSE). An accelerated solvent extraction 178 method was validated for determining carbon isotope ratios of Hexachlorocyclohexanes, DDT, and 179 their chlorinated metabolites (soil concentrations from 0.3 to 0.8 mg/kg) in two different types of soil 180 matrices (Ivdra et al. 2014). However, accelerated solvent extraction using three static cycles to 181 increase the recovery rate (>85%) was privileged in Wu et al. 2019a and further related studies. 182 accelerated solvent extraction enabled reproducible extraction of hexachlorocyclohexanes from

183 soil for C, H and Cl isotope analysis within the analytical precision of ±0.5%, ±5% and ±0.3%, 184 respectively. QuEChERS approaches (Anastassiades et al. 2003) involve an initial extraction with 185 acetonitrile followed by salt addition to induce phase separation. QuEChERS with acetonitrile 186 (Torrento et al. 2021) or modified for the use of ethylacetate (Alvarez-Zaldivar et al. 2018) or pentane:dichloromethane (3:1, v/v;Perèz et al. 2021) as extraction solvent, in pH buffered or 187 188 unbuffered media, were used to extract atrazine, metolachlor, alachlor, acetochlor or metalaxyl from soil samples without causing significant isotope fractionation for carbon ($\Delta \delta^{13}C \le 1\%$) and 189 nitrogen ($\Delta \delta^{15}$ N ≤ 0.5‰), despite variable extraction efficiencies. However, extraction from two 190 agricultural soils induced a systematic but reproducible isotope fractionation for CI ($\Delta \delta^{37}$ CI between 191 192 +2.5 and +3.5‰) using a QuEChERS extraction kit with acetonitrile (Torrento et al. 2021).

193 Overall, these results emphasize the versatility of QuEChERS approaches as a standard 194 preparation method for pesticide CSIA from soil samples and possible adaptations for specific 195 matrix-analyte combinations to reach more selective extraction. Acidic pesticides are usually 196 extracted from soil with solvents of medium polarity (ethyl acetate or methanol) in a solvent-water 197 mixture in buffered extraction (pH = 5), by citrate or acetate buffering, to enhance recoveries of 198 pesticides showing pH dependence (Tadeo et al. 2012). The effect of pH on extraction from soil 199 and matrix interferences should also be consirered. Indeed, humic acids at low pH values are 200 protonated and carry fewer negative charges (Melsbach et al., 2020). For less-polar pesticides, 201 such as chloroacetamides, triazines and their metabolites and acetamides, organic solvents like 202 methanol and acetonitrile are commonly used, alone or with mixtures with water (Perez-Rodriguez 203 et al., 2021). The solvent selection should be balanced with the soil matrix co-elution and reduction 204 of the matrix effect before CSIA. In this respect, acetonitrile has the advantages to extract less 205 lipophilic compounds from the soil, to facilitate the removal of residual water with drying agents, 206 and to form distinct partitioning phases with nonpolar solvents to facilitate further cleanup 207 (Anastassiades et al., 2003). Other methods, such as ultrasonic-assisted extraction, are faster 208 since several extractions can be performed simultaneously at relatively low-cost without 209 specialized laboratory equipment (Anastassiades et al., 2003). However, classical shaking or 210 QuEChERS approaches remain adaptative, low-cost and robust in routine extraction for pesticide 211 CSIA. Analyte extraction efficiency using QuEChERS usually increases with an additional step of 212 ultrasound radiation, in a water bath, probes or sonoreactors, enhancing solvent penetration into soil and mass transfer (e.g., Santalad et al. 2010). More selective extraction procedures may 213 214 simplify sample purification prior to a chromatographic analysis.

215 *Cleanup.* The pre-concentration step for pesticide CSIA and coextraction of matrix components 216 requires cleanup strategies to purify soil extracts (e.g., Tadeo et al. 2012) and reduce co-enriched 217 soil organic components of unknown isotopic composition. Among liquid-solid extraction 218 techniques, the most common procedure is solid phase extraction (SPE) with cartridges or 219 columns using sorbents, such as Florisil, silica, alumina, C18 or graphitized carbon black, 220 depending on the selected pesticides (Buszewski et al., 2012). A current trend is to develop 221 analytical methods where the extraction and purification are carried out simultaneously. Most soil 222 QuEChERS approaches developed for pesticide CSIA rely on dispersive solid-phase extraction, in 223 which a bulk amount of solid phase extraction sorbent is mixed with the extract rather than loading 224 the extract in a column or cartridge packed with the sorbent. Primary Secondary Amine (PSA) 225 sorbent was added to improve the cleanup of hexachlorocyclohexanes (Liu et al. 2020; Liu et al. 226 2021) chloroacetanilide (Perez-Rodriguez et al. 2021; Torrento et al. 2021; Perez-Rodriguez et al. 227 2021) and metalaxyl extracts (Masbou et al. 2018) with anhydrous magnesium or sodium sulfate 228 as a cleanup and drying agent to remove soil matrix, ensure pesticide recovery and reduce GC-229 background. In the future, stir bar sorptive extraction (SBSE) coated with IRMS 230 polydimethylsiloxane (PDMS) or newer materials may be used for additional clean up and 231 enrichment following QuEChERS extraction of soil matrices for more selective pesticide extraction 232 (Abdulra'uf and Tan 2014). For instance, molecularly imprinted polymers are synthetic polymeric 233 materials with specific cavities designed for a template molecule that can retain the target pesticide 234 (Faroog et al., 2018). Molecularly imprinted polymers could drastically reduce the matrix effect due 235 to dissolved organic carbon for accurate C and N-CSIA measurements of 1H-benzotriazol, a 236 typical corrosion inhibitor (Bakkour et al. 2018). In comparison with other methods, molecularly 237 imprinted polymers offer high thermo-chemical stability and reusability and can be used for solid 238 phase extraction as a highly selective sorbent for different types of pesticides in trace 239 concentrations in soils (Farooq et al. 2018).

240 **2.2 Isotope analysis of pesticides**

Isotope ratios in pesticides are measured as the ratio of the relative abundance of heavy and light isotopes in a sample (with R_{sample} = abundance of heavy isotope/abundance of light isotope) and equation 1:

244
$$\delta = \left(\frac{R_{sample}}{R_{standard}} - 1\right) \times 1000$$
 eq. 1

245 using the international R_{standards} given as e.g. in Aelion et al. (2010). The abundances of the heavy 246 and light isotopes are generally quantified by an isotope ratio mass spectrometer (IRMS). 247 Pesticides are often commercialized as formulations where the active molecule is mixed with 248 additives in a solid or liquid matrix to facilitate their use. In order to trace its fate, only the 249 abundances of isotopes from the active substance must be measured. Hence, it becomes obvious 250 that purification and chromatographic separation of the pesticide is essential before measurement 251 by IRMS. A large majority of isotope data on pesticides in this study (Supplementary Material 252 Annex 1) were measured on gas chromatography coupled to an IRMS (GC-IRMS). For carbon 253 isotope analysis of the polar herbicide glyphosate, the IRMS was coupled to liquid chromatography 254 (LC-IRMS). And finally in an intermediate number of cases where pesticides were available as 255 pure substances, the measurement was made on an elemental analyzer coupled to an IRMS (EA-256 IRMS). The latter is widely used in many labs since it permits measuring international standard 257 materials as references and calibrate the IRMS by multiple point calibration (Schimmelmann et al. 258 2016). The three analytical techniques are illustrated in Figure 1 and are presented in the following 259 in more detail.



260



265 **EA-IRMS:** The five elements H, O, C, N and S can be analyzed in elemental analyzers. However, 266 sulfur isotopes were analyzed in pesticides in only one article (Schürner et al. 2015), and also only 267 one single publication included oxygen isotopes in glyphosate (Sandy et al. 2013). Therefore, we 268 present here mainly H, C and N analysis. For hydrogen isotope analysis, the sample is pyrolyzed 269 at 1400°C in a stream of helium gas and the produced hydrogen gas is separated from other 270 gasses, including CO and H₂O, by a chromatographic column and transferred via an open split 271 system to the IRMS for the measurement of the abundance ratio ²H/¹H. The open split system was 272 developed when continuous-flow IRMS became available and permits an accurate dilution of the 273 sample. For C and N elemental analysis, the sample is combusted at 1000°C in an oxidation

furnace with a pulse of oxygen. Then, the nitrous oxides are reduced to N_{2} , and the latter is separated from CO₂ and other gases by the chromatographic column in the EA. The IRMS is first set to masses 28, 29, 30 to quantify the abundances of ¹⁴N₂, ¹⁴N¹⁵N and ¹⁵N₂ for the calculation of $\delta^{15}N$. The magnet of the IRMS is then switched by a peak jump to the masses 44-46 to quantify the CO₂ isotopologues with different masses. Correction for ¹⁸O is made with the method of (Santrock et al. 1985). The mass of sample needed for accurate EA-IRMS measurement is mostly between 0.2 to 0.5 milligrams of pure pesticide, which is weighted in tin or silver capsules.

281 GC-IRMS. This is the most widely applied method for pesticide isotope analysis, especially for real 282 field samples, capable of measuring isotope ratios of H, C or N (Fig. 1). The main differences 283 compared to EA-IRMS is that chromatographic separation of the compounds is made before 284 pyrolysis or combustion, and that the injected mass of compound is much lower. The mass of 285 carbon to reach reliable IRMS measurements of carbon stable isotope ratio is typically around 10-286 15 nanograms (Aelion et al. 2010). Knowing this value, the minimum injection amount into the GC 287 can be back calculated according to dilution factors in the injector. For the hydrogen analysis, 288 again a pyrolysis is operated at 1400°C to produce H₂. For N and C analysis, a combustion furnace is used, also with reduction of NOx compounds to N₂. The analysis of 15 N in pesticides requires, 289 however, a separation of N_2 from CO_2 , which is done by freezing out the CO_2 in a cooled trap. 290

291 LC-IRMS. This technique has so far only been applied to C isotope analysis in the polar pesticide 292 glyphosate (Fig. 1). The eluent was a 2.5 mM NaH₂PO₄ solution adjusted to pH 1.9. The reagents 293 to convert glyphosate to CO₂ were phosphoric acid and peroxodisulfate in a reactor kept at 99.9°C. 294 Then the CO_2 was flushed with He to the IRMS. Details are provided in (Kujawinski et al. 2013) 295 and Mogusu et al. (2015). This technique may also improve knowledge on the persistence of the 296 transformation product amino methyl phosphonic acid, or other mostly polar products, following 297 degradation processes of pesticides in soils. Apart from the three most-used techniques shown in 298 Figure 1, other techniques were used. They were:

GC-qMS. This technique couples a gas chromatograph to a quadrupole mass spectrometer and has been first described in 2007 (Sakaguchi-Soder et al. 2007) and first applied to the insecticide DDT in 2010 (Aeppli et al. 2010). It is only available to chlorine isotopes due to the large abundance of the ³⁷Cl isotope which contributes to about 24% of all chlorine atoms in the world. GC-qMS results have been compared in an interlaboratory study to continuous-flow IRMS (Bernstein et al. 2011) and proved to yield sufficient accuracy in isotope ratios when a correct bracketing was made using two standard compounds of different isotope ratios which lie slightly outside of the measurement range of the samples. The technique was recently applied to chlorineisotope analyses in acetochlor, metolachlor and atrazine (Ponsin et al. 2019; Torrento et al. 2021).

308 GC-MC-ICPMS. This technique consists of coupling a GC to a Multi-Collector Inductively Coupled 309 Plasma Mass Spectrometer, first described by (Van Acker et al. 2006). For pesticides it was only 310 used for chlorine isotope analysis. In the plasma, the masses of ³⁵Cl and ³⁷Cl are generated for the calculation of the isotope ratio, but care must be taken not to count also the mass of ³⁶ArH when 311 312 water vapor is reacting with argon gas (Horst et al. 2017). It has been applied in two studies looking at hexachlorocyclohexanes (Wu et al. 2019b), acetochlor and S-metolachlor (Lihl et al. 313 314 2019). GC-MC-ICPMS measurements of pesticides may open the way to analyze the stable 315 isotope ratios of heteroatoms, such as bromine or sulfur, in pesticide molecules.

Offline MS. Other studies converted pesticide samples by off-line methods to gaseous molecules
 containing chlorine or bromine and injected them to Dual-Inlet IRMS for the analysis of ratios of
 ³⁷Cl/³⁵Cl (Drenzek et al. 2002; Drenzek et al. 2004; Reddy et al. 2002) or ⁸¹Br/⁷⁹Br (Kuntze et al.
 2016).

320 **2.3 Isotopic compositions of pesticides from literature**

321 Isotopic compositions of pesticides and of key degradation products were compiled from the peer-322 reviewed literature (Figure 2 and Supplementary material Annex 1). The main criteria for selection 323 of isotopic compositions of pesticides were the existence of a reliable stable isotope composition of 324 the active molecule. Publications in which the isotope analysis by EA-IRMS targeted commercial 325 formulations, meaning the active molecule was not pure, were not included (e.g. (Annable et al. 326 2007). All hexachlorocyclohexane isomers were included although only the gamma isomer is a 327 pesticide. The method of analysis is provided along with the isotopic composition. Isotopic 328 compositions were found for 71 different compounds, in publications which appeared between 329 2002 and 2021. A total of 547 isotopic compositions were listed, with 337 values for ¹³C, 75 values for ³⁷Cl, 78 for ¹⁵N and 57 for ²H. One publication was found for ¹⁸O in glyphosate (Sandy et al. 330 2013), and one for ³⁴S in amethryn (Schürner et al 2015). 331



Number of compounds for isotopic compositions

332

Figure 2: Pesticide classes for which isotopic compositions are reported in literature (see
Supplementary material, Annex 1).

335 **2.4 Isotope fractionation values of pesticides**

Isotope fractionation values (enrichment factors) of pesticides and key degradation products were also compiled from the peer-reviewed literature (Supp. Mat. Annex 2). The criteria for selection included (i) the presentation of a well-described degradation pathway, (ii) a clear description of reaction conditions, and (iii) the quantification of isotope fractionation values from isotope data using the Rayleigh approach.

Isotope fractionation values were found for 32 compounds (Supp. Mat., Annex 2). A total of 263 isotope enrichment factors were retrieved, with 174 values for ¹³C, 19 values for ³⁷Cl, 50 for ¹⁵N, 13 for ²H, and 7 for ⁸¹Br, 2 for ³⁴S and 1 for ¹⁸O. The brominated compounds all concerned reactions acting on the fumigant ethylene dibromide. For ¹³C, all enrichments were normal (i.e., negative isotope fractionation values) except two for photooxidation of atrazine where inverse effects were observed. The most pronounced normal isotope effect (the most negative enrichment factor for ¹³C, -30.9 ‰) was observed for the abiotic reduction of ethylene dibromide, a small compound with

only two carbon atoms, both positioned at reactive positions of the molecule. This resulted in a
 strong enrichment in ¹³C, as observed for the reductive dechlorination of chloroethenes (see e.g.
 (Hunkeler et al. 2002)).

For ¹⁵N, both inverse and normal isotope effects were observed for biotic transformations. Inverse effects with positive isotope fractionation values were observed for the biotic degradation of triazine herbicides (Schürner et al. 2015; Chen et al. 2019), the aerobic degradation of bromoxynil (Knossow et al. 2020) and also for the acid hydrolysis of atrazine (Meyer et al. 2009; Masbou et al. 2018).

The most pronounced normal isotope effect for ¹⁵N was the alkaline hydrolysis of isoproturon, a substituted urea (Penning et al, 2008). In this compound family isotope effects were generally normal for ¹⁵N (SI, Annex 2). An inverse effect was also observed for ¹⁵N in atrazine photooxidation, like for carbon (Hartenbach et al. 2008).

360 **3. Applications of pesticide ME-CSIA**

361 **3.1 Source identification, Forensics**

Contaminations of surface waters, groundwaters or springs by pesticides are relatively frequent and call usually for actions to identify the polluters. The isotopic compositions of pesticides might help here for a better identification of the pollution source, especially when pesticides from different manufacturers have distinct isotopic signatures. For instance, a C and N CSIA of glyphosate (Mogusu et al. 2015) revealed that glyphosate is a molecule with a highly variable isotopic composition, depending on the commercial formulation (Fig. 3).



Figure 3: Isotopic compositions of A) glyphosate from 13 manufacturers. Data from Mogusu et al.
2015, including also data from Kujawinski et al. 2013. For list of manufacturers see Supp. Inf.
Annex A. B) atrazine from 7 manufacturers. Data from Chen et al (2017), Masbou et al. (2018),
Meyer et al. (2008, 2013), Ponsin et al. 2019, Torrento et al. 2019+2021. Schreglmann et al. 2013.
C) metolachlor from 5 manufacturers. Data from Alvarez-Zaldivar et al. (2018), Masbou et al.
(2018) and Torrento et al (2021).

The δ^{13} C ratios of glyphosate from different sources span from -33.7 to -24.6 ‰, while those for δ^{15} N range from -1.8 to +3.3 ‰. Figure 3 shows that δ^{13} C alone bears already a good potential for discrimination between two manufacturers, and that the dual element isotope assessment may improve further distinction among distinct glyphosate sources. While the authors did not discuss why the manufacturing process leads to such a high variability, they concluded that this variability would allow forensic investigations in cases of pollutions of glyphosate or its key metabolite, AMPA.

Another famous legacy pesticide, atrazine, also features highly variable isotopic composition (Fig. 3B). However, the δ^{13} C ratios of atrazine span only from -26.3 to -30 ‰ except for the atrazine of Ehrenstorfer which is at - 19 ‰, while δ^{15} N ratios range from -2.2 to -0.2 ‰. This highlights that the power for discrimination of distinct atrazine sources is far below that of glyphosate. However, stable chlorine isotope analysis has been developed for atrazine (Ponsin et al. 2019). Further data on the stable chlorine isotope composition may help in the future to establish a more powerful basis for forensic investigations of atrazine pollution.

In contrast to glyphosate and atrazine, *S*-metolachlor analyzed from five manufacturers shows relatively little isotopic variation for carbon and nitrogen (Figure 3C), suggesting that the chlorine isotope analysis must be included for forensic studies relying on *S*-metolachlor (Torrento et al. 2021). If isotopic signatures of *S*-metolachor in commercial formulations do not vary significantly, transformation processes involving bond breaking may be followed up in the environment, independent of the used commercial formulation. This may allow catchment-scale evaluation of metolachlor transformation.

The largest forensic investigation of a pesticide so far found in literature concerned 26 different pure samples of the insecticide lindane (γ -Hexachlorocyclohexane), including three analyses of stable isotopes of carbon, chlorine and hydrogen (Ivdra et al. 2017). In addition, other samples from commercial formulations, soils or wastes were also analyzed, as well as other 400 hexachlorocyclohexane isomers. Lindane is a banned persistent organic compound which is still 401 found in many soils worldwide. Stable isotope ratios of hexachlorocyclohexanes isomers and 402 lindane covered the ranges from -233‰ to +1‰ for δ^2 H, from -35.9‰ to -22.7‰ for δ^{13} C, and 403 from -6.69‰ to +0.54‰ for δ^{37} Cl.

404 To conclude this section, an example is presented where the stable isotope composition of a 405 pesticide was used to elucidate a criminal food intoxication case, presented by Kawashima (2015). 406 In December 2007 and January 2008, people suffered from food intoxication in two provinces of 407 Japan after eating frozen dumplings (gyosa) imported from China. These dumplings contained very 408 high concentrations (1490-19,290 ppm) of methamidophos, an organophosphorus pesticide. The 409 δ^{13} C of methamidophos from seven Japanese formulations and a Chinese one was analyzed by GC-C-IRMS. The isotope signatures of methamidophos ranged from -31.9 to -49.2 ‰, potentially 410 411 sufficient to distinguish most products without the addition of a second element. As this study 412 permitted to clearly point the Chinese source for the food contamination, a worker in the Chinese 413 food factory was arrested and later confessed the injection of stolen methamidophos into the food. 414 The study highlights the interest to have a good trackability of pesticide sources as it is relatively 415 easy for criminals to obtain pesticides and use them for poisoning.

416 3.2 Process identification by ME-CSIA

417 A common feature of diffuse agricultural pollution is that soil microorganisms cannot or only partly 418 degrade organic pollutants after a long period of ageing in soils. This has been observed for 419 instance for several banned or restricted agrochemicals, such as the s-triazine herbicides atrazine 420 and simazine (Stucki et al. 1995), racemic metolachlor, the insecticide chlordecone (Martin-Laurent 421 et al. 2014) or the soil fumigant 1,2-dibromoethane (Steinberg et al. 1987). All listed compounds 422 are known to be degraded to different extents by microorganisms in pure cultures and soil 423 microcosms under laboratory conditions. In contrast, in agricultural soils, degradation extent and 424 pathways of degradation could often not be evidenced based on conventional monitoring methods, 425 the molecules persist over decades, and they are still detected in the underlying groundwater or 426 nearby surface waters. This phenomenon has been called dormant pollution and resurrection 427 (Charles et al. 2021), and is also observable for some industrial pollutants, such as chlorinated 428 hydrocarbons (Zimmermann et al. 2020).

429 The collected information on isotope fractionation values can be used to predict changes of 430 isotopic ratios during the advancement of pesticide dissipation in soils. This is best illustrated in multi-elemental plots, like in section 3.1, for glyphosate, atrazine and metolachlor, for both carbon
and nitrogen isotopes. Changes in the isotopic ratios for each process were calculated using the
approximate Rayleigh equation (eq. 2), assuming a removal of 90% of the initial mass

434
$$\Delta \delta^{13} C = \varepsilon_C \ln (f) \qquad \text{eq. 2}$$

Where f is the remaining mass fraction in soil (0.1 in our calculations). Δ is the isotope shift with respect to the initial isotope composition. For the three compounds, we calculated an average isotope composition for all manufacturers, and drew calculated isotope shifts with arrows away from this average composition (Figure 4). It is worth noting that the slope of these arrows in Figure 439 4 corresponds to Lambda, which is generally used in the literature on ME-CSIA as (eq. 3, here for nitrogen and carbon isotopes):

441
$$\Lambda_{N/C} = \frac{\epsilon_N}{\epsilon_C}$$
 eq.3

Lambda (Λ) expresses the relative magnitude of isotope enrichment of two elements, but does not express absolute magnitude of enrichment, i.e., whether the final isotope ratios expected in samples which underwent a removal process are far away from the initial composition, or not. The Λ value may also be partly elusive since a process involving two inverse isotope effects for both elements may have the same Λ as a process with two normal isotope effects. Hence, tables of Λ were not printed in this review, but all Λ values can easily be calculated from the Excel sheet in annex 2.



- 450 **Figure 4:** Change of stable isotope ratios (¹⁵N vs. ¹³C) for selected processes acting on average
- 451 glyphosate (A), atrazine (B) and metolachlor (C). Grey shaded points represent variable
- 452 compositions from different manufacturers, already shown in Figure 3. Note that photolysis at
- 453 natural sunlight does not create an isotope shift for atrazine (Drouin et al, 2021). KMnO₄:
- 454 Potassium permanganate

For glyphosate, only the abiotic oxidation with KMnO₄ was characterized for fractionation of both stable carbon and nitrogen isotopes (Fig. 4A redrawn from (Mogusu et al. 2015)). This oxidation leads to a strong N enrichment, with a much higher shift than the range of variability of different δ^{15} N. The nitrogen isotope analysis may confirm the abiotic oxidation in soil even if the initial composition of the glyphosate was not known. In contrast, the carbon enrichment is small for abiotic oxidation, and carbon isotope analysis would not be able to confirm this removal process in soil, especially when the initial δ^{13} C of the glyphosate was not known.

462 The C and N isotope enrichment of atrazine was studied for various processes (Torrento et al. 463 2021; Drouin et al. 2021; Meyer et al. 2014; Meyer and Elsner 2013; Hartenbach et al. 2008), and 464 for a range of pure bacterial strains known to degrade atrazine (Schürner et al. 2015; Chen et al. 465 2017; Chen et al. 2019, Fig. 4B). Generally large carbon isotope shifts are expected after 90% 466 atrazine degradation, whereas nitrogen isotope shifts remain modest. However, nitrogen shifts are 467 either positive or negative, which can help for process identification. But the slopes of the isotope 468 shifts between various abiotic and biotic degradations of atrazine are not very distinctive, and for a 469 correct process identification one should add also stable chlorine isotope analysis (Ponsin et al. 470 2019).

471 For metolachlor removal processes, large C and N isotope shifts are illustrated in Fig. 4C, with 472 data from (Drouin et al. 2021; Masbou et al. 2018; Sun et al. 2021; Torrento et al. 2021). Photolysis 473 under UV lamp and alkaline hydrolysis result in well-distinct isotope fractionation patterns in Figure 474 4C, while acid hydrolysis, soil degradation and diffusion in water do not differ in their slope Λ . They 475 differ, however, in the magnitude of carbon isotope shifts: diffusion does almost not fractionate, 476 and soil biodegradation leads to lower carbon isotope shifts than acid hydrolysis. To tease apart 477 those two processes, additional analyses are needed. It is also worth noting that enantiomers of 478 chiral pesticides, such as metolachlor, have identical physico-chemical properties, and transport 479 processes or abiotic reactions appear not to change enantiomer proportions (Bidleman et al. 480 2013). In contrast, degradation of chiral pesticides may be mainly microbiologically mediated and 481 enantioselective (Liu et al. 2005). Because degradation rates often differ among enantiomers and

result in deviation from racemic ratios, changes in the enantiomeric fraction (EF) of metolachlor may be used to follow degradation (Hegeman and Laane, 2002; Milosevic et al, 2013). In particular, combining CSIA and enantioselective analysis techniques, enantioselective stable isotope analysis (ESIA) may enhance the evaluation of sources and transformation processes of chiral micropollutants in the environment (Badea and Danet, 2015).

487 3.3 Identification and modeling of pesticide degradation in soil and across the soil-hillslope 488 water continuum

489 Isotope data derived from ME-CSIA require adapted modeling approaches to interpret and predict 490 the persistence and the transport of pesticides in soil and water. Reactive transport models can 491 support this deployment as they explore numerically and quantitatively the interactions of physical. 492 chemical and biological processes in surface and subsurface systems (Li et al. 2017, Prommer et 493 al. 2019). Historically, the first reactive transport models integrating isotopic signatures aimed at 494 interpreting monitored natural attenuation of legacy industrial pollutants in aquifers, e.g. (Hunkeler 495 et al. 2008; Thouement et al. 2019; Antelmi et al. 2021) or in wetlands e.g. (Alvarez-Zaldivara et al. 496 2016). Some reactive transport models incorporating isotope fractionation also derived the 497 physico-chemical conditions, including redox conditions, dissolved oxygen, temperature, or pH, to 498 predict pollutant degradation and associated isotopic fractionation across the aquifer domain 499 (D'Affonseca et al. 2011).

500 While soil reactive transport models evaluating pesticide persistence and transport exist since 501 decades (Gassmann 2021), models including pesticides CSIA or ME-CSIA data remain scarce. Indeed, ME-CSIA of pesticides was limited and no isotopic dataset existed in the early 2010. 502 503 Nevertheless, the virtual 2D hillslope reactive transport model HydroGeoSphere, including pesticide CSIA data (δ^{13} C and δ^{2} H) across the soil-hillslope-waters continuum (Lutz et al. 2013), 504 505 highlighted the potential to examine pesticide degradation on the catchment scale. Worthy of note, 506 pesticide transport was simplified in HydroGeoSphere by neglecting volatilization and sorption 507 processes, assuming soluble, non-volatile and non-sorptive pesticides. A main limitation of lumped 508 approaches, i.e., considering the catchment as an undivided entity (Surdyk et al. 2021), is to 509 neglect spatial information, including soil heterogeneity, water content or temperature, hampering 510 the capacity to identify degradation hot spot across the catchment (Grundmann et al. 2007). Combining carbon CSIA data (δ^{13} C) of the herbicide S-metolachlor in a river and a parsimonious 511 512 lumped agro-hydrological model may, however, improve interpretation of pesticide transport and 513 degradation at the catchment scale (Lutz et al. 2017). Complementarily to a soil-hillslope-waters

scale approach, modeling of water-sediment phase transfer or mass transfer limited transport
supported lab-scale biodegradation of pesticides with CSIA for complex systems, including
groundwater (Sun et al. 2021) or river-sediment interface (Droz et al. 2021).

517 There are several reasons explaining the limited development and use of soil reactive transport 518 models incorporating pesticide ME-CSIA data, which are also opportunities (Table 1). First, an 519 isotopic signature database of pesticides in commercial formulations is necessary for ME-CSIA in 520 soil reactive transport models to identify sources and processes (see sections 2.3; Table 1). If 546 521 isotopic compositions of pesticides were found in the literature, only 13% of them were associated 522 with commercial formulations. These isotopic compositions of applied pesticides can differ from 523 analytical standards often used for reference laboratory experiments (see section 2.3). Whenever 524 the range of δ^{13} C of the applied formulations (Figure 3) is larger than $\Delta\delta^{13}$ C during pesticide 525 degradation, it becomes difficult to evaluate the contribution of degradation in soil or water. At the 526 field scale, a unique commercial formulation may be used, allowing an evaluation of degradation (e.g., Alvarez-Zaldivar et al, 2018). At larger scale, e.g., from a few hectares to hundred km², a 527 survey of the main commercial formulations is required to constrain δ^{13} C and δ^{15} N initial values of 528 529 target pesticides and and to reduce the uncertainty to identify pesticide sources and dissipation.

530 In addition, estimating isotope fractionation values (e.g., ε_c) from reference laboratory experiments 531 under controlled conditions (e.g., a degradative process occurring under a specific redox condition) 532 is a preliminary requirement to evaluate in situ degradation (Centler et al. 2013; Fischer et al. 533 2016). However, the reference isotope fractionation values should be carefully selected prior to 534 interpreting and predicting *in situ* biodegradative processes in the field. It is thus essential that such 535 isotope fractionation values correspond to the established hydrochemical conditions in situ and the 536 corresponding degradation pathways. Bacterial isolates associated with micropollutant degradation 537 are less known than bacteria associated with degradation of industrial solvents (Elsner and Imfeld 538 2016). This severely limits the field interpretation of degradative processes and definition of 539 remediation strategies. It has been suggested that degradation of micropollutants may rather occur 540 via co-metabolism, where enzymes may no longer be harbored by a single microorganism only but 541 may be distributed in a range of microbial populations constituting the microbial community (Fenner et al. 2021). Therefore, microcosm experiments with the targeted soil and pesticide remain 542 543 an essential step identify the degradation mechanisms and to derive isotope fractionation values 544 associated with complex degrading microbial communities for in situ application of ME-CSIA 545 (Alvarez-Zaldivar et al. 2018). Additional characterization of transformation products of the targeted 546 pesticides may also help identify ongoing degradation pathways both in the reference microcosm547 experiments and at the field scale.

- 548 In addition, degradation is generally represented with a lumped soil half-life parameter, hampering
- 549 the incorporation of ME-CSIA data in soil reactive transport model codes (Table 1).

Table 1: Challenges, bottlenecks and opportunities to identify and predict pesticides degradation in

soil with CSIA and ME-CSIA in soil and across soil-hillslope-water continuum.

Challenge	Bottleneck	Opportunities
$\begin{array}{rllllllllllllllllllllllllllllllllllll$	Limited knowledge of isotopic signatures of pesticides in commercial products	 i) Isotopic signature database for commercial formulations of applied and legacy pesticides, ii) Characterization strategy from field to catchment scales
ProcesseidentificationsValidity of referencestableisotopefractionationfactors($\epsilon_{C}, \epsilon_{N}, \epsilon_{Cl,}$)?	Transposing reference stable isotope fractionation factors to interpret pesticide degradation in soil in relation with soil characteristics	Evaluating the impact of soil physico- chemical and microbial conditions on stable isotope fractionation for field estimation of pesticide degradation
Reactivity - Reactivity - Open the degradation black- box in soil reactive transport models by considering impact of soil bio-physico- chemical properties and by differentiating degradation pathways?	Degradation in soil reactive transport models as a lumped process with one soil half-life parameter	i) Introducing in soil reactive transport models the effect of soil temperature and moisture and the soil biological activity on degradation intensity, ii) considering different degradation pathways in soil reactive transport models

Transport - IntegrateMischaracterization ofpreferential flows inpreferential flow in soilsoil reactive transportreactive transportmodels forbetterand their impacts onpredictionofpesticidestransittime?

i) in situ measurements with geophysical techniques and tracer experiments for soil profiles, ii) geostatistical methods, pedotransfer functions or meta-models to parametrize preferential flow in soil reactive transport models

Therefore, the use of a single reference isotope fractionation factor (see sections 3.1 and 3.2) may bias the interpretation of pesticide degradation by neglecting multiple and possibly simultaneous degradative processes in soil. Here, the effect of soil temperature and moisture and the soil biological activity on degradation intensity may be introduced in soil reactive transport models to strengthen and expand the description of degradation within the catchment and across the hydrological or growing season (Gassmann 2021), even when degradation is considered as a lumped process.

559 Finally, a major interest of introducing CSIA data in soil reactive transport models is to prove, 560 characterize and quantify pesticide transformation. This, however, depends on the ability to 561 interpret the transport of water and pesticides by preferential and matrix flows (Jarvis et al. 2012; 562 Köhne et al. 2009a+b; Beven and Germann 2013; Gassmann et al. 2015; Weiler 2017). Including 563 preferential flow framework in soil reactive transport models may help to improve prediction of 564 transit times in soil. Here, CSIA may further help tease apart fast flows, mobilizing "fresh" pools of 565 pesticides with isotopic signatures mirroring that of the applied products, from the "aged" pools of 566 pesticides in soil, reflecting degradation in the soil matrix. In situ measurements of macropores, 567 e.g., earthworm channels, soil pipes and larger soil cracks, in the soil profile may help identifying 568 preferential flow and providing parameter values for soil reactive transport models using 569 geophysical techniques and tracer experiments (Weiler 2017). At the hillslope and catchment 570 scales, geostatistical methods, pedotransfer functions or meta-models, i.e., simplified statistical 571 models built with 1D soil reactive transport models such as MACRO (Lindahl et al. 2008), may 572 provide spatially distributed parameters. This, in turn, allows predicting preferential and soil matrix 573 flows in Horizontal 2D soil reactive transport models with associated pesticide loads (Jarvis et al. 574 2012).

575 4. Conclusions and perspectives

576 ME-CSIA was originally developed for industrial pollutants, which are frequently encountered in 577 groundwater at concentrations often in the mg/L range. These organic pollutants have often 578 relatively small molar masses and are therefore prone to undergo relatively large isotope 579 fractionations. ME-CSIA applied to pesticides was originally thought to be hardly possible, 580 especially since they occur mostly in the µg/L range as diffuse sources and are rather large 581 molecules which might exhibit little isotope fractionation. However, this review shows that large 582 current and future improvements of extraction techniques of pesticides from soils, combined with 583 lower limits of detection of analytical techniques, and new analytical methods make it possible to 584 apply ME-CSIA to study, follow and predict pesticide degradation in soil. It is shown also that this 585 opens ways to model the fate of pesticides on the catchment scale, which may help to design new 586 management practices.

587 While a lot of recent literature summarized in this work have significantly advanced science to use 588 ME-CSIA for a better understanding of the fate of pesticides at the field scale, prominent research 589 gaps still remain that must be tackled in the future.

590 First, our database of isotopic compositions of pesticides is strongly lacking analyses of pesticides 591 from commercial formulations available on the market worldwide. More work is thus needed to 592 systematically analyze commercial formulations from different manufacturers and to study their 593 variability from past to present. In the framework of the French project who is funding this research, 594 over 120 commercial formulations available on the French market of pesticides have been 595 collected and are currently analysed for better insight to be provided in a follow-up publication.

596 In addition, the database of isotope fractionation values of pesticides during degradation processes 597 has mainly been built on laboratory experiments performed at relatively high concentrations of 598 active molecules. Processes in real soils, in contrast, occur mostly at low concentrations. More 599 work is thus needed to investigate whether at low concentrations the same fractionation patterns 600 occur as at high concentrations. Depending on whether substrate uptake into the microbial cell, or 601 slow enzyme kinetics is rate-determining, the extent of stable isotope fractionation may also be 602 smaller than expected (Elsner and Imfeld, 2016). More reference biodegradation studies involving 603 bioavailability limitations and aging of residues in soil should be performed.

Third, the analytical techniques so far developed are not equally good for all elements in pesticide molecules. As shown in the supplementary material, isotope analysis of ¹³C is well developed and practiced in many labs. Fewer but still abundant analyses were made for ¹⁵N and ³⁷Cl. For these three isotopes, accuracy is generally better than ± 0.5 ‰. For ²H, less measurements were made so far. The robustness of ²H analysis is hampered by the low natural abundance of deuterium and the presence of water vapor in mass spectrometers and yields at best an accuracy of ± 5 ‰. Furthermore, the presence of chlorine atoms in a pesticide can cause problems during the pyrolysis of the molecule before mass spectrometry of the produced H₂. Only very few studies of isotope analysis of other elements in pesticides, such as ⁸¹Br (1 study, Kuntze et al. 2016), ¹⁸O (1 study, Sandy et al. 2013), or ³⁴S (1 study, Schürner et al. 2015) were found in literature. Future research may foster the development of stable isotope analysis for these elements to strengthen the ME-CSIA approach and enhance the distinction between transformation pathways in both laboratory and field settings.

- 617 More fundamentally, the mechanistic understanding of isotope fractionation and related processes 618 should still be improved. Fractionation by diffusion has been studied in soil air (e.g. Bouchard et al. 2008), water (Jeannottat and Hunkeler, 2012; Sun et al. 2021) or clay (Wanner and Hunkeler, 619 620 2015). More knowledge should be gained on isotope fractionation of pesticides during diffusion in 621 narrow soil pores, since this may be affected by Knudsen diffusion, where molecules are subject to 622 collisions with pore walls, in addition to collisions with other gas or water molecules. Also, the 623 fundamental understanding of isotope fractionation by sorption/desorption during multi-step 624 partitioning and volatilization should be improved. For instance, a recent study by Zamane et al 625 (2020) showed quite large isotope effects during volatilization of toluene and propan-2-ol from 626 sandy soil.
- Finally, while stable isotopes are good markers of different sources of pollutants, and markers of sinks, it remains difficult to tease apart both multiple sources and fractionating sinks. Combined source apportionment and degradation quantification of organic pollutants with ME-CSIA has been shown to be possible in groundwater (Lutz van Breukelen, 2014) but its adaptation to soils has not yet been developed.

632 Supplementary Material:

Annex 1: Table of isotopic compositions of pesticides from Literature. Annex 2: Table of
enrichment factors from Literature. Both tables are integrated in the EXCEL sheet given in the
supplementary material on the journal website.

- 636 Acknowledgment
- 637
- We thank the editorial board and reviewers of ECLE and Maria PRIETO-ESPINOZA for makinghelpful suggestions for the text.
- 640
- 641 **Declarations**
- 642

- Funding : This work is funded by the French National research Agency ANR through grant ANR 18-CE04-0004-01, project DECISIVE.
- **Authors' contributions** GI, PH, SP + FML initiated this work by writing the project, DG + PH collected the literature data, and all authors wrote parts of the text.
- **Conflicts of interest/Competing interests** No conflicts or competing interests declared
- **Ethics approval** (not applicable)
- **Consent to participate** (not applicable)
- **Consent for publication** (not applicable)
- 651 Availability of data and material (Supplementary Material)
- **Code availability** (not applicable)

654 **5 References**

- Abdulra'uf L, Tan G (2014) Review of SBSE technique for the analysis of pesticide residues in
 fruits and vegetables. Chromatographia 77:15-24. https://doi.org/10.1007/s10337-0132566-8
- Aelion CM, Höhener P, Hunkeler D, Aravena R (2010) Environmental isotopes in
 biodegradation and bioremediation. CRC Press (Taylor and Francis), Boca Raton.
- Aeppli C, Holmstrand H, Andersson P, Gustafsson O (2010) Direct compound-specific stable
 chlorine isotope analysis of organic compounds with quadrupole GC/MS using standard
 isotope bracketing. Anal Chem 82:420-426. https://doi.org/10.1021/ac902445f
- Allen-King R, Grathwohl P, Ball W (2002) New modeling paradigms for the sorption of
 hydrophobic organic chemicals to heterogeneous carbonaceous matter in soils,
 sediments, and rocks. Adv Water Res 25:985-1016. https://doi.org/10.1016/S0309 1708(02)00045-3
- Alvarez-Zaldivara P, Centler F, Maier U, Thullner, M, Imfeld, G (2016) Biogeochemical
 modelling of in situ biodegradation and stable isotope fractionation of intermediate
 chloroethenes in a horizontal subsurface flow wetland. Ecol Eng 90:170–179.
 https://doi.org/10.1016/j.ecoleng.2016.01.0370
- Alvarez-Zaldivar P, Payraudeau S, Meite F, Masbou J, Imfeld G (2018) Pesticide degradation
 and export losses at the catchment scale: insights from compound-specific isotope
 analysis (CSIA). Water Res 139:198-207. https://doi.org/10.1016/j.watres.2018.03.061
- Anastassiades M, Lehotay S, Stajnbaher D, Schenck F (2003) Fast and easy multiresidue
 method employing acetonitrile extraction/partitioning and "dispersive solid-phase
 extraction" for the determination of pesticide residues in produce. J AOAC Internat 86:412 431. https://doi.org/0.1093/jaoac/86.2.412
- Annable W, Frape S, Shouakar-Stash O, Shanoff T, Drimmie R, Harvey F (2007) Cl-37, N-15,
 C-13 isotopic analysis of common agro-chemicals for identifying non-point source
 agricultural contaminants. App Geochem 22:1530-1536.
 https://doi.org/10.1016/j.apgeochem.2007.03.049

- Antelmi M, Mazzon P, Höhener P, et al (2021) Evaluation of MNA in a chlorinated solvents contaminated aquifer using reactive transport modeling coupled with isotopic fractionation
 analysis. Water 13:2945. <u>https://doi.org/10.3390/w13212945</u>
- Badea S, Vogt C, Gehre M, Fischer A, Danet A, Richnow H (2011) Development of an
 enantiomer-specific stable carbon isotope analysis (ESIA) method for assessing the fate
 of alpha-hexachlorocyclohexane in the environment. Rapid Comm Mass Spectrom
 25:1363- https://doi.org/1372 10.1002/rcm.4987
- Badea S, Danet A (2015) Enantioselective stable isotope analysis (ESIA) a new concept to
 evaluate the environmental fate of chiral organic contaminants. STOTEN 514:459-46.
 https://doi.org/10.1016/j.scitotenv.2015.01.082
- Bakkour R, Bolotin J, Sellergren B, Hofstetter T (2018) Molecularly imprinted polymers for
 compound-specific isotope analysis of polar organic micropollutants in aquatic
 environments. Anal Chem 90:7292-7301. https://doi.org/10.1021/acs.analchem.8b00493
- 695 Bernstein A, Shouakar-Stash O, Ebert K, Laskov C, Hunkeler D, Jeannottat S (2011) 696 Compound-specific chlorine isotope analysis: comparison of а gas 697 chromatography/isotope ratio mass spectrometry and gas chromatography/guadrupole 698 mass spectrometry methods in an interlaboratory study. Anal Chem 83:7624-7634. https://doi.org/10.1021/ac200516c 699
- Beven K, Germann P (2013) Macropores and water flow in soils revisited. Water Resour Res
 49:3071-3092. https://doi.org/10.1002/wrcr.20156
- Bidleman, TF., Jantunen LM, Binnur Kurt-Karakus, P, Wong, F, Hung, H, Ma, J, Stern, G,
 Rosenberg, B (2013) Chiral chemicals as tracers of atmospheric sources and fate
 processes in a world of changing climate, Mass Spectrom. (Tokyo) 2.
 <u>https://doi.org/10.5702/massspectrometry.S0019</u>
- Bouchard D, Hunkeler D, Gaganis P, Aravena R, Höhener P, Kjeldsen P (2008) Carbon
 isotope fractionation during migration of petroleum hydrocarbon vapors in the unsaturated
 zone: field experiment at værløse airbase, denmark, and modeling. Environ Sci Technol
 42:596-601. <u>https://doi.org/10.1021/es070718f</u>

Brock TCM, Arts GHP, Maltby L, Van den Brink PJ (2009) Aquatic risks of pesticides,
ecological protection goals, and common aims in European Union legislation. Integr Envir
Assess Manag, 2, p e20-e46. https://doi.org/10.1002/ieam.5630020402Bruzzoniti M,
Checchini L, De Carlo R, Orlandini S, Rivoira L, Del Bubba M (2014) QuEChERS sample
preparation for the determination of pesticides and other organic residues in
environmental matrices: a critical review. Anal Bioanal Chem 406:4089-4116.
https://doi.org/10.1007/s00216-014-7798-4

- Buszewski B, Szultka M (2012) Past, present, and future of solid phase extraction: a review.
 Crit Rev Anal Chem 42:198-213. https://doi.org/10.1080/07373937.2011.645413
- 720 Carson R. (1962) Silent spring. Penguin.
- Centler F, Hesse F, Thullner M (2013) Estimating pathway-specific contributions to
 biodegradation in aquifers based on dual isotope analysis: Theoretical analysis and
 reactive transport simulations. J Contam Hydrol 152:97-116.
 <u>https://doi.org/10.1016/j.jconhyd.2013.06.009</u>
- 725 Charles M, Pierre S, Olivier E et al. (2021) Pesticide resurrection. Environ Chem Lett.
 726 https://doi.org/10.1007/s10311-021-01347-z
- Chen S, Yang P, Kumar J, Liu Y, Ma L (2017) Inconsistent carbon and nitrogen isotope
 fractionation in the biotransformation of Atrazine by Ensifer sp cx-t and Sinorhizobium sp
 k. Internat Biodeterioration & Biodegradation 125:170-176
 https://doi.org/10.1016/j.ibiod.2017.09.014
- Chen S, Zhang K, Jha R, Chen C, Yu H, Liu Y (2019) Isotope fractionation in atrazine
 degradation reveals rate-limiting, energy-dependent transport across the cell membrane of
 gram-negative Rhizobium sp. cx-z. Environ Pollut 248:857-864
 https://doi.org/10.1016/j.envpol.2019.02.078
- Chevallier M, Cooper M, Kümmel S, Barbance A, Le Paslier D, Richnow H, Saaidi PL, Adrian
 L (2018) Distinct carbon isotope fractionation signatures during biotic and abiotic reductive
 transformation of chlordecone. Environ Sci Technol 52:3615-3624. https://doi.org/ 10
 https://doi.org/.1021/acs.est.7b05394

- Correia-Sa L, Fernandes V, Carvalho M, Calhau C, Domingues V, Delerue-Matos C (2012)
 Optimization of QuEChERS method for the analysis of organochlorine pesticides in soils
 with diverse organic matter. J Separat Sci 35:1521-1530. https://doi.org/
 10.1002/jssc.201200087
- D'Affonseca FM, Prommer H, Finkel M, et al (2011) Modeling the long-term and transient
 evolution of biogeochemical and isotopic signatures in coal tar-contaminated aquifers.
 Water Resour Res 47:W05518. https://doi.org/10.1029/2010WR009108
- Drenzek N, Tarr C, Eglinton T, Heraty L, Sturchio N, Shiner V (2002) Stable chlorine and
 carbon isotopic compositions of selected semi-volatile organochlorine compounds. Org
 Geochem 33:437-444. https://doi.org/10.1016/S0146-6380(02)00004-9
- Drenzek N, Eglinton T, Wirsen C, Sturchio N, Heraty L, Sowers K (2004) Invariant chlorine
 isotopic signatures during microbial PCB reductive dechlorination. Environ Poll 128:445448. https://doi.org/10.1016/j.envpol.2003.09.006
- Droz B, Drouin G, Maurer L, Villette C, Payraudeau S, Imfeld G (2021) Phase Transfer and
 Biodegradation of Pesticides in Water-Sediment Systems Explored by Compound-Specific
 Isotope Analysis and Conceptual Modeling. Environ Sci Technol 55:4720-4728.
 <u>https://doi.org/10.1021/acs.est.0c06283</u>
- Drouin G, Droz B, Leresche F, Payraudeau S, Masbou J, Imfeld G (2021) Direct and indirect
 photodegradation of atrazine and S-metolachlor in agriculturally impacted surface water
 and associated C and N isotope fractionation. Environ Sci-Processes & Impacts 23:1791 1802.
- 760 https://doi.org/10.33774/chemrxiv-2021-sg66n
- Elsner M, Imfeld G (2016) Compound-specific isotope analysis (CSIA) of micropollutants in the
 environment: current developments and future challenges. Current Opinion in
 Biotechnology 41:60-72. https://doi.org/10.1016/j.copbio.2016.04.014
- Farooq S, Nie J, Cheng Y, Yan Z, Li J, Shah Bacha SA, Mushtaqa A, Zhang H (2018)
 Molecularly imprinted polymers' application in pesticide residue detection. Analyst, 2018,
 143, 3971.
- 767 https://doi.org/10.1039/C8AN00907D

- Fenner K, Canonica S, Wackett L, Elsner M (2013) Evaluating pesticide degradation in the
 environment: blind spots and emerging opportunities. Science 341:752-758
 https://doi.org/10.1126/science.1236281
- Fenner K, Elsner M, Lueders T, et al (2021) Methodological advances to study contaminant
 biotransformation: new prospects for understanding and reducing environmental
 persistence? ACS EST Water 1:1541-1554.
 https://doi.org/10.1021/acsestwater.1c00025
- Fischer A, Manefield M, Bombach P (2016) Application of stable isotope tools for evaluating
 natural and stimulated biodegradation of organic pollutants in field studies. Curr Opin
 Biotechnol 41:99–107. https://doi.org/10.1016/j.copbio.2016.04.026
- Gassmann M (2021) Modelling the fate of pesticide transformation products from plot to
 catchment scale-state of knowledge and future challenges. Front Environ Sci 9:717738.
 https://doi.org/10.3389/fenvs.2021.717738
- Gassmann M, Olsson O, Stamm C, Weiler, M, Kümmerer, K (2015) Physico-chemical
 characteristics affect the spatial distribution of pesticide and transformation product loss to
 an agricultural brook. Sci Total Environ 532:733-743.
 https://doi.org/10.1016/j.scitotenv.2015.06.068
- Grathwohl P, Reinhard M (1993) Desorption of trichloroethylene in aquifer material rate
 limitation at the grain scale. Environ Sci Technol 27:2360-2366.
 <u>https://doi.org/10.1021/es00048a008</u>
- Grundmann S, Fuss R, Schmid M, et al (2007) Application of microbial hot spots enhances
 pesticide degradation in soils. Chemosphere 68:511-517.
 https://doi.org/10.1016/j.chemosphere.2006.12.065
- Hartenbach AE, Hofstetter TB, Tentscher PR, Canonica S, Berg M, Schwarzenbach RP
 (2008) Carbon, hydrogen, and nitrogen isotope fractionation during light-induced
 transformations of Atrazine. Environ Sci Technol 42:7751-7756.
 https://doi.org/10.1021/es800356h
- Hegeman, WJ, Laane, RW (2002) Enantiomeric enrichment of chiral pesticides in the
 environment. Rev. Environ. Contam. Toxicol. 173:85-116.

- Holmstrand H, Andersson P, Gustafsson O (2004) Chlorine isotope analysis of submicromole
 organochlorine samples by sealed tube combustion and thermal ionization mass
 spectrometry. Anal Chem 76:2336-2342. https://doi.org/10.1021/ac0354802
- Horst A, Renpenning J, Richnow H, Gehre M (2017) Compound specific stable chlorine
 isotopic analysis of volatile aliphatic compounds using gas chromatography hyphenated
 with multiple collector inductively coupled plasma mass spectrometry. Anal Chem
 803 89:9131-9138. https://doi.org/10.1021/acs.analchem.7b01875
- Hunkeler D, Aravena R, Cox E (2002) Carbon isotopes as a tool to evaluate the origin and fate
 of vinyl chloride: laboratory experiments and modeling of isotope evolution. Environ Sci
 Technol 36:3378-3384. https://doi.org/10.1021/es011479d
- Hunkeler D, Meckenstock RU, Sherwood Lollar B, Schmidt TC, Wilson J (2008) A guide for
 assessing biodegradation and source identification of organic ground water contaminants
 using compound specific isotope analysis (CSIA). Report EPA 600/R-08/148
 www.epa.gov/ada
- 811 Ivdra N, Herrero-Martin S, Fischer A (2014) Validation of user- and environmentally friendly
 812 extraction and clean-up methods for compound-specific stable carbon isotope analysis of
 813 organochlorine pesticides and their metabolites in soils. J Chromato A 1355:36-45.
 814 https://doi.org/10.1016/j.chroma.2014.06.014
- 815 Ivdra N, Fischer A, Herrero-Martin S, Giunta T, Bonifacie M, Richnow H (2017) Carbon,
 816 hydrogen and chlorine stable isotope fingerprinting for forensic investigations of
 817 hexachlorocyclohexanes. Environ Sci Technol 51:446-454.
 818 https://doi.org/10.1021/acs.est.6b03039
- Jarvis NJ, Moeys J, Koestel J, Hollis JM (2012) Preferential Flow in a Pedological Perspective.
 Elsevier Academic Press Inc, San Diego.
- Jeannottat S, Hunkeler D (2012) Chlorine and carbon isotopes fractionation during
 volatilization and diffusive transport of trichloroethene in the unsaturated zone. Environ Sci
 Technol 46:3169-3176. https://doi.org/10.1021/es203547p

- Kalathoor R, Zeiner M, Schmidt B. et al. (2015) First evidence for covalent linkage of acidic
 metabolites of metalaxyl and DDT as non-extractable pesticide residues in soil and
 sediment. Environ Chem Lett 13, 431-437. https://doi.org/10.1007/s10311-015-0514-6
- Kawashima H (2015) The measurement of stable carbon isotope ratios of eight
 Methamidophos samples. J Forensic Sci 60:1360-1364. https://doi.org/10.1111/15564029.12825
- Knossow N, Siebner H, Bernstein A (2020) Isotope analysis method for the herbicide
 Bromoxynil and its application to study photo-degradation processes. J Haz Mat
 388:122036.
- 833 https://doi.org/10.1016/j.jhazmat.2020.122036
- Köhne J, Köhne S, Simunek J (2009a) A review of model applications for structured soils: a)
 water flow and tracer transport. J Contam Hydrol 104:4-35.
 https://doi.org/10.1016/j.jconhyd.2008.10.002
- Köhne JM, Köhne S, Simunek J (2009b) A review of model applications for structured soils: b)
 Pesticide transport. J Contam Hydrol 104:36-60.
 https://doi.org/10.1016/j.jconhyd.2008.10.003
- 840 Kujawinski DM, Wolbert JB, Zhang LJ, Jochmann MA, Widory D, Baran N (2013) Carbon 841 isotope ratio measurements of Glyphosate and AMPA by liquid chromatography coupled 842 isotope ratio mass spectrometry. Anal Bioanal Chem 405:2869-2878. to 843 https://doi.org/10.1007/s00216-012-6669-0
- Kuntze K, Kozell A, Richnow H, Halicz L, Nijenhuis I, Gelman F (2016) Dual carbon-bromine
 stable isotope analysis allows distinguishing transformation pathways of ethylene
 dibromide. Environ Sci Technol 50:9855-9863. https://doi.org/10.1021/acs.est.6b01692
- Lewis KA, Tzilivakis J, Warner D, Green A (2016) An international database for pesticide risk assessments and management. Hum Ecol Risk Assess 22:1050-1064. https://doi.org/10.1080/10807039.2015.1133242
- Li L, Maher K, Navarre-Sitchler A, et al (2017) Expanding the role of reactive transport models in critical zone processes. Earth-Sci Rev 165:280-301. <u>https://doi.org/10.1016/j.earscirev.2016.09.001</u>

- Lichtfouse, E (2000) Compound-specific isotope analysis. Application to archaelogy, biomedical sciences, biosynthesis, environment, extraterrestrial chemistry, food science, forensic science, humic substances, microbiology, organic geochemistry, soil science and sport. Rapid Comm Mass Spectrom, 14:1337- 1344. https://doi.org/10.1002/1097-0231
- Lihl C, Renpenning J, Kümmel S, Gelman F, Schürner H, Daubmeier M (2019) Toward improved accuracy in chlorine isotope analysis: synthesis routes for in-house standards and characterization via complementary mass spectrometry methods. Anal Chem 91:12290-12297. https://doi.org/10.1021/acs.analchem.9b02463
- Lindahl AML, Soderstrom M, Jarvis N (2008) Influence of input uncertainty on prediction of within-field pesticide leaching risks. J Contam Hydrol 98:106-114. https://doi.org/10.1016/j.jconhyd.2008.03.006
- Liu, W, Gan J, Schlenk, D, Jury, WA (2005) Enantioselectivity in environmental safety of current chiral insecticides, Proc National Acad Sci, 102:701-706. https://doi.org/10.1073/pnas.0408847102
- Liu Y, Liu J, Renpenning J, Nijenhuis I, Richnow H (2020) Dual C-Cl isotope analysis for
 characterizing the reductive dechlorination of alpha- and gamma-hexachlorocyclohexane
 by two Dehalococcoides mccartyi strains and an enrichment culture. Environ Sci Technol
 54:7250-7260. https://doi.org/10.1021/acs.est.9b06407
- 871 Liu X, Li W, Kümmel S, Merbach I, Sood U, Gupta V (2021) Soil from a 872 hexachlorocyclohexane contaminated field site inoculates wheat in a pot experiment to 873 facilitate the microbial transformation of beta-hexachlorocyclohexane examined by 874 compound-specific isotope analysis. Environ Sci Technol 55:13812-13821. 875 https://doi.org/10.1021/acs.est.1c03322
- Lutz S, van Meerveld HJ, Waterloo MJ, et al (2013) A model-based assessment of the potential use of compound-specific stable isotope analysis in river monitoring of diffuse pesticide pollution. Hydrol Earth Syst Sci 17:4505-4524. https://doi.org/10.5194/hess-17-4505-2013
- Lutz S, Van Breukelen B (2014) Combined source apportionment and degradation quantification of organic pollutants with CSIA: 1. Model derivation. Environ Sci Technol

- 48:6220-6228.
- 883 https://doi.org/10.1021/es405400w
- Lutz S, van der Velde Y, Elsayed O, Imfeld G, Lefrancq M, Payraudeau S (2017) Pesticide fate on catchment scale: conceptual modelling of stream CSIA data. Hydrol Earth System Sci 21:5243-5261. https://doi.org/10.5194/hess-21-5243-2017
- Martin-Laurent F, Sahnoun MM, Merlin C, Vollmer G, Lubke M (2014) Detection and
 quantification of chlordecone in contaminated soils from the French west indies by GC-MS
 using the C-13(10)-chlordecone stable isotope as a tracer. Environ Sci and Pollut Res
 21:4928-4933. https://doi.org/10.1007/s11356-013-1839-y
- Masbou J, Meite F, Guyot B, Imfeld G (2018) Enantiomer-specific stable carbon isotope
 analysis (ESIA) to evaluate degradation of the chiral fungicide metalaxyl in soils. J Haz
 Mat 353:99-107. https://doi.org/10.1016/j.jhazmat.2018.03.047
- Mateo-Sagasta, J, Zadeh, SM, Turral, H, Burke, J (2017) Water pollution from agriculture: a
 global review. Executive summary IWMI FAO (book)
 <u>https://www.fao.org/3/ca0146en/CA0146EN.pdf</u>
- Melsbach A, Torrento C, Ponsin V, Bolotin J, Lachat L, Prasuhn V, Hofstetter T, Hunkeler D,
 Elsner M (2020) Dual-element isotope analysis of desphenylchloridazon to investigate its
 environmental fate in a systematic field study: a long-term lysimeter experiment. Environ
 Sci Technol 54:3929-3939, https://doi.org/10.1080/10256016.2020.1812599
- Meyer A, Penning H, Lowag H, Elsner M (2008) Precise and accurate compound specific
 carbon and nitrogen isotope analysis of Atrazine: critical role of combustion oven
 conditions. Environ Sci Technol 42:7757-7763. <u>https://doi.org/10.1021/es800534h</u>
- Meyer, A, Penning, H, Elsner, M (2009) C and N Isotope fractionation suggests similar
 mechanisms of microbial atrazine transformation despite involvement of different enzymes
 (AtzA and TrzN). Environ Sci Technol, 43: 8079-8085.
- 907 Meyer A, Elsner M (2013) C-13/C-12 and N-15/N-14 isotope analysis to characterize
 908 degradation of Atrazine: evidence from parent and daughter compound values. Environ
 909 Sci Technol 47:6884-6891. https://doi.org/10.1021/es305242q

- Meyer A, Dybala-Defratyka A, Alaimo P, Geronimo I, Sanchez A, Cramer C (2014)
 Cytochrome p450-catalyzed dealkylation of Atrazine by Rhodococcus sp strain n186/21
 involves hydrogen atom transfer rather than single electron transfer. Dalton Transactions
 43:12175-12186. https://doi.org/10.1039/c4dt00891j
- Milosevic N, Qiu S, Elsner M, Einsiedl F, Maier M, Bensch H (2013) Combined isotope and
 enantiomer analysis to assess the fate of phenoxy acids in a heterogeneous geologic
 setting at an old landfill. Water Res 47:637-649.
 https://doi.org/10.1016/j.watres.2012.10.029
- Mogusu E, Wolbert J, Kujawinski D, Jochmann M, Elsner M (2015) Dual element (N-15/N-14,
 C-13/C-12) isotope analysis of Glyphosate and AMPA by derivatization-gas
 chromatography isotope ratio mass spectrometry (GC/IRMS) combined with LC/IRMS.
 Anal Bioanal Chem 407:5249-5260. https://doi.org/10.1007/s00216-015-8721-3
- Penning H, Elsner M (2007) Intramolecular carbon and nitrogen isotope analysis by
 quantitative dry fragmentation of the phenylurea herbicide isoproturon in a combined
 injector/capillary reactor prior to GC separation. Anal Chem 79:8399-8405.
 https://doi.org/10.1021/ac071420a
- Penning H, Cramer C, Elsner M (2008) Rate-dependent carbon and nitrogen kinetic isotope
 fractionation in hydrolysis of isoproturon. Environ Sci Technol 42:7764-7771.
 https://doi.org/10.1021/es801101c
- 929 Perez-Rodriguez P, Schmitt A, Gangloff S, Masbou J, Imfeld G (2021) Plants affect the 930 dissipation and leaching of anilide pesticides in soil mesocosms: insights from compound-931 specific 308: isotope analysis (CSIA). Agri Ecosys Environ 107257. 932 https://doi.org/10.1016/j.agee.2020.107257
- Ponsin V, Torrento C, Lihl C, Elsner M, Hunkeler D (2019) Compound-specific chlorine isotope
 analysis of the herbicides Atrazine, Acetochlor, and Metolachlor. Anal. Chem 91:1429014298. https://doi.org/10.1021/acs.analchem.9b02497
- Prommer H, Sun J, Kocar BD (2019) Using reactive transport models to quantify and predict
 groundwater quality. Elements 15:87-92. https://doi.org/10.2138/gselements.15.2.87

- Reddy C, Drenzek N, Eglinton T, Heraty L, Sturchio N, Shiner V (2002) Stable chlorine
 intramolecular kinetic isotope effects from the abiotic dehydrochlorination of DDT. Env Sci
 Poll Research 9:183-186. https://doi.org/10.1007/BF02987486
- Reinnicke S, Bernstein A, Elsner M (2010) Small and reproducible isotope effects during
 methylation with trimethylsulfonium hydroxide (TMSH): a convenient derivatization method
 for isotope analysis of negatively charged molecules. Anal Chem 82:2013-2019.
 . https://doi.org/10.1021/ac902750s
- Reinnicke S, Simonsen A, Sorensen S, Aamand J, Elsner M (2012) C and N isotope
 fractionation during biodegradation of the pesticide metabolite 2,6-dichlorobenzamide
 (BAM): potential for environmental assessments. Environ Sci Technol 46:1447-1454.
 https://doi.org/10.1021/es203660g
- Roch F, Alexander M (1997) Inability of bacteria to degrade low concentrations of toluene in
 water. Env Tox Chem 16:1377-1383. https://doi.org/10.1002/etc.5620160707
- 951 Sakaguchi-Soder K, Jager J, Grund H, Matthaus F, Schuth C (2007) Monitoring and
 952 evaluation of dechlorination processes using compound-specific chlorine isotope analysis.
 953 Rapid Communications in Mass Spectrometry 21:3077-3084.
 954 https://doi.org/10.1002/rcm.3170
- Sandy E, Blake R, Chang S, Jun Y, Yu C (2013) Oxygen isotope signature of UV degradation
 of Glyphosate and phosphonoacetate: tracing sources and cycling of phosphonates. J
 Haz Mat 260:947-954. https://doi.org/10.1016/j.jhazmat.2013.06.057
- Santalad A, Zhou L, Shang F, Fitzpatrick D, Burakham R, Srijaranai S (2010) Micellar
 electrokinetic chromatography with amperometric detection and off-line solid-phase
 extraction for analysis of carbamate insecticides. J Chromato A 1217:5288-5297.
 https://doi.org/10.1016/j.chroma.2010.06.024
- 962Santrock J, Studley S, Hayes J (1985) Isotopic analyses based on the mass-spectrum of963carbon-dioxide. Anal Chem 57:1444-1448. https://doi.org/10.1021/ac00284a060
- Schimmelmann A, Qi H, Coplen T, Brand W, Fong J, Meier-Augenstein W (2016) Organic
 reference materials for hydrogen, carbon, and nitrogen stable isotope-ratio
 measurements: caffeines, n-alkanes, fatty acid methyl esters, glycines, l-valines,

967polyethylenes,andoils.AnalChem88:4294-4302.968https://doi.org/10.1021/acs.analchem.5b04392

- Schürner H, Seffernick J, Grzybkowska A, Dybala-Defratyka A, Wackett L, Elsner M (2015)
 Characteristic isotope fractionation patterns in s-triazine degradation have their origin in
 multiple protonation options in the s-triazine hydrolase TRZN. Environ Sci Technol
 49:3490-3498. https://doi.org/10.1021/es5055385
- 973 Schreglmann K, Hoeche M, Steinbeiss S, Reinnicke S, Elsner M (2013) Carbon and nitrogen
 974 isotope analysis of atrazine and desethylatrazine at sub-microgram per liter
 975 concentrations in groundwater. Anal Bioanal Chem 405:2857-2867.
 976 https://doi.org/10.1007/s00216-012-6616-0
- 977 Steinberg SM, Pignatello JJ, Sawhney BL (1987) Persistence of 1,2-dibromoethane in soils 978 entrapment in intraparticle micropores. Environ Sci Technol 21:1201-1208.
 979 <u>https://doi.org/10.1021/es00165a007</u>
- Stokstad, E., 2018. France's decade-old effort to slash pesticide use failed. Will a new attempt
 succeed? Science (80-october 17). <u>https://doi.org/10.1126/science.aav6762</u>
- Stucki G, Yu CW, Baumgartner T, Gonzalezvalero JF (1995) Microbial Atrazine mineralization
 under carbon limited and denitrifying conditions. Water Res 29:291-296.
 https://doi.org/10.1016/0043-1354(94)E0101-B
- Sun F, Mellage A, Gharasoo M, Melsbach A, Cao X, Zimmermann R, Griebler C, Thullner M,
 Cirpka OA, Elsner, M (2021) Mass-transfer-limited biodegradation at low concentrations.
 Evidence from reactive transport modeling of isotope profiles in a bench-scale aquifer.
 Environ Sci Technol 55:7386-7397. https://doi.org/10.1021/acs.est.0c08566
- Surdyk N, Gutierrez A, Baran N, Thiery D (2021) A lumped model to simulate nitrate
 concentration evolution in groundwater at catchment scale. J Hydrol 596:125696.
 https://doi.org/10.1016/j.jhydrol.2020.125696
- Tadeo J, Perez R, Albero B, Garcia-Valcarcel A, Sanchez-Brunete C (2012) Review of sample
 preparation techniques for the analysis of pesticide residues in soil. J AOAC International
 95:1258-1271. https://doi.org/10.5740/jaoacint.sge_tadeo

- Thouement H, Kuder T, Heimovaara T, van Breukelen B (2019) Do CSIA data from aquifers
 inform on natural degradation of chlorinated ethenes in aquitards? J Contam Hydrol 226:
 103520. https://doi.org/10.1016/j.jconhyd.2019.103520
- 998Torrento C, Bakkour R, Glauser G, Melsbach A, Ponsin V, Hofstetter TB (2019) Solid-phase999extraction method for stable isotope analysis of pesticides from large volume1000environmentalwatersamples.TheAnalyst144:2898.1001https://doi.org/10.1039/C9AN00160C
- 1002Torrento C, Ponsin V, Lihl C, Hofstetter T, Baran N, Elsner M (2021) Triple-element1003compound-specific stable isotope analysis (3d-csia): added value of cl isotope ratios to1004assess herbicide degradation. Environ Sci Technol 55:13891-13901.1005https://doi.org/10.1021/acs.est.1c03981
- 1006 Van Acker M, Shahar A, Young E, Coleman M (2006) GC/multiple collector-ICPMS method for
 1007 chlorine stable isotope analysis of chlorinated aliphatic hydrocarbons. Anal Chem
 1008 78:4663-4667. https://doi.org/10.1021/ac0602120
- 1009Vogel TM (1996) Bioaugmentation as a soil bioremediation approach. Curr Opin Biotechnol10107:311-316. https://doi.org/10.1016/s0958-1669(96)80036-x.
- 1011 Vogt C, Dorer C, Musat F, Richnow HH (2016) Multi-element isotope fractionation concepts to
 1012 characterize the biodegradation of hydrocarbons from enzymes to the environment. Curr
 1013 Opin Biotechnol 41:90-98. https://doi.org/10.1016/j.copbio.2016.04.027
- Wanner P, Hunkeler D (2015) Carbon and chlorine isotopologue fractionation of chlorinated
 hydrocarbons during diffusion in water and low permeability sediments. Geochim
 Cosmochim Acta 157:198-212. https://doi.org/10.1016/J.GCA.2015.02.034
- 1017 Weiler M (2017) Macropores and preferential flow-a love-hate relationship. Hydrol Process
 1018 31:15-19. https://doi.org/10.1002/hyp.11074
- 1019Wu L, Yao Y, Trebse P, Zhang N, Richnow H (2014) Compound specific isotope analysis of1020organophosphoruspesticides.Chemosphere111:458-446.1021https://doi.org/10.1016/j.chemosphere.2014.04.037

- Wu L, Liu Y, Liu X, Bajbaj A, Sharma M, Lal R (2019a) Fractionation approach to characterize
 the reactive transport processes governing the fate of hexachlorocyclohexanes at a
 contaminated site in India. Environ Int 219:132.
 https://doi.org/10.1016/j.envint.2019.105036
- Wu L, Moses S, Liu Y, Renpenning J, Richnow H (2019b) A concept for studying the
 transformation reaction of hexachlorocyclohexanes in food webs using multi-element
 compound-specific isotope analysis. Anal Chim Acta 1064:56-64.
 https://doi.org/10.1016/j.aca.2019.03.030
- Zamane S, Gori D, Höhener P (2020) Multistep partitioning causes significant stable carbon
 and hydrogen isotope effects during volatilization of toluene and propan-2-ol from
 unsaturated sandy aquifer sediment. Chemosphere 251:126345.
 https://doi.org/10.1016/j.chemosphere.2020.126345
- 1034Zimmermann J, Halloran L, Hunkeler D (2020) Tracking chlorinated contaminants in the1035subsurface using compound-specific chlorine isotope analysis: a review of principles,1036current challenges and applications.Chemosphere 244:125476.1037https://doi.org/10.1016/j.chemosphere.2019.125476