

A caged tris(2-pyridylmethyl)amine ligand equipped with a C triazole –H hydrogen bonding cavity

Gege Qiu, Donglin Diao, Leo Chaussy, Sabine Michaud-Chevallier, A. Jalila

Simaan, Paola Nava, Alexandre Martinez, Cédric Colomban

▶ To cite this version:

Gege Qiu, Donglin Diao, Leo Chaussy, Sabine Michaud-Chevallier, A. Jalila Simaan, et al.. A caged tris(2-pyridylmethyl)amine ligand equipped with a C triazole –H hydrogen bonding cavity. Dalton Transactions, 2022, 51 (28), pp.10702-10706. 10.1039/d2dt00607c . hal-03789423

HAL Id: hal-03789423 https://amu.hal.science/hal-03789423

Submitted on 29 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A caged tris(2-pyridylmethyl)amine ligand equipped with a C_{triazole}-H hydrogen bonding cavity.

Gege Qiu,^{‡a} Donglin Diao,^{‡a} Leo Chaussy,^a Sabine Michaud-Chevallier, ^a Paola Nava, ^a Alexandre Martinez^{*a} and Cédric Colomban^{*a}

A CAPPED BIOINSPIRED LIGAND BUILT FROM A TRIS(2-PYRIDYL-METHYL)AMINE (TPA) UNIT AND SURMOUNTED BY A TRIAZOLE-BASED INTRAMOLECULAR H-BONDING SECONDARY SPHERE, WAS PREPARED. THE RESULTING CAGE PROVIDES A WELL-DEFINED CAVITY COMBINING A HYDROPHOBIC NATURE WITH H-BONDING PROPERTIES. ITS COORDINATING PROPERTIES WERE EXPLORED USING ZN(II) AND CU(II) METAL IONS.

The binding cavities found in metalloproteins govern reaction's selectivity and efficiency. In their hydrophobic channels, destabilizing (like steric repulsion) and stabilizing (like hydrogen bonding, H-bonding) forces, allow for specific enzyme-substrate interactions, substrate positioning and activation/stabilization of highly reactive intermediates.¹ H-bonding is particularly important in metalloenzymes involved in dioxygen processing, such as copper-containing oxygenases and oxidases. The postulated oxidative active species of these systems is the highly reactive mononuclear cupric superoxide ($Cu^{II}-O2^{\bullet-}$).² Many efforts have been dedicated to the development of artificial Cu ligands able to generate and stabilize such metastable intermediate. Among them, the tris(2-pyridylmethyl)amine (**TPA**) ligand has been widely used as scaffold for mimicking the first coordination sphere in structural and functional models of copper,³ but also iron,⁴ mono-oxygenases. Interestingly, incorporating intramolecular H-bonding secondary spheres to the **TPA** ligand, was reported as an efficient strategy to stabilize either mononuclear hydroperoxo [(L)Cu^{II}-OOH]^{+,5} binuclear peroxodicopper [{(L)Cu^{II}}₂(O2²⁻)],⁶ or end-on superoxo [(L)Cu^{II}-O2^{•-}]⁷ copper-dioxygen intermediates.

On another side, synthetic supramolecular chemistry is a powerful tool to build cage-like second coordination sphere around bioinspired catalysts.⁸ In particular, the archetypal **TPA** ligand has been equipped with well-defined cavities by mean of its covalent substitution,^{9, 10} or host-guest encapsulation into an H-bonded capsule.¹¹ In this context, **TPA**-based hemicryptophanes are organic cages built from a bowl-shaped cyclotriveratrylene (CTV) cap, connected to the tripodal ligand via three linkers. We have recently demonstrated that **TPA**- hemicryptophanes displaying methylene or phenyl linkers, could respectively control the helical arrangement of the ligand,¹² and lead to enhanced oxidation catalysts.¹³ However, the hydrophobic cavities found in such cages were devoid of H-bonding groups that could allow substrate positioning or intermediate stabilization.

Despite these progresses, the preparation of **TPA**-based complexes combining hydrophobic cavities with intramolecular H-bonding units at their secondary sphere is still needed. Designing and discovering new methodologies to prepare such advanced model complexes is in fact crucial to better reproduce the key structural properties of metalloenzymes. Beside their connecting benefits, triazole bridges are particularly interesting due to their H-bonding donor properties. For instance, a triazolo organic cage has been recently reported as the most efficient chloride-binding receptor to date, by mean of $C_{triazole}$ -H H-bonding interactions.¹⁴ We therefore envisioned that the covalent substitution of the **TPA** by another C_3 symmetrical cap, using triazole spacers, will represent an efficient strategy to construct functionalized cavities.

In this communication, we report the preparation of unprecedented bioinspired complexes displaying a hydrophobic cavity offering three H-bonding triazoles, aiming at reproducing the metalloenzyme's functionalized hydrophobic channels. We design the hemicryptophane **Hm-TriA-TPA** where the archetypal **TPA** ligand is linked to a northern CTV cap, via three triazole bridges, resulting in an H-bond donor decorated cavity.

Hm-TriA-TPA was prepared in an eight-step synthetic strategy (Scheme 1). The cage's walls were first prepared (Scheme 1a), before generating the southern **TPA** and the CTV cap in a final intramolecular macrocyclization closing the structure (Scheme 1b). The aryl propargyl ether derivative **2** was prepared in twosteps by alkylation of the starting vanillyl alcohol with propargyl bromide, followed by the protection of the resulting alcohol **1** with THP. **2** was then connected to the pyridine derivative **4** by a triazole link formed in a Cu-catalyzed azide-alkyne cycloaddition reaction (CuAAC). The CuAAC reaction between equimolar amounts of the propargyl **2**, and the azide **4** precursors, catalyzed by CuSO₄ (10 mol%) in the presence of the sodium ascorbate reducing agent (10 mol%), resulted in the formation of the triazole **5** in 83% yield. Precursor **7** was then prepared in two steps by reduction of **5** into the alcohol **6** followed by it mesylation. The addition of ammonia to **7** in the presence of Cs₂CO₃, at 90°C in THF, afforded the open **TPA** derivative **8** in 73% yield. Formation of **Hm-TriA-TPA** was finally achieved in a 29% yield, via the intramolecular cyclization of **8** in CH₃CN, catalyzed by the Lewis acid scandium triflate [Sc(OTf₃)], under diluted conditions.



Scheme 1. (a) Synthesis of the pyridine-triazole precursor 5. (b) synthesis of the targeted hemicryptophane Hm-TriA-TPA.

The ¹H NMR spectrum of **Hm-TriA-TPA** indicates a C_3 symmetrical structure on average, in CDCI₃, at 298K (Fig. 1a). Identical, sharp and well-defined signals could be observed for the protons belonging to the northern CTV unit (H_h, H_i, H_j and H_{k,k'}), the -CH₂- links (H_{e,e'} and H_{g,g'}), the C_{triazole}-H bonds (H_f) and the southern **TPA** (H_{a,a'}, H_b, H_c and H_d). 2D-NMR experiments (see the ESI) were used to assign these resonances. Slow diffusion of Et₂O to a CH₂Cl₂ solution of **Hm-TriA-TPA** afforded single crystals suitable for X-ray diffraction. The XRD structure of **Hm-TriA-TPA** confirms the endohedral functionalization of the **TPA** unit by the bowl-shaped CTV via the three triazole bridges having their C_{triazole}-H bonds pointing toward the inside of the cavity (Fig. 1b). It should be noted that, in the X-ray structure of the cage, a pyridine unit of the **TPA** reside inside the cavity. This C₁ symmetrical conformation observed in the solid state, contrasts with the symmetrical ¹H NMR spectrum of **Hm-TriA-TPA** in solution. Fast conformational exchanges between in-out orientation of the pyridines could explain this behavior in solution at 298K.



Figure 1. (a) ¹H NMR spectra (CDCl₃, 400 MHz) of **Hm-TriA-TPA** along with (b) view of its X-ray crystal structure. Only the hydrogen atom belonging to the three triazole units have been included for clarity.

We have then investigated the ability of our caged ligand to form metallo-complexes in solution via coordination at its TPA unit. Binding of the air-stable and diamagnetic zinc triflate salt Zn^{II}(OTf)₂, was monitored by ¹H-NMR in CD₃CN at 298K (Fig. 2 and Fig. S6 ,ESI). The ¹H-NMR spectra of Hm-TriA-TPA in the presence of 0.5 equiv. of the zinc salt reveal two sets of signals for each protons of the cage, that could be attributed to the presence of ZnII(Hm-TriA-TPA)(OTf)2 and Hm-TriA-TPA in a 1:1 ratio (Fig. 2b). Interestingly, in the presence of a stoichiometric amount of the metal salt, the resonances belonging to the free cage fully disappear to the profit of the zinc complex signals, indicating full complexation. Upon metalation, a strong down-field shift occurs on the protons of the **TPA**'s pyridines (H_b , H_c and H_d , \Box_{ppm} : 0,35-0,7 ppm) that remain equivalent. The triazole bridges appear less affected with a modest upfield-shift observed for the Ctriazole-H bond (H_f, □ppm<0,03 ppm). Overall, the ¹H-NMR analysis of Zn^{II}(Hm-TriA-TPA)(OTf)₂ attests for retained C₃ symmetry of the caged ligand (on average) with identical and sharp signals for every resonances. Altogether, these observations unambiguously confirm the coordination of the Zn^{II} metal ion at the **TPA** unit with retention of the endohedral functionalization of the resulting complex.¹² This was further supported by the optimized DFT (Density Functional Theory) structure that clearly reveals a C₃ symmetrical caged Zn(II) complex in a trigonal bipyramidal geometry with an apical molecule of acetonitrile (Fig. 2e, Fig. S7, ESI). Finally, identical spectra were observed upon addition of a second equivalent of Zn^{II}(OTf)₂ (Fig. 2d), ruling out the possibility of a second metal-binding event occurring at the triazole crown.15



Figure 2. Monitoring of the ¹H NMR spectra (CD₃CN, 400 MHz) of **Hm-TriA-TPA** (a), upon addition of 0.5 equiv. (b), 1.0 equiv. (c) and 2.0 equiv. (d) of $Zn(OTf)_2$ (depicted region: 6.7 – 8.1 ppm). (e) DFT-optimized structure (PBE0-D3/def2-TZVP, COSMO, see ESI for full computational details) of $Zn^{II}(Hm-TriA-TPA)(OTf)_2$.

We have next investigated the possibility of preparing Cu^{II} complexes at our caged ligand. The **Cu^{II}(Hm-TriA-TPA)(OTf)**₂ complex was prepared by reacting the ligand with stoichiometric amounts of Cu^{II}(OTf)₂ in CH₃CN, at room temperature (see ESI). Formation of the targeted complex, in a 1:1 stoichiometry, was evidenced by High-Resolution Mass Spectrometry analysis (ESI-HRMS, Fig. S8, ESI). Furthermore, crystallization by slow diffusion of diethyl ether to a CH₃CN solution of the complex affords single crystals suitable for X-ray diffraction, allowing for determination of its solid-state structure. The structure of the **Cu^{II}(Hm-TriA-TPA)(OTf)**₂ complex shows a pentacoordinated copper center in a distorted square pyramidal geometry (Fig. 3a). The distorted tetragonal plane comprises the tertiary amine (N1), two pyridines (N2 and N3), and one triazole (N5) in *trans* to the tertiary amine with similar Cu-N bond length ranging from 1.978 Å to 2.055 Å. The coordination sphere is completed by an apical pyridine (N4) with a Cu-N distance of 2.225 Å. Importantly, a well-defined cavity, described by the northern CTV unit and the tris-triazole crown could be observed just above the Cu^{II} center confirming its endohedral functionalization. The latter is occupied by a non-bonded guest molecule of acetonitrile (solvent).

Could the Cu-triazole bond in Cu^{II}(Hm-TriA-TPA)(OTf)₂ be replaced by anion coordination at the Cu^{II}-TPA core ?

Anion binding at Cu-**TPA** derivatives has been reported to result in trigonal-bipyramidal Cu(II) complexes with axial binding of the anion in *trans* to the tertiary amine.¹⁶ Among them, azido adducts [(Cu^{II}-**TPA**^X)(N₃⁻)]⁺ have been described as electronic and structural analogues of cupric superoxide intermediates [(Cu^{II}-**TPA**^X)(O₂^{•-})]⁺. The N₃⁻ anion binds to Cu^{II}-**TPA** complexes with very similar bond lengths and angles (Cu^{II}-**TPA**^X)(O₂^{•-})]⁺. The N₃⁻ anion binds to Cu^{II}-**TPA** complexes with very similar bond lengths and angles (Cu^{II}-**TPA**^X)(O₂^{•-})]⁺. The N₃⁻ anion binds to Cu^{II}-**TPA** complexes (Cu^{II}-**TPA**^X)(N₃⁻)]⁺ have been extensively studied by electronic and vibrational spectroscopies.¹¹ They display typical azido \rightarrow Cu^{II} LMCT transitions, and \Box (N-N) stretching frequency. On this basis, we explored the possibility of binding N₃⁻ to our caged Cu(II) complex. UV-vis analysis of **Cu^{II}(Hm-TriA-TPA)(OTf)**₂, in acetone, reveal typical d-d transition centered at 634 nm (0,5 mM, \Box = 139 M⁻¹ cm⁻¹, Fig. S9, ESI). The UV-vis monitored addition of tetrabutylammonium azide NBu₄N₃ to a 0,5 mM solution of the caged complex, in acetone, results in the appearance of an intense absorption band centered at 415 nm (LMCT, \Box = 2068 M⁻¹cm⁻¹) and a d-d band at 713 nm (\Box = 249 M⁻¹ cm⁻¹, Fig. 3b). The final spectrum was identical to that of the azido adduct prepared using NaN₃ and isolated by precipitation (see ESI).



Figure 3. (a) Views of the X-ray crystal structure of $Cu^{II}(Hm-TriA-TPA)(OTf)_2$ (b) UV-vis monitoring of the formation of the azido adduct $Cu^{II}(Hm-TriA-TPA)(N_3)(OTf)$, upon addition of NBu_4N_3 (0 to 1.0 equiv.) to a 0,5 mM solution of $Cu^{II}(Hm-TriA-TPA)(OTf)$ in acetone. (c) spectroscopic features (UV-vis and IR) of the azido complexes stabilized by ^{Me3}TPA and Hm-TriA-TPA.

These UV-vis data are consistent with the formation of the azido complex $Cu^{II}(Hm-TriA-TPA)(OTf)(N_3)$.^{2, 7, 11} In TPAcomplexes bearing intramolecular H-bonding donors, the potential presence of stabilizing interactions with the azido moiety has been associated to a blue-shift of the LMCT band, as well as a blue-shift of the \Box (N-N) stretching frequency.^{2,17} Value of the LMCT band observed in the case of **Hm-TriA-TPA** was therefore compared with the one of its corresponding open model ligand ^{Me3}TPA devoid of intramolecular H-bonding groups (Fig 3c). Compared to the open **TPA**-based complex $Cu^{II}(M^{e3}TPA)(OTf)(N_3)$ ($\Box_{max} = 429nm$, Fig. S10, ESI), the $Cu^{II}-N_3^-$ LMCT bands in our triazolecontaining cage shift to higher energy ($\Box_{max} = 415nm$). Finally, to further support the stabilization offer by Hm-TriA-TPA, the two azido complexes have been isolated and they antisymmetric N_3^- IR stretch compared to the "base" ligand ^{Me3}TPA (Fig. 3c, Fig. S11 and S12, ESI). These findings are therefore consistent with a bonding stabilization of the azido adduct within the triazole-containing cavity.

Conclusions

In summary, the preparation of an organic cage where the canonical **TPA** ligand is surmounted by a Hbonding hydrophobic cavity offering three triazole units is described. We demonstrate that this cage can coordinate zinc(II) and copper(II) metal ions at its **TPA** unit with an endo-functionalization of the complex, in both solution and solid state. These are the first examples of bioinspired complexes equipped with a tristriazole decorated cavity mimicking the functionalized (H-bonding) hydrophobic channels of metalloenzymes. Finally, we found that the azidocopper(II) adduct $[(L)Cu^{II}-N_3^-]$ can be prepared upon addition of N₃ to the caged Cu(II) complex. Spectroscopic analysis of this structural analogue of the cupric superoxide intermediates $[(L)Cu^{II}-O_2^{\bullet-}]$, suggest stabilization of the azido adduct, within the C_{triazole}-H based cavity. We envision that our strategy might finds applications toward the development of non-enzymatic catalysts able to stabilize reactive intermediates and/or control substrate positioning by their H-bonding hydrophobic cavity. Future work will focus on the use of **TPA** ligands equipped with our triazolefunctionalized cavity to generate, stabilize, and explore the reactivity of end-on superoxocopper(II) complexes upon dioxygen activation.

Notes and references¹⁸

¹ Z. Wojdyla and T. Borowski, *Chem. Eur. J.*, 2022, DOI: 10.1002/chem.202104106

²² M. A. Ehudin, A. W. Schaefer, S. M. Adam, D. A. Quist, D. E. Diaz, J. A. Tang, E. I. Solomon and K. D. Karlin, *Chem. Sci.*, 2019, **10**, 2893-2905

³ S. Y. Quek, S. Debnath, S. Laxmi, M. van Gastel, T. Krämer and J. England, *J. Am. Chem. Soc.* 2021, **143**, 19731–19747

⁴ M. Borrell, E. Andris, R. Navrátil, J. Roithová and M. Costas, *Nat. Commun.*, 2019, **10**, 901

⁵ A. Wada, M. Harata, K. Hasegawa, K. Jitsukawa, H. Masuda, M. Mukai, T. Kitagawa, and Hisahiko Einaga, *Angew. Chem. Int. Ed.* 1998, **37**, 798-799

⁶ E. W. Dahl, H. T. Dong and N. K. Szymczak, *Chem. Commun.*, 2018, **54**, 892-895

⁷ (a) M. Bhadra, J. Yoon C. Lee, R. E. Cowley, S. Kim, M. A. Siegler, E. I. Solomon, and K. D. Karlin, *J. Am. Chem. Soc.* 2018, **140**, 9042–9045, (b) D. E. Diaz, D. A. Quist, A. E. Herzog, A. W. Schaefer, I. Kipouros, M.

Bhadra, E. I. Solomon, and K. D. Karlin, Angew. Chem. Int. Ed. 2019, 58, 17572–17576

⁸ (a) S. C. Bete, and M. Otte, Angew. Chem. Int. Ed. 2021, 60, 18582 –18586, (b) C. Colomban, V. Martin-Diaconescu, T. Parella, S. Goeb, C. García-Simón, J. Lloret-Fillol, M. Costas, and X. Ribas, Inorg. Chem., 2018, 57, 3529–3539

⁹ C. Bravin, E. Badetti, G. Licini, C. Zonta, *Coordination Chemistry Reviews*, 2021, **427**, 213558

¹⁰ N. Le Poul, B. Colasson, G. Thiabaud, D. Jeanne Dit Fouque, C. Iacobucci, A. Memboeuf, B. Douziech,

J. Řezáč, T. Prangé, A. de la Lande, O. Reinaud and Y. Le Mest, Chem. Sci., 2018, 9, 8282-8290

¹¹ T. Zhang, L. Le Corre, O. Reinaud, and B. Colasson, *Chem. Eur. J.* 2021, **27**, 434-443

¹² G. Qiu, C. Colomban, N. Vanthuyne, M. Giorgi and A. Martinez, *Chem. Commun.*, 2019, **55**, 14158–14161.

¹³ S. A. Ikbal, C. Colomban, D. Zhang, M. Delecluse, T. Brotin, V.Dufaud, J. P. Dutasta, A. B. Sorokin and A. Martinez, *Inorg. Chem.*, 2019, **58**, 7220-7228

¹⁴ Yun Liu, Wei Zhao, Chun-Hsing Chen and Amar H. Flood, *Science*, 2019, **365**, 159-161

¹⁵ Identical ¹H-NMR spectra was obtain after isolation of the Zn complex by precipitation with Et₂O (Fig. S6, ESI)

¹⁶ C. M. Moore and N. K. Szymczak, *Chem. Commun.*, 2015, **51**, 5490-5492

¹⁷ A. Wada, Y. Honda, S. Yamaguchi, S. Nagatomo, T. Kitagawa, K. Jitsukawa, and H. Masuda, *Inorg. Chem.* 2004, **43**, 5725-5735