



HAL
open science

Forces in T cell Immunology: examples, role and consequences

Farah Mustapha, Kheya Sengupta, Pierre-Henri Puech

► To cite this version:

Farah Mustapha, Kheya Sengupta, Pierre-Henri Puech. Forces in T cell Immunology: examples, role and consequences. Immunology and Cancer Biology, 2022, <https://videleaf.com/mechanical-forces-in-t-cell-biology/>. hal-03975398

HAL Id: hal-03975398

<https://amu.hal.science/hal-03975398>

Submitted on 21 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author Edited Manuscript

Forces in T cell Immunology: examples, role and consequences.

Running title: **Forces in T cell Immunology.**

Farah Mustapha ^{1,2,3,4,5}, Kheya Sengupta ^{4,5,*}, Pierre-henri Puech ^{1,2,3,4,*}

Affiliations:

Laboratoire Adhésion et Inflammation (LAI)

¹ Aix Marseille University, LAI UM 61, Marseille, F-13288, France.

² Inserm, UMR_S 1067, Marseille, F-13288, France.

³ CNRS, UMR 7333, Marseille, F-13288, France.

CENTURI

⁴ Turing Center for Living systems, Marseille, France

Centre Interdisciplinaire de Nanoscience de Marseille (CINAM)

⁵ CNRS - AMU UMR 7325, Marseille, F-13288, France

* To whom correspondence should be addressed : sengupta@cinam.univ-mrs.fr ; pierre-henri.puech@inserm.fr

Short abstract

Cell/cell interactions are a paradigm in cell biology since they establish, through biochemical and mechanical contacts, the structures of the living matter, allowing the cells core functions to take place, with broad impacts in tissue structure and function. While the biochemical aspects of these interactions have been intensively studied for decades, the mechanical aspects, known as mechano-transduction, have only more recently attracted the interest of biologists. Its importance is increasingly recognized as being instrumental to finely regulate cellular functions. Lymphocytes, either T or B, live and function as single cells, and do not assemble in a continuous tissue, while they need to transiently contact each other in a one-to-one manner to allow their efficient activation and subsequent action. These contacts are at the origin of some of the forces, and forces can strongly affect, and even potentiate, the effector functions of immune cells. In the present review, we will present the actors and scales, and discuss the modes of operations of this mechanobiology in the particular frame of immunological cells.

Keywords: Immune system, T cell, Forces, Mechanics, Mechanotransduction, TCR, Integrins

Looking back on history

In all forms of life, survival depends on the ability to adapt to environmental stresses, including mechanical stimuli such as external physical forces. It is a requirement so fundamental that it is at the core of all biological

designs; virtually all organisms have evolved structures from the macro (organs, tissues) to the micro (cells) and even down to the nanoscale (molecular assemblies, single proteins) that are not only sensitive, but also responsive to forces.

The biological effects of these forces are perhaps most evident in the context of physical structure and activity—the skeleton provides structural support to sustain the force of gravity. The skin provides a protective barrier that is maintained upon the application of external stretch. Even the simplest of physiological functions, such as respiration and circulation, require the generation of forces. This could explain why the earliest understanding and quantifications of these forces were focused on the organism and organ levels. In 1917, biologist D’Arcy Thompson published his book ‘On Growth and Form’, in which he discussed how mechanical forces contribute to the shape and size of living organisms (Thompson 1992). Near contemporaries of Thompson, Cecil Murray and Julius Wolff, proved respectively that shear stress controls the size of blood vessels (Murray 1926) and that mechanical loading increases the thickness and density of bone (Teichtahl et al. 2015).

This goes to show that the study of the interplay between physical forces and biological function dates back to well before the term ‘mechanobiology’ was even coined. Today, there is a general consensus that cells constantly sense the various mechanical cues (e.g. force, stress, strain, rigidity, topology and adhesiveness) of their micro-environment, *via* a process called ‘mechanosensing’. They then translate these cues into biochemical signals such as modified binding affinity, altered phosphorylation state, and/or a conformational change; a process called ‘mechanotransduction’.

These features are ubiquitous among different cell types and find themselves at the core of many physiological functions; in particular, it has been demonstrated that they are instrumental for key moments of immune cell life and function (Huse 2017). For decades, immunological research had focused on identifying the networks of secreted ligands, cell surface receptors, intracellular signaling pathways, and transcriptional factors mediating immune response (Murphy, Travers, and Walport 2008). These networks have been predominantly regarded as chemical in nature, largely because the individual molecules that make them up have been characterized by their non-covalent molecular interactions and/or enzymatic activity. Though this chemical description may not be incorrect, it neglects the influence of physical cues, in particular mechanical forces on molecular kinetics (Bell 1978) and signaling networks, as well as the influence of signaling networks on the mechanical environment within and outside the cell. Such an oversight becomes particularly relevant when studying immune cells whose lives are intensely “physical”: regularly deforming, migrating through tight interstitial spaces, adhering under shear flow, and forming stable interfaces (known as immunological synapses) with other cells (Zhang et al. 2020). Effectively, this means that the receptor-ligand interactions that govern immune cell function are likewise being subjected to and influenced by the same mechanically tempestuous microenvironment. And, given that several cell surface proteins (e.g. integrins) are known to be strongly connected to the actin cytoskeleton, which

is in turn connected to other intracellular proteins, this makes the molecular machinery involved in signal transduction ideal for relaying physical information about the extracellular environment into the cell, as well as translating biochemical signals inside the cell into physical forces exerted against the environment (Limozin and Puech 2019; Malissen and Bongrand 2015).




Fig. 1 Origins of forces, and orientation, at cellular and molecular scales in T cell recognition and function

Exemplifying the importance of mechanosensing and mechanotransduction in their development and function are T cells, key players of the adaptive immune system (Rossy, Laufer, and Legler 2018). Broadly speaking, T cells can be divided into three categories; Cytotoxic T cells that directly kill virally infected cells and cancer cells, and Helper and Regulatory T cells that activate and tune the effector functions of other cells in the immune system. In either case, T cells carry out the formidable task of identifying a particular cognate peptide bound to the major histocompatibility complex (pMHC) (Fig. 2A), against a very noisy environmental background of endogenous self-peptides MHCs, many of which involve the same MHC molecule (Huppa and Davis 2013). They do so even though the T cell receptors (TCRs) are cross-reactive and typically low in affinity when measured in isolation. One would expect that such high-fidelity decisions would be time consuming, however, T cells scan numerous APCs in a very short time (~ a few minutes) so that the immune system can react fast enough and avoid any potential significant damage to the body. The ability of T cells to perform their function properly while simultaneously abiding by all these constraints has baffled the scientific community for many years. Over the last decade, mechano-sensing/transduction has been proposed to be the missing puzzle piece in our understanding of T cell function (Pageon et al. 2018; Rushdi et al. 2020). Different players may have different roles, as we will exemplify further on.

Integrins: the prototypes of mechanosensitive cell surface proteins

As in any architectural structure, if mechanical load is to be transmitted across the cell surface into the cell, the simplest manner to do so would be through pliable structural elements that are physically interconnected (He and Bongrand 2012). Given that integrins link either the ECM or integrin ligands on other cells (through their extracellular domains) to the actin-cytoskeleton (through their cytoplasmic tails and adapter molecules) (Fig. 2B), they represent excellent candidates for both mechanosensing and mechanotransduction. In fact, the demonstration that integrins are indeed mechanoreceptors was made almost three decades ago in a series of elegant experiments using magnetic twisting cytometry, where twisting ligand-bearing beads bound to $\beta 1$ integrins caused endothelial cells to stiffen (N. Wang, Butler, and Ingber 1993).




Fig. 2. Key mechanosensory molecules for T cells. A: TCR interacts with peptide bearing MHC of an APC and directs the specificity of the adaptive immune response through signaling via the CD3 phosphorylatable cytoplasmic tails. B: integrins can modulate they extension and interaction with the cytoskeleton depending on forces acting on them (outside-in signaling) or in response to eg. T cell activation through the TCR (inside out-signaling) (Springer and Dustin 2012).

T cells specifically rely heavily on integrins, whether it is for adhesion during trafficking from the bloodstream, migration within tissues, immune synapse formation, or for signaling and cell polarization (Gérard et al. 2021). Lymphocyte function-associated antigen-1 (LFA-1) is the predominant integrin on T cells, binding intercellular adhesion molecule-1 and -2 (ICAM-1 and ICAM-2) on partner cells (antigen presenting cells or endothelial cells) (Murphy, Travers, and Walport 2008). Like other members of the integrin family, LFA-1 is a heterodimer comprising one α and one β chain, each containing a long, stalk-like extracellular domain, a transmembrane helix, and a short intracellular tail responsible for interacting with cytoplasmic signaling and cytoskeletal proteins (Luo, Carman, and Springer 2007).

The affinity of LFA-1 to ICAMs, however, is intimately coupled to its conformation, which is in turn is set by the cell activation status (Dustin and Springer 1989). In the cell resting state, LFA-1 exhibits a low affinity, bent conformation in which its ligand binding pockets are oriented towards the plasma membrane. In the presence of

activating TCR signals during immune synapse formation, specific protein complexes (e.g. talin and kindlins) assemble on the cytoplasmic tails of the α and β chains and drive them apart (Kim et al 2011). The conformational change induces the extension of the extracellular domain, thus allowing ligand recognition. Although this extended conformation is capable of ligand binding, it can only do so at intermediate affinity. In fact, TCR signaling alone is insufficient to unlock the full binding potential of LFA-1 (Feigelson et al. 2010).

Only under applied tangential force, originating from the actin cytoskeleton (further elaborated later on), and transferred to integrins via interactions between cytoskeletal adaptors such as talin and the tail of the β subunit, does LFA-1 reach peak binding affinity (~ 100 fold increase) (Astrof et al. 2006; Friedland, Lee, and Boettiger 2009), a clear signature of catch-bond behavior. Catch bonds are an unusual kinetic behavior of ligand receptor interactions where the exertion of a physical force on a molecular complex counter-intuitively prolongs its bond lifetime, in contrast to the so called ‘ordinary’ slip bonds, where force intuitively shortens bond lifetime. Indeed, similar to other integrins (Kong et al. 2009), LFA-1 binding with ICAM-1 behaves as a catch bond (W. Chen, Lou, and Zhu 2010)).

Interestingly enough, the engagement of LFA-1 alone does not generate any measurable forces or intracellular signaling (Husson et al. 2011). This observation suggests that the mechano-sensing/transduction capacity of T cells could not be limited to conventional adhesion molecules such as integrins.

TCR is at the core of early activating contacts and responds to forces

In the event that a cognate pMHC on APCs is encountered, TCR signaling will rapidly convert the ligand-binding event to the phosphorylation of up to 10 immunoreceptor tyrosine-based activation motif elements (ITAMS) in the cytoplasmic tails of the associated CD3 complexes. The ensuing signaling cascade ultimately results in developmental decisions, effects, or functions (Smith-Garvin, Koretzky, and Jordan 2009). Unfortunately, our current knowledge of this signaling cascade far exceeds our limited understanding of how it is initiated upon TCR-pMHC binding.

The TCR-pMHC interaction is probably among the weakest protein-protein interactions that can initiate an effective biological response (Chakraborty and Weiss 2014). The affinity of a TCR binding to a pMHC is only around 10^{-4} - 10^{-6} M (Rudolph, Stanfield, and Wilson 2006), about 1000 times weaker than a typical antigen/antibody binding (10^{-6} - 10^{-10} M (Sundberg and Mariuzza 2002)). Aside, shape-complementarity at the TCR-pMHC interface has been shown to be extremely poor (Rossjohn et al. 2015). Despite that, the TCR is still capable of discriminating as few as one to ten non-self antigens in a sea of endogenous self antigens that are presented by the same self-MHC molecule on the APC surface, and even a single pMHC is thought to be sufficient to trigger an efficient TCR signaling and subsequent T cell activation (Huang et al. 2013). All of this

begs the question: How can a seemingly weak interaction simultaneously achieve such level of specificity and sensitivity?

In an attempt to answer this question, in 2008, Ma and colleagues proposed a ‘receptor deformation model’ for TCR signaling. This model proposes that TCR signaling is initiated by significant conformational changes of the TCR/CD3 complex, induced by a pulling force originating from the cytoskeleton of the T cell and transmitted through pMHC-TCR binding interactions with enough strength to resist rupture (Z. Ma, Janmey, and Finkel 2008). Essentially, providing a mechanistic explanation to the specificity and sensitivity of the TCR.

A year later, Kim and colleagues provided the first concrete proof that the TCR behaves as a mechanosensor (Kim et al. 2009). They used optical tweezers (OT) and nuclear magnetic resonance (NMR) techniques to characterize the distinct functional consequences of several anti-CD3 monoclonal antibodies (mAbs) binding to T cells. In parallel, they quantified Ca^{2+} levels as a measure of T cell activation. The NMR cross-correlation analysis showed that agonist Abs (i.e. those capable of triggering calcium fluxes) bind CD3 in a *diagonal* fashion, in comparison to CD3 Abs that do not trigger downstream signaling which bind CD3 in an upright mode (perpendicular to cell membrane plane). Interestingly enough, perpendicularly binding Abs were still capable of activating T cells but only when a significant *tangential* force, of ~50 piconewtons (pN), i.e. ~10-12 times the thermal agitation limit, was applied by OT. Based on these observations, the authors proposed a model in which external tangential forces generated following pMHC ligation during the scanning of the APC by the T cell, allow TCRs to mechanically sense and then transduce the first activation signals.

TCR/pMHC can exhibit complex behaviors: the catch bond proposition

In 2014, Liu and colleagues connected yet another piece of the puzzle (B. Liu et al. 2014). Using biomembrane force probes (BFPs), they showed that the lifetime of the bond between a TCR and its specific pMHC was *prolonged* by the application of a ~ 10pN force, indicative of catch-bond behavior. Such complex response was also associated with more robust and long-lived cellular calcium fluxes, suggesting that catch bond formation may be required for stronger T cell activation. By contrast, the affinity of non-specific TCR-pMHC bonds peaked at zero force, indicative of slip-bond behavior. OT experiments using DNA tethers further revealed that it is in fact the FG loop of the constant domain of the β chain that allosterically controls the V domain modules’ catch bond lifetime and peptide discrimination, through a force-driven conformational transition (Das et al. 2015). Collectively, these findings demonstrate that by eliciting antigen-specific catch bonds, external forces may amplify the power of T cell antigen discrimination by separating agonist pMHCs that induce catch bonds from non-specific pMHCs that exhibit only slip bonds.




Fig. 3: Different behaviors of bonds, which have been proposed to depend on the peptide presented by a MHC. Slip bonds, whose lifetime only decreases when submitted to increasing forces, vs. catch bonds for which a range of forces sees an increasing lifetime, up to an optimum (which has been estimated to be ~ 10 pN for TCR/pMHC) before decreasing as for a catch bond (B. Liu et al. 2014).

While catch bonds have been observed in a broad range of molecules, TCR-pMHC catch bonds are still enigmatic, as their origin is still a matter of debate especially; numerous reports employing purely acellular systems have demonstrated that, outside the cellular context, the TCR does not exhibit catch bond behavior (Limozin et al. 2019). Aside, how can a tangential (to the membrane) force applied to the TCR-pMHC bond make it stronger? The same group attempted to answer this question using an integrated approach of steered molecular dynamics (SMD) simulation, MTs, and BFPs (Wu et al. 2019). Their results showed that forces acting on the TCR-agonist pMHC complex induced a conformational change in the MHC that subsequently increased the length of the complex. Specifically, the increased force experienced by the TCR-agonist pMHC bond uncoupled the α -chain $\beta 2$ -microglobulin ($\beta 2m$) interdomain interaction, resulting in a 5–10 nm extension of the MHC. They proposed that such pronounced extension would not only stabilize the TCR- agonist pMHC bond but also promote the formation of new interactions after forces rupture the preexisting ones. Based on these results, the authors hypothesized that, in the case of agonist pMHCs, the forces acting on the TCR-agonist pMHC complex would induce a conformational change in the MHC, ultimately stabilizing the complex and creating a catch bond. The catch bond would then endow the TCR with the power to sensitively discriminate between peptides (self and non-self), plus, the increased chance of bond formation would make T cell activation easier. Nevertheless, this still does not explain the discrepancy observed by (Limozin et al. 2019) and more work will be needed to clarify (i) if the catch bond behavior is needed for the T cell to perform its action and (ii) if it is a hallmark of TCR-cognate pMHC bonds, where is it originating from.

A very recent view : not only the TCR but also the pre-TCR is mechanosensitive

Even before the TCR, force-based discrimination, is thought to be conserved in its developmental precursor, the pre-TCR, for the selection of efficient TCRs. Early thymic progenitors (ETPs; uncommitted thymic cells retaining some myeloid, NK and little if any B lineage potential) enter the corticomedullary junction of the thymus as double-negative cells (DN, stages DN1 to 4), lacking the expression of both CD4 and CD8, as well as the full T cell receptor (whether TCR $\alpha\beta$ or TCR $\gamma\delta$). For the $\alpha\beta$ T cell lineage, a surrogate preT- α chain (denoted

pT α , which lacks V α of final TCR $\alpha\beta$) is expressed on the surface of DN2 cells in place of the α chain seen in the final $\alpha\beta$ TCR. Shortly after, the cells enter the DN3 stage where they synthesize the TCR β chain and express it on their surface in association with the pT α chain, forming what is known as the pre-TCR receptor (Carpenter and Bosselut 2010). Signaling through this pre-TCR marks the first major checkpoint in early thymic development, referred to as β -selection, whereby only DN cells with productive TCR β are selected to continue their development. The question here is how does pre-TCR signaling occur?

Initially, pre-TCR signaling was thought to be ligand-autonomous (Irving, Alt, and Killeen 1998; Yamasaki and Saito 2007) and purely dependent on pT α charge-based receptor oligomerization (Smelty et al. 2010). That theory was readily discredited by Mallis and colleagues (Mallis et al. 2015) who showed through nuclear magnetic resonance in solution and BFP experiments that the pre-TCR, just like its mature form, and through the β chain alone, is capable of recognizing its respective pMHC (albeit with a broader specificity than its final TCR $\alpha\beta$ form), as well as triggering calcium fluxes. Using OT, the same group later showed that this pre-TCR-pMHC recognition occurs specifically through the V β hydrophobic patch, in partnership with the C β FG loop of the TCR β , and that the recognition is in-fact force-sensitive (Das et al. 2016). Indeed, the pre-TCR-pMHC interaction, similar to the TCR-pMHC one, was shown to exhibit features of catch bond behavior. Diminishing bond strength and/or bond lifetime (through mutating either the V β or the C β FG loop) negatively impacted pre-TCR ligand discrimination and ultimately reduced post-DN3 thymocyte proliferation and developmental progression (Li et al. 2021).

These observations show that only under force is pre-TCR signaling induced during β -selection. In this sense, the β repertoire is tuned prior to the $\alpha\beta$ repertoire final tuning, with mechanotransduction through the β subunit serving as the first checkpoint towards ensuring a functional TCR. As for the diminished ligand specificity of the pre-TCR in comparison with that of the final $\alpha\beta$ TCR, it is possible that the broader ligand focus allows the β chain to interact with multiple self-pMHC ligands in the pMHC-rich stromal environment, affording DN3 growth/survival advantage to pMHC binding competent preTCRs and imprinting self-reactivity in the developing repertoire. Thus, DN progression selects for a self-reactive repertoire early in development. The V β patch may contribute to this behavior, relaxing peptide specificity requirements and functioning as a surrogate V α domain whose replacement at the double positive (DP) stage (signaling through the pre-TCR marks the end of DN3 stage and the transition into the DP stage where the cells stop β chain rearrangement, undergo a period of proliferation, and begin to express both CD4 and CD8) where by an actual V α domain then imposes more precise peptide recognition. Negative selection, that corresponds to the final selection before T cells leave the thymus where only DP T cells that bind self antigens at low affinity survive, therein purges high pMHC self-reactivity while maintaining a low self-pMHC bias.

Sensing and exerting forces on a cellular level: the role of the actomyosin cytoskeleton

Moving up from the molecular to the cellular level, mechanical forces play a very important role in T cell function. However, before diving into that, one should first address how forces are generated and sensed on the cellular scale. Ultimately, mechano-sensing/transduction, on any scale, and force exertion are tightly linked processes. Mechanically induced conformational changes, just as those described for activating the pMHC-TCR and LFA-1-ICAM bonds, only occur under the influence of force. Ergo, mechanotransduction necessitates that the cell exerts and receives forces from its environment. Conversely, force exertion is itself regulated by feedback from mechanosensing pathways, as we will see later on.

Cells exert forces against their environments via dynamic cytoskeletal remodeling; the cytoskeleton is a polymer network composed of three distinct biopolymers: actin, microtubules, and intermediate filaments. Typically, it is the filamentous actin (F-actin) cytoskeleton that bears the brunt of the mechanical load; It is a highly dynamic structure that undergoes continuous reorganization in response to external mechanical cues. This feature is what enables the cell to rapidly change its elastic properties and what consequently endows it with the capacity to apply forces against a substrate and move (Ananthakrishnan and Ehrlicher 2007). The classical model for F-actin dependent force exertion involves myosin motors consuming chemical energy in the form of ATP and walking on actin filaments in a general three-step process of binding, power stroke, and unbinding. This process is continuously repeated and leads to the generation of a contractile force (actomyosin contractile force) (Salvi and DeMali 2018) . Although actomyosin contractility was initially characterized in muscle cells, it is now clear that it is a universal mechanism for force generation in most eukaryotic cells, fueling a wide range of processes including adhesion, division and motility. With that being said, it is important to note that actin polymerization alone, in the absence of myosin motors, does also generate force. However, such protrusive forces are far less characterized, most likely because they are easily masked by the long-lasting, contractile ones (Ananthakrishnan and Ehrlicher 2007).

Whether it is protrusive or contractile, in order for forces to propagate from the cytoskeleton onto the extracellular environment (substrate or cell), both parties have to be linked through *adhesive* contact points. The most characterized of such contact points are focal adhesions (FAs); FAs constitute large protein assemblies in which transmembrane adhesion receptors (e.g. integrins) and F-actin are bridged via a specialized layer of cytoplasmic scaffolding proteins (e.g. paxillin, vinculin, talin...) (C. S. Chen, Tan, and Tien 2004). The size, composition, and structure of such adhesion sites are directly dependent on the mechanical forces that they are subjected to, whether it is from actomyosin contractility or from the extracellular environment. This explains why FAs are readily observed for fibroblasts cultured on stiff supports, while similar prominent contacts are harder to detect in-vivo, where the ECM is much more compliant (Prager-Khoutorsky et al. 2011). Interestingly, the process of building FAs from initial adhesion receptors is intricately coupled to the activity of intracellular signaling cascades, not through their possession of enzymatic activity, rather, their capacity to recruit specific, “classical” adhesome signaling components to the growing FAs (Welf et al. 2020). For example, in the case of

integrin mediated adhesions, the focal adhesion kinase (FAK) recruited to the FA site regulates diverse downstream signaling pathways, including those promoting cell growth and survival (Schlaepfer et al. 1994).

It has to be underlined that, unlike large adherent cells such as the fibroblasts mentioned above, many immune cells, among which the T cells, do *not* form distinct FAs-like structures in vitro or in vivo. Rather, they form *transient* adhesive contacts that contain cell surface receptors, F-actin, and cytoplasmic proteins such as the ones typically found in FAs. These contacts likely serve as sites for force exertion during migration and cell-cell interactions (Huse 2017).

The most straightforward way in which cellular forces could contribute to T cell function is through enabling their migration and trafficking. Typically, as a cell moves on a substrate (whether it is the ECM or simply a cover slide), it experiences external forces, mainly the viscous force/resistance from the surrounding medium and cell-substrate interaction forces, as well as internal forces that are generated by the cytoskeleton. In T cells, as in most animal cells, the cytoskeleton is the essential component in creating these motility-driving forces, and in coordinating the entire process of movement: First, a cell propels the membrane forward by growing the actin network at its leading edge, creating an F-actin rich lamellipodium. Second, it adheres to the substrate (for example through integrin adhesions in T cells) at the leading edge and deadheres (releases) at the cell body and rear of the cell (also known as uropod). Finally, the cell propels forward by the F-actin retrograde flow generated against the adhesive contacts present at the base of the leading edge of the cell; retrograde flow describes the variable movement of actin filaments rearward with respect to the substrate, generally in the direction opposite to cell movement (Ananthakrishnan and Ehrlicher 2007), and it is caused by actin polymerization against the plasma membrane, which drives the growing fibers backwards, and myosin contractility, which collapses the leading edge F-actin network into linear bundles (Huse 2017).

Aside from motility, cellular forces come into play at different time points in T cell activation. To begin with, the most basic requirement for T cell activation is for the TCR to interact with the pMHC. This may seem trivial to point out, however, there are physical barriers that make this interaction not as straightforward; The TCR-pMHC bonds (10-15 nm) are much smaller than individual TCR and APC glyocalyx proteins, such as the T cell receptor tyrosine phosphatases CD45 (28-50 nm) and CD148 (47-55 nm), and even LFA-ICAM bonds (45-50 nm for the couple). Though models such as the kinetic segregation one (Davis and Merwe 2006) were originally put forth to explain how the T cell overcomes these barriers, there still remains several key issues that the model does not account for (Comrie and Burkhardt 2016). Recently, Cai et al. combined time-resolved lattice light-sheet microscopy and quantum dot-enabled synaptic contact mapping microscopy to show how highly dynamic T cell F-actin-rich microvilli colocalized with TCR MCs, and in the absence of external stimulus, scanned the entire area of opposing cells and surfaces (coated with antagonist/agonist pMHCs and ICAM-1) before and during antigen recognition, at a time frame (≈ 1 min) similar to that recorded for T cell-APC contacts in vivo (Cai et al. 2017). These observations, coherent with earlier ones (Brodovitch, Bongrand, and Pierres 2013),

suggest that T cell microvilli, with an average length of 380 nm, can promote TCR signaling by surpassing the size-related restrictions, penetrating the glycocalyx, and bringing the TCR into close proximity with the pMHCs. Additionally, one could imagine that the applied F-actin protrusive forces would further stabilize low affinity TCR-pMHC bonds, and with the microvilli containing pre-clustered TCRs, it would provide an easy access platform for signal amplification explaining the high sensitivity of T-cells to low numbers of pMHC antigens.

After TCR engagement, is actin polymerization at the T cell-APC contact zone commences. The membrane deformation resulting from such polymerization forces allows the T cell to spread over the APC. This spreading process is critical as it not only allows the T cell to scan a larger area of the APC and thus increase the efficiency of antigen sampling (Brodovitch et al. 2015), but it also exerts force on the receptor-ligand pairs engaged, such as the mechanosensitive TCR-pMHC and LFA-1-ICAM-1/2 bonds, further enhancing peptide discrimination and TCR activation.

As the T cell reaches its maximal spreading area, the same actin polymerization forces, combined with myosin contractility, create retrograde flow. Forces originating from this retrograde flow organize the various receptor microclusters (MCs; upon ligand binding, TCRs coalesce into signaling microclusters containing >10 receptors each) and signaling molecules present at the T cell-APC contact zone, and order them into the infamous spatially symmetric bullseye structure of the IS (Blumenthal and Burkhardt 2020). To be more specific, the TCR MCs are swept towards the center of the contact by retrograde F-actin centripetal flow at the periphery and then by myosin II dependent actin arcs closer to the center, leading to the formation of the cSMAC (central supramolecular activation cluster) surrounded by a ring of integrins (LFA-1/ICAM bonds) in the pSMAC (peripheral supramolecular activation cluster). The interruption of F-actin centripetal flow eradicates TCR MC signaling within seconds, further confirming that force exertion is imperative for maintaining proper TCR activation. Interestingly enough though, the same actin machinery described above may also break TCR-pMHC bonds, allowing the serial engagement of the same pMHC with the other TCRs present in the TCR MC, consequently augmenting TCR signaling.

Once the IS is established, it has to be maintained for a long enough period of time (up to hours) to enable the proper activation of the T cell. This is a particularly difficult task as T cells are already highly motile cells and the T cell-APC interaction occurs in non-static conditions. By monitoring the T cell cytoskeletal organization during their interaction with both APCs and APC mimetic surfaces, Kumari et al. found that antigen recognition triggered the formation of actin foci (by the help of Wiskott–Aldrich syndrome protein) at the T cell-APC/substrate contact that, with the assistance of myosin II contractility, generated and sustained intracellular tension within the T cell that maintained the stability and symmetry of the IS for the activation time frame (Kumari et al. 2020).

Finally, in an elegant series of experiments combining pMHC and ICAM-1 coated on beads beared by deformable micropipettes and on micropillar arrays, Basu et al. demonstrated that mechanical forces at the IS potentiate cytotoxic T cell (CTL) cytotoxicity: CTLs destroy target cells by secreting a mixture of the protein perforin and granzyme proteases where perforin forms pores in the target cell membrane that enable granzymes to access the cytoplasm and induce apoptosis (Basu et al. 2016). Specifically, their study revealed that altering the membrane tension of the CTLs using pharmacological drugs or osmotic shock strongly perturbed the pore-forming activity of perforin. Similarly, altering the membrane tension of the target cell by changing substrate stiffness modulated CTL killing, with cells on stiffer substrates exhibiting a higher sensitivity to perforin-induced pore formation. Taken together, these results point towards a model in which forces at the IS promote CTL killing by straining the target cell membrane, thus facilitating the formation of perforin pores. Considering that several reports have correlated transformation and malignancy with cellular softening, this work puts forth a very compelling hypothesis of whether tumor cells modulate their mechanical properties to relief forces at the IS and thus evade the immune system (Huse 2017; Basu and Huse 2017).

T cells can detect substrate mechanics and react to it

Just as we do when we use our fingers to apply pressure on an object, T cells exert forces to probe their mechanical environment, particularly stiffness. Pioneering work by Judokusumo et al. initially documented this property by stimulating naïve CD4⁺ mouse T cells with polyacrylamides of different rigidities, and functionalized with activating antibodies against CD3 and CD28 (Judokusumo et al. 2012). Their experiments revealed that T cells exhibited stronger activation, quantified as IL-2 secretion, with increasing substrate rigidities (over the range of 10-200 KPa), and that this mechano-sensing/transduction ability was largely affiliated with the TCR/CD3 complex rather than CD28. Intriguingly, this “stiffness sensitivity” property was observed only when the anti-CD3 antibody was immobilized onto the surface of the gel, rather than added as a soluble solution, and it was lost upon myosin inhibition. These observations are in accordance with the now commonly accepted idea that antigen receptors pull against their ligands for optimal signaling. Conversely, similar experiments done by O’Connor et al. on polydimethylsiloxane substrates with the same functionalization but using a different rigidity range (100-200 KPa), showed that naïve CD4⁺ human T cells were stimulated and proliferated more on softer substrates in comparison to stiffer ones (O’Connor et al. 2012). Taken together, these studies suggest a possible biphasic response to stiffness sensitivity. Another crucial piece of information came from Tabdanov et al. who employed a combination of activating anti-CD3 antibody and ICAM-1 functionalized flat micropatterned PDMS substrates (5 KPa- 2000 KPa) and micropillar arrays to delineate the contributions of both the TCR/CD3 complex and LFA-1 in stimulated CD4⁺ human T cell activation (Tabdanov et al. 2015). In these experiments, early T cell activation, measured by the total phosphor-tyrosine levels, was weaker on soft substrates than on rigid ones. Though this stiffness sensitivity was observed in the absence of LFA-1 engagement, it was enhanced by its presence. Even more interestingly, their results also highlighted a

mechanical cooperation between the TCR/CD3 and LFA-1-ICAM-1 systems, whereby actin nucleation downstream of TCR signaling sustained the growth of the LFA-1 dependent actin network, which in turn provided the cytoskeletal tension to allow mechanical sensing, T-cell spreading and enhanced TCR activation.

Similar experiments were later repeated but on substrates with stiffnesses of more physiologic relevance in terms of T cell function, considering that APCs display a stiffness range between \approx 200 Pa and 2 KPa (Bufi et al. 2015). Notable of which were those performed by Hui et al. (Hui and Upadhyaya 2017; Hui et al. 2015) who used poly-l-lysine-antiCD3-coated soft polyacrylamide gels (1- 5 KPa) to demonstrate the contributions of actin polymerization and myosin contractility, as well as dynamic microtubules, to force generation and maintenance during enhanced green fluorescent protein (eGFP)-actin expressing Jurkat T cell activation (quantified as phosphotyrosine signaling). Their results, similar to what was originally documented by Judokusumo et al., showed that T cells exhibited higher levels of activation on stiffer gels in comparison to softer ones.

Though these studies are difficult to directly compare because they differ in substrate chemistry, antibody/protein immobilization, stiffness ranges, and more importantly the T cell types/subtypes used, they do overall reveal that T cells possess the inherent ability to sense stiffness. This, at least partly, explains their modified behaviors in mechanically distinct interactions, whether it is different APCs that have been activated by different stimuli and present a varying repertoire of agonist/non-agonist pMHCs, or endothelial cells in blood vessels, or infected/tumor cells inside tissues. Even if the change in stiffness between these surfaces may seem quite modest and inconsequential, it is nevertheless sensed and responded to by T cells. Wahl et al. recently proposed a model in which increased substrate stiffness heightens the TCR-pMHC resistance to cytoskeletal forces and thus increases T cell spreading and activation. That is to a certain limit, beyond which the tension on the bonds becomes too high and breaks them, which consequently decreases spreading and weakens T cell activation (Wahl et al. 2019).

Ligand mobility is felt by T cells and may influence force transmission

Aside from stiffness sensitivity, T cells have also been shown to be sensitive to ligand mobility (Dillard et al. 2014). The interaction between a T cell and an APC necessitates extensive cytoskeletal and lipid membrane composition changes for both cells, as to allow for the spatial ligand/receptor re-ordering mentioned above. In an innovative approach, Mossman et al. investigated the impact of ligand mobility on T cell signaling by creating “artificial APCs” where nanofabricated 10–20 nm high chromium barriers were assembled on pMHC and ICAM-1 coated supported lipid bilayers (Mossman et al. 2005); set up as is, the bilayer would allow for free lipid diffusion, however, the barriers would block the movement of proteins with larger cytoplasmic domains, and more importantly, TCR MCs. Interestingly, trapping the TCR clusters in the in the IS periphery (as opposed to their natural position in the cSMAC of the IS) augmented early TCR-associated phosphotyrosine signaling and cytoplasmic Ca²⁺ levels in the spatially constrained IS in comparison to the native ones. In a similar

approach, but playing on the lipid bilayer composition instead of using chromium barriers for limiting ligand mobility, Hsu et al. revealed that tyrosine phosphorylation and persistent elevation of cytoplasmic Ca^{2+} was in fact more pronounced for T cells (Jurkat and naïve or stimulated $CD4^+$ murine) on mobile membranes than on less mobile ones (Hsu et al. 2012). Though these two studies seem contradictory, the immobilization of the TCRs differed between the two systems; in the former, the chromium barriers completely trapped the TCR clusters in the periphery, on the other hand, the latter still permitted the diffusion of TCRs however at a slower rate. This could underline a complex mechanism, potentially reliant on the spatial and temporal parameters of ligand constriction, by which T cell sensitivity to ligand mobility impacts T cell activation. However, this would require further experimentation to decipher.

More recently, pioneering work done by Bukhardt and colleagues (Comrie et al. 2015; Comrie and Burkhardt 2016) revealed that dendritic cell (DC) maturation- a process characterized by an increase in DC cortical stiffness- induced a dramatic actin-dependent decrease in ICAM-1 mobility. The reported decrease in ICAM-1 mobility helped generate a counterforce that drove the centripetal flow of the actomyosin network in the T cells spreading over the APC. This flow, in turn, recruited LFA-1 to the IS, maintained it in a high affinity conformation, and consequently promoted efficient binding to ICAM-1. One could imagine that since LFA-1 connects the extra- and intra-cellular compartments, similar to other integrins, the tension on LFA-1 will also affect the dynamics of the underlying T cell actin network (Jankowska et al. 2018); since the TCR is thought to be interacting with said network, this will indirectly influence tension on the TCR, potentially modulating TCR signaling (He and Bongrand 2012). This work is of particular importance as firstly, it explains how LFA-1 reaches peak binding affinity necessary for proper T cell activation, and secondly, it suggests that cells can regulate intercellular communication by altering the physical status of the signaling molecules in question, rather than just their expression level or spatial localization.

How to pass on the message ?

Although the influence of mechanical forces on the specificity and sensitivity of antigen recognition by the TCR is coming to light, how information regarding TCR-antigen binding is relayed into the cell still remains unclear (Harrison, Fang, and Huang 2019).

As mentioned above, TCR signaling propagates across the membrane through the CD3 intracellular domains, specifically through ITAM phosphorylation. In their unphosphorylated state, ITAM chains have been shown to be buried in the hydrophobic interior of the membrane, hence inaccessible to Src kinases. Ligand binding by the TCR has been recently proposed to induce conformational change in the CD3 chains, extending them and exposing their ITAMs to phosphorylation (Lee et al. 2015). Although there are currently no definitive studies directly linking mechanical forces applied onto the TCR protein to this $CD3_{\zeta}$ conformational change (eg. are the

forces needed pushing or pulling on the complex to unlock it like you do with an umbrella?), a recent study using fluorescence resonance energy transfer (FRET) showed that the TCR, under force, is able to decipher structural subtle differences between peptides by different bond conformations, independent of binding affinity and kinetics. Peptide potency then appears to directly regulate the amount of conformational change, which in turn dictates the degree of dissociation of the CD3 (ζ chain) from the inner membrane leaflet and consequently the exposure of its ITAMs to phosphorylation (Sasmal et al. 2019).




Fig.4. Typical times scales for the signal propagation and consequences. The question marks are underlying the interactions between different cell biology “modules” (Puech and Bongrand 2021) which are still to be clarified for the entire process of T cell mechanotransduction and activation to be fully understood.

Another important question to address is the link between the different scales (Fig. 4), particularly the TCR and the actin cytoskeleton. As mentioned previously, in adherent cells, the maintenance, growth and signaling through FAs are completely dependent on cytoskeletal forces. As such, FAs act as mechano-sensors/transducers bridging transmembrane adhesion receptor binding and actin flow with cell signaling. It is intriguing to imagine that the TCR MCs serve similar purposes. Using novel ratiometric tension probes, Ma et al. have demonstrated that TCRs undergoing clustering within the first few minutes stimulation experience tension in the pN range (Y. Liu et al. 2016; V. P.-Y. Ma et al. 2016). It is thus highly likely that TCR clustering is stabilized by the underlying F-actin network or even through direct tethering of the TCR complex to cortical actin. Interestingly, the force-sensing protein lymphocyte-specific Crk-associated substrate (Cas-L) has recently been proposed by Santos et al. to mechanically link TCR MCs to the underlying actin network (Santos et al. 2016). Their experiments showed that Cas-L participates in a positive feedback loop, whereby upon TCR triggering, Cas-L localized the TCR MCs undergoes actin-polymerization dependent activation (through phosphorylation), leading to Ca^{2+} signal amplification, regulation of TCR microcluster transport, inside-out integrin signaling, as well as actomyosin contraction (Santos et al. 2016).

Concluding remarks

Besides the mechano-sensitive/transductive abilities of the TCR and integrins, there are several other membrane receptors, ion channels, cytoskeletal proteins, and transcriptional factors that are thought to be also affected by physical forces. For example, pulling forces on bound Notch receptors during endocytosis of Notch ligands induce a conformational change in Notch that ultimately drives early thymic progenitors to commit to the T cell lineage (X. Wang and Ha 2013; Luca et al. 2017) and mechanical stretch of the membrane during IS formation activates Piezo channels, thereby triggering Ca^{2+} flux and regulating TCR signaling (C. S. C. Liu et al. 2018).

What is even more interesting is that these different elements do not function in isolation, but rather as parts of a complex mechanical signaling network with cross-talks and feedback loops, that ultimately regulates T cell mechanics, gene expression, and behavior.

The challenge is, now that some of the key elements have been described separately, to understand how these mechano-signaling components and pathways are intertwined and integrated across time and length scales, and in different intra-cellular compartments, to shape the T cell response (Limozin and Puech 2019; Puech and Bongrand 2021). To take the TCR and LFA-1 as an example, Bernard and colleagues attempted to decipher the mechanical link between these two molecules by imaging T cells on anti-TCR Ab micropatterned soluble lipid bilayers (SLBs) (Benard et al. 2018). Their results showed that the TCRs do in fact aggregate into microclusters that colocalize with the anti-TCR Ab patterns, however, the clusters do not move (by the means of retrograde actin flow) to the center of the contact area, as seen during the formation of the central supramolecular activating complex of the IS. Only upon the addition of ICAM-1 to the SLBs, do the TCR microclusters centralize with the actin and form a peripheral ring around them. This study, in addition to many others (Tabdanov et al. 2015; Verma, Kelleher, and Alerts 2018; Y. Chen et al. 2017), supports a model in which the actin cytoskeleton couples the TCR and LFA-1 in a positive feedback loop that coordinates IS formation and growth. It also puts forward a very exciting concept of the actin cytoskeleton acting as a mechanical intermediate that integrates force-dependent signals coming from different receptor-ligand interactions, and then coordinates outgoing responses over large distances (Comrie and Burkhardt 2016).

Over the last decade, a sturdy foundation has emerged for measuring and interpreting mechanical forces in T cells. Nevertheless, the field remains in its infancy and more work, both experimental and theoretical, will still be needed to fully connect the dots, and in particular the behaviors at different time and space scales, from molecular to cellular worlds. However, what is becoming more and more apparent is that forces play quite an important role in the T cell response and can no longer be overlooked. It is our hope that the literature and arguments presented in this review would raise awareness to this emerging area of research in T-cell biology. What also appears clearer is that these concepts apply to all and every immune cell type, with basic similar

phenomena and subtle differences, for lymphocytes such as T cells, B cells, NK cells but also APC cells such as dendritic cells or macrophages (Huse 2017; Zhang et al. 2020).

Aknowledgments & fundings

Part of this work was supported by institutional grants from INSERM, CNRS and Aix-Marseille University to the LAI and CINAM.

FM was supported by a PhD grant from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No713750, with the financial support of the Regional Council of Provence- Alpes-Côte d'Azur and with of the A*MIDEX (n° ANR- 11-IDEX-0001-02), funded by the Investissements d'Avenir project funded by the French Government, managed by the French National Research Agency (ANR)

References

- Ananthakrishnan, Revathi, and Allen Ehrlicher. 2007. 'The Forces behind Cell Movement'. *International Journal of Biological Sciences* 3 (5): 303–17. <https://doi.org/10.7150/ijbs.3.303>.
- Astrof, Nathan S., Azucena Salas, Motomu Shimaoka, JianFeng Chen, and Timothy A. Springer. 2006. 'Importance of Force Linkage in Mechanochemistry of Adhesion Receptors'. *Biochemistry* 45 (50): 15020–28. <https://doi.org/10.1021/bi061566o>.
- Basu, Roshni, and Morgan Huse. 2017. 'Mechanical Communication at the Immunological Synapse'. *Trends in Cell Biology* 27 (4): 241–54. <https://doi.org/10.1016/j.tcb.2016.10.005>.
- Basu, Roshni, Benjamin M Whitlock, Julien Husson, Judy Lieberman, Lance C Kam, Morgan Huse, Roshni Basu, et al. 2016. 'Cytotoxic T Cells Use Mechanical Force to Potentiate Target Cell Killing Cytotoxic T Cells Use Mechanical Force to Potentiate Target Cell Killing'. *Cell* 165 (1): 100–110. <https://doi.org/10.1016/j.cell.2016.01.021>.
- Bell, G. I. 1978. 'Models for the Specific Adhesion of Cells to Cells.' *Science* 200 (4342): 618–27.
- Benard, Emmanuelle, Jacques A Nunès, Laurent Limozin, and Kheya Sengupta. 2018. 'T Cells on Engineered Substrates : The Impact of TCR Clustering Is Enhanced by LFA-1 Engagement' 9 (September): 1–12. <https://doi.org/10.3389/fimmu.2018.02085>.
- Blumenthal, Daniel, and Janis K. Burkhardt. 2020. 'Multiple Actin Networks Coordinate Mechanotransduction at the Immunological Synapse'. *The Journal of Cell Biology* 219 (2). <https://doi.org/10.1083/jcb.201911058>.
- Brodovitch, Alexandre, Pierre Bongrand, and Anne Pierres. 2013. 'T Lymphocytes Sense Antigens within Seconds and Make a Decision within One Minute'. *Journal of Immunology (Baltimore, Md.: 1950)* 191 (5): 2064–71. <https://doi.org/10.4049/jimmunol.1300523>.
- Brodovitch, Alexandre, Eugene Shenderov, Vincenzo Cerundolo, Pierre Bongrand, Anne Pierres, and Philip Anton van der Merwe. 2015. 'T Lymphocytes Need Less than 3 Min to Discriminate between Peptide MHCs with Similar TCR-Binding Parameters'. *European Journal of Immunology*, n/a-n/a. <https://doi.org/10.1002/eji.201445214>.
- Bufi, Nathalie, Michael Saitakis, Stéphanie Dogniaux, Oscar Buschinger, Armelle Bohineust, Alain Richert, Mathieu Maurin, Claire Hivroz, and Atef Asnacios. 2015. 'Human Primary Immune Cells Exhibit Distinct Mechanical Properties That Are Modified by Inflammation'. *Biophysical Journal* 108 (9): 2181–90. <https://doi.org/10.1016/j.bpj.2015.03.047>.
- Cai, En, Kyle Marchuk, Peter Beemiller, Casey Beppler, Matthew G. Rubashkin, Valerie M. Weaver, Audrey Gérard, et al. 2017. 'Visualizing Dynamic Microvillar Search and Stabilization during Ligand Detection by T Cells'. *Science* 356 (6338): eaal3118–eaal3118. <https://doi.org/10.1126/science.aal3118>.
- Carpenter, Andrea C., and Rémy Bosselut. 2010. 'Decision Checkpoints in the Thymus'. *Nature Immunology* 11 (8): 666–73. <https://doi.org/10.1038/ni.1887>.
- Chakraborty, Arup K, and Arthur Weiss. 2014. 'Insights into the Initiation of TCR Signaling.' *Nature Immunology* 15 (9): 798–807. <https://doi.org/10.1038/ni.2940>.
- Chen, Christopher S., John Tan, and Joe Tien. 2004. 'Mechanotransduction at Cell-Matrix and Cell-Cell Contacts'. *Annual Review of Biomedical Engineering* 6 (1): 275–302. <https://doi.org/10.1146/annurev.bioeng.6.040803.140040>.
- Chen, Wei, Jizhong Lou, and Cheng Zhu. 2010. 'Forcing Switch from Short- to Intermediate- and Long-Lived States of the AA Domain Generates LFA-1/ICAM-1 Catch Bonds'. *Journal of Biological Chemistry* 285 (46): 35967–78. <https://doi.org/10.1074/jbc.M110.155770>.
- Chen, Yunfeng, Lining Ju, Muaz Rushdi, Chenghao Ge, and Cheng Zhu. 2017. 'Receptor-Mediated Cell Mechanosensing'. *Molecular Biology of the Cell* 28 (23): 3134–55. <https://doi.org/10.1091/mbc.E17-04-0228>.
- Comrie, William A., and Janis K. Burkhardt. 2016. 'Action and Traction: Cytoskeletal Control of

- Receptor Triggering at the Immunological Synapse'. *Frontiers in Immunology* 7 (March). <https://doi.org/10.3389/fimmu.2016.00068>.
- Comrie, William A., Shuixing Li, Sarah Boyle, and Janis K. Burkhardt. 2015. 'The Dendritic Cell Cytoskeleton Promotes T Cell Adhesion and Activation by Constraining ICAM-1 Mobility'. *The Journal of Cell Biology* 208 (4): 457–73. <https://doi.org/10.1083/jcb.201406120>.
- Das, Dibyendu Kumar, Yinnian Feng, Robert J. Mallis, Xiaolong Li, Derin B. Keskin, Rebecca E. Hussey, Sonia K. Brady, et al. 2015. 'Force-Dependent Transition in the T-Cell Receptor β -Subunit Allosterically Regulates Peptide Discrimination and PMHC Bond Lifetime'. *Proceedings of the National Academy of Sciences* 112 (5): 1517–22. <https://doi.org/10.1073/pnas.1424829112>.
- Das, Dibyendu Kumar, Robert J. Mallis, Jonathan S. Duke-Cohan, Rebecca E. Hussey, Paul W. Tetteh, Mark Hilton, Gerhard Wagner, Matthew J. Lang, and Ellis L. Reinherz. 2016. 'Pre-TCRs Leverage V β CDRs and Hydrophobic Patch in Mechanosensing Thymic Self-Ligands'. *Journal of Biological Chemistry* 291 (49): jbc.M116.752865-jbc.M116.752865. <https://doi.org/10.1074/jbc.M116.752865>.
- Davis, Simon J, and P. Anton van der Merwe. 2006. 'The Kinetic-Segregation Model: TCR Triggering and Beyond.' *Nat Immunol* 7 (8): 803–9. <https://doi.org/10.1038/ni1369>.
- Dillard, Pierre, Rajat Varma, Kheya Sengupta, and Laurent Limozin. 2014. 'Ligand-Mediated Friction Determines Morphodynamics of Spreading T Cells'. *Biophysical Journal* 107 (11): 2629–38. <https://doi.org/10.1016/j.bpj.2014.10.044>.
- Dustin, M. L., and T. A. Springer. 1989. 'T-Cell Receptor Cross-Linking Transiently Stimulates Adhesiveness through LFA-1.' *Nature* 341 (6243): 619–24. <https://doi.org/10.1038/341619a0>.
- Feigelson, Sara W, Ronit Pasvolsky, Saso Cemerski, Ziv Shulman, Valentin Grabovsky, Tal Ilani, Adi Sagiv, et al. 2010. 'Occupancy of Lymphocyte LFA-1 by Surface-Immobilized ICAM-1 Is Critical for TCR- but Not for Chemokine-Triggered LFA-1 Conversion to an Open Headpiece High-Affinity State.' *Journal of Immunology (Baltimore, Md. : 1950)* 185 (12): 7394–7404. <https://doi.org/10.4049/jimmunol.1002246>.
- Friedland, Julie C., Mark H. Lee, and David Boettiger. 2009. 'Mechanically Activated Integrin Switch Controls Alpha5beta1 Function'. *Science (New York, N.Y.)* 323 (5914): 642–44. <https://doi.org/10.1126/science.1168441>.
- Gérard, Audrey, Andrew P. Cope, Claudia Kemper, Ronen Alon, and Robert Köchl. 2021. 'LFA-1 in T Cell Priming, Differentiation, and Effector Functions'. *Trends in Immunology* 42 (8): 706–22. <https://doi.org/10.1016/j.it.2021.06.004>.
- Harrison, Devin L., Yun Fang, and Jun Huang. 2019. 'T-Cell Mechanobiology: Force Sensation, Potentiation, and Translation'. *Frontiers in Physics* 7 (April). <https://doi.org/10.3389/fphy.2019.00045>.
- He, Hai-Tao, and Pierre Bongrand. 2012. 'Membrane Dynamics Shape TCR-Generated Signaling'. *Frontiers in Immunology* 3: 90. <https://doi.org/10.3389/fimmu.2012.00090>.
- Hsu, Chih-Jung, Wan-Ting Hsieh, Abraham Waldman, Fiona Clarke, Eric S. Huseby, Janis K. Burkhardt, and Tobias Baumgart. 2012. 'Ligand Mobility Modulates Immunological Synapse Formation and T Cell Activation'. *PloS One* 7 (2): e32398. <https://doi.org/10.1371/journal.pone.0032398>.
- Huang, Jun, Mario Brameshuber, Xun Zeng, Jianming Xie, Qi-jing Li, Yueh-hsiu Chien, Salvatore Valitutti, and Mark M Davis. 2013. 'A Single Peptide-Major Histocompatibility Complex Ligand Triggers Digital Cytokine Secretion in CD4(+) T Cells.' *Immunity* 39 (5): 846–57. <https://doi.org/10.1016/j.immuni.2013.08.036>.
- Hui, King Lam, Lakshmi Balagopalan, Lawrence E. Samelson, and Arpita Upadhyaya. 2015. 'Cytoskeletal Forces during Signaling Activation in Jurkat T-Cells'. *Molecular Biology of the Cell* 26 (4): 685–95. <https://doi.org/10.1091/mbc.E14-03-0830>.

- Hui, King Lam, and Arpita Upadhyaya. 2017. 'Dynamic Microtubules Regulate Cellular Contractility during T-Cell Activation'. *Proceedings of the National Academy of Sciences* 114 (21): E4175–83. <https://doi.org/10.1073/pnas.1614291114>.
- Huppa, Johannes B., and Mark M. Davis. 2013. 'The Interdisciplinary Science of T-Cell Recognition'. In *Advances in Immunology*, 119:1–50. Elsevier. <https://doi.org/10.1016/B978-0-12-407707-2.00001-1>.
- Huse, Morgan. 2017. 'Mechanical Forces in the Immune System'. *Nature Reviews Immunology* 17 (11): 679–90. <https://doi.org/10.1038/nri.2017.74>.
- Husson, Julien, Karine Chemin, Armelle Bohineust, Claire Hivroz, and Nelly Henry. 2011. 'Force Generation upon T Cell Receptor Engagement'. Edited by Javed N. Agrewala. *PLoS ONE* 6 (5): e19680. <https://doi.org/10.1371/journal.pone.0019680>.
- Irving, B. A., F. W. Alt, and N. Killeen. 1998. 'Thymocyte Development in the Absence of Pre-T Cell Receptor Extracellular Immunoglobulin Domains'. *Science (New York, N.Y.)* 280 (5365): 905–8. <https://doi.org/10.1126/science.280.5365.905>.
- Jankowska, Katarzyna I., Edward K. Williamson, Nathan H. Roy, Daniel Blumenthal, Vidhi Chandra, Tobias Baumgart, and Janis K. Burkhardt. 2018. 'Integrins Modulate T Cell Receptor Signaling by Constraining Actin Flow at the Immunological Synapse'. *Frontiers in Immunology* 9 (January). <https://doi.org/10.3389/fimmu.2018.00025>.
- Judokusumo, Edward, Erdem Tabdanov, Sudha Kumari, Michael L. Dustin, and Lance C. Kam. 2012. 'Mechanosensing in T Lymphocyte Activation'. *Biophysical Journal* 102 (2): L5-7. <https://doi.org/10.1016/j.bpj.2011.12.011>.
- Kim, Sun Taek, Koh Takeuchi, Zhen-Yu J Sun, Maki Touma, Carlos E Castro, Amr Fahmy, Matthew J Lang, Gerhard Wagner, and Ellis L Reinherz. 2009. 'The Alphabeta T Cell Receptor Is an Anisotropic Mechanosensor'. *J Biol Chem* 284 (45): 31028–37. <https://doi.org/10.1074/jbc.M109.052712>.
- Kong, Fang, A Paul Mould, Martin J Humphries, and Cheng Zhu. 2009. 'Demonstration of Catch Bonds between an Integrin and Its Ligand'. *Cell* 185 (7): 1275–84. <https://doi.org/10.1083/jcb.200810002>.
- Kumari, Sudha, Michael Mak, Yeh-Chuin Poh, Mira Tohme, Nicki Watson, Mariane Melo, Erin Janssen, Michael Dustin, Raif Geha, and Darrell J Irvine. 2020. 'Cytoskeletal Tension Actively Sustains the Migratory T-cell Synaptic Contact'. *The EMBO Journal* 39 (5). <https://doi.org/10.15252/emj.2019102783>.
- Lee, Mark S., Caleb R. Glassman, Neha R. Deshpande, Hemant B. Badgandi, Heather L. Parrish, Chayasith Uttamapinant, Philipp S. Stawski, Alice Y. Ting, and Michael S. Kuhns. 2015. 'A Mechanical Switch Couples T Cell Receptor Triggering to the Cytoplasmic Juxtamembrane Regions of CD3ζζ'. *Immunity* 43 (2): 227–39. <https://doi.org/10.1016/j.immuni.2015.06.018>.
- Limozin, Laurent, Marcus Bridge, Pierre Bongrand, Omer Dushek, Philip Anton van der Merwe, and Philippe Robert. 2019. 'TCR-PMHC Kinetics under Force in a Cell-Free System Show No Intrinsic Catch Bond, but a Minimal Encounter Duration before Binding'. *Proceedings of the National Academy of Sciences of the United States of America* 116 (34): 16943–48. <https://doi.org/10.1073/pnas.1902141116>.
- Limozin, Laurent, and Pierre-Henri Puech. 2019. 'Membrane Organization and Physical Regulation of Lymphocyte Antigen Receptors: A Biophysicist's Perspective'. *The Journal of Membrane Biology* 252 (4–5): 397–412. <https://doi.org/10.1007/s00232-019-00085-2>.
- Liu, Baoyu, Wei Chen, Brian D. Evavold, and Cheng Zhu. 2014. 'Accumulation of Dynamic Catch Bonds between TCR and Agonist Peptide-MHC Triggers T Cell Signaling'. *Cell* 157 (2): 357–68. <https://doi.org/10.1016/j.cell.2014.02.053>.
- Liu, Chinky Shiu Chen, Deblina Raychaudhuri, Barnali Paul, Yogaditya Chakrabarty, Amrit Raj Ghosh, Oindrila Rahaman, Arindam Talukdar, and Dipyaman Ganguly. 2018. 'Cutting Edge:

- Piezo1 Mechanosensors Optimize Human T Cell Activation'. *The Journal of Immunology* 200 (4): 1255–60. <https://doi.org/10.4049/jimmunol.1701118>.
- Liu, Yang, Lori Blanchfield, Victor Pui-Yan Ma, Rakieb Andargachew, Kornelia Galior, Zheng Liu, Brian Evavold, and Khalid Salaita. 2016. 'DNA-Based Nanoparticle Tension Sensors Reveal That T-Cell Receptors Transmit Defined PN Forces to Their Antigens for Enhanced Fidelity'. *Proceedings of the National Academy of Sciences* 113 (20): 5610–15. <https://doi.org/10.1073/pnas.1600163113>.
- Luca, Vincent C., Byoung Choul Kim, Chenghao Ge, Shinako Kakuda, Di Wu, Mehdi Roein-Peikar, Robert S. Haltiwanger, Cheng Zhu, Taekjip Ha, and K. Christopher Garcia. 2017. 'Notch-Jagged Complex Structure Implicates a Catch Bond in Tuning Ligand Sensitivity'. *Science* 355 (6331): 1320–24. <https://doi.org/10.1126/science.aaf9739>.
- Luo, Bing-Hao, Christopher V. Carman, and Timothy A. Springer. 2007. 'Structural Basis of Integrin Regulation and Signaling'. *Annual Review of Immunology* 25: 619–47. <https://doi.org/10.1146/annurev.immunol.25.022106.141618>.
- Ma, Victor Pui-Yan, Yang Liu, Lori Blanchfield, Hanquan Su, Brian D. Evavold, and Khalid Salaita. 2016. 'Ratiometric Tension Probes for Mapping Receptor Forces and Clustering at Intermembrane Junctions'. *Nano Letters* 16 (7): 4552–59. <https://doi.org/10.1021/acs.nanolett.6b01817>.
- Ma, Zhengyu, Paul A Janmey, and Terri H Finkel. 2008. 'The Receptor Deformation Model of TCR Triggering'. *The FASEB Journal* 22: 1002–8. <https://doi.org/10.1096/fj.07-9331hyp>.
- Malissen, Bernard, and Pierre Bongrand. 2015. 'Early T Cell Activation: Integrating Biochemical, Structural, and Biophysical Cues'. *Annual Review of Immunology* 33 (1): 539–61. <https://doi.org/10.1146/annurev-immunol-032414-112158>.
- Mallis, Robert J., Ke Bai, Haribabu Arthanari, Rebecca E. Hussey, Maris Handley, Zhenhai Li, Loice Chingozha, et al. 2015. 'Pre-TCR Ligand Binding Impacts Thymocyte Development before A β TCR Expression'. *Proceedings of the National Academy of Sciences* 112 (27): 201504971–201504971. <https://doi.org/10.1073/pnas.1504971112>.
- Mossman, Kaspar D, Gabriele Campi, Jay T Groves, and Michael L Dustin. 2005. 'Altered TCR Signaling from Geometrically Repatterned Immunological Synapses.' *Science* 310 (5751): 1191–93. <https://doi.org/10.1126/science.1119238>.
- Murphy, Ken, Paul Travers, and Mark Walport. 2008. *Janeway's Immunobiology 7th Edition*. Edited by Garland Science.
- Murray, C. D. 1926. 'The Physiological Principle of Minimum Work: I. The Vascular System and the Cost of Blood Volume'. *Proceedings of the National Academy of Sciences of the United States of America* 12 (3): 207–14. <https://doi.org/10.1073/pnas.12.3.207>.
- O'Connor, Roddy S, Xueli Hao, Keyue Shen, Keenan Bashour, Tatiana Akimova, Wayne W Hancock, Lance C Kam, and Michael C Milone. 2012. 'Substrate Rigidity Regulates Human T Cell Activation and Proliferation.' *Journal of Immunology (Baltimore, Md. : 1950)* 189 (3): 1330–39. <https://doi.org/10.4049/jimmunol.1102757>.
- Pageon, Sophie V, Matt A Govendir, Daryan Kempe, and Maté Biro. 2018. 'Mechanoimmunology : Molecular-Scale Forces Govern Immune Cell Functions' 29: 1919–26. <https://doi.org/10.1091/mbc.E18-02-0120>.
- Prager-Khoutorsky, Masha, Alexandra Lichtenstein, Ramaswamy Krishnan, Kavitha Rajendran, Avi Mayo, Zvi Kam, Benjamin Geiger, and Alexander D. Bershadsky. 2011. 'Fibroblast Polarization Is a Matrix-Rigidity-Dependent Process Controlled by Focal Adhesion Mechanosensing'. *Nature Cell Biology* 13 (12): 1457–65. <https://doi.org/10.1038/ncb2370>.
- Puech, Pierre-Henri, and Pierre Bongrand. 2021. 'Mechanotransduction as a Major Driver of Cell Behaviour: Mechanisms, and Relevance to Cell Organization and Future Research'. *Open Biology* 11 (11): 210256. <https://doi.org/10.1098/rsob.210256>.

- Rossjohn, Jamie, Stephanie Gras, John J. Miles, Stephen J. Turner, Dale I. Godfrey, and James McCluskey. 2015. 'T Cell Antigen Receptor Recognition of Antigen-Presenting Molecules'. *Annual Review of Immunology* 33: 169–200. <https://doi.org/10.1146/annurev-immunol-032414-112334>.
- Rossy, Jérémie, Julia M. Laufer, and Daniel F. Legler. 2018. 'Role of Mechanotransduction and Tension in T Cell Function'. *Frontiers in Immunology* 9 (November). <https://doi.org/10.3389/fimmu.2018.02638>.
- Rudolph, Markus G, Robyn L Stanfield, and Ian A Wilson. 2006. 'How TCRs Bind MHCs, Peptides, and Coreceptors.' *Annu Rev Immunol* 24: 419–66. <https://doi.org/10.1146/annurev.immunol.23.021704.115658>.
- Rushdi, Muaz, Kaitao Li, Zhou Yuan, Stefano Travaglino, Arash Grakoui, and Cheng Zhu. 2020. 'Mechanotransduction in T Cell Development, Differentiation and Function'. *Cells* 9 (2). <https://doi.org/10.3390/cells9020364>.
- Salvi, Alicia M., and Kris A. DeMali. 2018. 'Mechanisms Linking Mechanotransduction and Cell Metabolism'. *Current Opinion in Cell Biology* 54 (October): 114–20. <https://doi.org/10.1016/j.ceb.2018.05.004>.
- Santos, Luís C, David A Blair, Sudha Kumari, Michael Cammer, Thomas Iskratsch, Olivier Herbin, Konstantina Alexandropoulos, Michael L Dustin, and Michael P Sheetz. 2016. 'Actin Polymerization-Dependent Activation of Cas-L Promotes Immunological Synapse Stability'. *Immunology and Cell Biology* 94 (10): 981–93. <https://doi.org/10.1038/icb.2016.61>.
- Sasmal, Dibyendu K., Wei Feng, Sobhan Roy, Peter Leung, Yanran He, Chufan Cai, Guoshuai Cao, et al. 2019. 'TCR–PMHC Bond Conformation Controls TCR Ligand Discrimination'. *Cellular & Molecular Immunology*, September. <https://doi.org/10.1038/s41423-019-0273-6>.
- Schlaepfer, David D., Steven K. Hanks, Tony Hunter, and Peter van der Geer. 1994. 'Integrin-Mediated Signal Transduction Linked to Ras Pathway by GRB2 Binding to Focal Adhesion Kinase'. *Nature* 372 (6508): 786–91. <https://doi.org/10.1038/372786a0>.
- Smelty, Philippe, Céline Marchal, Romain Renard, Ludivine Sinzelle, Nicolas Pollet, Dominique Dunon, Thierry Jaffredo, Jean-Yves Sire, and Julien S. Fellah. 2010. 'Identification of the Pre-T-Cell Receptor Alpha Chain in Nonmammalian Vertebrates Challenges the Structure-Function of the Molecule'. *Proceedings of the National Academy of Sciences of the United States of America* 107 (46): 19991–96. <https://doi.org/10.1073/pnas.1010166107>.
- Smith-Garvin, Jennifer E, Gary A Koretzky, and Martha S Jordan. 2009. 'T Cell Activation'. *Annu Rev Immunol* 27: 591–619. <https://doi.org/10.1146/annurev.immunol.021908.132706>.
- Springer, Timothy A, and Michael L Dustin. 2012. 'Integrin Inside-out Signaling and the Immunological Synapse'. *Current Opinion in Cell Biology* 24 (1): 107–15. <https://doi.org/10.1016/j.ceb.2011.10.004>.
- Sundberg, Eric J., and Roy A. Mariuzza. 2002. 'Molecular Recognition in Antibody-Antigen Complexes'. *Advances in Protein Chemistry* 61: 119–60. [https://doi.org/10.1016/s0065-3233\(02\)61004-6](https://doi.org/10.1016/s0065-3233(02)61004-6).
- Tabdanov, Erdem, Sasha Gondarenko, Sudha Kumari, Anastasia Liapis, Michael L. Dustin, Michael P. Sheetz, Lance C. Kam, and Thomas Iskratsch. 2015. 'Micropatterning of TCR and LFA-1 Ligands Reveals Complementary Effects on Cytoskeleton Mechanics in T Cells'. *Integrative Biology (United Kingdom)* 7 (10): 1272–84. <https://doi.org/10.1039/c5ib00032g>.
- Teichtahl, Andrew J., Anita E. Wluka, Pushpika Wijethilake, Yuanyuan Wang, Ali Ghasem-Zadeh, and Flavia M. Cicuttini. 2015. 'Wolff's Law in Action: A Mechanism for Early Knee Osteoarthritis'. *Arthritis Research & Therapy* 17 (1): 207. <https://doi.org/10.1186/s13075-015-0738-7>.
- Thompson, D'Arcy Wentworth. 1992. *On Growth and Form*. Revised ed. édition. New York: Dover Publications Inc.

- Verma, Navin Kumar, Dermot Kelleher, and Email Alerts. 2018. 'Not Just an Adhesion Molecule: LFA-1 Contact Tunes the T Lymphocyte Program'. <https://doi.org/10.4049/jimmunol.1700495>.
- Wahl, Astrid, Céline Dinet, Pierre Dillard, Aya Nassereddine, Pierre-Henri Puech, Laurent Limozin, and Kheya Sengupta. 2019. 'Biphasic Mechanosensitivity of T Cell Receptor-Mediated Spreading of Lymphocytes'. *Proceedings of the National Academy of Sciences* 116 (13): 5908–13. <https://doi.org/10.1073/pnas.1811516116>.
- Wang, N., J. P. Butler, and D. E. Ingber. 1993. 'Mechanotransduction across the Cell Surface and through the Cytoskeleton'. *Science (New York, N.Y.)* 260 (5111): 1124–27. <https://doi.org/10.1126/science.7684161>.
- Wang, X., and T. Ha. 2013. 'Defining Single Molecular Forces Required to Activate Integrin and Notch Signaling'. *Science* 340 (6135): 991–94. <https://doi.org/10.1126/science.1231041>.
- Welf, Erik S., Christopher E. Miles, Jaewon Huh, Etai Sapoznik, Joseph Chi, Meghan K. Driscoll, Tadamoto Isogai, et al. 2020. 'Actin-Membrane Release Initiates Cell Protrusions'. *Developmental Cell* 55 (6): 723-736.e8. <https://doi.org/10.1016/j.devcel.2020.11.024>.
- Wu, Peng, Tongtong Zhang, Baoyu Liu, Panyu Fei, Lei Cui, Rui Qin, Huaying Zhu, et al. 2019. 'Mechano-Regulation of Peptide-MHC Class I Conformations Determines TCR Antigen Recognition'. *Molecular Cell* 73 (5): 1015-1027.e7. <https://doi.org/10.1016/j.molcel.2018.12.018>.
- Yamasaki, Sho, and Takashi Saito. 2007. 'Molecular Basis for Pre-TCR-Mediated Autonomous Signaling'. *Trends in Immunology* 28 (1): 39–43. <https://doi.org/10.1016/j.it.2006.11.006>.
- Zhang, Xuexiang, Tae-Hyung Kim, Timothy J Thauland, Hongjun Li, Fatemeh Sadat Majedi, Chau Ly, Zhen Gu, Manish J Butte, Amy C Rowat, and Song Li. 2020. 'Unraveling the Mechanobiology of Immune Cells'. *Current Opinion in Biotechnology* 66 (December): 236–45. <https://doi.org/10.1016/j.copbio.2020.09.004>.